

## Effect of Initial Quality and Compositional Parameters on Total Polar Compounds Formation during Continuous Heating of Olive Oil

R. Farhoosh<sup>1\*</sup>, M. H. Haddad Khodaparast<sup>1</sup>, A. Sharif<sup>1</sup>, and S. Alavi Rafiee<sup>1</sup>

### ABSTRACT

The present study was conducted to determine the relative contribution of initial quality and compositional parameters to the rate of changes ( $RC_{TPC}$ ) in the content of total polar compounds (TPC), which is considered to be one of the most reliable indicators of heated oils quality, of olive oil during continuous heating at 180 °C. The ratio between monounsaturated and polyunsaturated fatty acids showed the highest relative contribution (44.86%) to the  $RC_{TPC}$ . The content of total phenolic compounds indicated a considerable contribution (30.76%) to the  $RC_{TPC}$  whereas the content of total tocopherols was found to be actually ineffective (0.01%). Peroxide value was another important parameter that showed a marked contribution (20.51%) in this regard. The contributions of acid value and initial TPC content to the  $RC_{TPC}$  of olive oil were 3.85 and 0.01%, respectively.

**Keywords:** Alteration rate, Oil quality, Relative contribution, Stability.

### INTRODUCTION

Olive oil is a major source of dietary fat in the Mediterranean diet, used as salad dressing and frying oil. It accounts for less than 2% of total vegetable oil production, but its consumption is also becoming increasingly popular in non-producer countries. Several studies support a relationship between olive oil and a lower incidence of some important diseases, including cancer (Assmann *et al.*, 1997) and cardiovascular diseases (Covas, 2007). Health properties, oxidative stability, and sensory quality of olive oil are associated with prominent and well-balanced chemical composition (Bendini *et al.*, 2007). High content of oleic acid in olive oil serves to retard penetration of fatty acids into arterial walls (Charbonnier, 1982). The oils that are much higher in monounsaturated fatty acids

(MUFA) and lower in saturated fatty acids (SFA) are preferred because of the proved beneficial effects of MUFA on serum cholesterol levels. The interesting properties of olive oil are also related to the presence of minor components, especially natural antioxidants such as tocopherols and, particularly, phenolic compounds (Owen *et al.*, 2000).

Fatty acid composition and significant amounts of stability- and health-promoting components make olive oil a very interesting option among heating oils and fats used for frying purposes. Numerous analytical methods have been described to measure the changes in heated oils, but determination of polar compounds is considered to be one of the most reliable indicators of heated oils quality. By definition, total polar compounds (TPC) content is the sum of the materials that are not triglycerides (Ruiz-Mendez *et al.*, 1997). Regulatory agencies in European

<sup>1</sup> Department of Food Science and Technology, Ferdowsi University of Mashhad, Faculty of Agriculture, Mashhad, Islamic Republic of Iran.

\* Corresponding author; e-mail: rfarhoosh@um.ac.ir



countries have established a level of 24-27% TPC for the rejection of heated oils (Paul and Mittal, 1997).

The rate of changes in TPC content of olive oil may be a function of many initial quality and compositional parameters. Peroxide value (PV) and acid value (AV) are two of the most frequently determined quality indices, which indicate the levels of total hydroperoxides and hydrolytic rancidity, respectively. Therefore, they are considered to be guides to initial quality of olive oils (Lovaas, 1992). The ratio between MUFA and polyunsaturated fatty acids (PUFA) is a measure of olive oils tendency to undergo autoxidation. Phenolic compounds and tocopherols are well-known effective chain-breaking antioxidants naturally occurring in edible fats and oils (Frankel, 1998). Moreover, the initial amount of TPC in vegetable oils can have a pronounced contribution to create the off-flavour compounds as well as a marked effect on the primary oxidation (Farhoosh and Pazhouhanmehr, 2009).

The aim of this study was to determine the relative contribution of compositional parameters, including PV, AV, MUFA/PUFA ratio, total tocopherols (TT) content, total phenolics (TP) content, and TPC content, to the rate of changes in TPC content ( $RC_{\text{TPC}}$ ) of olive oil during continuous heating process.

## MATERIALS AND METHODS

### Materials

Four deodorized olive oil samples (1, 2, 5, and 6 as shown in Table 1) and three virgin olive oil samples (3, 4, and 7 as shown in Table 1) were purchased from local shops and were stored at 4°C until analysis. Fatty acid methyl ester (FAME) standards and all chemicals and solvents used in this study were of analytical reagent grade and supplied by Merck and Sigma Chemical Companies.

### Oil Oxidation

The oil samples (200 g) were placed in a fryer (Kenwood DF280, Havant, Hampshire, UK) and maintained at 180°C for 16 hours with no stirring. The surface-to-volume ratio of the oil samples in the fryer was about 0.7 cm<sup>-1</sup>. After intervals of 4 hours of heating, samples were removed and analyzed.

### Chemical Analyses

The fatty acid composition of the oils was determined by gas-liquid chromatography. Fatty acids were transesterified into their corresponding FAMES by vigorous shaking of a solution of oil in hexane (0.3 g in 7 mL) with 2 mL of 7N methanolic potassium hydroxide at 50°C for 10 minutes. The FAMES were identified using an HP-5890 chromatograph (Hewlett-Packard, CA, USA) equipped with a CP-FIL 88 (Supel Co., Inc., Bellefonte, PA) capillary column of fused silica, 60 m in length × 0.22 mm ID, 0.2 μm film thickness, and a flame ionization detector (FID). Nitrogen was used as carrier gas with a flow rate of 0.75 mL min<sup>-1</sup>. The oven temperature was maintained at 198°C, and that of the injector and the detector at 250°C (Farhoosh *et al.*, 2008). The fatty acids in the sample were identified using standard methyl esters of fatty acids by comparing the retention times. The results were expressed as relative area percent.

The spectrophotometric method described by Shantha and Decker (1994) was used to determine the PV. The AV was determined according to the AOCS Official Method Cd 3d-63 (1993). The TT content was determined according to the colorimetric method described by Wong *et al.* (1988). The TP content was determined spectrophotometrically using Folin-Ciocalteu's reagent according to the method described by Capannesi *et al.* (2000). A calibration curve of gallic acid in methanol was performed in concentration range 0.04-0.40 mg mL<sup>-1</sup>. The TPC content

was determined according to the economical micro method developed by Schulte (2004).

### Statistical Analysis

All determinations were carried out in triplicate, and data were subjected to analysis of variance (ANOVA). ANOVA and regression analyses were performed according to the MSTATC and SlideWrite software. To standardize the compositional parameters, each parameter was subtracted from its mean among the olive oils, and the difference was divided by the corresponding standard deviation. Significant differences between means were determined by Duncan's multiple range tests. *P* values less than 0.05 were considered statistically significant.

## RESULTS AND DISCUSSION

The compositional parameters of the olive oil samples are shown in Table 1. As can be seen, the fatty acid compositions were statistically different. The contents of SFA, MUFA, and PUFA ranged from 15.67 to 20.19% (mainly palmitic acid, C16:0, and stearic acid, C18:0), from 59.13 to 76.10% (mainly oleic acid, C18:1), and from 7.67 to 20.21% (mainly linoleic acid, C18:2), respectively. These contents were in accordance with those reported in the literature for olive oils (Firestone *et al.* 1996). From information stated above, the MUFA/PUFA ratio varied significantly amongst the olive oils studied and widely ranged from 2.93 to 9.92.

A wide range of the PV (8.4-13.4 meq kg<sup>-1</sup>) and AV (0.10-2.00 mg g<sup>-1</sup>) was observed amongst the olive oil samples, indicating the olive oils had no similar initial qualities. The olive oils experimented contained statistically different contents of TT (44 - 457 ppm) and TP (12-203 ppm), leading to their different oxidative stabilities.

Table 2 shows the TPC content of the olive oils during the heating process at 180°C. Lumley (1988) showed that the TPC content

of fresh oils normally ranges between 0.4 and 6.4%, although other different amounts of TPC content (up to 14%) have been reported in crude or refined vegetable oils (Ruiz-Mendez *et al.*, 1997). Depending on the fatty acid composition, initial quality, and technology used, crude oils may undergo decreases or even increases in their TPC contents during refining. Considering that the polar fraction in edible fats and oils includes essentially alteration compounds, the higher the TPC contents, the lower will be the expected quality of the oils. The olive oils tested in this study contained initial TPC contents ranging from 3.96 to 11.59% (Table 2). As can be seen, the TPC contents linearly increased with high determination coefficients ( $R^2 > 0.98$ ) during the heating process. Assuming that the limit of acceptance for the TPC content is 24%, the time required to reach this limit was calculated as a measure of oxidative stability ( $t_{24}$ ). A higher  $t_{24}$  value means greater stability over the heating process. Except for the samples 2 and 3, all the oils studied reached the discarding range of TPC content during the heating process. The olive oil samples 1-7 indicated the  $t_{24}$  values of 10.23, 19.41, 23.96, 7.75, 8.26, 10.99, and 14.49 h, respectively (Figure 1).

It is obvious that the  $RC_{TPC}$  is a function of all initial quality and compositional parameters showed in Table 1, whereas the  $t_{24}$  value is highly affected by the initial TPC content (Table 2). Therefore, the inverse of percentage increase in the TPC content after 16 h of heating ( $\%TPC_{16h}^{-1}$ ) was considered as a quantitative measure of the  $RC_{TPC}$ . The  $\%TPC_{16h}^{-1}$  was correlated with a unified variable defined as  $\sum_{i=1}^6 (C_i \times SCP_i)$ , in which  $C_i$  and  $SCP_i$  were coefficients and the standardized (and inversed in the cases needed, Table 3) compositional parameters, respectively. The  $C_i$  values in the regression models were determined based on reaching the maximum possible determination coefficient. A very good correlation was found between the  $RC_{TPC}$  and the unified variable consisting

**Table 1.** The initial quality and compositional parameters of the olive oil samples\*.

Parameters	Olive oil samples						
	1	2	3	4	5	6	7
<b>Fatty acids</b>							
16:0	15.68±0.26 a	12.50±0.09 d	14.58±0.11 b	14.63±0.00 b	13.08±0.13 c	12.40±0.05 d	10.63±0.01 e
16:1	1.98±0.03 a	1.26±0.06 c	1.66±0.06 b	1.92±0.04 a	1.27±0.02 c	1.10±0.00 d	1.13±0.11 cd
18:0	3.70±0.08 d	4.56±0.12 b	3.38±0.03 e	3.52±0.10 de	4.90±0.11 a	4.82±0.06 a	4.27±0.08 c
18:1	56.70±0.51 g	68.40±0.37 b	60.91±0.10 f	63.25±0.38 e	65.03±0.13 d	67.59±0.11 c	74.62±0.37 a
18:2	19.02±0.05 a	10.34±0.10 g	16.91±0.18 b	13.87±0.08 c	12.59±0.24 e	11.01±0.15 f	6.84±0.19 h
18:3	1.20±0.05 a	0.89±0.00 d	1.13±0.04 a	1.11±0.10 ab	0.89±0.13 cd	1.00±0.04 bc	0.84±0.09 d
20:0	0.81±0.04 b	0.84±0.07 abc	0.65±0.11 c	0.76±0.05 bc	0.93±0.01 a	0.88±0.04 ab	0.77±0.06 bc
20:1	0.45±0.09 a	0.49±0.01 a	0.43±0.01 a	0.42±0.01 a	0.51±0.07 a	0.42±0.04 a	0.35±0.16 a
22:0	0.24±0.07 a	0.25±0.06 a	0.16±0.03 a	0.25±0.07 a	0.25±0.04 a	0.26±0.04 a	0.23±0.11 a
Others	0.24±0.02 c	0.48±0.10 ab	0.21±0.04 c	0.29±0.06 bc	0.57±0.04 a	0.53±0.02 c	0.32±0.11 b
SFA <sup>d</sup>	20.19±0.22 a	17.89±0.04 c	18.60±0.03 b	18.91±0.15 b	18.90±0.04 b	18.09±0.03 c	15.67±0.13 d
MUFA <sup>b</sup>	59.13±0.39 g	70.14±0.31 b	63.00±0.15 f	65.59±0.33 e	66.81±0.08 d	69.11±0.15 e	76.10±0.64 a
PUFA <sup>c</sup>	20.21±0.10 a	11.23±0.10 f	18.04±0.13 b	14.98±0.18 c	13.48±0.11 d	12.01±0.11 c	7.67±0.28 g
MUFA/PUFA	2.93±0.02 g	6.25±0.01 b	3.49±0.05 f	4.38±0.04 e	4.96±0.02 d	5.75±0.03 c	9.92±0.05 a
PV <sup>d</sup>	13.2±1.8 a	8.4±0.5 c	11.3±0.7 a	9.3±0.7 bc	9.9±0.2 b	10.4±1.2 abc	13.4±1.7 a
AV <sup>e</sup>	1.86±0.02 b	0.30±0.02 d	2.00±0.02 a	1.92±0.15 ab	0.10±0.02 e	0.16±0.02 e	0.66±0.02 c
TT content <sup>f</sup>	388±67 b	102±1 d	457±24 a	44±11 e	144±6 c	109±15 d	382±12 b
TP content <sup>g</sup>	165±8 b	12±1 e	203±18 a	100±12 d	13±3 e	15±2 e	138±3 c


\* Mean±SD (standard deviation) within a row with the same lowercase letters are not significantly different at  $P < 0.05$ .

<sup>a</sup> Saturated fatty acids (%); <sup>b</sup> Monounsaturated fatty acids (%); <sup>c</sup> Polyunsaturated fatty acid (%); <sup>d</sup> Peroxide value (meq O<sub>2</sub> per kg oil); <sup>e</sup> Acid value (mg KOH per g oil); <sup>f</sup> Total tocopherols (mg α-tocopherol per kg oil); <sup>g</sup> Total phenolics (mg gallic acid per kg oil).

**Table 2.** Total polar compounds (TPC, %) content of the olive oil samples during the heating process at 180°C\*.

Time (Hour)	Olive oil samples						
	1	2	3	4	5	6	7
0	3.98±0.19 Ec	4.74±0.28 Db	3.96±0.12 Dc	11.59±1.97 Ca	8.15±1.05 Ea	5.41±0.35 Db	11.52±1.60 Ca
4	10.27±1.50 Dc	9.76±1.12 Cc	8.03±1.10 Cc	15.98±1.27 Ca	12.95±1.16 Dabc	11.27±1.47 Cbc	14.45±1.21 BCab
8	16.98±2.31 Cab	13.63±2.10 ABCb	11.05±1.85 BCb	22.47±2.82 Ba	21.43±2.66 Ca	18.00±2.03 Bab	18.91±1.97 ABab
12	26.44±2.77 Bab	17.28±2.28 ABc	13.90±1.70 ABC	33.75±2.83 Aa	32.17±3.03 Ba	26.52±1.86 Aab	22.44±1.89 Ab
16	38.53±2.06 Aa	20.07±2.14 Acd	17.47±2.11 Ad	38.48±1.58 Aa	42.43±3.78 Aa	32.70±2.89 Aab	24.75±2.64 Abc

\* Mean±SD (standard deviation) within a row with the same lowercase letters are not significantly different at  $P < 0.05$ . Means±SD within a column with the same uppercase letters are not significantly different at  $P < 0.05$ .  
The corresponding linear equations: Sample 1,  $y = 2.13x + 2.19$ ,  $R^2 = 0.979$ ; Sample 2,  $y = 0.96x + 5.46$ ,  $R^2 = 0.989$ ; Sample 3,  $y = 0.82x + 4.30$ ,  $R^2 = 0.996$ ; Sample 4,  $y = 1.79x + 10.14$ ,  $R^2 = 0.976$ ; Sample 5,  $y = 2.20x + 5.87$ ,  $R^2 = 0.982$ ; Sample 6,  $y = 1.75x + 4.81$ ,  $R^2 = 0.996$ ; Sample 7,  $y = 0.86x + 11.52$ ,  $R^2 = 0.990$ .




**Figure 1.** The time required to reach total polar compounds (TPC) content of 24% ( $t_{24}$ ) for the olive oil samples (1–7) during the heating process at 180°C. Means with the same lowercase letters are not significantly different at  $P < 0.05$ .

of the compositional parameters (Figure 2).

The relative contributions of compositional parameters to the  $RC_{TPC}$  of olive oils were calculated from the  $C_i$  values obtained from the regression models in Figure 2 (Table 3). Amongst the parameters studied, the MUFA/PUFA ratio showed the highest contribution (44.86%) to the  $RC_{TPC}$  of the olive oils. This clearly means that less monounsaturated and/or more polyunsaturated olive oils are more prone to produce polar compounds during heating processes. In other words, the MUFA/PUFA ratio can be considered a good indicator of olive oils stability.

The TP content, as Table 3 shows, played an important role in the  $RC_{TPC}$  of olive oils, so that its relative contribution to the production of polar compounds was 30.76%. The contribution of phenolic compounds to the olive oils stability and antioxidant activity has been estimated to be higher than that of other compounds (Gutierrez *et al.*, 2001; Pellegrini *et al.*, 2001), and a number of researchers have demonstrated a positive linear relationship between oil stability and the TP content (Gutfinger, 1981; Baldioli *et al.*, 1996). It has been shown that the major phenolic compounds in olive oils are oleuropein, hydroxytyrosol (3-dihydroxyphenylethanol), and tyrosol (4-



**Figure 2.** Linear relationship between the inverse of percentage increase in total polar compounds (TPC) content after 16 h of heating at 180°C ( $\%TPC_{16h}^{-1}$ ) and the unified variable consisting of the compositional parameters ( $\sum_{i=1}^6(C_i \times SCP_i)$ ).

hydroxyphenylethanol). The phenolic compounds presented in olive oils are strong metal chelators (Chimi *et al.*, 1991) and radical scavengers (Tuck and Hayball, 2002). Typically, hydroxytyrosol is a superior antioxidant and radical scavenger to oleuropein and tyrosol (Tuck and Hayball 2002).

In addition to phenolic compounds, tocopherols are a group of the most important natural antioxidants in edible fats and oils. Autoxidation studies have demonstrated that tocopherols not only inhibit the formation of hydroperoxides (Lampi *et al.*, 1999) in many substrates, but also reduce the total amount of decomposition products from hydroperoxides (Lampi *et al.*, 1999; Kulas *et*

*al.*, 2001; Makinen *et al.*, 2001; Kulas *et al.*, 2002). Nevertheless, it was interesting to find that these antioxidative compounds were actually ineffective on the production rate of polar compounds (0.01%) in our study. It should be noted that research on the antioxidant activity of tocopherols is conflicting sometimes, and this may be due to different experimental conditions and antioxidants concentrations (Schieberle and Grosch, 1981; Koskas *et al.*, 1984).

The PV was another important parameter that showed considerable contribution (20.51%) to the  $RC_{TPC}$  of olive oils. This means that olive oils with higher initial PVs are more prone to produce the polar compounds. Measurement of AV is important to determine the degree of hydrolytic

**Table 3.** The coefficients of compositional parameters ( $C_i$  values) obtained from the regression model and the relative contribution (%) of each parameter to the rate of changes in total polar compounds (TPC) content ( $RC_{TPC}$ ) of olive oil samples.

Parameters	$RC_{TPC}: [1000 \times (\%TPC_{16h}^{-1})]$	
	$C_i$ value	Relative contribution (%)
$(PV^{-1})_{Std}^a$	16	20.51
$(AV^{-1})_{Std}^b$	3	3.85
$MUFA/PUFA_{Std}^c$	35	44.86
$TT_{Std}^d$	0.01	0.01
$TP_{Std}^e$	24	30.76
$(TPC^{-1})_{Std}$	0.01	0.01

<sup>a</sup> Peroxide value; <sup>b</sup> Acid value; <sup>c</sup> Monounsaturated/Polyunsaturated fatty acids ratio; <sup>d</sup> Total tocopherols, <sup>e</sup> Total phenolic compounds.

deterioration (Cecchi, 2003) and its level is a valuable determinant of oils quality. Surprisingly, however, the AV had a partial role in the RC<sub>TPC</sub> of olive oils, so that its relative contribution to the production of polar compounds was 3.85%. A previous study indicated that the initial TPC content has a pronounced contribution to create the off-flavor compounds as well as a marked effect on the primary oxidation (Farhoosh and Pazhouhanmehr, 2009). However, our findings in the present study revealed that the initial TPC content had almost no influence on the production rate of polar compounds (0.01%, Table 3).

### CONCLUSIONS

In the present study, the effect of two groups of initial quality (PV, AV, and TPC content) and compositional parameters (TT content, TP content, and MUFA/PUFA ratio) on the rate of changes in the TPC content of olive oil was investigated during heating process. The parameters in the first group are function of handling, storage, and/or processing conditions, while the parameters in the second group are indigenous properties of edible fats and oils. Our study showed that the most effective initial quality and compositional parameters during heating process of olive oil were MUFA/PUFA ratio (44.86%), TP content (30.76%), and PV (20.51%), respectively.

### ACKNOWLEDGEMENTS

The research was funded by Ferdowsi University of Mashhad under Project Grant AGRI/47266/2009.

### REFERENCES

1. Assmann, G., de Backer, G., Bagnara, S., Betteridge, J., Crepaldi, G., Fernandez-Cruz, A., Godtfredsen, J., Jacotot, B., Paoletti, R., Renaud, S., Ricci, G., Rocha, E., Trautwein, E., Urbinati, G. C., Varela, G. and Williams, C. 1997. International Consensus Statement on Olive Oil and the Mediterranean Diet: Implications for Health in Europe. The Olive Oil and the Mediterranean Diet Panel. *Eur. J. Can. Prev.*, **6**: 418-421.
2. AOCS. 1993. Official Methods and Recommended Practices of the American Oil Chemists' Society. AOCS Press, Champaign.
3. Baldioli, M., Servili, M., Perretti, G. and Montedoro, G. F. 1996. Antioxidant Activity of Tocopherols and Phenolic Compounds of Virgin Olive Oil. *J. Am. Oil Chem. Soc.*, **73**: 1589-1593.
4. Bendini, A., Cerretani, L., Carrasco-Pancorbo, A., Gomez-Caravaca, A. M., Segura-Carretero, A., Fernandez-Gutierrez, A. and Lercker, G. 2007. Phenolic Molecules in Virgin Olive Oils: A Survey of Their Sensory Properties, Health Effects, Antioxidant Activity and Analytical Methods. An Overview of the Last Decade. *Molecules*, **12**: 1679-1719.
5. Capannesi, C., Palchetti, I., Mascini, M. and Parenti, A. 2000. Electrochemical Sensor and Biosensor for Polyphenols Detection in Olive Oils. *Food Chem.*, **71**: 553-562.
6. Cecchi, H. M. 2003. Fundamentos Teoricos e Praticos em Analise de Alimentos. Editora da Unicamp, Campinas.
7. Charbonnier, A. 1982. Main Conclusion Drawn from the International Symposium on the Recent Medical Researches on the Value of the Olive Oil to Health, Paris, PP. 1-4.
8. Chimi, H., Cillard, J., Cillard, P. and Rahamani, M. 1991. Peroxyl and Hydroxyl Radical Scavenging Activity of Some Natural Phenolic Antioxidants. *J. Am. Oil Chem. Soc.*, **68**: 307-312.
9. Covas, M. I. 2007. Olive Oil and the Cardiovascular System. *Pharmacol. Res.*, **55**: 175-186.
10. Farhoosh, R., Niazmand, R., Rezaei, M. and Sarabi, M. 2008. Kinetic Parameter Determination of Vegetable Oil Oxidation under Rancimat Test Conditions. *Eur. J. Lipid Sci. Technol.*, **110**: 587-592.
11. Farhoosh, R. and Pazhouhanmehr, S. 2009. Relative Contribution of Compositional Parameters to the Primary and Secondary Oxidation of Canola Oil. *Food Chem.*, **114**: 1002-1006.
12. Firestone, D., Fedeli, E. and Emmons, E. W. 1996. Olive Oil. In: "Bailey's Industrial Oil



- and Fat Products", (Ed.): Hui, Y. H.. John Wiley and Sons, New York, PP. 241-247.
13. Frankel, E. N. 1998. *Lipid Oxidation*. The Oily Press, Dundee.
  14. Gutfinger, T. 1981. Polyphenols in Olive Oils. *J. Am. Oil Chem. Soc.*, **58**: 966-968.
  15. Gutierrez, F., Arnaud, T. and Garrido, A. 2001. Contribution of Polyphenols to the Oxidative Stability of Virgin Olive Oil. *J. Sci. Food Agric.*, **81**: 1463-1470.
  16. Koskas, J. P., Cillard, J. and Cillard, P. 1984. Autoxidation of Linoleic Acid and Behavior of its Hydroperoxides with and without Tocopherols. *J. Am. Oil Chem. Soc.*, **61**: 1466-1469.
  17. Kulas, E. and Ackman, R. G. 2001. Properties of  $\alpha$ -,  $\gamma$ - and  $\delta$ -tocopherols in Purified Fish Oil Triacylglycerols. *J. Am. Oil Chem. Soc.*, **78**: 361-367.
  18. Kulas, E., Olsen, E. and Ackman, R. G. 2002. Effect of  $\alpha$ -,  $\gamma$ - and  $\delta$ -tocopherol on the Distribution of Volatile Secondary Oxidation Products in Fish Oil. *Eur. J. Lipid Sci. Technol.*, **104**: 520-529.
  19. Lampi, A. M., Kataja, L., Kamal-Eldrin, A. and Piironen, V. 1999. Antioxidant Activities of  $\alpha$ - and  $\gamma$ -tocopherols in the Oxidation of Rapeseed Oil Triacylglycerols. *J. Am. Oil Chem. Soc.*, **76**: 749-755.
  20. Lovaas, E. A. 1992. Sensitive Spectrophotometric Method for Lipid Hydroperoxide Determination. *J. Am. Oil Chem. Soc.*, **69**: 777-783.
  21. Lumley, I. D. 1988. Polar Compounds in Heated Oils. In: "Frying of Food: Principles, Changes, New Approaches", (Eds.): Varela, G., Bender, A. E. and Morton, I. D.. Ellis Horwood Ltd., Chichester, PP. 166-173.
  22. Makinen, M., Kamal-Eldrin, A., Lampi, A. M. and Hopia, A. 2001.  $\alpha$ -,  $\gamma$ - and  $\delta$ -tocopherols as Inhibitors of Isomerization and Decomposition of Cis, Trans Methyl Linoleate Hydroperoxides. *Eur. J. Lipid Sci. Technol.*, **103**: 286-291.
  23. Owen, R. W., Mier, W., Giacosa, A., Hull, W. E., Spiegelhalder, B. and Bartsch, H. 2000. Phenolic Compounds and Squalene in Olive Oils: The Concentration and Antioxidant Potential of Total Phenols, Simple Phenols, Secoiridoids, Lignans and Squalene. *Food Chem. Toxicol.*, **38**: 647-659.
  24. Paul, S. and Mittal, G. S. 1997. Regulating the Use of Degraded Oil/Fat in Deep-fat/Oil Food Frying. *Crit. Rev. Food Sci. Nutr.*, **37**: 635-662.
  25. Pellegrini, N., Visioli, F., Buratti, S. and Brighenti, F. 2001. Direct Analysis of Total Antioxidant Activity of Olive Oil and Studies on the Influence of Heating. *J. Agric. Food Chem.*, **49**: 2532-2538.
  26. Ruiz-Mendez, M. V., Marquez-Ruiz, G. and Dobarganes, M. C. 1997. Relationships between Quality of Crude and Refined Edible Oils Based on Quantitation of Minor Glyceridic Compounds. *Food Chem.*, **60**: 549-554.
  27. Schieberle, P. and Grosch, W. 1981. Decomposition of Linoleic Acid Hydroperoxides II. Breakdown of Methyl 13-hydroperoxy-cis-9-trans-11-octadecadienoate by Radicals and Copper II ions. *Zeit. Leben. u.-Forsch. A.*, **173**: 192-198.
  28. Schulte, E. 2004. Economical Micromethod for Determination of Polar Components in Frying Fats. *Eur. J. Lipid Sci. Technol.*, **106**: 772-776.
  29. Shantha, N. C. and Decker, E. A. 1994. Rapid, Sensitive, Iron-based Spectrophotometric Methods for Determination of Peroxide Values of Food Lipids. *J. AOAC Int.*, **77**: 421-424.
  30. Tuck, K. L. and Hayball, P. J. 2002. Major Phenolic Compounds in Olive Oil: Metabolism and Health Effects. *J. Nutr. Biochem.*, **13**: 636-644.
  31. Wong, M. L., Timms, R. E. and Goh, E. M. 1988. Colorimetric Determination of Total Tocopherols in Palm Oil, Olein and Stearin. *J. Am. Oil Chem. Soc.*, **65**: 258-261.



## اثر پارامترهای ساختاری و کیفی اولیه بر شکل گیری کل ترکیبات قطبی طی حرارت دهی مداوم روغن زیتون

ر. فرهوش، م. ح. حدادخداپرست، ع. شریف و س. علوی رفیعی

### چکیده

پژوهش حاضر به قصد تعیین سهم نسبی پارامترهای ساختاری و کیفی اولیه در سرعت تغییرات میزان کل ترکیبات قطبی به عنوان یکی از مطمئن ترین شاخصهای کیفیت روغنهای حرارت دیده طی حرارت دهی پیوسته روغن زیتون در دمای ۱۸۰ درجه سانتیگراد به اجرا در آمد. نسبت اسیدهای چرب تک غیراشباع به چند غیراشباع حائز بالاترین سهم نسبی (۴۴/۸۶ درصد) در این خصوص بود. سرعت تغییرات میزان کل ترکیبات قطبی به طور قابل ملاحظه‌ای (۳۰/۷۶ درصد) تحت تاثیر میزان کل ترکیبات فنلی قرار گرفت اما میزان کل ترکیبات توکوفرولی در این خصوص بواقع بی-تاثیر بود (۰/۰۱ درصد). عدد پراکسید، پارامتر حائز اهمیت دیگری بود که توانست سهم مشخصی از سرعت تغییرات میزان کل ترکیبات قطبی را به خود اختصاص دهد (۲۰/۵۱ درصد). سهم عدد اسیدی و میزان کل ترکیبات قطبی اولیه در این خصوص به ترتیب ۳/۸۵ و ۰/۰۱ درصد تعیین شد.