# Genetic Diversity of Synthetic Alfalfa Generations and Cultivars Using Tetrasomic Inherited Allozyme Markers

M. Valizadeh<sup>1\*</sup>, M. Mohayeji<sup>1</sup>, N. Yasinzadeh<sup>1</sup>, S. Nasrullazadeh<sup>1</sup>, and M. Moghaddam<sup>1</sup>

## ABSTRACT

Enzyme electrophoresis was used to measure genetic variation within, and divergence among, three generations of recently bred synthetic alfalfa generations (Syn<sub>1</sub>, Syn<sub>2</sub>, and Syn<sub>3</sub>) originating from a polycross of 12 selected parents and several cultivars. Three isozyme loci, exhibiting tetrasomic inheritance in 10-day seedlings, were detected from five enzymatic systems analyzed by polyacrylamide slab gel electrophoresis for about 100 individuals of each alfalfa population. Very high levels of heterozygosity (ranging from 0.521 to 0.699) were observed within alfalfa populations, using polymorphic loci. The reduction in heterozygosity was about 5% from Syn1 to Syn2 and from Syn2 to Syn3. The last open pollinated generation was found to be in W-H equilibrium as well as Gharayonja, a native ecotype under examination, using  $\chi^2$ -test. Application of Wright's Fstatistics revealed that the estimated overall inbreeding coefficient, (F<sub>IT</sub>), of 9.4% was mainly related to inbreeding or double reduction in alfalfa (F<sub>IS</sub>= 8.61%) rather than random genetic drift or population differentiation ( $F_{ST}$ = 1.6%). Therefore, due to very large intra-population diversity, the breeding program of the synthetic alfalfa did not generate a large variety differentiation. However, the use of seedling allozymic loci can be applied successfully for estimation of the population genetic parameters.

Keywords: Alfalfa, Allozymes, Genetic diversity, Tetrasomic inheritance.

## **INTRODUCTION**

Alfalfa (Medicago sativa L.), is the most important forage crop grown in the temperate regions. It is cultivated over 32 million hectares worldwide (Michaud et al., 1988) and over 680 thousands hectares in Iran (Anonymous, 2007), especially in the Northwest region, which is considered as one of the two alfalfa origins (Sinskaya, 1950). In Iran, cultivated alfalfa mainly consists of several adaptive landraces including Gharayonja and Hamadani as well as occasionally introduced cultivars such as Ranger or Maopa.

Tetrasomic inheritance (2n=4x=32) in autotetraploid alfalfa (Stanford, 1951; Demarly, 1954) and pronounced inbreeding depression (McCoy and Bingham, 1988), have caused the genetic improvement of alfalfa to

be less than other major crops. Improved varieties are synthetic cultivars, usually obtained through three or four generations of open pollinated reproduction of polycross seeds of selected parents (Tysdal and Crandal, 1948; Flajulot *et al.*, 2005).

One of the most useful measures of population structure is F-statistics (Wright, 1951), which describes the amount of inbreeding-like effects within subpopulations  $(F_{IS})$ , among subpopulations  $(F_{ST})$ , and within the entire population (F<sub>IT</sub>) (Flajulot et al., 2005). In contrast to diploids, the correlation individuals of genes between within population with respect to genes between populations (F<sub>ST</sub>) may vary among loci due to double reduction (Wricke and Weber, 1986) and appropriate number of loci may be required for estimating demographic

<sup>&</sup>lt;sup>1</sup> Department of Crop Production and Plant Breeding, Faculty of Agriculture, University of Tabriz, Islamic Republic of Iran.

<sup>\*</sup> Corresponding author, e-mail: valizadeh@hotmail.com

parameters of autotetraploid populations. The allozymic markers showing tetrasomic inheritance patterns can be used to perform genetic diversity analysis.

The objective of this work was to measure population structure of three generations belonging to a newly produced synthetic alfalfa (Syn<sub>1</sub>, Syn<sub>2</sub>, and Syn<sub>3</sub>), some parental populations, and improved varieties by allozymic loci.

## **MATERIALS AND METHODS**

## **Plant Material**

A total of 600 alfalfa seedlings were analyzed from six populations of alfalfa including three generations of a synthetic alfalfa (Syn<sub>1</sub>, Syn<sub>2</sub> and Syn<sub>3</sub>), one parental landrace, Ghara-yonja, and two cultivars, namely, Ranger and Sea-river, released in the USA and Australia, respectively. Syn1 was obtained by polycrossing of 12 selected ecotypes in 2002 arranged in an isolated latinsquare design after a period of four years of screening 30 parental ecotypes from Northwest of Iran by progeny test, at the Research Station of University of Tabriz (Valizadeh et al., 2002). Syn<sub>2</sub> and Syn<sub>3</sub> were produced by isolated open pollination in 2004 and 2005, respectively. Gara-yonja, which was collected from Maragheh region in East-Azarbaijan Province, is the most common landrace grown in Northwest of Iran.

## **Enzyme Extraction and Electrophoresis**

Hypocotyls and cotyledons of 10-day old seedlings were crashed with separate mortar and pestle in a tris-HCl extraction buffer pH 7.5 (tris 50 mM, sucrose 5%, ascorbic acid 50 mM, sodium metabisulfite 20 mM, PEG 2% and 2ME 0.1% before use) with a ratio of 0.5 mg  $\mu$ l<sup>-1</sup> and centrifuged at 4°C and 8,000g for 20 minutes using small Eppendof tubes. Enzyme extracts (supernatant) were immediately absorbed onto 3×5 mm wicks cut from Whatman 3 MM filter paper and

loaded onto 7% horizontal slab acrylamide gel. Electrophoresis was performed using Poulik gel buffer and tris 32 mM, boric acid 0.6 mM, and EDTA-Na<sub>2</sub> 0.1 mM electrode buffer at pH 8.8. Five enzymatic systems were stained on sliced gels: Esterase (Est), glutamate oxaloacetate transaminase (Got), catalase (Cat) and malate dehydrogerase (Mdh), according to Soltis and Soltis (1990) and peroxidase (Pox) according to Olson and Varner (1993). Putative loci were designated sequentially, with the most anodally migrating isozyme designated A, the next one B, etc. Likewise, alleles were designated sequentially with the most anodally migrating allele designated 1.

#### **Data Analysis**

Some standard genetic parameters including proportion of polymorphic loci (P) and genetic diversity, expected or heterozygosity (He= $1 - \Sigma p_i^4$ ), were calculated for about 100 individuals in each alfalfa population. A  $\chi^2$  test was performed to verify the existence of equilibrium in populations with one locus and two allele models, by keeping the most common allele and pooling the other alleles, when necessary. To determine whether the allele frequencies were significantly different, a  $\chi^2$  test was performed according to Workman and Niswander (1970):

$$\chi^2 = \frac{2NV_{(p)}}{\overline{pq}}$$

Where, *N* is the total sample size, *p* and *q* are the weighted averages of the two allele frequencies, and  $V_{(p)}$  is the weighted variance (See Hederick (2005) for more details). Due to the limited number of loci, no statistical analysis was used for the parameters estimations.

Nei's (1972) genetic distance measures were calculated for all pairs of populations to estimate genetic divergence among the populations. Wrights *F*-statistics (Wright, 1951), which is based on the decrease in the proportion of heterozygosity, was used to evaluate the amount of inbreeding-like effects



**Figure 1**. Banding pattern of esterase for 12 alfalfa seedlings, showing two polymorphic loci (Est-A and Est-B) and one monomorphic locus (Est-C). The allele's nomenclature and the genotypes of some individuals are indicated.

within each population  $(F_{IS})$ , among populations  $(F_{ST})$ , and within the entire germplasm under study, using Autotet software.

## **RESULTS AND DISCUSSION**

Eight putative loci were resolved from five enzymatic systems. These were *Got-A*, *Got-B*, *Mdh-A*, *Cat-A*, *Est-A*, *Est-B*, *Pox-B*, and *Pox-C*. The four former loci were monomorphic and the four latter were polymorphic. Therefore, the proportion of polymorphic loci (P) was estimated to be 50%. This proportion, as a measure of genetic variation, was higher than the 26% reported by Nevo (1978) for 15 plant species. This high (P) value could be attributed to the tetrasomic inheritance as well as self-incompatibility in alfalfa.

JAST

The *Est-A*, *Est-B*, and *Pox-C* loci exhibited unambiguous tetrasomic inheritance (Figures 1 and 2) for two, three and three allozymes, respectively, in all populations and were used to study the populations genetic structure. The polymorphic *Pox-B* locus was not included in this study due to faint or inconsistent staining. A summary of the allele frequencies for three polymorphic loci are given in Table 1. All populations shared the same highest allelic



Figure 2. Banding pattern of peroxidase for 17 alfalfa seedlings, showing one unambiguous polymorphic locus (Pox-C). The allele's nomenclature and the genotypes of some individuals are indicated.



Alfalfa	$\mathbf{N}^{a}$	Est-B		Est-A		Pox-C			
population		$b_1$	$b_2$	$b_3$	$a_1$	$a_2$	<i>c</i> <sub>1</sub>	<i>c</i> <sub>2</sub>	<i>C</i> <sub>3</sub>
Ranger	69	0.149	0.366	0.486	0.062	0.938	-	-	-
Sea-river	70	0.143	0.500	0.357	0.104	0.896	-	-	-
Ghara-yonja	94	0.181	0.352	0.468	0.188	0.812	0.283	0.042	0.675
Syn <sub>1</sub>	100	0.214	0.339	0.446	0.120	0.880	0.303	0.037	0.660
Syn <sub>2</sub>	102	0.172	0.360	0.468	0.127	0.873	0.281	0.071	0.648
Syn <sub>3</sub>	111	0.243	0.279	0.478	0.113	0.887	0.291	0.051	0.658
Average		0.183	0.366	0.450	0.119	0.881	0.288	0.052	0.660

 Table 1. Allele frequencies for three polymorphic loci in the studied populations of alfalfa.

<sup>a</sup> Number of individuals (seedlings) analyzed for each population and each enzyme system.

- : Data not available

Table 2. Observed (expected) levels of heterozygosity in alfalfa populations based on polymorphic loci.

Alfalfa - population		Meen			
	Est-B	Est-A	Pox-C	heterozygosity	
	(3 alleles)	(2 alleles)	(3 alleles)	neterozygosity	
Ranger	0.843 (0.926)**	0.198 (0.225)*	-	0.521 (0.575)	
Sea-river	$0.846 (0.921)^{ns}$	$0.320 (0.354)^{*}$	-	0.583 (0.637)	
Ghara-yonja	0.872 (0.936) <sup>ns</sup>	$0.526 (0.564)^{*}$	$0.690 (0.790)^{\rm ns}$	0.696 (0.753)	
$Syn_1$	$0.906 (0.945)^{\rm ns}$	0.366 (0.400)**	$0.780~{(0.820)}^{*}$	0.684 (0.722)	
Syn <sub>2</sub>	$0.853~{(0.938)}^{*}$	$0.368 (0.420)^{\rm ns}$	$0.760~{(0.810)}^{**}$	0.660 (0.721)	
Syn <sub>3</sub>	$0.833 (0.938)^{*}$	$0.333 (0.380)^{ns}$	$0.730 (0.810)^{ns}$	0.632 (0.709)	

Data not available

ns, \*, \*\* Non-significant difference from equilibrium and significant at 5 and 1% levels of probability, based on two alleles model of  $\chi^2$ - test.

frequencies at three loci:  $b_3$  at Est-B,  $a_2$  at Est-A and  $c_3$  at *Pox-C*, with the exception of  $b_2$  at Est-B for Sea-river, a cultivar that originated from Australia. The observed and expected values of heterozygosity based on these loci are shown in Table 2. Very high levels of heterozygosity were found within alfalfa population, especially for Est-B and Pox-C loci. The observed mean heterozygosities (Ho) ranged from 0.521 in the improved variety Ranger to 0.696 in Ghara-yonja landrace. The  $Syn_1$  generation exhibited the highest *Ho* of 0.684. But, the two successive open pollinated generations showed reduction in heterozygosity. The reduction was about 5% from Syn<sub>1</sub> to Syn<sub>2</sub> and also from Syn<sub>2</sub> to Syn<sub>3</sub>. The expected mean heterozygosity (He), as a measure of genetic diversity, ranged from 0.575 to 0.753, indicating high withinpopulation variability. This result was in agreement with the He level of 0.665 to 0.717

in some cultivars evaluated by using seven SSR loci (Flajoulot *et al.*, 2005). Mengoni *et al.* (2000), Brummer *et al.* (1991), and Puppilli *et al.* (1996) also confirmed the high heterozygosity of alfalfa populations by using RAPD, SSR, or RFLP markers.

Results obtained on genetic distances (Nei, 1972) between pairs of alfalfa populations are shown in Table 3. Apparently, cultivar Seariver, originating from Australia, was a more distant population, but, Ranger did not show any appreciable divergence from other alfalfa populations.

Estimation of  $F_{ST}$ ,  $F_{IS}$ , and  $F_{IT}$  for three generations of Syn<sub>1</sub>, Syn<sub>2</sub> and Syn<sub>3</sub> and one of their 12 polycrossed parents (Ghara-yonja) by three polymorphic loci (Table 4) indicated that, in spite of small differences in the effect of the three allozymic loci on the above mentioned parameters, similar results were obtained from all loci. Therefore, the overall

Table 3. Nei's genetic distances between alfalfa populations based on analyzed polymorphic loci.

	Ranger	Sea-river	Ghara-yonja	Syn <sub>1</sub>	Syn <sub>2</sub>	Syn <sub>3</sub>
Ranger	-	$0.0224^{*}$	0.0075	0.0051	0.0021	0.0115
Sea-river		-	$0.0260^{*}$	$0.0250^{*}$	$0.0211^{*}$	$0.0486^{**}$
Ghara-yonja			-	0.0042	0.0019	0.0071
Syn <sub>1</sub>				-	0.0012	0.0026
Syn <sub>2</sub>					-	0.0058
Syn <sub>3</sub>						-
Mean distance	0.0097	0.0281	0.0093	0.0076	0.0058	0.0151

\* and \*\*: Significant at 5 and 1% levels of probability, according to Workman and Niswander (1970)  $\chi^2$ -values.

 Table 4. Average F-statistics in alfalfa populations (three synthetic generations and Ghara-yonja landrace) obtained from three allozymic loci.

Loci –	<i>F</i> components				
LOCI	$F_{IS}$	$F_{ST}$	$F_{IT}$		
Pox-C	0.0796	0.0022	0.0817		
Est-B	0.0801	0.0080	0.0874		
Est-A	0.0986	0.0245	0.1156		
Average	0.0861	0.0116	0.0948		

inbreeding coefficient ( $F_{IT}$ ) was mainly due to inbreeding effect ( $F_{IS}$ ) or double reduction in alfalfa rather than random genetic drift or population differentiation ( $F_{ST}$ ). The mean values of  $F_{IT}$ ,  $F_{IS}$ , and  $F_{ST}$  were estimated as 0.0948, 0.0861 and 0.0116, respectively.

Accordingly, due to very large intrapopulation diversity in alfalfa, the three generations of synthetic alfalfa did not generate a large amount of differentiation compared to Ghara-yonja landrace. Flajoulot *et al.* (2005) also reported low amount of  $F_{ST}$ parameter (between 0.001 and 0.009), but significant for some pairs of alfalfa populations.

#### REFERENCES

- Anonymous. 2007. Agricultural Statistics. Information and Statistical Center, Ministry of Jihad-Keshavarzi (Agriculture), Teheran, Iran.
- Brummer, E. C., Kochert, G. and Bouton, J. H. 1991. RFLP Variation in Diploid and Tetraploid Alfalfa. *Theor. Appl. Genet.*, 88: 89-96.
- 3. Demarly, Y. 1954. Etude de l' Heredite de la Bigarrure de la Fleur Chez la Luzerne. *Ann. Amelior. Plants*, **4:** 5-20.
- Flajoulot, S., Ronfort J., Boudin, P., Barre, P., Huguet, T., Hugyhe, C. and Julier, B. 2005.

Genetic Diversity among Alfalfa (*Medicago sativa*) Cultivars Coming from a Breeding Program, Using SSR Markers. *Theor. Appl. Genet.*, **111:** 1420-1429.

- Hederick, P. W. 2005. *Genetic of Populations*. 3<sup>rd</sup> Edition, Jones and Bartlett Publishers, Massachusetts, USA, PP: 97-109.
- McCoy, T. J. and Bingham, E. T. 1988. Cytology and Cytogenetics of Alfalfa. In: "Alfalfa and Alfalfa Improvement". (Eds.): Hanson et al., ASA, CSSA. Madison, WI, USA, 29: 737-776.
- Mengoni, A., Gori, A. and Bazzicalupo, M. 2000. Use of RAPD and Micro Satellite (SSR) Variation to Assess Genetic Relationships among Populations of Tetraploid Alfalfa (*Medicago sativa*). *Plant Breed.*, **119**: 311-317.
- Michaud, R., Lehman, W. F. and Rumbauch, M. D. 1988 World Distribution and Historical Development. In: "Alfalfa and Alfalfa Improvement". (Eds.): Hanson et al., Agronomy Monographs. ASA, CSSA, Madison, WI, USA, 29: 25-91.
- Nei, M. 1972. Analysis of Gene Diversity in Subdivided Populations. *Proc. Natl. Acad. Sci.*, 70: 3321-3323.
- Nevo, E. 1978. Genetic Variation in Natural Populations: Patterns and Theory. *Theor. Pop. Biol.*, 13: 121-177.
- Olson, P. D. and Varner, J. E. 1993. Hydrogen Peroxides and Lignification. *Plant J.*, 4: 887-892.



- Pupilli, F., Businelli, S., Paolocci, F., Scotti, C., Damiani, F. and Arcioni, S. 1996. Extent of RFLP Variability in Tetraploid Population of Alfalfa (*Medicago sativa*). *Plant Breed.*, **115**: 106-112.
- Sinskaya, G. N. 1950. Flora of Cultivated Plants of USSR XIII, Perennial Leguminous Plants. Part I: "*Medic, Sweet Clover*". Fenugreek, Moscow.
- Soltis, D. E. and Soltis, P. S. 1990. *Isozymes in Plant Biology*. Chapman and Hall, London, 259 PP.
- 15. Stanford, E. H. 1951. Tetrasomic Inheritance in Alfalfa. *Agron. J.*, **43**: 222-225.
- Tysdal, H. M. and Crandal, B. H. 1948. The Polycross Progeny Performance as an Index of Combining Ability of Alfalfa Clones. *J. Amer. Soc. Agron.*, 40: 293-306.

- Valizadeh, M., Moghaddam, M., Talebi, P., Kazemi, M. H., Monirifar, H. and Hassanpanah, D. 2002. Breeding and Introduction of Suitable Alfalfa Cultivars in East-Azarbaijan. University of Tabriz Research Affairs Pub., Tabriz, Iran. pp 120
- Workman, P. L. and Niswander, J. D. 1970. Population Studies on Southern Indian Tribes. II. Local Genetic Differentiation in the Papago. *Am. J. Hum. Genetic.*, **22**: 24-29.
- Wricke, G. and Weber, W. E. 1986. Quantitative Genetics and Selection in Plant Breeding. Walter de Gruyer. Berline, pp. 406(p.30-38).
- Wright, S. 1951. The Genetical Structure of Populations. Ann. Eugen., 15: 323-354.

## بررسی تنوع ژنتیکی نسلهای سنتتیک و ارقام یونجه با استفاده از وراثت تتراسومیک نشانگرهای آلوزیمی

م. ولیزاده، م. مهیجی، ن. یاسینزاده ، ص. نصرالهزاده و م. مقدم

## چکیدہ

برای اندازه گیری تنوع ژنتیکی درون و بین ارقام یونجه و نیز سه نسل از یک جمعیت سنتیک (Syn<sub>1</sub> ، Syn<sub>1</sub>) و <sub>۲</sub> Syn<sub>7</sub>) به دست آمده از تلاقی پلی کراس بین ۱۲ والد، از الکتروفورز آنزیم ها استفاده شد. از بین پنج سیستم آنزیمی مورد مطالعه در حدود ۱۰۰ گیاهچه ۱۰ روزه از هر جمعیت با الکتروفورز در ژل افقی پلی اکریلامید، سه مکان ژنی ایزوزیمی تشخیص داده شد که وراثت تتراسومیک نشان دادند. با بررسی این مکانهای پلی مورف، سطح بسیار بالایی از هتروزیگوسی (بین ۲۵/۰ تا ۲۹/۰) در داخل جمعیتهای یونجه مشاهده شد. کاهش میزان هتروزیگوسی از متروزیگوسی (بین ۲۵/۰ تا ۲۹۹۹) در داخل جمعیتهای یونجه مشاهده شد. کاهش میزان هتروزیگوسی از متروزیگوسی (بین ۲۵/۰ تا ۲۹۹۹) در داخل جمعیتهای یونجه مشاهده شد. کاهش میزان هتروزیگوسی از ۲۵/۱ یونجه میاه در داخل جمعیتهای یونجه مشاهده شد. کاهش میزان هتروزیگوسی از ۲۵/۰ یونجه ۲۹/۰ به ۲۹/۰ به داخل جمعیتهای یونجه مشاهده شد. کاهش میزان هتروزیگوسی از ۲۵/۱ یونجه میران دادند. با بردن آزمون <sup>۲</sup> هاب کردید که نسل اخیر (Syn بالایی از هتروزیگوسی (بین ۲۵/۰ تا ۲۹۹۹ به جرود ۵/۰) در داخل جمعیتهای یونجه مشاهده شد. گردید که نسل اخیر (Syn بالایی از میه Syn به ۲۹/۰) به تعاد معیتهای یونجه مشاهده شد. آماره F<sub>1</sub> واریت نشان داد که میزان ضریب خویشامیزی کل /۲/۰ به جرال بر آورده شده عمدتا (/۲/۱ به ۲۹/۹) به تعایزی یا کاهش دو گانه قابل انتساب است تا رانش تصادفی ژنتیکی یا تمایز جمعیتی (/۲/۱ اینه) بنابراین، به دلیل وجود تنوع بسیار بالای درون - جمعیتی، یونجه سنتیک نتوانست تمایز واریته ای چشمگیری نولید کند. با وجود این، استفاده از مکانهای آلوزیمی میتواند به شکل موفقیت آمیزی برای بر آورد پارامترهای ژنتیکی جمعیتها و ارقام یونجه بکار رود.