

## Effects of Some Indigenous Plant Extracts on *Meloidogyne javanica* Infesting Eggplant and Pepper under Greenhouse Condition

I. Kepenkci<sup>1,2,\*</sup> and H. D. Saglam<sup>3</sup>

### ABSTRACT

Among the major pests of vegetables are Root-Knot Nematodes (*Meloidogyne* spp.) (RKNs), which cause loss of production due to galling and reduction in root development and shoot growth. Herein, the efficacy of plant extracts of *Capsicum frutescens*, *Hyoscyamus niger*, *Melia azedarach*, *Xanthium strumarium* and *Achillea wilhelmsii* were evaluated at 3 concentrations (3, 6, and 12%) against *Meloidogyne javanica* on pepper and eggplant. Experiments were carried in pots under greenhouse condition, using pepper cv. Charleston and eggplant cv. Kemer as assay plants that are commonly cultivated in Turkey. Approximately 3,000 eggs of *M. javanica* were used for hatching test and 1,000 J2 of *M. javanica* were used for mortality test. Approximately, 5 mL of plant extracts were added by a syringe into the soil. Each experiment was arranged in a Randomized Block Design (RBD) with 5 replications. The control (+) pots received water containing *M. javanica* and the control (-) pots received only water. At the end of the experiment, plants heights and weights were measured. According to the results, all plant extracts showed a different level of nematicidal activity at 3, 6, and 12% concentrations. The plant extracts of *H. niger*, and *X. strumarium* at 12% concentration were found more efficient than *M. azedarach*, *C. frutescens* and *A. wilhelmsii* extract on egg hatching, on both pepper and eggplant plants. J2s mortality experiments showed that 12% concentration of *H. niger*, *M. azedarach* and *X. strumarium* were more effective against *M. javanica* than the other treatments, on both pepper and eggplant plants. In general, there was no significant difference was found among nematode mortality and growth parameters (such as plant height, the fresh and dry weights of the above-ground parts of the plants, fresh and dry weights of roots of both pepper and eggplant plants). Accordingly, using *H. niger* and *X. strumarium* plant extracts can provide effective methods of *M. javanica* control.

**Keywords:** Hatching test, Eggplant, Nematicidal effect, Pepper.

### INTRODUCTION

Plant parasitic nematodes, especially Root-Knot Nematodes (RKNs) from the genus *Meloidogyne*, are widely distributed and cause significant yield losses in a wide range of crops (Davis, 2005; Luc *et al.*, 2005). *M. incognita*, *M. arenaria*, *M. javanica* and *M. hapla* are the most commonly found root-

knot nematode species in Turkey (Kepenkci, 2012). Current management of nematodes has been attempted using plant resistance, crop rotation, cultural practices, biological control, or using chemical nematicides (Chitwood, 2002; Khan *et al.*, 2007, 2008, 2012). Traditionally, synthetic nematicides are used to control *Meloidogyne* spp. These synthetic nematicides increase production costs and have potentially

<sup>1</sup> Department of Plant Protection, Faculty of Agriculture, Gaziosmanpasa University, 60250 Tokat, Turkey.

<sup>2</sup> Department of Plant Protection, Faculty of Agriculture, Kyrgyz Turkish Manas University, 720038, Bishkek-Kyrgyzstan.

<sup>3</sup> Department of Plant Protection, Faculty of Agriculture, Ahi Evran University, 40200 Kırsehir, Turkey.

\*Corresponding author; e-mail: kepenkci@gmail.com



negative impacts on the human health and the environment including on non-target organisms. New strategies are needed to substitute traditional chemicals such as antagonistic plants or plant extracts against plant-parasitic nematodes (Chitwood, 2002; Akhtar, 2004). Many plants including Brassicaceae, Asteraceae, Myrtaceae and Rutaceae families' member plants contain nematicidal compounds (Sukul, 1992; Andres *et al.*, 2012). The use of plant extracts as an alternative to synthetic pesticides for control of RKNs is becoming important and, in recent years, studies on plant extracts have accelerated (Lee, 2011, Ntalli *et al.*, 2011; Andres *et al.*, 2012; Oka, 2012). *Azadirachta indica* is well known as a pesticide and controlling insect, mite, nematode, and plant diseases (Agbenin *et al.* 2005; Bashir, 2013; Anwar, 2015; Benelli, 2015). *Tagetes* spp. includes a-Terthienyl and this content shows highly nematicidal activities against plant-parasitic nematodes, especially *Meloidogyne* spp. (Ploeg, 1999). Nematicidal effect of garlic has been studied against *Meloidogyne* spp. (Bekhiet *et al.*, 2010; El-Nagdi and Youssef, 2013). Therefore, the objective of this study was to determine the efficacy of plant extracts from *Capsicum frutescens*, *Hyoscyamus niger*, *Melia azedarach*, *Xanthium strumarium* and *Achillea wilhelmsii* as alternative nematicides against *Meloidogyne javanica* on pepper and eggplant under greenhouse condition.

## MATERIALS AND METHODS

### Nematodes

The egg masses of *M. javanica* were collected from tomato roots infected with the nematode (SC-2121 variety susceptible to RKNs). RKNs eggs were extracted from roots using a 0.575% NaOCl solution and the eggs were collected using the modified technique described by McClure *et al.* (1973). Eggs were washed by rinsing with tap water through a 75  $\mu\text{m}$  sieves, collected

on a 26  $\mu\text{m}$  sieve and transferred into the distilled water. The egg suspension was poured on to a cotton-wool filter and incubated at  $26\pm 2^\circ\text{C}$ . Emerged J2s were collected daily for up to 4 days and stored at  $4^\circ\text{C}$  until used for the experiment. The population density of J2 and eggs of *M. javanica* were determined from 3 replications of one mL subsample of an inoculum suspension. A total number of J2 and eggs were calculated by multiplying the mean number of nematodes per subsamples by the number of subsamples in the total volume.

### Plant Material

Five selected indigenous plants were collected from different regions of Anatolia, Turkey. These were *Capsicum frutescens* (Solanaceae), *Hyoscyamus niger* (Solanaceae), *Melia azedarach* (Meliaceae), *Xanthium strumarium* (Asteraceae) and *Achillea wilhelmsii* (Asteraceae).

### Preparation of Plant Extracts

Plants leaves were picked from their branches and spread on polythene in the laboratory for ten days to air dry. After that, they were dried at  $80^\circ\text{C}$  for 3-4 days. The dried plants parts were ground to fine particles using a blender. Ethanol was added to the ground plant powder and shaken on a rotary shaker at 120 rpm for 48 hours. The solution was filtered to remove solids and the material was vacuum concentrated in a rotary evaporator at  $50\text{-}60^\circ\text{C}$  to obtain the corresponding organic crude extracts (Brauer and Davkota, 1990; Lee *et al.*, 2008). Each plant extract was prepared in  $200\text{ g } 200\text{ mL}^{-1}$  and was used immediately in greenhouse tests. Suspensions of concentrations of 3, 6, and 12% were prepared with distilled water (Orisajo *et al.*, 2007).

## Greenhouse Pot Experiments

According to treatment, approximately 3,000 eggs (egg hatching test) and 1,000 J2s (mortality test) of *M. javanica* were applied to the root of pepper and eggplants (Adekunle and Akinlua, 2007; Liman *et al.*, 2010). The whole plant extracts were applied at the same time with *M. javanica* on pepper (*C. annuum*, Charleston variety) and eggplant (*Solanum melongena*, Kemer variety) roots. Each experimental unit consisted of the plastic pot (10×10 cm) containing 800 cm<sup>3</sup> sterilized loamy sand/pot (sterile moist loamy soil, 80% sand, 15% silt, and 5% clay) and a seedling of pepper or eggplant. Eggs or J2s of *M. javanica* were applied to 2 cm depth of the soil surface. Approximately 5 mL of extract was added by a syringe into the soil. The control (+) pots received water containing *M. javanica* and the control (-) pots received only water. Each experiment was arranged in a Randomized Block Design (RBD) with 5 replications. Greenhouse conditions were recorded by a data logger (HOBO-Onset computer cooperation, USA). During the experiment, the average temperature and the humidity were recorded at 25.04±4 °C and 30.14±10%, respectively. Nine weeks after inoculation, plants were uprooted and their roots gently washed with tap water. Galled roots were placed in an aqueous solution of phloxine B (0.15 g L<sup>-1</sup> tap water) for 15-20 minutes. After weighing, egg masses were counted. To facilitate counting of egg masses, they were stained red with phloxine B (Fenner, 1962; Dickson and Struble, 1965; Holbrook *et al.*, 1983). Root systems were rinsed in tap water to remove the residual stain from the roots, and egg masses were counted under a dissecting microscope. Plant height was measured from the base to the terminal bud. Fresh roots were weighed, then dried at 70°C for 48 hours in the incubator and weighed again. Data recorded included the total number of egg masses for each plant, plant height, the fresh and dry weight of the plant shoots and roots.


## Statistical Analysis

Data were analyzed by analysis of variance, and means were compared using Duncan's multiple range test (SPSS, 1999). Differences were reported at  $P \leq 0.05$ .

## RESULTS

### Egg Hatching Test

The concentrations of 3, 6, and 12% of plant extracts displayed varied nematicidal effects on egg hatching of *M. javanica*. On eggplant, 12% concentration of *X. strumarium* showed the highest nematicidal effect (11.13±1.3). This was followed by 12% of *H. niger* on eggplant (15.53±2.4), 12% *X. strumarium* on pepper (17.83±1.3) and 12% of *H. niger* on pepper (19.53±3.4) (Figure 1-A). Their differences were found statistically significant ( $P \leq 0.05$ ). Also, 3 and 6% concentrations of *C. frutescens* plant extracts influenced *M. javanica* both on pepper and eggplant, although this effect was not statistically important ( $P \leq 0.05$ ). Although the low effect was observed in 3% and 6% concentrations of *H. niger*, the high effect was observed in 12% concentration. The 3, 6, and 12% concentrations of *M. azedarach* plant extract affected nematode both on pepper and eggplant, although these effects were not statistically important ( $P \leq 0.05$ ). Treatments 3, 6, and 12% concentrations of *A. wilhelmsii* plant extracts had less influence on *M. javanica* on both pepper and eggplant (Figure 1-A). The concentration of 12% of *X. strumarium* and *H. niger* were highly effective against *M. javanica* compared with the other extracts. As the concentration increased, the nematicidal effects also increased against *M. javanica* on pepper and eggplant.



**Figure 1.** Effect of selected concentrations (3, 6 and 12%) of some indigenous plant extracts on *Meloidogyne javanica* reproduction in greenhouse pepper and eggplants. [(A) Total number of egg masses; (B) Plant height; (C) Fresh; (D) Dry weight of the aboveground parts of plants; (E) Fresh, and (F) Dry weights of roots].

## J2 Mortality Test

Different effects were observed in 3, 6 and 12% concentrations of all plant extracts against 2. stage of *M. javanica*. The highest effects were observed at the concentration of 12% of *X. strumarium* ( $19.3 \pm 2.3$ ) on eggplant. This was followed by 12% of *M. azedarach* ( $21.2 \pm 2.2$ ) on pepper, 12% of *H. niger* ( $22.5 \pm 2.1$ ) on eggplant, 12% of *M. azedarach* ( $33.3 \pm 3.8$ ) on eggplant and 12% of *X. strumarium* ( $42.3 \pm 2.1$ ) on pepper,


respectively (Figure 2-A). Their differences were statistically significant ( $P \leq 0.05$ ). The concentrations of 3, 6, and 12% of *C. frutescens* and *A. wilhelmsii* plant extracts were less effective against *M. javanica* on both pepper and eggplant. Their extracts displayed lower nematicidal effect, therefore, this impact was not significantly important. Additionally, the 3% concentration of *H. niger* had a low nematicidal effect on both pepper and eggplant besides that 6% concentration of *H. niger* had lower effect against *M. javanica*

on eggplant (Figure 2-A). The increase in *M. azedarach* and *X. strumarium* concentrations led to increase in the nematicidal effect against *M. javanica* on pepper and eggplant.

### Effect of Plant Extracts on Plant Growth

At the end of the egg hatching and J2

mortality test experiments, growth of pepper and eggplant plants were measured on plant height, the fresh and dry weight of the aboveground plant parts, and fresh and dry weight of roots. There was no significant difference between the control and plant extract treated plants in parameters such as plant height, the fresh and dry weight of the aboveground parts of plants, and fresh and dry weight of roots, both in pepper and



**Figure 2.** Effect of selected concentrations (3, 6 and 12%) of some indigenous plant extracts on *Meloidogyne javanica* reproduction (mortality test, 1000 J2s were applied) in greenhouse pepper and eggplants. Symbols A to F are defined in Figure 1.



eggplant plants from both egg hatching and mortality tests [Figures 1 (B-F) and 2 (B-F)] ( $P < 0.05$ ). The 12% concentration of *X. strumarium* and *H. niger* plants extracts caused a significant reduction in root galling and nematode population on pepper and eggplant, even though the plant height, the fresh and dry weight of the aboveground plant parts, and fresh and dry weight of roots were not increased by using plant extracts [Figures 1 (B-F) and 2 (B-F)] ( $P < 0.05$ ).

## DISCUSSION

In this study, the nematicidal activities of *C. frutescens*, *H. niger*, *M. azedarach*, *X. strumarium* and *A. wilhelmsii* were evaluated on egg hatching and J2 mortality of *M. javanica* in pots of pepper and eggplant plants under greenhouse condition. The plant extracts used in our study showed a different level of nematicidal effect in a concentration-dependent manner. *C. frutescens* contains phytochemicals such as capsaicin and capsaicinoids, which are used to control plant diseases (*Aspergillus flavus*, *A. niger*, *Penicillium* sp. and *Rhizopus* sp.) (Soumya and Nair, 2012) and storage insects (*Callosobruchus maculatus* and *Sitophilus zeamais*) (Oni, 2011) and were found highly effective to them. In this study, *C. frutescens* was identified as having low nematicidal activities against *M. javanica* on pepper and eggplant plants. Kepenekci *et al.* (2016) applied *C. frutescens* extracts against *M. javanica* on tomato and found low nematicidal activity against nematode. Our study supports this finding. *H. niger* is known to be rich in tropane alkaloids. Tropane alkaloids of hyoscyamine and scopolamine show the insecticidal effect by affecting the activity of the neurotransmitter acetylcholine (Roddick, 1991; Shonle and Bergelson, 2000). Acetylcholine mechanism is enzymatic and inhibition of acetylcholinesterase is the target for the control of plant-parasitic nematodes by tropane alkaloids. The concentration 12% of *H. niger* showed higher nematicidal effect

on eggplant and pepper infected by *M. javanica*. Kepenekci *et al.* (2016) also reported similar results on tomatoes infected by *M. javanica* both *in vivo* and *in vitro* studies. *M. azedarach* has potent active ingredient azadirachtin. Azadirachtin is used for controlling insects, diseases, and weeds effectively. (Isman, 2006; El-Ghany *et al.*, 2015; Phuagphong *et al.*, 2015). Salgado *et al.* (2003) reported that 3 different (methanol, chloroform, and ethyl acetate) extracts of *M. azedarach* against *Meloidogyne exigua* were used and methanol and chloroform extracts of *M. azedarach* caused 94.3 and 82.5% mortality, while this rate was 48.7% for ethyl acetate. Ntalli *et al.* (2010) stated that melia methanol extracts higher than 0.08% showed higher nematicidal activity. In our study, the concentration of 6 and 12% of *M. azedarach* impact on mortality of J2 of *M. javanica* was higher both on eggplant and pepper plants. In contrast, all concentrations of *M. azedarach* extracts were had a lower effect on egg hatching on *M. javanica*. Chaudhary *et al.* (2013) reported that the nematicidal activities of ethanol and aqueous extracts of *X. strumarium* were tested against egg hatching and mortality of J2 of *M. incognita* and hot water and ethanol extracts showed lower effects on nematode, while Kepenekci *et al.* (2016) reported that *X. strumarium* demonstrated highly active nematicidal activities on tomato against *M. javanica*. The extract of *A. wilhelmsii* is known as an insecticide. It shows fumigant toxicity against stored pests (Asghari *et al.*, 2010). All concentration of *A. wilhelmsii* extracts displayed lower nematicidal activities in both egg hatching and mortality experiment on eggplant and pepper. According to the results of this study, *X. strumarium*, *M. azedarach* and *H. niger* present good inhibitory effect on nematode egg hatching and juvenile reproduction. The use of *M. azedarach*, *H. niger* and *X. strumarium* extract are suggested as potential substitutes for synthetic nematicides used in the management of RKNs disease in the greenhouse vegetable production. Further

studies will be conducted in the field conditions for determining the nematicidal ability of plant extracts. Therefore, the use of indigenous plant extracts should be considered in integrated pest management strategies. It is suggested that further trials be conducted in the field on the basis of the promising results from these studies. It is interesting to note that high concentration of the plant extracts showed a greater effect on egg hatching. Hence, we suggest that high concentrations of plants extracts are required for effective nematode control.

### ACKNOWLEDGEMENTS

We thank Dr. F. D. Erdoğan, Dr. T. Katı Çekengil and Dr. E. Evlice [Department of Entomology (Nematology Lab.), Plant Protection Central Research Institute, Ankara-Turkey] for helping us during this study. Also, we thank Plant Protection Central Research Institute, Ankara, Turkey which supplied the plant extracts.

### REFERENCES

1. Adekunle, O. K. and Akinlua, A. 2007. Nematicidal Effects of *Leucaena leucocephala* and *Gliricidia sepium* Extracts on *Meloidogyne incognita* Infecting Okra. *J. Agr. Sci.*, **2**: 53-63.
2. Agbenin, N.O., Emechebe, A. M., Marley, P. S. and Akpa, A. D. 2005. Evaluation of Nematicidal Action of some Botanicals on *Meloidogyne incognita* in Vivo and in Vitro. *J. Agr. Rural Dev. Trop. (JARTS)*, **106**(1): 29-39.
3. Akhtar, M. 2004. Current Options in Integrated Management of Plant-Parasitic Nematodes. *Integrated Pest Manage. Rev.*, **2**: 187-197.
4. Andres, M. F., Coloma, A.G., Sanz, J., Burillo, J. and Sainz, P. 2012. Nematicidal Activity of Essential Oils: A Review. *Phytochem. Rev.*, **11**: 371-390.
5. Anwar, W., Haider, M. S., Aslam, M., Shahbaz, M., Khan, S. N. and Bibi, A. 2015. Assessment of Antifungal Potentials of some Aqueous Plant Extracts and Fungicides against *Alternaria alternata*. *J. Agric. Res.*, **53**: 1.
6. Asghari, J., Khani, A. and Rakhshani, E. 2010. Insecticidal Activity of Flower Essential Oil from *Achillea wilhelmeii* (Asteraceae) against Two Stored Product Pests. *Proceedings of the 19<sup>th</sup> Iranian Plant Protection Congress*, July 31-August 3, 2010, Iranian Reserch Institute of Plant Protection, Tehran.
7. Bashir, M., Gogi, M. D., Ashfaq, M., Afzal, D. M., Khan, M. A. and Ihsan, M. 2013. The Efficacy of Crude Aqueous Extracts of some Plants as Grain Protectants against the Stored Grain Mite, *Rhizoglyphus tritici*. *Turk. J. Agr. For.*, **37**(5): 585-594.
8. Bekhiet, M. A., Kella, A., El-Gindi, A. Y. and Hammad, E. A. 2010. Effect of Certain Inorganic Acids and Garlic Cloves Oil for Controlling the Root-Knot Nematode *Meloidogyne javanica* Infecting Banana Plant. *Egypt. J. Agronematol.*, **9**: 202-214.
9. Benelli, G., Bedini, S., Cosci, F., Toniolo, C., Conti, B. and Nicoletti, M. 2015. Larvicidal and Ovideterrent Properties of Neem Oil and Fractions against the Filariasis Vector *Aedes albopictus* (Diptera: Culicidae): A Bioactivity Survey across Production Sites. *Parasitol. Res.*, **114**(1): 227-236.
10. Brauer, M. and Davkota, B. 1990. Control of *Thaumatopoea pycnandra* (Den. & Schiff) by Extracts of *Melia azedarach* L. (Meliaceae). *J. Appl. Entomol.*, **110**: 128-135.
11. Chaudhary, K. K., Haile, A. Ayresea, Z. G., Semereab, G. and Weldegergish, T. 2013. Nematicidal Activity of Eritrean Weeds Plants against Root-kKnot Nematode. *Nematropica*, **43**(2): 207-215.
12. Chitwood, D. J. 2002. Phytochemical Based Strategies for Nematode Control. *Annu. Rev. Phytopathol.*, **40**: 221-249.
13. Davis, R. F. 2005. Effect of the Southern Root-Knot Nematode on Watermelon Yield. In: "Georgia Vegetable Research-Extension Report 2004", (Eds.): Kelly, W. T. and Langston, D. R. University of Georgia, Tifton, GA, USA.
14. Dickson, D. W. and Struble, F. B. 1965. A Sieving-Staining Technique for Extraction of Egg Masses of *Meloidogyne incognita* from Soil. *Phytopathol.*, **55**: 497.
15. El-Ghany, A., Roushdy, M. M. and Mohamed, A. A. 2015. Efficacy of Certain



- Plant Extracts as Safe Fungicides against Phytopathogenic and Mycotoxigenic Fungi. *Agric. Biol. Sci. J.*, **1(3)**: 71-75.
16. El-Nagdi, W. M. A. E. and Youssef, M. M. A. 2013. Comparative Efficacy of Garlic Clove and Castor Seed Aqueous Extracts against the Root-Knot Nematode, *Meloidogyne incognita* Infecting Tomato Plants. *J. Plant Prot. Res.*, **53(3)**: 285-288.
  17. Fenner, L. M. 1962. Determination of Nematode Mortality. *Plant Dis. Rep.*, **46**: 383.
  18. Holbrook, C. C., Knauff, D. A. and Dickson, D. W. 1983. A Technique for Screening Peanut for Resistance to *Meloidogyne arenaria*. *Plant Dis.*, **57**: 957-958.
  19. Isman, M. B. 2006. Botanical Insecticides, Deterrents, and Repellents in Modern Agriculture and an Increasingly Regulated World. *Annu. Rev. Entomol.*, **51**: 45-66.
  20. Kepenekci, İ. 2012. Nematoloji (Bitki Paraziti ve Entomopatogen Nematodlar) [Genel Nematoloji (Cilt-I) ISBN 978-605-4672-11-0, Taksonomik Nematoloji (Cilt-II) ISBN 978-605-4672-12-7]. Eğitim Yayın ve Yayınlar Dairesi Başkanlığı. *Tarım Bilim Serisi Yayın.*, **3**: 1155.
  21. Kepenekci, İ., Erdoğan, D. and Erdoğan, P. 2016. Effects of Some Plant Extracts on Root-Knot Nematodes *In Vitro* and *In Vivo* Conditions. *Turk. J. Entomol.*, **40**: 3-14.
  22. Khan, Z., Kim, Y. H., Kim, S. G. and Kim, H. W. 2007. Observations on the Suppression of Root-Knot Nematode (*Meloidogyne arenaria*) on Tomato by Incorporation of Cyanobacterial Powder (*Oscillatoria chlorina*) into Potting Field Soil. *Bioresour. Technol.*, **98**: 69-73.
  23. Khan, Z., Kim, S. G., Jeon, Y. H., Khan, H. U., Son, S. H. and Kim, Y. H. 2008. A Plant Growth Promoting Rhizobacterium, *Paenibacillus polymyxa* GBR-1, Suppresses Root-Knot Nematode. *Bioresour. Technol.*, **99**: 3016-3023.
  24. Khan, Z., Son, S. H., Akhtar, J., Gautam, N. K. and Kim, Y. H. 2012. Plant Growth-Promoting Rhizobacterium, *Paenibacillus polymyxa* Induced Systemic Resistance in Tomato (*Lycopersicon esculentum*) against Root-Knot Nematode (*Meloidogyne incognita*). *Indian J. Agr. Sci.*, **82**: 613-617.
  25. Lee, T. O., Khan, Z., Kim, S. G. and Kim, Y. H. 2008. Amendment with Peony (*Paeonia suffruticosa*) Root Bark Powder Improves the Biocontrol Efficacy of *Trichoderma harzianum* against *Rhizoctonia solani*. *J. Microbiol. Biotechnol.*, **18**: 1537-1543.
  26. Lee, H. M., Khan, Z., Kim, S. G., Baek, N. I. and Kim, Y. H. 2011. Evaluation of Biocontrol Potential of Some Medicinal Plant Materials Alone and in Combination with *Trichoderma harzianum* against *Rhizoctonia solani* AG 2-1. *Plant Pathol. J.*, **27**: 68-77.
  27. Liman, B., Ibrahim, M., Ibrahim, N. T. and Rabah, A. B. 2010. Effect of Mahogany (*Khaya senegalensis* L.) Leaf Extract on Root-Knot Nematode of Tomatoes (*Lycopersicon esculentum* L.). *Nig. J. Basic Appl. Sci.*, **18**: 272-276.
  28. Luc, M., Sikora, R. A. and Bridge, J. 2005. *Plant Parasitic Nematodes in Subtropical and Tropical Agriculture*. CABI Publishing, Wallingford, UK, 871 PP.
  29. McClure, M. A., Kruk, T. H. and Misaghi, I. 1973. A Method for Obtaining Quantities of Clean *Meloidogyne* Eggs. *J. Nematol.*, **5**: 230.
  30. Ntalli, N. G., Menkissoglu-Spiroudi, U. and Giannakou, I. O. 2010. Nematicidal Activity of Powder and Extracts of *Melia azedarach* Fruits against *Meloidogyne incognita*. *Ann. Appl. Biol.*, **156**: 309-317.
  31. Ntalli, N.G., Ferrari, F., Giannakou, I. and Menkissoglu-Spiroudia, U. 2011. Synergistic and Antagonistic Interactions of Terpenes against *Meloidogyne incognita* and the Nematicidal Activity of Essential Oils from Seven Plants Indigenous to Greece. *Pest Manage. Sci.*, **67**: 341-351.
  32. Oka, Y. 2012. Nematicidal Activity of *Verbesina encelioides* against the Root-Knot Nematode *Meloidogyne javanica* and Effects on Plant Growth. *Plant Soil*, **355(1-2)**: 311-322.
  33. Oni, M. O. 2011. Evaluation of Seed and Fruit Powder of *Capsicum annum* and *C. frutescens* for Control of *Callosobruchus maculatus* (Fab.) in Stored Cowpea and *Sitophilus zeamais* in Stored Maize. *Int. J. Biol.*, **3(2)**: 185 -188.
  34. Orisajo, S. B., Okeniyi, M. O., Fademi, O. A. and Dongo, L. N. 2007. Nematicidal Effects of Water Leaf Extracts of *Acalypha ciliata*, *Jatropha gossypifolia*, *Azadirachta indica* and *Allium ascalonicum* on *Meloidogyne incognita* Infection on Cacao Seedlings. *J. Res. Biosci.*, **3**: 49-53.



35. Phuagphong, P., Nawanopparatsakul, S. and Kitcharoen, N. 2015. Effect of *Azadirachta indica* A. Juss var *Indica*, *Nicotiana tabacum* L., and *Derris elliptica* (Roxb.) on Growth of Duckweed. *Adv. Mat. Res.*, **1060**: 211-214.
36. Ploeg, A. 1999. Greenhouse studies on the effect of marigolds (*Tagetes* spp.) on four *Meloidogyne* species. *J. Nematol.*, **31**: 62-9.
37. Roddick J. 1991. The Importance of the Solanaceae in Medicine and Drug Therapy. In: "*Solanaceae III: Taxonomy, Chemistry, Evolution*", (Eds.): Hawkes J. G., Lester R. N., Nee, M., Estrada, R. N. The Royal Botanic Gardens, Kew, London, UK, PP. 7-23.
38. Salgado, S. L. M., Campos, V. P., Cardos M. D. G. and Salgado, A. P. S. 2003. Hatching and Mortality of Second-Stage Juveniles of *Meloidogyne exigua* in Essential Plant Oils. *Nematol. Brasil.*, **27**: 17-22
39. Shonle, I. and Bergelson, J. 2000. Evolutionary Ecology of the Tropane Alkaloids of *Datura stramonium* L.(Solanaceae). *Evol.*, **54(3)**: 778-788.
40. Soumya, S. L. and Nair, B. R. 2012. Antifungal Efficacy of *Capsicum frutescens* L. Extracts against Some Prevalent Fungal Strains Associated with Groundnut Storage. *J. Agric. Technol.*, **8(2)**: 739-750.
41. SPSS. 1999. *SPSS for Windows, Release 10.0.1*. SPSS, Chicago, IL, USA.
42. Sukul, N. C. 1992. Plant Antagonist to Plant-Parasitic Nematodes. *Ind. Rev. Life Sci.*, **2**: 23-52.

## اثرات عصاره های برخی گیاهان بومی روی *Meloidogyne javanica* آلوده کننده بادمجان و فلفل در شرایط گلخانه

ی. کپنسکی، و.ه. د. ساگلام

### چکیده

نماتدهای گره ریشه (RKNs) (*Meloidogyne* spp.) از شمار آفات عمده سبزیجات است که می تواند به لحاظ ایجاد گال (galling) و کاهش توسعه ریشه و رشد اندام هوایی منجر به تلف شدن تولید شود. در این پژوهش، موثر بودن عصاره گیاهی *Hyoscyamus*، *Capsicum frutescens*، *Achillea wilhelmsii* و *Xanthium strumarium*، *Melia azedarach niger* در ۳ غلظت (۳٪، ۶٪ و ۱۲٪) بر علیه *Meloidogyne javanica* روی گیاه فلفل و بادمجان ارزیابی شد. آزمایش ها در گلدان در شرایط گلخانه روی فلفل کولتیوار چارلستون و بادمجان کولتیوار *Kemer* که در ترکیه به طور گسترده کشت می شود اجرا شد. برای آزمون از تخم بیرون آمدن (*hatching*) تقریباً ۳۰۰۰ تخم *M. javanica* و برای آزمون مرگ و میر (*mortality*)، *J2* ۱۰۰۰ از *M. javanica* استفاده شد. حدود ۵ میلی لیتر از عصاره گیاهی با سرنگ به خاک افزوده شد. طرح هر آزمایش بلوک های تصادفی (RBD) با ۵ تکرار بود. به گلدان های شاهد (+) آب حاوی *M. javanica* و به گلدان های شاهد (-) فقط آب داده شد. در آخر آزمون، بلندی و وزن گیاه اندازه گیری شد. بر اساس نتایج، همه عصاره های گیاهی در غلظت های ۳، ۶ و ۱۲٪ سطح متفاوتی از فعالیت ضد نماتدی نشان دادند. در آزمون بیرون آمدن از تخم در هر دو گیاه فلفل و بادمجان، اثر عصاره های گیاهی *H. niger* و *X.*



*A. wilhelmsii*, *C. frutescens* و *M. azedarach* در غلظت ۱۲٪ بیشتر از *strumarium* بود. نیز، در آزمون مرگ و میر *J2* در هر دو گیاه فلفل و بادمجان، عصاره *M. javanica* موثرتر از دیگر تیمارها بود. به طور کلی، هیچ اختلاف معناداری بین مرگ و میر نماتدها و پارامترهای رشد مانند بلندی گیاه، وزن تر و خشک اندام هوایی گیاهان و وزن تر و خشک ریشه های فلفل و بادمجان وجود نداشت. به این قرار، استفاده از عصاره گیاهان *H. niger* و *X. strumarium* می تواند روش موثری برای کنترل *M. javanica* فراهم کند.