# Studying Genetic Diversity in *Zymoseptoria tritici*, Causal Agent of Septoria Tritici Blotch, by Using ISSR and SSR Markers

M. Dalvand<sup>1\*</sup>, D. Zafari<sup>1</sup>, M. J. Soleimani Pari<sup>1</sup>, R. Roohparvar<sup>2</sup>, and S. M. Tabib Ghafari<sup>3</sup>

#### ABSTRACT

Seventy-five isolates of the fungus Zymoseptoria tritici, collected from seven wheat producing provinces in Iran, were studied using the molecular markers ISSR and SSR to determine its structure and genetic variations within its populations. Results indicated that the populations of the causal agent of this disease had a relatively high level of genetic diversity, with the total average genetic diversity (Ht) of 0.34 in the studied samples. When the genetic diversity was divided by the genetic diversity within and between the tested fungal populations, intra-population and inter-population genetic diversities were, respectively, 69 and 31%. Moreover, results showed that there was a strong gene flow between the studied provinces. The maximum genetic diversity among the studied provinces was observed in Ardebil and Khuzestan Provinces, and cluster analysis also revealed that the fungal populations of these two provinces had the greatest similarity with each other.

Keywords: Gene flow, Intra-population genetic diversity, Molecular markers, Wheat.

#### **INTRODUCTION**

The fungus Zymoseptoria tritici (Desm) (Quaedvlieg et al., 2011) (teleomorph Mycosphaerella graminicola), which was previously named Septoria tritici, is the causal agent of Septoria Tritici Blotch (STB), which is one of the most important pathogenic agents in Dothideomycetes Class.

Knowledge of genetic diversity in various strains of pathogenic agents is of great importance in breeding wheat cultivars for producing cultivars resistant to STB (McDonald *et al.*, 1995).

Numerous studies have shown that the population structure of *Zymoseptoria tritici* 

has high genetic diversity. Czembor and Arseniuk (1999) studied several species of *Septoria* genera and demonstrated that SSRs and ISSRs were the most sensitive markers for investigating DNA polymorphism in this subgenus. The genetic structure of *Zymoseptoria tritici* has been previously studied using markers including SCA, SRRs, RFLP, AFLP, and RAPD (Kema *et al.*, 2002; Linde *et al.*, 2002; Talbot, 2015).

Schnieder *et al.* (2001) studied isolates of this fungus in Germany and showed that diversity of these isolates was related to their geographical separation. Razavi and Hoghes (2004) studied 90 isolates collected from western Canada using the RAPD marker and found that these isolates exhibited high

<sup>&</sup>lt;sup>1</sup>Bu-Ali Sina University, Hamadan, Islamic Republic of Iran.

<sup>&</sup>lt;sup>2</sup>Seed and Plant Improvement Institute, Agricultural Research Education, and Extension Organization (AREEO), Karaj, Islamic Republic of Iran.

<sup>&</sup>lt;sup>3</sup> Department of Seed and Plant Improvement Research, Safiabad Agricultural Research, Education and Natural Resources Center, AREEO, Dezful, Islamic Republic of Iran.

<sup>\*</sup> Corresponding author; e-mail: md5576@yahoo.com

molecular diversity. It was also shown in a study carried out in Kansas (United States of America) that isolates on micro-plot, macro-plot, and state-wide scales showed diversity of 98 percent (Kabbage *et al.*, 2008). In a study conducted in Argentina in 2010 using ISSRs, 81 haplotype groups were identified from 126 isolates of *Mycosphaerella graminicola* collected in bread wheat from various parts of the country (Castillo *et al.*, 2010).

Knowledge of genetic structure of fungi is necessary to adopt management strategies for controlling this disease. Considering the greater efficiency and speed of the ISSR and SSR markers, this research aimed to use them to study genetic diversity in this fungus.

# MATERIALS AND METHODS

## **Collection and Purification of Isolates**

During travels to various regions in Iran (in the years 2014 and 2015), wheat fields were visited and samples of leaves with symptoms of STB were collected from Ilam, Khuzestan, Hamadan, Lorestan, Markazi, Golestan, and Ardebil Provinces and transferred to the laboratory. The infected leaves were dried first to isolate the causal agent of the disease when the suitable opportunity arose. For long-term storage of these colonies, their yeast-like spores were collected and transferred to Eppendorf tubes and kept at -20°C. In total, 75 isolates were selected for study (Table 1).

## **DNA Extraction**

The CTAB method (Doyle and Doyle, 1990) was used with minor alterations to extract DNA from the fungal samples. About 25-50 mg of the isolated fungal spores were weighed out and poured into a sterile porcelain mortar that was cooled in freezer at -20°C beforehand, and were turned into a fine powder by adding liquid

1308

nitrogen and crushing them. Five  $\mu$ L of the extracted DNA were run on 0.8% agarose gel to study DNA quality. The extracted DNA was diluted to the concentration of 20 ng  $\mu$ L<sup>-1</sup> that was further used in Polymerase Chain Reaction (PCR).

#### **ISSR-PCR**

Ten ISSR primers supplied by the SinaClon Company (Iran) were selected according to previous data (Czembor and Arseniuk, 1999) and they used to amplify DNA fragments (Table 2).

For each PCR, 13  $\mu$ L of PCR Master Mix, 1  $\mu$ L of DNA, 1  $\mu$ L of the primer, and 10  $\mu$ L of twice-sterilized distilled water were used. The PCR master mix was supplied by either the SinaClon Company of (Iran) or the Ampliqon Company (Denmark) and contained 1.5 mM of MgCl<sub>2</sub>, 0.8 mM dNTPs, and 0.4 unit of *Taq* polymerase.

Amplification was carried out in a thermocycler (Bio-Rad) and the PCR cycle was as follows: initial denaturation for 2 minutes at 94°C, then, 40 cycles of denaturation for 2 minutes at 94°C, annealing 40 seconds at the tested temperature of 48-53°C and elongation for 1 minute at 72°C. The final elongation cycle was extended to 10 minutes.

#### SSR-PCR

Four pairs of primers supplied by the Macrogen Inc. (South Korea) were used to amplify SSR markers (Owen *et al.*, 1998) (Table 3) and PCR mixture was composed with the same ingredients as above but only master mix supplied by the Ampliqon Company (Denmark) was used for the amplification.

Amplification was carried out in the thermocycler (Bio-Rad) and was programed for initial denaturation for 2 minutes at 94°C, 35 cycles of denaturation for 2 minutes at 94°C, annealing for 35 seconds at the tested temperatures of 52-56°C, and elongation for

No	Name	Province	Origin	Host	No	Name	Province	Origin	Host
	of		(Location)			of		(Location)	
	isolate					isolate			
1	IR1	Ardebil	Moghan	Bread Wheat	39	IR39	Ardebil	Moghan	Bread Wheat
2	IR2	Khuzestan	Zeydoon	Bread Wheat	40	IR40	Ardebil	Moghan	Bread Wheat
3	IR3	Khuzestan	Dezful	Bread Wheat	41	IR41	Lorestan	Aleshtar	Bread Wheat
4	IR4	Khuzestan	Dezful	Bread Wheat	42	IR42	Khuzestan	Sardasht	Dry land Wheat
5	IR5	Khuzestan	Shushtar	Bread Wheat	43	IR43	Khuzestan	Sardasht	Dry land Wheat
6	IR6	Khuzestan	Zeydoon	Bread Wheat	44	IR44	Khuzestan	Shushtar	Bread Wheat
7	IR7	Khuzestan	Zeydoon	Bread Wheat	45	IR45	Khuzestan	Zeydoon	Bread Wheat
8	IR8	Khuzestan	Zeydoon	Bread Wheat	46	IR46	Lorestan	Noorabad	Bread Wheat
9	IR9	Ardebil	Moghan	Bread Wheat	47	IR47	Khuzestan	Zeydoon	Bread Wheat
10	IR10	Ardebil	Moghan	Bread Wheat	48	IR48	Khuzestan	Sardasht	Dry land Wheat
11	IR11	Khuzestan	Shushtar	Bread Wheat	49	IR49	Khuzestan	Sardasht	Bread Wheat
12	IR12	Khuzestan	Safiabad	Bread Wheat	50	IR50	Golestan	Gorgan	Bread Wheat
13	IR13	Khuzestan	Dezful	Triticale	51	IR51	Khuzestan	Zeydoon	Bread Wheat
14	IR14	Lorestan	Noorabad	Bread Wheat	52	IR52	Ilam	Ilam	Bread Wheat
15	IR15	Khuzestan	Safiabad	Durum Wheat	53	IR53	Ilam	Ilam	Bread Wheat
16	IR16	Golestan	Araghi Mahale	Bread Wheat	54	IR54	Golestan	Araghi Mahale	Bread Wheat
17	IR17	Ardebil	Moghan	Bread Wheat	55	IR55	Ilam	Mehran	Bread Wheat
18	IR18	Ardebil	Moghan	Bread Wheat	56	IR56	Ilam	Mehran	Bread Wheat
19	IR19	Lorestan	Noorabad	Bread Wheat	57	IR57	Ilam	Mehran	Bread Wheat
20	IR20	Khuzestan	Safiabad	Durum Wheat	58	IR58	Ilam	Mosian	Bread Wheat
21	IR21	Khuzestan	Safiabad	Triticale	59	IR59	Ilam	Mosian	Bread Wheat
22	IR22	Khuzestan	Andimeshk	Dry land Wheat	60	IR60	Hamadan	Jokar	Bread Wheat
23	IR23	Khuzestan	Shush	Bread Wheat	61	IR61	Hamadan	Jokar	Bread Wheat
24	IR24	Khuzestan	Safiabad	Durum Wheat	62	IR62	Hamadan	Jokar	Bread Wheat
25	IR25	Ardebil	Moghan	Bread Wheat	63	IR63	Hamadan	Jokar	Bread Wheat
26	IR26	Khuzestan	Sardasht	Dry land Wheat	64	IR64	Hamadan	Jokar	Bread Wheat
27	IR27	Ardebil	Moghan	Bread Wheat	65	IR65	Hamadan	Jokar	Bread Wheat
28	IR28	Markazi	Arak	Bread Wheat	66	IR66	Hamadan	Jokar	Bread Wheat
29	IR29	Ardebil	Moghan	Bread Wheat	67	IR67	Hamadan	Jokar	Bread Wheat
30	IR30	Khuzestan	Safiabad	Durum Wheat	68	IR68	Hamadan	Jokar	Bread Wheat
31	IR31	Khuzestan	Shushtar	Bread Wheat	69	IR69	Hamadan	Jokar	Bread Wheat
32	IR32	Golestan	Araghi Mahale	Bread Wheat	70	IR70	Hamadan	Jokar	Bread Wheat
33	IR33	Lorestan	Noorabad	Bread Wheat	71	IR71	Khuzestan	Shushtar	Bread Wheat
34	IR34	Khuzestan	Safiabad	Bread Wheat	72	IR72	Ardebil	Moghan	Bread Wheat
35	IR35	Lorestan	Noorabad	Bread Wheat	73	IR73	Golestan	Gorgan	Bread Wheat
36	IR36	Lorestan	Noorabad	Bread Wheat	74	IR74	Ardebil	Moghan	Bread Wheat
37	IR37	Khuzestan	Sardasht	Dry land wheat	75	IR75	Khuzestan	Shushtar	Bread Wheat
38	IR 38	Khuzestan	Sardasht	Dry land wheat					

Table 1. Codes, names, origins, and host of collection of Zymoseptoria tritici isolates used in this study.

 Table 2. ISSR markers used in this study.

No.	ISSR marker	Marker sequence $5' \rightarrow 3'$	Annealing	Number of	PIC
			temperature	polymorphic	
			(°C)	bands	
1	(GACA)4	GACAGACAGACAGACA	48	6	0.36
2	(AAC)7	AACAACAACAACAACAACAACAAC	52	7	0.41
3	(ATC)7	ATCATCATCATCATCATCATC	52	8	0.43
4	(AC)9	ACACACACACACACACAC	53	8	0.48
5	(AAG)7	AAGAAGAAGAAGAAGAAGAAGAAG	52	11	0.41
6	(AG)9	AGAGAGAGAGAGAGAGAGAG	53	9	0.46
7	(AGC)5	AGCAGCAGCAGCAGC	52	8	0.47
8	(CAG)5	CAGCAGCAGCAGCAG	52	8	0.48
9	(GTG)5	GTGGTGGTGGTGGTG	52	12	0.48
10	(GACAC)3	GACACGACACGACAC	48	11	0.47



No	SSR marker	Primer sequence $5' \rightarrow 3'$ (forward)	Primer sequence $3' \rightarrow 5'$ (reverse)	Annealing temperature (°C)
1	ST1E7	GATCTCGAGCAGGGCGGAAGT	TCACACGCTGGTCTGTGAATC	54
2	ST2E4	GAAGATCAACAGCAT GGGCGG	CTCCAGAGGGATCACAAAGGC	56
3	ST1D7	ATCCTCCATTCACTACTGCAT	TGTGGAACAGGAATAGGCTTG	56
4	ST1G7	ATGCTGAGAAGTTCGGTGAGG	CGTTCTTCCACCTCCTCCAACACT	52

Table 3. SSR markers used in this study.

1 minute at 72°C.The final elongation cycle was extended to 10 minutes.

After the amplification, PCR products were electrophoresed on 1.5% agarose gel at 90 volts for 1 hour (although it is probably insensitive to  $\pm 10\%$  variation in fragment size). After electrophoresis, gels were photographed on a UV transilluminator.

#### **Analysis of Molecular Data**

The gel images were evaluated for presence (score 1) or absence (score 0) of specific amplicons. The PopGene version 1.32 (Yeh et al., 1999) and GenAlex version 6.501 (Peakall and Smouse, 2006) software were applied to calculate the number and percentage of polymorphic markers, Polymorphism Information Content (PIC) of each primer, Nei's genetic diversity (h), Shannon's Index (I), Nei's genetic Distance (D), total genetic diversity (Ht), interpopulation diversity (Gst), and intrapopulation diversity (Hst), (Yeh et al., 1999). The NTSYSpc version 2.02e software was applied for analysis of the relationship genetic among isolates according to Dice's similarity coefficient and the Unweighted Pair Group Method with Arithmetic mean (UPGMA) (Rohlf, 2010).

#### **RESULTS AND DISCUSSION**

All of the 10 ISSR primers produced polymorphic bands that could be scored, and the lengths of the amplified DNA fragments varied from 100 to 3,000 bp, but only bands 100 to 1,500 bp long were used in analyses and weak bands were removed. Only two of the SSR primers (ST1D7 and ST1G7) produced distinct band pattern. In total, 91 polymorphic bands were considered in further analyses. The *PIC* index, varied from 0.36 to 0.48 (Table 3).

#### **Results of Cluster Analysis**

Based on molecular data, the diversity of isolates was distributed in a completely homogeneous manner within and between the sampled populations (Figure 1). Based on this dendrogram, the 75 studied samples were put within their distinctly related populations.

Results of the cluster analysis indicated that two clusters had the maximum number of isolates. One of them included most of the isolates from Lorestan, Hamadan, Markazi, and Ilam Provinces, and the other most of the isolates from Khuzestan and Ardebil Provinces.

#### **Genetic Diversity of the Populations**

Statistical analysis of genetic diversity within and between populations of *Zymoseptoria tritici* in Iran indicated that the maximum number of polymorphic loci belonged to the isolates from Khuzestan Province, with eighty-nine loci (97.8%). This was followed by the isolates from Ardebil Province with eighty-seven loci (95.6%) and the minimum to the isolates



**Figure 1.** Dendrogram for 75 isolates of *Zymoseptoria tritici*, based on DNA polymorphism revealed by ISSR and SSR molecular markers using Unweighted Pair Group Method of Arithmetic average (UPGMA) from pairwise comparisons employing the Dice genetic similarity coefficient.

JAST

from Ilam Province with thirty-nine loci (42.86%) (Table 4).

The largest genetic diversity based on Nei's genetic diversity (h) was that of Ardebil Province (0.3382) and the smallest that of Ilam Province (0.166).

Results of this research showed that the fungus causing STB in Iran had relatively high genetic diversity (Table 5). The average total genetic diversity was 0.34, of which 0.23 (equivalent to 69%) was that of intrapopulation diversity and the rest (31%) was that of inter-population diversity. Therefore, most of the genetic diversity was contributed by intra-population diversity.

#### **Analysis of Molecular Variance**

Analysis of Molecular Variance (AMOVA) showed significant variance within and between populations and was highly related (91%) to variation within populations (Table 6).

# Grouping of Populations Based on Genetic Diversity

Genetic grouping of populations based on Nei's genetic diversity showed that the maximum similarity existed between the populations collected in Khuzestan and Ardebil Provinces (Table 7 and Figure 2).

The word population cannot be an accurate definition for pathogens because, allelically, pathogens in various regions are basically combinations of various isolates that can behave as a population (Boeger *et al.*, 1993). Nevertheless, study of genetic diversity in pathogenic agents attracts the interest of plant pathologists in many ways. Therefore, researchers search for causes of and influential factors in pathogenicity of

Table 4. Genetic diversity of the studied population Zymoseptoria tritici using Nei's genetic diversity index and Shannon index.

Population	Nei's genetic	Shannon	Observed	Effective	Number of	Polymorphic
	diversity	index	alleles	alleles	polymorphic	percentage
	index		means	means	loci	
Khuzestan	0.3376	0.5081	1.9780	1.5678	89	97.8%
Lorestan	0.2741	0.4082	1.7473	1.4686	68	74.73%
Ilam	0.1660	0.2438	1.4286	1.2913	39	42.86%
Markazi	0	0	1	1	0	0
Ardebil	0.3382	0.5060	1.9560	1.5791	87	95.60%
Golestan	0.2777	0.4067	1.7033	1.4913	64	70.33%
Hamedan	0.2383	0.3595	1.7253	1.4061	44	72.53%
Means	0.3517	0.5258	2	1.6030		

Table 5. The genetic diversity indices of all populations *Zymoseptoria tritici*.

	Total genetic diversity(Ht)	Inter populations genetic diversity (Hs)	Gst	Nm
Total Populations indices	0.3377	0.2331	0.3087	1.1197

Table 6. Analysis of Molecular Variance (AMOVA) for 75 Zymoseptoria tritici isolates.

Source	Df	SS	MS	Est Var	%	Pr > F
Among Pops	5	212.911	42.582	2.047	9%	<0.0001**
Within Pops	68	1366.562	20.096	20.096	91%	< 0.0001**
Total	73	1579.473		22.144	100%	

\*\* Significant at P = 0.01.

khuzestan	lorestan	Ilam	Ardebil	Golestan	Hamadan	
****	0.9378	0.8309	0.7404	0.9663	0.9063	khuzestan
0.0643	****	0.8158	0.6997	0.9204	0.8915	lorestan
0.1853	0.2036	****	0.643	0.8072	0.8506	Ilam
0.3006	0.3571	0.4417	****	0.7483	0.7203	Ardebil
0.0343	0.0829	0.2142	0.2900	****	0.9092	Golestan
0.0983	0.1149	0.1619	0.3280	0.0952	****	Hamadan

**Table 7.** Pairwise population matrix of mean pop binary genetic distance.<sup>a</sup>

<sup>a</sup> Nei's genetic identity (above diagonal) and genetic distance (below diagonal).



Figure 2. Dendrogram (UPGMA) of Zymoseptoria tritici isolates based on Nei's (1978) Genetic distance.

pathogens (Pryor, 2000). Various researchers have previously studied diversity in pathogenicity of *Z. tritici* isolates (Gurung *et al.*, 2011).

Relatively similar results have been obtained in research carried out on this subject. For example, in the United States, genetic diversity reported from various states varied from 0.31 to 0.56 (Gurung et al., 2013). Furthermore, Razavi and Hughes (2004) reported that genetic diversity in 90 isolates collected in Canada was 0.44, of which 88% was related to intra-population and 12% to inter-population diversity. In another study conducted by Linde et al. (2002) on samples from various countries including Switzerland and the United States, it was found that genetic diversity varied from 0.44 to 0.50, of which 77% was related to intra-population and 23% to interpopulation diversity.

Although the diversity obtained in the present research was lower compared to some countries, yet it was relatively high. In

the mentioned studies, it was mentioned that gene flow was the reason for the higher share of intra-population diversity compared to inter-population diversity. In addition, in the present research, the higher intrapopulation diversity compared to interpopulation diversity, as well as the high Nm index (1.1197), suggested the occurrence of a strong gene flow between the studied provinces. This result is in contrast to a previous study conducted in Iran. Abrinbana et al. (2010) found an Nm of 0.78 and noted the considerable contrast with other regional (Schneider et al., 2001) and global (Linde et al., 2002) scales. Of course, populations of these two studies are not the same and are different in terms of the host type and the locations of the isolate collection.

In general, there are three ways in which gene flow happens in *Z. tritici*: formation of a sexual reproduction cycle, presence of intermediate hosts such as some grasses that can play the role of a green bridge, and movement of infected seeds and straw from one place to another by people.

Formation of the sexual reproduction cycle and ascospore production have not been reported in Iran, but existence of a strong gene flow in this fungus and of its sexual and asexual reproduction that were previously reported in Iran (Abrinbana *et al.*, 2010) raise the probability of the formation of the sexual reproduction cycle. This requires further research.

In general, we cannot ignore the role played by humans in the transmission of pathogenic agents to remote places through transportation of infected straw and seeds. Although it has not been proven that infected seeds could give rise to infected seedlings (Schnieder *et al.*, 2001), we assumed that infected grains might have contributed to such a gene flow (Consolo *et al.* 2009).

In grouping the various populations through genetic diversity, it was found that the greatest similarity existed between populations collected from Khuzestan and Ardebil Provinces (Figure 3). Similar results were obtained by Abrinbana *et al.* (2010) on genetic diversity of this fungus through using mating types, so that the fungal population in Khuzestan Province had the greatest similarity to that of Fars Province, followed by that of Ardebil Province.

Study of percentages of polymorphism indicated that the maximum polymorphism was also observed in these provinces. In general, in this research, the higher the degree of similarity, the greater was the degree of relatedness between the this populations. One reason for phenomenon can be the fact that fungal populations that have higher degrees of polymorphism naturally possess a greater number of genes to exchange. Moreover, geographical conditions, especially relative humidity in the air, and agricultural practices such as irrigation of most areas under cultivation in places including Khuzestan Province and the Dasht-e Moghan area allow the annual incidence of the disease. This, by itself, can be effective in

expanding, displacing, and creating diversity in the causal agent of the disease and can influence diversity in pathogenicity.

The importance that the effect of diversity in pathogenicity has on genetic diversity must not be ignored either, because many of these loci are associated with pathogenicity genes, and any factor that is involved in selection of pathogenicity genes, including the type of the involved host, also influences genetic diversity. In general, STB is a sporadic disease in most Iranian provinces, and is strongly dependent on environmental conditions, especially relative humidity and farming practices. Therefore, in many provinces, this disease appears once every few years. This, in itself, strongly influences the increases in the volume and diversity of the population and allows permanent exchange with other populations. Moreover, gene drift must not be ignored when interpreting small populations that are different from other populations.

### REFERENCES

- Abrinbana, M., Mozafiri, J., Shamsbakhsh, M. and Mehrabi, R. 2010. Genetic Structure of *Mycosphaerella graminicola* Populations in Iran. *Plant Path.*, **59:** 829-838.
- Boeger, J. M., Chen, R. S. and McDonald, B. A. 1993.Gene Flow between Geographic Populations of *Mycosphaerella graminicola* (Anamorph *Septoria tritici*) Detected with Restriction Fragment Length Polymorphism Markers. *Phytopathol.*, 83: 1148-1154.
- 3. Castillo, N., Cordo, C. and Simón, M. R. 2010. Molecular Variability among Isolates of *Mycosphaerella graminicola*, the Causal Agent of *Septoria tritici* Blotch, in Argentina. *Phytoparasitica*, **38**: 379–389.
- Consolo, V. F., Albani, C. M., Berón, C. M., Salerno, G. L. and Cordo, C. A. 2009. A Conventional PCR Technique to Detect *Septoria tritici* in Wheat Seeds. *Aus. Plant Pathol.*, 38(3): 222-227.
- 5. Czembor, P. C. and Arseniuk, E. 1999. Study of Genetic Variability among

Monopycnidial and Monopycnidiospore Isolates Derived from Single Pycnidia of *Stagonospora* ssp. and *Septoria tritici* with the Use of RAPD-PCR, MP-PCR and Rep-PCR Techniques. *Int. J. Phytopathol.*, 147:539–546.

- Doyle, J.J. and Doyle, J. L. 1990. Isolation of Plant DNA from Fresh Tissue. *Focus*, 12: 13–15.
- Gurung, S., Goodwin, S. B., Kabbage, M., Bockus, W. W. and Adhikari, T. B. 2011. Genetic Differentiation at Microsatellite Loci among Populations of *Mycosphaerella graminicola* from California, Indiana, Kansas, and North Dakota. *Phytopath.*, **101:** 1251-1259.
- Kabbage, M., Leslie, J. F., Zeller, K. A., Hulbert, S. H. and Bockus, W. W. 2008. Genetic Diversity of *Mycosphaerella* graminicola, the Causal Agent of Septoria tritici Blotch, in Kansas Winter Wheat. J. Agric. Food Environ. Sci., 2: 1-9.
- Kema, G. H. J., Goodwin, S. B., Hamza, S., Verstappen, E. C. P., Cavaletto, J. R. and van der Lee, T. A. J. 2002. A Combined AFLP and RAPD Genetic Linkage Map of *Mycosphaerella* graminicola, the Septoria tritici Leaf Blotch Pathogen of Wheat. Genet., 161: 1497–1505.
- Linde, C. C., Zhan, J. and McDonald, B. A .2002. Population Structure of *Mycosphaerella graminicola*: From Lesions to Continents. *Phytopath*, **92(9)**: 946-955.
- McDonald, B. A., Pettway, R. E., Chen, R. S., Boeger, J. M. and Martinez, J. P. 1995. The Population Genetics of *Septoria tritici* (Teleomorph *Mycosphaerella graminicola*). *Can. J. Bot.*, **73(S1)**: 292-301.
- Nei, M. 1978. Estimation of Average Heterozygosity and Genetic Distance from a Small Number of Individuals. *Genet.*, 89: 583–590
- Owen, P. G., Pei, M., Karp, A., Royle, D. J. and Edwards, K. J. 1998. Isolation and Characterization of Microsatellite Loci in

the Wheat Ppathogen *Mycosphaerella* graminicola. Mol. Eco., **7(11)**: 1611-1612.

- Peakall, R. O. D. and Smouse, P. E. 2006. GENALEX 6: Genetic Analysis in Excel. Population Genetic Software for Teaching and Research. *Mol. Eco. Res.*, 6(1): 288-295.
- 15. Pryor, B. M. and Gilbertson, R. L. 2000. Molecular Phylogenetic Relationships amongst *Alternaria* Species and Related Fungi Based upon Analysis of Nuclear ITS and mt SSU rDNA Sequences. *Mycol.*, **104(11)**: 1312-1321.
- Quaedvlieg, W., Kema, G. H. J., Groenewald, J. Z., Verkley, G. J. M., Seifbarghi, S., Razavi, M., Gohari, A. M., Mehrabi, R. and Crous, P. W. 2011. *Zymoseptoria* gen. Nov.: A New Genus to Accommodate *Septoria*-Like Species Occurring on Graminicolous Hosts. *Persoonia: Mol. Phylogenet. Evol.*, 26: 57-69.
- Razavi, M. and Hughes, G. R. 2004. Molecular Variability of *Mycosphaerella* graminicola as Detected by RAPD Markers. Int. J. Phytopathol., 152: 543-548.
- Rohlf, F. J. 2010. NTSYSpc: Numerical Taxonomy and Multivariate Analysis System Version 2.21j. Exeter Software, Setauket, New York.
- 19. Schnieder, F., Koch, G., Jung, C. and Verreet, J. A. 2001. Genotypic Diversity of the Wheat Leaf Blotch Pathogen *Mycosphaerella graminicola* (Anamorph) *Septoria tritici* in Germany. *Eur. Int. J. Phytopathol*, **107**: 285–290.
- 20. Talbot, N. J. 2015. Taming a Wild Beast: Developing Molecular Tools and New Methods to Understand the Biology of *Zymoseptoria tritici. Fungal Genet. Biol.*, **79**: 193–195.
- 21. Yeh, F. C., Yang, R. C. and Boyle, T. 1999. Popgene Version 1.31, Microsoft Window-Based Freeware for Population Genetic Analysis. Centre for International Forestry Research, University of Alberta.

# بررسی تنوع ژنتیکی قارچ Zymoseptoria tritici عامل بیماری سپتوریوز بر گی گندم با استفاده از نشانگرهای ISSR وSSR

م. دالوند، د. ظفری، م. ج. سلیمانی پری، ر. روح پرور، و س. م. طبیب غفاری

# چکیدہ

برای بررسی ساختار و ژنتیک جمعیت قارچ Zymoseptoria tritici گندم خیز ایران جمع آوری شده بودند توسط برگی گندم ۷۵ ایزوله این قارچ که از 7استان گندم خیز ایران جمع آوری شده بودند توسط نشانگرهای مولکولی ISSR و SSRمورد مطالعه قرار گرفتند .نتایج این بررسی نشان داد جمعیت قارچ عامل بیماری از تنوع ژنتیکی نسبتاً بالایی برخوردار است بطوریکه میانگین تنوع ژنتیکی کل (Ht) در نمونه هی مورد بررسی ۳۳/۰ بود و وقتی تنوع ژنتیکی به تنوع ژنتیکی داخل و بین جمعیتهای قارچ مورد آزمایش تقسیم شد مشخص شد ۶۹٪ تنوع در داخل جمعیتها و ۳۱٪ تنوع در بین جمعیتها وجود دارد .همچنین نتایج این تحقیق نشان از وقوع جریان ژنی قوی در بین استانهای مورد بررسی داشت.در بین استانهای مورد بررسی بیشترین تنوع ژنتیکی دراستانهای اردبیل و خوزستان مشاهده گردید که تجزیه خوشه ای نیز نشان داد جمعیتهای این دو استان بیشترین شباهت را با هم دارند.