In vitro Propagation of Caprifig and Female Fig Varieties (*Ficus carica* L.) from Shoot-tips

Ch. Bayoudh^{1, 2*}, R. Labidi², A. Majdoub¹, and M. Mars^{1, 2}

ABSTRACT

Fig trees are threatened by the attack of Fig Mosaic Disease (FMD) on leaves and fruits caused by viruses of several genera. Shoot-tip culture is a convenient method for viral sanitation. For this purpose, a reliable protocol for rapid in vitro propagation was developed with shoot-tips of three major Tunisian local fig (Ficus carica L.) varieties Zidi (ZDI), Soltani (SNI), Bither Abiadh (BA) and one rare and recalcitrant caprifig Assafri (ASF). For each in vitro step, four Murashige and Skoog (MS) media with different combinations of plant regulators were used. The best initiation of shoot-tips with sizes 0.5, 1 and 1.5 mm was obtained on medium M_3 containing 0.2 mg L 1 Benzyle Amino Purine (BAP), 0.1 mg L⁻¹ 1-NaphthaleneAcetic Acid (NAA) and 0.1 mg L⁻¹ Gibberellic acid (GA₃). The variety (SNI) showed the highest shoot-tip initiation potentialities for the establishment step with 100% of explant development rate. The shoot multiplication and plantlet development were provided by medium M_6 with 0.5 mg L⁻¹ BAP and 0.1 mg L⁻¹ NAA. The highest average of leaf number increase (92 leaves per plant) and proliferation rate (16.91 branches per plant) were reached on M₆. The best rooting rate (83.34%) was favored by medium M_{11} with half-strength MS and 1 mg L⁻¹ Indole-3-Butyric Acid (IBA). Ex vitro rooting of fig plantlets was successfully performed on moist peat with success rate of 90%. The acclimatized fig vitroplants showed high establishment rates (92.1%) and rapid growth on substrates S1 composed by peat without symptoms of virus diseases or morphological abnormalities.

Keywords: Acclimatization, Culture media, Explant size, Micropropagation.

INTRODUCTION

Fig tree (Ficus carica L.) has gained more interests in recent years for its economic importance and medicinal virtues. In Tunisia, fig tree is with remarkable genetic diversity and all cultivars are well adapted to local conditions (Mars et al., 2008; 2009). However, several abiotic and biotic constraints limit the development of this fruit crop and contribute to the decrease in revenue and the gradual disappearance of plantations (Saddoud et al., 2011). The lack of selected pollinators according to the needs of female cultivars, and the difficulties of their multiplication in fields, are major constraints (Gaaliche *et al.*, 2013). Virus diseases, especially fig mosaic disease, are actually a threat and spread in all areas of production (Saddoud *et al.*, 2007); which inhibits the development of healthy orchards and limits the production (Ashihara *et al.*, 2004). Improvement of farming techniques (Fraguas *et al.*, 2004), the sanitation of local varieties (Mars *et al.*, 2008), and the largescale production of good quality and healthy fig plants and pollinators by meristem or shoot tip culture (Pontikis and Melas, 1986; Gella *et al.*, 1998; Günver and Ertan, 1998)

¹ Regional Research Center for Horticulture and Organic Agriculture (CRRHAB), B. P. 57, 4042 Chott Mariem, Sousse, Tunisia.

^{*} Corresponding author; e-mail: chokribayoudh@gmail.com

² Research Unit on Agro-biodiversity, High Agronomic Institute, IRESA-University of Sousse, 4042 Chott-Mariem, Tunisia.

seem to be basic requirements for successful commercial orchards (Pasqual and Ferreira, 2007). This requires the mastery of rapid micropropagation of local figs. Tissue culture is a basic method for the propagation of plant material with high multiplication rates. In Tunisia, *in vitro* protocols for rapid regeneration of virus-free fig trees were not developed. Therefore, the main objective of this study was the establishment and optimization of a reliable protocol for *in vitro* regeneration of virus-free fig plants from three important local female varieties and one selected caprifig (pollinator).

MATERIALS AND METHODS

Four to five cm length shoots were taken from fig mosaic symptomatic female adult trees of three local varieties Zidi (ZDI), Soltani (SNI), Bither Abiadh (BA) and the pollinator Assafri (ASF) (Bayoudh et al., 2012). All samples were washed under running tap water for 15 minutes. Then, the samples were kept for 25 minutes in 10% NaOCl solution with 2 drops of Tween-20, followed by three rinses of 10 minutes in sterile distilled water. Shoot-tips of 0.5, 1.0 and 1.5 mm were excised from the aseptic explants (Figure 1) and were immediately placed on 4 Murashige and Skoog (MS) (1962) basal salts supplemented with 30 g L⁻ sucrose, 7 g \hat{L}^{-1} Agar, 90 mg L^{-1} Phloroglucinol (PG) (Bio Basic Canada Inc., Ontario) and different growth regulators for the study of their initiation (Figure 2).



Figure 1. Shoot-tip of fig (M) (G: 32x).



Figure 2. Initiation of fig shoot-tip on medium $M_3(a, b)$.

The initiation media were:

 M_1 : MS+0.2 mg L⁻¹ BAP+0.1 mg L⁻¹ NAA+0.1 mg L⁻¹ Kin+PG; M₂: MS+0.2 mg L⁻¹ BAP+0.1 mg L⁻¹ NAA+0.1 mg L⁻¹ IsoPentyl Adenosine (IPA)+PG; M₃: MS+0.2 mg L⁻¹ BAP+0.1 mg L⁻¹ NAA+0.1 mg L⁻¹ GA₃+PG, and M₄: MS+0.2 mg L⁻¹ BAP+0.1 mg L⁻¹ 2,4-dichlorophenoxy acetic acid (2,4-D)+PG.

For the proliferation and the elongation of the *in vitro* shoots, the following media were tested:

Rooting of plantlets was studied on:

 $M_{9}: MS+1 mg L^{-1} IBA; M_{10}: MS+1 mg L^{-1} NAA; M_{11}: MS^{1/2} (Half strenth of MS)+1 mg L^{-1} IBA, and M_{12}: MS^{1/2}+1 mg L^{-1} NAA.$

The pH of all media was adjusted to 6.0 prior to autoclaving for 20 minutes at 121° C and 1.1 kg cm⁻². Cultures were maintained in a growth chamber at $25\pm1^{\circ}$ C under 16 hours photoperiod with a light intensity of 40 µmol m⁻² s⁻¹ provided by cool white fluorescent lamps.

Recorded Parameters for Proliferated and Acclimatized Plants

The number of shoots per each explant was counted at the multiplication stage.

Well- developed fig plantlets of female varieties SNI and ZDI and the pollinator ASF were potted on moist peat and covered by transparent plastic film for ex vitro rooting. For the acclimatization step, wellrooted vitroplants were washed and placed on two substrates: (S_1) Composed by only peat, and (S_2) By a 2:1 mixture of peat and perlite. Survival rates (after four and ten weeks) were recorded for all studied Plant during varieties. growth acclimatization was studied for Smyrna type varieties ZDI and SNI. On each substrate, we determined the average increases in height, leaf number, and proliferation rate of acclimatized plants after 6 weeks. The survived plants were transferred to larger pots under greenhouse covered by insect proof sheet.

Data Analysis

All experiments were carried out in a factorial statistical scheme and a completely randomized design. Each treatment was replicated three times. Data were submitted to analysis of variance (ANOVA), and means were compared by Duncan test, using software SPSS (SPSS Inc., Ver.11.0).

RESULTS AND DISCUSSION

Effect of Explant Size and Media Composition

Shoot-tip size is a very important factor in the elimination of viruses from plants (Verbeek *et al.*, 1995; Panattoni *et al.*, 2013). In our attempts of fig sanitation via shoot-tip culture, we studied the effect of three shoot-tip sizes (0.5, 1, and 1.5 mm) (Bayoudh *et al.*, 2013) on their survival and development on medium M₃. For all studied varieties, shoot-tip size appeared to significantly affect their evolution rates (Table 2). For the pollinator ASF, the highest rate of recovery (96%) was obtained with shoot-tip size 0.5 mm. For the other varieties, this rate was obtained with

larger sizes: 1 mm for ZDI and BA (79 and 73.33%, respectively) and 1.5 mm for SNI (95.22%). It seems that the size of excised shoot-tip is not only important to produce virus-free plants of many crops, but also to determine the survivability of the explants in culture (Malaurie et al., 1995; Manganaris et al., 2003). In this context, Parmar et al. (2013) mentioned that in vitro shoot growth of Clerodendrum inerme was higher with larger size explants than the smaller one. This may be due to the high number of leaf primordia in large shoot-tips. Sahraroo et al. (2009) encountered many difficulties to regenerate Ficus carica plantlets from 0.2 to 0.4 mm meristems since they failed to grow on the media. For all varieties, abundant necroses were observed at the bases of large explant sizes and the highest necrosis rate (76.45%) was recorded at the base of 1.5 mm shoot-tips of the male variety ASF.

Establishment Media

Effect of establishment media was studied on small size shoot-tips (0.5 mm) of varieties ZDI, BA, SNI and pollinator ASF. These explants were cultured and initiated on the establishment media M_1 , M_2 , M_3 and M_4 . During the first 2 weeks of establishment, explants browning were noticed. Its negative effects on the explants were reduced by PG, which acts as antioxidant and increases growth rate of the cultures (Hepaksoy and Aksoy, 2005). Similar behavior of small fig shoot tip explants on establishment media was observed by Gella et al. (1998) and Dhage et al. (2012), who reported that explant browning was avoided by the use of the antioxidants. After eight weeks of culture, growth and emergence of shoot apices were highly dependent to the establishment composition of medium. Significant interaction was noted between the 4 varieties and the establishment media concerning shoot-tip necrosis rate. The variety Soltani (SNI) showed better growth and less necrosis and calli at the bases of explants than the other three varieties (Table 1). For SNI, the highest rate of shoot-tip survival and

Variety	Size (mm)	% Necrosis	% Callus at explant bases	% Development
ZDI ^{<i>a</i>}	0.5	46.67 ± 21.66	23.89 ± 3.89	61.11 ± 11.11
	1	36.82 ± 4.55	33.39 ± 13.34	79.00 ± 11.59
	1.5	64.31 ± 12.87	55.10 ± 17.5	70.49 ± 3.32
BA b	0.5	58.50 ± 11.64	31.10 ± 1.1	67.77 ± 19.28
	1	43.33 ± 16.66	33.33 ± 8.81	73.33 ± 8.81
	1.5	70.00 ± 5.77	60.00 ± 5.77	56.67 ± 12.01
SNI ^c	0.5	$5.71 \pm 3,68$	$4.28 \pm 2,97$	$90.00 \pm 6,54$
	1	31.42 ± 5.53	7.14 ± 5.65	55.71 ± 2.97
	1.5	6.18 ± 3.03	4.28 ± 2.97	95.22 ± 3.07
ASF^{d}	0.5	72.67 ± 8.19	37.67 ± 16.89	96.00 ± 4.0
	1	69.79 ± 4.01	48.36 ± 20.54	92.63 ± 3.68
	1.5	76.45 ± 10.50	46.65 ± 10.18	87.96 ± 7.23
Statistical	significance:			
Size e	effect	NS	NS	*
Intera	ction Variety×Size	**	**	**

Table 1. Effect of shoot-tip size of different varieties of fig on necrosis, callus, and development rates.

^{*a*} Zidi; ^{*b*} Bither Abiadh; ^{*c*} Soltani, ^{*d*} Assafri. *: Significant differences (Duncan, P \leq 0.05), **: Highly significant differences (Duncan, P \leq 0.01), NS: Not significant differences. The values are compared vertically.

Fable 2. Effect of four establishment media on shoo	t-tip explants dev	velopment of different	varieties of fig.
--	--------------------	------------------------	-------------------

Variety ^{<i>a</i>}	Medium	% Necrosis	% Callus at explant bases	% Development
	M_1	53.33 ± 3.33	39.17 ± 0.83	59.17 ± 12.27
ZDI^{a}	M_2	53.33 ± 12.01	77.59 ± 7.21	16.67 ± 3.33
	M ₃	46.67 ± 21.66	23.89 ± 3.89	61.11 ± 11.11
	M_4	75.17 ± 13.10	62.20 ± 23.19	31.83 ± 16.53
	M ₁	78.33 ± 11.66	38.33 ± 7.26	41.67 ± 22.04
\mathbf{BA}^{b}	M_2	66.67 ± 17.63	58.33 ± 13.64	73.33 ± 14.52
	M ₃	58.50 ± 11.64	31.10 ± 1.1	67.77 ± 19.28
	M_4	53.33 ± 18.55	40.00 ± 0	66.67 ± 18.55
	M ₁	$5.00 \pm 2,67$	$8.75 \pm 5,80$	100 ± 0
SNI ^c	M_2	0 ± 0	0 ± 0	100 ± 0
	M ₃	$5.71 \pm 3,68$	$4.28 \pm 2,97$	$90 \pm 6,54$
	M_4	$2.22 \pm 2,22$	0 ± 0	$96.66 \pm 3,33$
	M_1	40.00 ± 11.54	40.00 ± 20.0	86.67 ± 13.33
ASF^{d}	M_2	65.77 ± 22.9	81.30 ± 13.53	66.30 ± 16.15
	M ₃	72.67 ± 8.19	37.67 ± 16.89	96.00 ± 4.0
	M_4	61.67 ± 3.34	20.37 ± 15.15	83.77 ± 11.11
Statistical s	ignificance:			
Media effec	et	**	NS	NS
Interaction	Varieties×Media	**	NS	*

^{*a*} Zidi; ^{*b*} Bither Abiadh; ^{*c*} Soltani, ^{*d*} Assafri. * Significant differences (Duncan, P \leq 0.05), ** Highly significant differences (Duncan, P \leq 0.01), NS: Not significant differences. The values are compared vertically.

evolution (100%) was achieved on the media M₁ and M₂. However, on M₂, no necrosis or callus formation at the bases of shoot-tips occurred. Zidi (ZDI) showed low evolution rates on all initiation media and the lowest rate (16.67%) was obtained with M₂. The majority of explants were susceptible to necrosis on all culture media. The highest rate of necrosis (78.33%) was recorded with (BA) on M₁. Assafri (ASF) explants gave the highest rate of callus at their bases (81.3%) on M₂. Among the four tested establishment media, M_2 and M_3 were the most adequate for shoot-tip development of ZDI, BA, SNI and ASF, but M₃ supplemented with GA₃ was the best one sine it produced lower rates of callus at the explant bases. Similar results were reported by Fraguas et al. (2004).

Proliferation Media

Plantlets of the four tested varieties ZDI, BA, SNI and ASF, obtained during the

initiation phase, were transferred to the multiplication media M_5 , M_6 , M_7 , and M_8 containing BAP and Kinetin. Statistical analysis of shoot development and growth on all proliferation media for 10 weeks showed that shoot proliferation was significantly dependent on varieties and culture media (Table 3).

results Similar were obtained by Hepaksoy and Aksoy (2006) and Mustafa and Taha (2012). The variety ZDI presented the highest proliferation potentialities: average number of branches per plant (16.91), average number of leaves per plant (92) and average increase in plant height (16.33 mm) (Table 3). The highest rates of shoot proliferation for varieties BA (7.08 branches per plant), ZDI (16.91 branches per plant) and pollinator ASF (6.83 branches per plant) were obtained with the medium M_6 containing BAP, IBA, and GA₃. On this medium, the shoots were well developed,

Table 3. Effect of four proliferation media on in vitro plant growth of four genotypes of fig.

Variety	Medium	Average increase of plant height (mm)	Number of leaves	Number of newly formed shoots
SNI ^a	M ₅	9.01 ± 1.37 a	48.41 ± 9.21 a	9.00 ± 1.17 a
	M_6	9.60 ± 1.75 ab	30.58 ± 4.84 a	7.17 ± 1.36 a
	M_7	5.88 ± 1.27 b	5.91 ± 3.15 b	1.08 ± 0.31 b
	M_8	6.93 ± 2.28 ab	10.33 ± 6.45 c	1.67 ± 1.02 c
\mathbf{BA}^{b}	M_5	9.79 ± 0.82 a	34.31 ± 5.18 a	6.83 ± 0.90 a
	M_6	7.83 ± 1.03 ab	26.77 ± 5.80 a	7.08 ± 1.16 a
	M_7	9.59 ± 1.00 b	8.70 ± 4.79 b	3.67 ± 1.02 b
	M_8	5.24 ± 1.71 ab	7.36 ± 2.60 c	1.42 ± 0.33 c
ZDI^{c}	M_5	17.0 ± 0.82 a	66.91 ± 7.25 a	13.41 ± 1.49 a
	M_6	11.92 ± 0.97 ab	92.00 ± 8.83 a	16.91 ± 1.98 a
	M_7	10.54 ± 2.05 b	61.17 ± 13.35 b	9.91 ± 1.82 b
	M_8	16.33 ± 2.4 ab	26.67 ± 4.30 c	1.91 ± 0.39 c
ASF^{d}	M_5	6.94 ± 0.99 a	28.58 ± 4.14 a	6.75 ± 0.85 a
	M_6	7.60 ± 1.38 ab	28.41 ± 8.31 a	6.83 ± 1.43 a
	M_7	4.74 ± 1.35 b	7.75 ± 2.13 b	1.33 ± 0.65 b
	M_8	3.74 ± 1.87 ab	6.41 ± 3.82 c	1.50 ± 0.85 c
Statistical signific	ance:			
Variety effect		**	**	**
Media effect		*	**	**
Interaction Varieties×Media		*	**	**

^{*a*} Soltani, ^{*b*} Bither Abiadh; ^{*c*} Zidi; ^{*d*} Assafri. *: Significant differences (Duncan, P \leq 0.05). **: Highly significant differences (Duncan, P \leq 0.01), The values are compared vertically. Means with different letter in a row are statistically different (Duncan, P \leq 0.05).

greenish and with small amount of basal calli (Figure 3). BAP is a very critical cytokinin for proliferation. It was reported to stimulate multiple shoot formation in vitroplants (Saidi et al., 2007; Singh et al., 2010; Karimpour et al., 2013; Zuraida et al., 2014). Also, the proliferation medium M₆ contained GA₃, which is considered to be essential not only to improve the growth of the explants, but to increase significantly the number of nodes and leaves (Rostami and Shahsava, 2012). For SNI, the best proliferation rate (9.0 branches per plant) was provided by the medium M_5 . The lowest rates were, generally, obtained by the medium M8 containing Kinetin as reported by Mustafa and Taha (2012). On medium M₈, the shoots were stunted, with callus and some vellowish leaves. This may be due to reaction effects of Kinetin and NAA on plant respiratory metabolism (Akemine et al., 1975). The best plantlets elongation rate and increase in number of leaves were mainly provided by media M₅ and M₆ and the lowest values were recorded on M7 and M_8 . It appeared that BAP (M_5 and M_6) had significant higher effects on the vitroplants. Similar results have been reported by Kumar et al. (1998) and Nobre and Romano (1998)



Figure 3. Proliferation and elongation of fig shoots on medium M₆.

for the regeneration of fig vitroplants from apical buds on media containing BAP.

Rooting Media

Single vigorous plantlets of each variety were transferred to different rooting media M_9 , M_{10} , M_{11} and M_{12} to induce and develop the root system (Figure 4). Results showed that plant rooting rate was significantly impacted by the variety and rooting media effects (Table 4). The highest percentages of rooted plants (83.34%) occurred for SNI on M_{10} and ASF on M_{11} . In addition, M_{11} contained half strength of MS and 1 mg L⁻ IBA gave best results for ZDI plants (75%). These results were in conformity with previous reports (Kumar et al., 1998; Yancheva et al., 2005; Soliman et al., 2010) where in vitro rooting of fig vitroplants was facilitated by 1 to 2 mg L⁻¹ IBA containing media. M₁₁ induced development of longest roots in ASF (52.4 mm) and variety SNI (39.2 mm) (Table 5). Furthermore, highest number of secondary roots of ZDI (11.5), ASF (9.7) and SNI (10.7) plantlets were recorded on M₁₁ medium. Therefore, the composition of this medium (with reduced nitrogen and auxins) was, globally, the most suitable for fig vitroplant rooting. Similar findings for other plant species were mentioned by many authors (Ebrahim et al.,



Figure 4. Multiple and long roots of fig vitroplant induced on medium M₁₁.

Variety	Medium	Rooting rate (%)	Root length (mm)	Principal root number/ plant	Secondary root number/plant
	M9	50.00 ± 15.07 b	26.8 ± 4.25 b	2.50 ± 0.67 b	2.50 ± 1.62 b
	M10	33.34 ± 14.21 b	35.5 ± 12.75 ab	4.00 ± 1.73 a	11.00 ± 5.80 a
ZDI^{a}	M11	75.00 ± 13.05 a	28.20 ± 5.61 a	2.55 ± 0.61 ab	11.50 ± 3.56 a
	M12	8.34 ± 8.34 b	18.60 ± 0.0 ab	1.00 ± 0 a	0 ± 0 b
	M9	8.34 ± 8.34 b	1.60 ± 0 b	1.00 ± 0 b	0 ± 0 b
	M10	16.67 ± 11.23 b	9.90 ± 0.29 ab	3.50 ± 1.45 a	7.00 ± 1.73 a
\mathbf{BA}^{b}	M11	58.34 ± 14.86 a	22.50 ± 5.49 a	2.30 ± 0.37 ab	5.70 ± 1.64 a
	M12	75.00 ± 13.05 b	22.50 ± 3.33 ab	3.80 ± 0.48 a	6.00 ± 1.39 b
	M9	41.66 ± 14.86 b	18.80 ± 2.80 b	1.60 ± 0.26 b	3.20 ± 1.22 b
	M10	50.00 ± 15.07 b	34.50 ± 5.18 ab	3.57 ± 0.57 a	8.57 ± 2.80 a
ASF^{c}	M11	83.34 ± 11.23 a	52.40 ± 6.32 a	3.40 ± 0.43 ab	9.70 ± 2.16 a
	M12	41.67 ± 14.86 b	37.10 ± 8.88 ab	3.37 ± 1.19 a	7.00 ± 2.73 b
	M9	58.34 ± 14.86 a	35.10 ± 5.91 b	2.50 ± 0.37 b	5.00 ± 1.0 b
	M10	83.34 ± 11.23 b	22.20 ± 3.68 ab	2.90 ± 0.68 a	10.40 ± 4.04 a
SNI ^c	M11	75.00 ± 13.05 a	39.23 ± 6.30 a	3.40 ± 0.78 ab	10.70 ± 1.64 a
	M12	50.00 ± 15.07 b	37.80 ± 4.03 ab	3.50 ± 1.05 a	2.00 ± 0.57 b
Statistical Significance:					
Media effec	t	**	*	*	*
Variety×media		NS	NS	NS	NS

Table 4. Fig plantlet root development.

^{*a*}Zidi; ^{*b*}Bither Abiadh; ^{*c*}Assafri, ^{*d*}Soltani. *: Significant differences (Duncan, P \leq 0.05), NS: Not significant differences. The values are compared vertically. Means with different letter in a row are statistically different (Duncan, P \leq 0.05).

Table 5. Development of ex vitro rooted fig plants on peat after six weeks.^a

Variety	Mean plant height (mm)	Leaf number/Plant
ZDI^{a}	126.10 ± 12.20	2.10 ± 0.69
SNI ^b	91.07 ± 12.79	1.91 ± 0.73
ASF ^c	104.59 ± 24.01	1.45 ± 0.54
Differences among varieties	NS	NS

^a Zidi, ^b Soltani, ^c Assafri. The values are compared vertically, NS: Not significant differences.

2007; Othmani et al., 2010 and Toppo et al., 2012). For (BA), the best rooting rate (75%) was obtained with M₁₂. Furthermore, on M_{12} , we obtained strong root system. The plants continued to grow well and to this proliferate medium and, on consequently, they were greenish and with an average of about 6 brunches per plant. Globally, a significant effect of rooting media for root length and number of principal and secondary roots of vitroplants was recorded. The highest average number of main roots (4) was obtained on M_{10} (Table 4).

Acclimatization Step

Ex vitro Rooting

After six weeks, we noticed that for all varieties, 90% of plants were successfully rooted. Throughout *ex vitro* rooting period, plantlets continued to grow properly. Strategy adopted in the present work was in agreement with *ex vitro* rooting of Blueberry (Guang-jie *et al.*, 2008). Bhatia *et al.* (2002) reported that best results for *ex vitro* Stackhousia plantlets reflects the potential

effects of combining *ex vitro* and hardening in one step, with view to reducing costs of vitroplants. The highest averages of increase in plant height (126.1 mm) and leaf number (2.1) were recorded for ZDI (Table 5). The rooted plants were transferred to larger pots and maintained under the insect proof greenhouse.

Effects of Substrates on Survival Rate

For all varieties, on the two substrates S_1 and S_2 , the survival rates recorded following the first four weeks ranged from 81.39 to 100%, while the survival rates recorded in the tenth week of transplantation extended from 63.15 to 92.1% (Table 6). This may be due essentially to the gradual loss of some acclimatized plants due to the relatively hard environment as compared to the in vitro conditions (Hajong et al., 2013). As reported by many authors (Demiralay et al., 1998; Kumar et al., 1998; Hepaksoy and Aksoy, 2005; Fraguas et al., 2012), the final survival rates of rooted acclimatized plants vary according to the varieties and the substrates. The highest final survival rate (92.1%) and the lowest final survival rate (63.15%)were recorded for ZDI, respectively, on S_1 and S_2 (Table 6). Generally, S_1 , which was composed of only peat, allowed the best survival rates. This substrate contained appropriate amount of nutrients (Fraguas et al., 2012); and had good water retention capacity, which played an important role in plant growth (Mengesha et al., 2013).

Effect of Substrates on Plant Growth

Plants of both varieties ZDI and SNI presented better growth on S_1 . The highest increase in plant height (15.48 mm) and branching number (0.7) were recorded with S_1 , whereas the highest leaf number (1.9) was obtained with S_2 (Table 7). These results of acclimatized plant growth are considered satisfactory if compared to those obtained by

lable 6. Survival rate	es (%) of acclimatized f.	ig plants on substrates S_1 and S_2 .				
	ZDIª	SNI	q	BA ^c		ASF^{d}
	Survival rate during	Final survival rate Survival rate during	Final survival	Survival rate during Fin	nal survival	Survival rate during Final survival

 85.96 ± 4.52 64.28 ± 3.26

*

NS

NS

SN

NS

89.47 ± 7.23

 86.27 ± 4.86

 96.07 ± 2.74 94.73 ± 5.71

 67.10 ± 5.42 66.17 ± 5.78

first 4 weeks

rate

 96.05 ± 2.24 81.39 ± 4.51

 92.10 ± 4.43 63.15 ± 8.03

 $\frac{\text{first 4 weeks}}{100 \pm 0}$ $\frac{100 \pm 0}{\text{NS}}$

first 4 weeks

rate

rate

first 4 weeks 96.49 ± 1.73 87.14 ± 3.15

	compared vertically.
	s. The values are
2	S: Not significant differences
211	differences (P≤ 0.01), N9
	Highly significant o
	Bither Abiadh; ^d Assafri. **:
substrates	^{<i>a</i>} Zidi; ^{<i>b</i>} Soltani, ^{<i>c</i>}

NS

*

Differences between

S₂: Peat+ Perlite

S₁: Peat

	ZD	οI ^α		SNI ^b		
	Plant height	Leaf	Branching	Plant height	Leaf	Branching
	(mm)	number	number	(mm)	number	number
S ₁ : Peat	$15.48 \pm 2,42$	1.54 ± 0.43	0.45 ± 0.15	11.36 ± 2.05	1.5 ± 0.8	0.7 ± 0.3
S ₂ : Peat+Perlite	13.89 ± 2.02	1.0 ± 0.38	0.18 ± 0.18	9.67 ± 2.16	1.9 ± 1.1	0.64 ± 0.27

Table 7. Growth of acclimatized plants on S_1 and S_2 .

^aZidi, ^bSoltani

Fraguas *et al.* (2012) who obtained 8.92 mm as the best plant height of plantlets acclimatized during 60 days on 5 different substrates.

Acclimatized plants transferred to the insect proof greenhouse showed good growth. They formed thick trunk, multiple branching and new leaves. No variations in growth or morphological characteristics were detected.

CONCLUSIONS

А suitable protocol for rapid multiplication of important local fig varieties of Smyrna type (Zidi and Soltani), San Pedro type (Bither Abiadh) and the rare and recalcitrant pollinator Assafri from shoottips was developed and optimized. Since this protocol is to be applied for sanitation of local fig varieties, the small shoot-tip size of 0.5 mm height was the most preferred size. The proliferation of fig shootlets was mainly provided by medium M₆ containing small amounts of BAP and NAA. With this optimized protocol, ex vitro rooting of vitroplants on humidified peat was successfully feasible and allowed high rooting rates, reached 90% with healthy roots, which resulted in reduction of plant micro-propagation period. The micropropagated plants were FMD-symptoms free and had normal morphological aspects. No abnormalities were noticed. Nevertheless, molecular analysis to test their genetic stability is needed. Following these tests, this optimized protocol for large-scale production of good quality and healthy female fig and pollinator plants will be applied for successful establishment of commercial production.

REFERENCES

- Akemine, T., Kikuta, Y. and Tagawa, T. 1. 1975. Effect Kinetin of and Naphthaleneacetic Acid Application on the Respiratory Metabolism during Callus Development in Potato Tuber Tissue Cultured In vitro. Journal of the Faculty of Agriculture, Hokkaido University, 58: 247-261.
- 2. Ashihara, W., Kondo, A., Shibao, M., Tanaka, H., Hiehata, K. and Izumi, K. 2004. Ecology and Control of Eriophyid Mites Injurious to Fruit Trees in Japan. *JARQ-Jpn. Agr. Res. Q.*, **38:** 31-41.
- Bayoudh, C., Barutcu, E., Labidi, R., Majdoub, A., Çaglayan, K. and Mars, M. 2012. Détection de Trois Virus Associés à la Mosaïque de Figuier (*Ficus carica* L.) Dans les Régions Côtières Centrales de la Tunisie. *Journées Internationales de l'A. T. Biotech.*, Décembre 2012, Mahdia, Tunisia, PP. 19-22.
- 4. Bayoudh, C., Elair, M., Labidi, R., Majdoub, A., Mahfoudhi, N. and Mars, M. 2013. In vitro Tissue Culture and Molecular Techniques as an Aid for Organic Healthy Fig (Ficus carica L.) Orchard Establishment. In International CTAB-ISOFAR-MOAN Symposium on Crop Protection Management in Mediterranean Organic Agriculture, 14-16 May, 2013, Sousse, Tunisia.
- 5. Bhatia, N. P., Bhatia, P. and Ashwath, N. 2002. *Ex vitro* Rooting of Micropropagated Shoots of *Stakhousia tryonii*. *Biologia Plantarum*, **45**: 441-444.
- 6. Demiralay, A., Yalçin-Mendi, Y., Aka-Kaçar, Y. and Çetiner, S. 1998. *In vitro* Propagation of *Ficus carica* L. var. Bursa

Siyahi through Meristem Culture. Acta Hort., **480:** 165-167.

- Dhage, S. S., Pawar, B. D., Chimote, V. P., Jadhav, A. S. and Kale, A. A. 2012. *In Vitro* Callus Induction and Plantlet Regeneration in Fig (*Ficus carica* L.). *J. Cell Tissue Res.*, 12: 3395-3400.
- Ebrahim, N., Shibli, R., Makhadmeh, I., Shatnawi, M. and Abu-Ein, A. 2007. *In vitro* Propagation and *in Vivo* Acclimatization of Three Coffee Cultivars (*Coffea arabica* L.) from Yemen. *World Appl. Sci. J.*, 2: 142-150.
- Fraguas, C. B., Pasqual, M. and Pereira, A. R. 2004. Multiplicaçao *In vitro* de *Ficus carica* L.: Efeito da Cinetina e do Acido Giberélico. *Ciênc. Agrotec.*, 28: 49-55.
- Fraguas, C. C., Moacir, P., Gomes de Araujo, A., Pereira, A. R. and Maoro de Castro, E. 2012. Acclimatization and Leaf Anatomy of Micropagated Fig Plantlets. *Rev. Bras. Frutic. Jaboticabal – SP*, 34: 1180-1188.
- Gaaliche, B., Majdoub, A., Trad, M. and Mars, M. 2013. Assessment of Pollen Viability, Germination, and Tube Growth in Eight Tunisian Caprifig (*Ficus carica* L.) Cultivars. *ISRN Agronomy*, VOL??? 1-4. <u>http://dx.doi.org/10.1155/2013/207434</u>
- Gella, R., Marin, J. A., Corrales, M. L. and Toribio, F. 1998 Elimination of Fig Mosaic from Shoot Tip Cultures by Thermotherapy. *Acta Hort.*, 480: 173-177.
- Guang-jie Z., Zhan-bin, W. and Dan, W. 2008. In vitro Propagation and Ex vitro Rooting of Blueberry Plantlets. Plant Tissue Cult. Biotech., 18: 187-195. doi. 10.3329/ptcb.v18i2.3650
- 14. Günver, G. and Ertan, E. 1998. A Study on the Propagation of Figs by the Tissue Culture Techniques. *Acta Hort.*, 480: 169-172.
- Hajong, S., Kumaria, S. and Tandon, P. 2013. Effect of Plant Growth Regulators on Regeneration Potential of Axenic Nodal Segments of *Dendrobium chrysanthum* Wall. ex Lindl. J. Agr. Sci. Tech., 15: 1425-1435.
- Hepaksoy, S. and Aksoy, U. 2005. In vitro Propagation of Ficus carica cv. Sariop Clone Selected for Its High Performance. Acta Hort., **798**: 199-204.
- 17. Hepaksoy, S. and Aksoy, U. 2006. Propagation of *Ficus carica* L. Clones by *In*

vitro Culture. Biologia Plantarum, 50: 433-436.

- Karimpour, S., Davarynejad, G. H., Bagheri, A. and Tehranifar, A. 2013. *In vitro* Establishment and Clonal Propagation of Sebri Pear Cultivar. *J. Agr. Sci. Tech.*, 15: 1209-1217.
- Kumar, V., Radha, A. and Kumar Chita, S. 1998. *In vitro* Plant Regeneration of Fig (*Ficus carica* L. cv. Gular) Using Apical Buds from Mature Trees. *Plant Cell Rep.*, 17: 717-720.
- Malaurie, B., Thouvenel, J. C. and Pungu, O. 1995. Influence of Meristem-tip Size and Location on Morphological Development in *Diowcorea cayenensis*-D. *Rotundata comlex* 'Grosse Caille' and One Genotype of D. *praehensilis. Plant Cell Tiss. Organ. Cult.*, 42: 215-218.
- Manganaris, G. A., Economou, A. S., Boubourakas, I. N. and Katis, N. I. 2003. Elimination of PPV and PNRSV through Thermotherapy and Meristem Tip Culture in Nectarine. *Plant Cell Rep.*, 22: 195-200.
- Mars, M., Chatti, K., Saddoud, O., Salhi-Hannachi, A., Trifi, M. and Marrakchi, M. 2008. Fig Cultivation and Genetic Resources in Tunisia: An Overview. *Acta Hort.*, **798**: 27-32.
- 23. Mars, M., Gaaliche, B., Ouerfelli, I. and Chouat, S. 2009. Systèmes de Production et Ressources Génétiques du Figuier (*Ficus carica* L.) à Djebba et Kesra, Deux Villages de Montagne au Nord Ouest de la Tunisie. *Revue des Régions Arides*, 22: 33-45.
- Mengesha, A., Ayenew, B. and Tadesse, T. 2013. Acclimatization of *In vitro* Propagated Pineapple (*Ananas comosuss* L., var. Smooth Cayenne) Plantlets to *Ex vitro* Condition in Ethiopia. *Am. J. Plant Sci.* (AJPS), 4: 317-323. doi: 10.4236/ajps.2013.42042.
- 25. Murashige, T. and Skoog, F. 1962. A Revised Medium for Rapid Growth and Bioassays with Tobacco Tissue Cultures. *Physiol. Plant.*, **15:** 473-497.
- 26. Mustafa, N. S. and Taha, R. A. 2012. Influence of Plant Growth Regulators and Subculturing on *In vitro* Multiplication of Some Fig (*Ficus Carica* L.) Cultivars. *J. Appl. Sci. Res.*, 8: 4038-4044.
- 27. Nobre, J. and Romano, A. 1998. *In vitro* Cloning of *Ficus carica* L. Adult Trees. *Acta Hort.*, 480: 161-164.

- Othmani, A., Rhouma, S., Bayoudh, C., Mzid, R., Drira, N. and Trifi, M. 2010. Regeneration and Analysis of Genetic Stability of Plantlets as Revealed by RAPD and AFLP Markers in Date Palm (*Phoenix dactylifera* L.) cv. Deglet Nour. *Intl. Res. J. Plant Sci. (IRJPS)*, 1: 048-055.
- 29. Panattoni, A., Luvisi, A. and Triolo, E. 2013. Elimination of Viruses in Plants: Twenty Years of Progress. *Span. J. Agric. Res.*, **11:** 173-188.
- Parmar, V. R., Patel, H. A. and Jasrai, Y. T. 2013. Developing Normal Plants of *Clerodendron* from Viral Infected Stock through Meristem Culture. *Cibtech. J. Bio-Protocols*, 2: 1-5. http://www.cibtech.org/cjbp.htm
- Pasqual, M. and Ferreira, E. A. 2007. Micropropagation of Fig Tree (*Ficus carica* sp). In: "Protocols for Micropropagation of Woody Trees and Fruits", (Eds.): Mohan Jain, S. and Häggman, H.. Springer, PP. 409-416.
- 32. Pontikis, C. A. and Melas, P. 1986. Micropropagation of *Ficus carica* L. *HortSci.*, **21:** 153.
- 33. Rostami, A. A. and Shahsava, A. R. 2012. In vitro Micropropagation of Olive (Olea europaea L.) 'Mission' by Nodal Segments. J. Biol. Environ. Sci., 6: 155-159. http://jbes.uludag.edu.tr
- 34. Saddoud, O., Chatti, K., Salhi-Hannachi, A., Mars, M., Rhouma, A., Marrakchi, M. and Trifi, M. 2007. Genetic Diversity of Tunisian Fig (*Ficus caraica* L.) as Revealed by Nuclear Microsatellites. *Hereditas*, 144: 149-157.
- 35. Saddoud, O., Baraket, G., Chatti, K., Trifi, M., Marrakchi, M., Mars, M. and Salhi-Hannachi, A. 2011. Using Morphological Characters and Simple Sequence Repeat (SSR) Markers to Characterize Tunisian Fig (*Ficus caraica* L.) Cultivars. *Acta Biol. Cracov. Bot.*, **53**: 7-14.

- 36. Sahraroo, A., Babalar, M., Ebadi, A., Habibi, M. K. and Khadivi-Khub, A. 2009. Influence of Apical Meristem Culture and Thermotherapy on Production of Healthy Fig Plants. *Hortic. Environ. Biotechnol.*, **50**: 45-50.
- Saidi, R., Lamarti, A. and Badoc, A. 2007. Micropropagation du Caroubier (*Ceratonia siliqua*) par Culture de Bourgeons Axillaires Issus de Jeunes Plantules. *Bull. Soc. Pharm. Bordeaux*, 146: 113-129.
- Singh, J. and Tiwari, K. N. 2010. Highfrequency *In vitro* Multiplication System for Commercial Propagation of Pharmaceutically Important *Clitoria ternatea* L.: A Valuable Medicinal Plant. *Ind. Crop. Prod.*, **32:** 534–538.
- Soliman, H. I., Gabr, M. and Abdallah, N. 2010. Efficient Transformation and Regeneration of Fig (*Ficus carica* L.) *via* Somatic Embryogenesis. *GM Crop.*, 1: 47-58. doi: 10.4161/gmcr.1.1.10632.
- 40. Toppo, D. D., Singh, G., Purshottam, D. K. and Misra, P. 2012. Improved *In vitro* Rooting and Acclimatization of *Jatropha curcas* Plantlets. *Biomass Bioener.*, 44: 42-46.
- Verbeek, M., Van Dijk, P. and Van Well, P. M. A. 1995. Efficiency of Eradication of Four Viruses from Garlic (*Allium sativum*) by Meristem-tip Culture. *Eur. J. Plant Pathol.*, 101: 231-239.
- Yancheva, S. D., Golubowicz, S., Yablowicz, Z., Perl, A. and Flaishman, M. A. 2005. Efficient Agrobacterium-mediated Transformation and Recovery of Transgenic Fig (*Ficus carica* L.) Plants. *Plant Sci.*, 168: 1433–1441.
- Zuraida, A. R., Fatin, L. I. K. and Ayu, N. O. 2014. *In vitro* Plant Propagation for Rapid Multiplication of *Melicope lunu-ankenda*: A Olant Species of High Medicinal Value. *Int. J. Pharm. Bio. Sci.*, 5: 1148-1156.



تکثیر درون شیشه ای بَراَنجیر و ارقام انجیر ماده (.*Ficus carica* L) از نوک شاخه ها

چ. بایوده، ر. لابیدی، ۱. مجدوب، و م. مارس

چکیدہ

برگ ها و میوه های درختان انجیر در تهدید حمله مرض موزائیک انجیر(FMD) قرار دارند که توسط ویروس هایی از چند جنس ایجاد می شود. کشت نوک شاخه (یا ساقه) روش راحتی برای بهسازی و عاری کردن از ویروس است. به این منظور، دستورالعملی قابل اعتماد برای تکثیر سریع درون شیشه ای نوک شاخه های سه رقم محلی و اصلی انجیر تونس (.Ficus carica L) به نام های زیدی (ZDI)، سلطانی (SNI) و بیذر ابیاد (BA) و یک بَراَنجیر گرده افشان مقاوم و کمیاب به نام عصافری (ASF) تهیه شد. در هر گام از این فرایند درون شیشه ای، چهار بستر رشد Murashige and Skoog (MS) با ترکیب های مختلف تنظیم کننده های رشد گیاه به کار گرفته شد. بهترین فرایند آغازین نوک شاخه ها با اندازه های ۱٬۰٫۵ ، و ۱/۵ میلی متر روی بستر M3 به دست آمد که حاوی ۲/۲ میلی گرم در لیتر بنزیل آمینو پورین (BAP)، ۰/۱ میلی گرم در لیتر نفتالین استیک اسید (NAA) و ۰/۱ میلی گرم در لیتر جیبرالیک اسید (GA₃) بود. رقم SNI با نرخ رشد ۱۰۰٪ ریز نمونه بیشترین فرايند آغازين نوک شاخه ها را براي مرحله استقرار نشان داد. تکثير ساقه و رشد گياهچه در بستر M₆ بود که حاوی ۰/۵ میلی گرم در لیتر BAP و ۰/۱ میلی گرم در لیتر NAA بود. بالاترین میانگین افزایش تعداد برگ (۹۲ برگ در هر گیاه) و شاخه زنی(۱۶/۹۱ شاخه در هر گیاه) روی بسترM₆ رخ داد. بهترین نرخ ریشه زنی (۸۳/۳۴٪) در بستر M₁₁ثبت شد که حاوی نیم–قدرت MS و ۱ میلی گرم در ليتر ايندول-۳-بوتيريک اسيد (IBA) بود. ريشه زايي گياهچه هاي انجير در شرايط طبيعي(مزرعه اي) روی پیت مرطوب با موفقیت ۹۰٪ انجام شد. گیاهچه های ساز گارشده انجیر های درون شیشه ای نرخ استقرار زیاد (۹۲/۱٪) و رشدی سریع روی بستر S1 که حاوی پیت بود نشان دادند و علایم امراض ويروسي يا نابهنجاري هاي شكلي در آنها مشاهده نمي شد.