

## Evaluation of Water Stress Memory in Compensation Response of Cotton (*Gossypium hirsutum* L.) during Subsequent Water Deficiency

M. Habibi<sup>1</sup>, E. Faghani<sup>2\*</sup>, M. Rezaei<sup>1</sup>, M. A. Mahmood Janlou<sup>1</sup>, and M. H. Razzagi<sup>3</sup>

### ABSTRACT

This research was carried out to provide suitable cotton seed for seed propagation in dryland. In this study, the potential of cotton seeds that have been stressed for the third-consecutive year was investigated to evaluate water stress memory responses. The experiment was arranged in split-plot factorial design with four irrigation levels of  $W_0$  (No-irrigation),  $W_1$  (33% FC),  $W_2$  (66% FC), and  $W_3$  (100% FC), as the main factor, and five seed treatments (four third-stressed seeds, i.e.  $S_{21}$  to  $S_{24}$ , and registered seed), as a sub-plot. Seeds of cotton were grown under different levels of water-stress exposure for three crop-seasons. As results showed,  $S_{32}$  received water stress signal in both  $W_0$  and  $W_3$  conditions through physiological mechanisms change. Seeds of  $S_{32}$  accumulated the lowest ABA and the highest calcium in exposure to  $W_0$  and  $W_3$ . Enhancement to superoxide dismutase and Aspartate peroxidase activity in leaves of  $S_{32}$  in exposure to  $W_0$  and  $W_1$  is another memorial stress mechanism for scarce water acclimation. The highest-potential thirty-boll weight, thirty-fiber weight, and first-harvesting yield were obtained from  $S_{32}$  against  $W_0$ ,  $W_1$ , and  $W_2$ . Also, the seeds of  $S_{32}$  had the most seedling vigor and germination percentage in exposure to  $W_0$ ,  $W_1$ , and  $W_2$ . It can be concluded that stress memory, via modification of physiology and morphology of plant behavior, helps plants to tolerate water deficiency when subjected to recurrent drought.

**Keywords:** Compensation mechanism, Drought tolerant, Seed, Yield.

### INTRODUCTION

Drought is a common abiotic stress that severely limits production of cotton (*Gossypium hirsutum* L.) grown in rain-fed farms of Iran (Zare *et al.*, 2014). Severe water stress may result in plant death, whereas mild water stress, in a short period of time, may improve drought tolerance by inducing stress memory that is introduced as drought imprinting, drought priming (as training and conditioning), hardening, and drought adaptation (Wojtyla *et al.*, 2020). In fact, plant

stress “memory” occurring in subsequent exposures to plants makes them more resistant to future exposure (Tombesi *et al.*, 2018). The main habitat of cotton cultivation was the moderate climate in the north of Iran, but these days, the cotton planted area has been reduced (Kolahi *et al.*, 2021). Indeed, irrigated cotton farmers have moved to dryland (Faghani *et al.*, 2018). On the other hand, episodic water shortage influences cotton physiological traits and crop production. Plants often remember environmental stress that they have encountered during growth to harvesting time (Wojtyla *et al.*, 2020). This ability of plants

<sup>1</sup> Department of Biology, Faculty of Basic Sciences, Gorgan Branch, Islamic Azad University, Gorgan, Islamic Republic of Iran.

<sup>2</sup> Department of Agronomy, Cotton Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Gorgan, Islamic Republic of Iran.

<sup>3</sup> Department of Agricultural Engineering, Golestan Agriculture and Natural Resources Research and Education Center, Agricultural Research, Education and Extension Organization (AREEO), Gorgan, Islamic Republic of Iran.

\*Corresponding author; e-mail: elhamfaghanibio@gmail.com



reactions refers to plant memory. Sometimes these physiological changes to manage environmental stress are temporary. However, when stress is continued, physiological modifications are stable and plants memorize them, and they transfer this information to the following generations as a stress memory (Sun *et al.*, 2021). Moreover, modern plant physiologists emphasize “memorial plant” and their behavior (Wojtyla *et al.*, 2020). There is limited information about stress memory and plant behavior against drought stress (Crisp *et al.*, 2016). Therefore, recognizing cotton metabolic reactions to reproduce seed for dry field requires management practices in order to achieve potential yield and drought tolerant seed. Assimilation, which is important to growing bolls and subtending leaves, comes from “main stem leaves” (Schubert *et al.*, 1986). Severity drought stress was declined by the accumulation of such compounds as abscisic acid (ABA), as a fundamental hormonal signal and enzymatic antioxidants adjustment in double-stressed reaction was declined. In exposure to water-stress consecutively, ABA can induce starch mobilization, which is an essential signal for metabolic adjustment against stress (Kempa *et al.*, 2008). Wang *et al.* (2019) results suggest that ABA mainly acts in cotton seed after ripening. Therefore, ABA plays an important role in seed germination and dormancy. Then, it reduces the damage effects to seed germination against abiotic stress by changing ABA regulation (Zhang *et al.*, 2014). The most common special role of Calcium (Ca) is known as adaptation against water stress. Under drought-stress condition, an increase in ABA induced the accumulation of intracellular Ca, which is known as a secondary messenger (Nayyar and Kushal, 2002). ABA and Ca increase the activities of Superoxide Dismutase (SOD) and Ascorbate Peroxidase (APX) against drought stress. The fundamental scavenging enzymatic defense antioxidants, in exposure to water deficiency stress, are SOD and APX (Ighodaro and Akinloye, 2018). ROS scavenging enzymes, such as SOD and APX, play an important role in cellular parts and act as a detoxifier for

cellular survival (Wahid *et al.*, 2007). Prolonged water stress exposure during subsequent drought stress leads to further acclimation by activating ABA, Ca, antioxidant enzyme, and morphological trait changes (Kolahi *et al.*, 2020). Antioxidant enzyme and the accumulation of ABA and Ca compound influence seed germination and seedling vigor. Seed vigor is important to evaluate the potential for rapid, uniform emergence, and development of normal seedlings under a wide range of field conditions (Alvarenga and Marcos-Filho, 2014). Therefore, yield components, such as thirty boll weight and fiber weight, were affected in exposure to subsequent drought stress. In cotton, water deficit at the filling stage influences crop production (Foulkesa *et al.*, 2007). Due to unlimited growth, cotton is often known as a drought-resistant crop and compensates for the loss of yield by activating the drought tolerance mechanism, including physiological and biochemical changes (Niu *et al.*, 2018). Evidence has indicated that plants adapt through changing their physiology and morphology characteristics in reaction to the previous occurrence of water stress (Kolahi *et al.*, 2021). Then, some physiological mechanisms store this information from previous water stress exposure by activating memory on plant behavior during subsequent water deficiency stress.

On the basis of the aforementioned points, the aim of this research was to investigate the role of three season’s exposures to water stress in activating water stress memorial reaction in cotton seed production in order to improve acclimatization and adaptive capacity to water deficiency condition, and also its transference to the next generations.

## MATERIALS AND METHODS

### Plant Material

Field experiments were carried out at Hashemabad Cotton Research Station. This station site is at the southeast corner of the Caspian Sea (36° 51' N latitude, 54° 16' E


longitude, and 13.3 m above mean sea level). The soil texture was sandy clay silt (6, 6.8, 3) throughout the 0.5 m soil profile. Water content at field capacity and wilting point were 28.1 and 14.1% by soil volume.

Figure 1 shows the diagram that summarizes three years of research. In the first year, once water stress exposure was done, (2016), registered seeds were prepared from the market for the purpose of the study. Water treatments were no-irrigation [rain-fed], 33, 66, and 100% FC ( $W_0$ ,  $W_1$ ,  $W_2$ , and  $W_3$ , respectively). The seeds, i.e., Golestan cultivar, were named  $S_{10}$ ,  $S_{11}$ ,  $S_{12}$ ,  $S_{13}$ , and  $S_{14}$  and were grown under  $W_0$ ,  $W_1$ ,  $W_2$ , and  $W_3$  treatments.

In the second year (2017), the second water stress exposure was done. Seed treatments were chosen from the ones exposed to stress once ( $S_{11}$ ,  $S_{12}$ ,  $S_{13}$ , and  $S_{14}$ ) and the certified seeds were obtained from the market ( $S_{10}$ ). These cultivated seeds were subjected to irrigation treatments similar to the first year. Then, the harvested seeds were named  $S_{20}$  (from the market),  $S_{21}$ ,


$S_{22}$ ,  $S_{23}$ , and  $S_{24}$  respectively, and were used for the following crop season.

The third water-stress exposure was done in the third year of the experiment (2018). Five seed treatments, i.e.,  $S_{20}$  (purchased from the market),  $S_{21}$ ,  $S_{22}$ ,  $S_{23}$ , and  $S_{24}$ , were cultivated. Then, the registered seeds, i.e., ( $S_{21}$  and  $W_0$ ), ( $S_{22}$  and  $W_1$ ), ( $S_{23}$  and  $W_2$ ), and ( $S_{24}$  and  $W_3$ ), were labeled as  $S_{30}$ ,  $S_{31}$ ,  $S_{32}$ ,  $S_{33}$ , and  $S_{34}$  according to exposure to four irrigation treatments (Non-irrigation [rain-fed] and 33, 66, and 100% FC) (Figure 1). This experiment was done in split-plot factorial design with three replications. Different irrigation treatments were started from the first square formation to the boll opening (in average 50 days) (Figure 2). Different irrigation treatments were set in the main plots, and seeds with two water stress exposures were cultivated in the sub-plots. In this experiment, seeds spacing were 20 cm (inter row) and 80 cm (intra rows) (Figures 3-a and -b). The amount of irrigation was measured by a water flow meter, and the time of irrigation was



\*  $S_{11}$  means S (Seeds), the first number (1) used for first generation, and the second number (1) Used for the plot in which seed generated (first irrigation treatment or  $W_1$ )

**Figure 1.** Diagram of conducting three years of research.



**Figure 2.** Sowing to harvest schedule and irrigation treatments for seed productions from double water stress exposure to the third water stress exposure experiments (2018 crop-season).



(a)



(b)

**Figure 3.** (a) Cotton seed cultivation and different irrigation treatments in farm. (b) Sampling cotton leaves of different seeds from third water stress exposure under different irrigations.

determined by gravimetric determination of soil moisture.

### Biochemical Studies

The seed and leaf samples were collected for biochemical analyses and were kept in liquid nitrogen to be carried out at the plant physiology laboratory. The following biochemical traits were evaluated:

#### Calcium and ABA Assay

Ca was extracted using EDTA method based on Motsara and Roy (2008). After burning 5 g of powdered seed samples in the muffle furnace at 550°C for about 12 hours, ashes were dissolved in HCl.

ABA content of seeds were estimated using HPLC (Unicam-Crystal-200, UK) according to Kelen *et al.* (2004) method. In brief, 100 mg seed dry weight was homogenized with 10 mL, 80% methanol that contained PVP. The pH of extraction was adjusted to 2.8 with H<sub>3</sub>PO<sub>4</sub>. The acidic solution was passed through a Diamonsic, C<sub>18</sub>, 5 µm cartridge to trap and concentrate ABA. The acidic solution was passed through HPLC 184 Eurospher-100 C18 column (250×4 mm ID; 5 µm particle size, 185 Knauer, Germany) by filtering a 0.45 µm membrane 183 syringe filter (Whatman, England). The isocratic mobile phase consisted of 100% methanol and 0.2% acetic acid (50:50 v/v) at a flow rate of 0.7 mL/min. ABA peak was estimated using a standard solution and the signal was monitored at 257 nm at 40°C.

#### Antioxidant Enzyme

##### SOD Activity Enzyme

SOD was analyzed using Beyer and Fridovich (1987) method. In doing so, about 1 g fresh weight was grounded with a mixture of 1 mL of 50 mM potassium

phosphate and 1 mM EDTA (pH 7.5) buffer. The homogenates solutions were centrifuged at 4°C for 15 minutes at 43,000×g. The supernatants were used for the estimation of enzymatic activities and protein concentrations. One enzyme unit of SOD was determined by spectrophotometer at 560 nm (Hitachi model U- 2000) based on the amount of enzyme reduction required to cause 50% inhibition of the rate of NBT (Nitroblue tetrazolium) was measured.

##### APX Enzyme Activity

APX assay was carried out following Nakano and Asada (1981). One g seed fresh weight was homogenized in the extracted solution [50 mM Na<sub>2</sub>HPO<sub>4</sub>/KH<sub>2</sub>PO<sub>4</sub> (pH 7.0), 1 mM EDTA, 5% polyvinylpyrrolidone (PVP-40) and 1 mM ASA]. The homogenate was centrifuged at 105,000×g for 10 minutes at 4°C. Oxidation of ASA was read at 300 nm (extinction coefficient 0.74 mM<sup>-1</sup>cm<sup>-1</sup>). APX was figured as the amount of enzyme oxidizing 1 µmol of ASA per minute.

##### Germination Percentage Test (Gmax)

Fifty Fuzzy seeds were placed on a filter paper (25×38 cm) and between double layered rolled germination papers in three replications. Each roll had been soaked with deionized water for 24 h. Paper rolls with seeds were put in a germination chamber (Paradise, Iran) at a constant temperature of 30°C with 250 mol m<sup>-2</sup>s<sup>-1</sup> light intensity (diurnal cycle was 8 hours light and 16 hours darkness). After nine days, the germination percentage (%) was assessed using the following equation:

$$G_{\max} = 100 \times \frac{\text{Germinated seed number}}{9\text{th day}/50}$$

##### Cool Warm Vigor Index (CWVI)

The testing procedure consisted of the following steps:



The cool germination test was estimated at 18°C. After seven days, the number of normal seedlings greater than 1.5 inches long was counted (AOSA, 1993). The standard warm germination test was studied (alternating 20°C for 16 hours and 30°C for 8 hours, respectively). Then, the number of normal seedlings that were greater than 1.5-inch-long were counted after four days. Finally, the results of the two tests were added.

The following rating scale is currently being used based on AOSA (1993):

CWVI	Rating
160+	Excellent
140-159	Good
120-139	Fair
<120	poor

### Statistical Analysis

The collected data were analyzed based on the average of three plants as one replicate


for each treatment. SAS package (9; SAS Institute, Cary, NC, USA) was utilized to analyze data. Specifically, analysis of variance was estimated by Proc-Mixed model using Fisher's Least Significant Difference (LSD) test at the 5% level (SAS, 2001).

## RESULTS


### Biochemical Studies

#### ABA and Calcium Concentration

As the results demonstrated, the highest and the lowest ABA levels were obtained in S<sub>34</sub> and S<sub>32</sub> against W<sub>0</sub> water stress exposure with 46.4 and 19.4 ng g<sup>-1</sup> FW, respectively. Also, in W<sub>2</sub> treatment, ABA content in seeds of S<sub>34</sub> was the highest. In W<sub>1</sub> treatment, ABA accumulation in the seeds of S<sub>32</sub> increased to 41.8 ng g<sup>-1</sup> FW compared to other seed treatments (Figure 4). As Figure 3 displays, ABA concentration seeds of S<sub>31</sub> in the third



**Figure 4.** ABA content trend in cotton seed under different irrigation treatments [W<sub>0</sub>; Rain-fed (without irrigation); W<sub>1</sub> (33% FC); W<sub>2</sub> (66% FC) and W<sub>3</sub> (100% FC)] and seed sources [(registered seeds); (S<sub>21</sub> and W<sub>0</sub>); (S<sub>22</sub> and W<sub>1</sub>); (S<sub>23</sub> and W<sub>2</sub>); and (S<sub>24</sub> and W<sub>3</sub>) were labeled as S<sub>30</sub>, S<sub>31</sub>, S<sub>32</sub>, S<sub>33</sub>, and S<sub>34</sub> in the third water stress exposure. (Different letters represent significantly different).



**Figure 5.** Calcium content trend in cotton seed under different irrigation treatments [ $W_0$ ; Rain-fed (without irrigation);  $W_1$  (33% FC);  $W_2$  (66% FC) and  $W_3$  (100% FC)] and seed sources [(registered seeds); ( $S_{21}$  and  $W_0$ ); ( $S_{22}$  and  $W_1$ ); ( $S_{23}$  and  $W_2$ ); and ( $S_{24}$  and  $W_3$ )] were labeled as  $S_{30}$ ,  $S_{31}$ ,  $S_{32}$ ,  $S_{33}$ , and  $S_{34}$  in the third water stress exposure. (Different letters represent significantly different).

exposure to  $W_3$  irrigation was the most compared to other seed treatments. As the result of Ca analysis showed,  $S_{32}$  against  $W_0$  and  $W_3$  had the highest ( $151.35 \text{ mg } 100 \text{ g}^{-1}$ ) and the lowest ( $87.85 \text{ mg } 100 \text{ g}^{-1}$ ) Ca content in matured seed, respectively. Being exposed to  $W_0$  for the third time, Ca content in seeds of  $S_{34}$  decreased compared to other seed treatments (Figure 5). As the data showed, there was a reverse relationship between the accumulation of ABA and Ca in cotton seed. Also,  $S_{31}$  received drought-signal in treatment  $W_1$  and  $W_3$  irrigations due to having more ABA and low Ca content compared to other seed treatments, which might have weak germination for the next crop season. Generally, in  $W_0$ ,  $S_{32}$  seeds stored the most Ca and the least ABA, but in  $W_1$ , seeds of  $S_{32}$  had the greatest ABA and the lowest Ca (Figures 4 and 5).

### Antioxidant Enzyme

The effect of drought stress on the activities of antioxidant enzymes participating in the scavenging of ROS is


shown by evaluating SOD and APX activities.

### SOD Activity

SOD activity, as a radical scavenger enzyme, plays an important role against water shortage. However, on the basis of the results of the present study, SOD activity in seeds was more than its activity in cotton leaves (Figure 6).

In water stress exposure, SOD activity in cotton-seeds showed a significant variation among seed source and irrigation treatments. The values of SOD in cotton seed varied from 32.7 to  $47.4 \text{ U g}^{-1}$  total protein (Figure 6). Among these treatments, the highest SOD activity was found in  $S_{30}$  ( $47.7 \text{ U g}^{-1}$  total protein) seeds in  $W_0$  treatment, but the lowest SOD activity was related to  $S_{33}$  in  $W_3$  and  $W_0$  treatments (Figure 6). In  $W_0$  to  $W_2$ , SOD activity in the seeds of  $S_{32}$  was the least (Figure 6).

Different seed irrigating results were observed in a wide range of variations of SOD activity of leaves (Figure 6). The most



**Figure 6.** SOD activity content trend in cotton seed and different irrigation treatments [W<sub>0</sub>; Rain-fed (without irrigation); W<sub>1</sub> (33% FC); W<sub>2</sub> (66% FC) and W<sub>3</sub> (100% FC)] and seed sources [(registered seeds); (S<sub>21</sub> and W<sub>0</sub>); (S<sub>22</sub> and W<sub>1</sub>); (S<sub>23</sub> and W<sub>2</sub>); and (S<sub>24</sub> and W<sub>3</sub>) were labeled as S<sub>30</sub>, S<sub>31</sub>, S<sub>32</sub>, S<sub>33</sub>, and S<sub>34</sub> in the third water stress exposure. (Different letters represent significantly different).

SOD activity in the leaves of S<sub>32</sub> was obtained in W<sub>3</sub> with 0.19 U g<sup>-1</sup> total protein. Nevertheless, in W<sub>0</sub>, W<sub>1</sub>, and W<sub>3</sub> irrigation treatments, S<sub>32</sub> leaves had the most SOD activity (Figure 6). Moreover, SOD activity in the leaves of S<sub>32</sub> under W<sub>0</sub> and W<sub>1</sub> irrigation were 17 and 16.6%, respectively, more than S<sub>30</sub> (control) (Figure 6). In W<sub>2</sub> and W<sub>3</sub>, S<sub>32</sub> leaves had, respectively, 25.5% lower and 29.5% more SOD activity than S<sub>30</sub> (Figure 6). Also, SOD activity in the leaves of S<sub>32</sub> under W<sub>0</sub> and W<sub>1</sub> irrigations were 7 and 4.4% lower than that in S<sub>30</sub>.

### APX Activity


As the data showed, APX activity was lower than SOD activity in seeds and leaves. The present study indicated that the most significant APX activity was observed in the seeds of S<sub>31</sub> W<sub>3</sub> (Figure 7). Also, APX activity in the seeds of S<sub>32</sub> W<sub>0</sub> was the lowest in comparison to other treatments in this condition. In the third W<sub>2</sub> irrigation, APX activity in the seeds of S<sub>32</sub> was the

most while APX activity in S<sub>33</sub> seeds was the lowest, compared to others. The assessment of the APX activity in W<sub>3</sub> revealed that the S<sub>31</sub> and S<sub>32</sub> seeds had the highest while the seeds of S<sub>33</sub> had the lowest APX activity (Figure 7). Evaluation of APX activity of leaves against several irrigation treatments and seeds gathered from the third water stress exposure indicated that the highest APX activity was related to the seeds of S<sub>31</sub> against W<sub>2</sub> irrigation. (Figure 7). APX activity in the leaves of S<sub>32</sub> was about 84% lower than the leaves of S<sub>31</sub> in W<sub>2</sub> exposure (Figure 7).

### Seed Gmax and Seedling Vigor

Gmax and the vigor of seeds, which were harvested after the third water stress exposure, were estimated in each irrigation plot. The data showed that the maximum Gmax (100%) was related to S<sub>32</sub> and S<sub>33</sub> in W<sub>2</sub>, and S<sub>33</sub> in W<sub>3</sub> irrigation. To state it differently, the least Gmax, which was 80%, was obtained from S<sub>30</sub> against W<sub>2</sub> and S<sub>31</sub> in W<sub>3</sub>. Seedling vigor resulting in W<sub>0</sub> stress appeared to reverse the





**Figure 7.** APX activity content trend in cotton seed and leaves under different irrigation treatments [ $W_0$ ; Rain-fed (without irrigation);  $W_1$  (33% FC);  $W_2$  (66% FC) and  $W_3$  (100% FC)] and seed sources [(registered seeds); ( $S_{21}$  and  $W_0$ ); ( $S_{22}$  and  $W_1$ ); ( $S_{23}$  and  $W_2$ ); and ( $S_{24}$  and  $W_3$ ) were labeled as  $S_{30}$ ,  $S_{31}$ ,  $S_{32}$ ,  $S_{33}$ , and  $S_{34}$  in the third water stress exposure. (Different letters represent significantly different).

G max result (Table 1). It means that, in exposure to  $W_0$  water stress, seeds with high potential G max had weak seedling vigor and vice versa. Also, in exposure to  $W_1$ , G max of  $S_{32}$  with 100% had the maximum seedling vigor index, i.e. 176, and  $S_{34}$  with 86% Gmax showed the lowest seedling vigor index, i.e. 138. In addition,  $W_2$  water exposure, seed of  $S_{32}$  and  $S_{33}$  in subjected to  $W_3$  irrigation represented 100% Gmax (Table 1).  $S_{32}$  and  $S_{33}$  in exposed to  $W_2$  irrigation were classified in the 'excellent' class. Seeds of  $S_{32}$  and  $S_{33}$  in exposure to  $W_1$  (with 176 and 152 seed vigor indexes, respectively) were set in the 'excellent' and 'good' classes, and seeds of  $S_{32}$  and  $S_{33}$  in exposure to  $W_0$  (with seedling vigor index 154 and 158, respectively) were classified in the 'good' group.

### Yield

Water stress significantly affected the weight and fiber of thirty-boll.  $S_{32}$  in  $W_1$  and  $W_2$  had the highest thirty-boll weight with 164 g and fiber of thirty boll weight with

61.66 g per plot in the first harvest. However, the minimum thirty-boll weight and fiber at the first harvest were obtained from  $S_{31}$  against  $W_3$  and  $S_{33}$  in  $W_0$  with 99.7 g and 31.53 g per plot, respectively.

Boll weight trait was affected differently among deficit irrigated conditions and seed treatments. The most significant boll weight at the first harvest was observed in  $S_{32}$   $W_1$  (Table 1). Additionally, thirty-boll weight and first harvest yield in  $S_{32}$  were increased in all irrigation treatments, i.e.,  $W_0$ ,  $W_1$ , and  $W_2$  and were more than other seed treatments (Table 1). The highest yield in the first harvest was gained from  $S_{32}$   $W_2$  with 1581.3 g per plot, while in 6.2-fold was more in comparison to  $S_{32}$  under  $W_0$  treatment. As the results showed, in  $W_0$ , the first harvest yield of  $S_{32}$  produced 2.9- and 4.1-fold of  $S_{30}$  and  $S_{31}$  (Table 1).

### DISCUSSION

Recently, dryland areas have been extended. Although facing long-term water shortage in cotton causes loss in yield, the impact of



**Table 1.** Irrigation treatments [ $W_0$ ; Rain-fed (without irrigation);  $W_1$  (33% FC);  $W_2$  (66% FC) and  $W_3$  (100% FC)] and seed sources [(registered seeds); ( $S_{21}$  and  $W_0$ ); ( $S_{22}$  and  $W_1$ ); ( $S_{23}$  and  $W_2$ ); and ( $S_{24}$  and  $W_3$ )] were labeled as  $S_{30}$ ,  $S_{31}$ ,  $S_{32}$ ,  $S_{33}$ , and  $S_{34}$  effects on first harvest, thirty- Boll- and thirty-fiber weight, germination (Gmax) and seedling vigor in exposed to third water stress condition. (Different letters represent significantly different).

WS Treatment	Thirty-Boll Fiber Weight (First harvest) (g 3 m <sup>-2</sup> )	Thirty-Boll Weight (First harvest) (g 3 m <sup>-2</sup> )	Yield (First harvest) (g 3 m <sup>-2</sup> )	Germination (percentage)	Seedling Vigor (Index)
$W_0S_{30}$	38.90 ±7.60 g-i	121.00 ±17.05 c-h	55.3 ± 6.00 h	88 ±3.25 a-c	150 ±1.62 i-l
$W_0S_{31}$	42.80 ±1.94 d-i	134.33 ±5.78 a-f	42.0 ± 21.35 h	96 ±3.25 ab	164 ±3.25 e-g
$W_0S_{32}$	46.33 ±0.88 c-h	136.33 ±1.85 a-e	217.3 ± 78.28 gh	94 ±1.62 ab	154 ±1.62 g-j
$W_0S_{33}$	33.53 ±8.13 i	104.10 ±26.70 f-h	126.0 ± 66.74 gh	92 ±3.25 a-c	158 ±1.62 f-i
$W_0S_{34}$	39.73 ±6.36 f-i	122.00 ±22.32 c-h	198.3 ± 75.98 gh	98 ±1.62 ab	146 ±1.62 j-m
$W_1S_{30}$	49.70 ±2.40 b-f	144.47 ±7.29 a-d	526.0 ± 86.53 d-g	90 ±4.87 a-c	160 ±3.25 e-i
$W_1S_{31}$	50.83 ±1.75 b-e	151.00 ±4.04 a-c	260.3 ± 172.34 gh	96 ±3.25 ab	162 ±4.87 e-h
$W_1S_{32}$	58.86 ±2.25 ab	164.00 ±6.25 a	1297.3 ± 33.94 ab	96 ±3.25 ab	176 ±3.25 b-d
$W_1S_{33}$	47.13 ±8.23 c-h	131.33 ±19.00 b-f	717.0 ± 119.65 d-f	94 ±1.62 ab	152 ±0.0 k-j
$W_1S_{34}$	56.13 ±1.27 a-c	154.77 ±2.42 ab	788.7 ± 104.18 de	86 ±4.87 bc	138 ±1.62 m
$W_2S_{30}$	55.80 ±1.74 a-c	143.33 ±4.05 a-d	1219.3 ± 192.04 a-c	80 ±6.50 c	138 ± 4.87 m
$W_2S_{31}$	51.66 ±1.76 a-d	129.00 ±2.00 b-h	1308.7 ± 81.71 ab	94 ±1.62 ab	178 ±1.62 bc
$W_2S_{32}$	61.66 ±1.20 a	149.67 ±3.28 a-c	1581.3 ± 95.69 a	100 ±0.0 a	162 ±4.87 e-h
$W_2S_{33}$	51.00 ±2.00 a-d	126.67 ±5.84 b-h	912.7 ± 82.28 b-d	100 ±0.0 a	192 ±3.25 a
$W_2S_{34}$	53.73 ±3.41 a-c	127.80 ±5.77 b-h	1496.0 ± 72.53 a	98 ±1.62 ab	170 ±4.87 c-e
$W_3S_{30}$	47.00 ±1.52 c-h	115.43 ±4.16 d-h	846.0 ± 136.72 cd	98 ±1.62 ab	166 ±1.62 d-f
$W_3S_{31}$	36.83 ±0.84 hi	99.77 ±2.47 h	399.3 ± 41.95 d-h	80 ±0.0 c	142 ±1.62 k-l
$W_3S_{32}$	40.06 ±0.96 e-i	101.13 ±0.94 gh	260.3 ± 34.72 f-h	90 ±8.12 a-v	140 ±0.0 ml
$W_3S_{33}$	47.73 ±1.82 c-g	117.67 ±2.60 d-h	698.7 ± 109.09 d-f	100 ±0.0 a	182 ±1.62 ab
$W_3S_{34}$	47.33 ±1.20 c-h	112.77 ±10.63 e-h	872.7 ± 43.63 cd	94 ±1.62 ab	166 ±1.62 df

drought-stress memory is induced by using some morphological parameters and physiological mechanisms to compensate for the effects of damage. Moreover, the heritable memory, which was known as “plant stress memory”, helps plants to react efficiently in exposure to stresses, not only for the current crop season but also for the next generations. Developing stress tolerance in plants for achieving high yield potential and stability affects modification of physiological mechanism.

#### ABA and Calcium Concentration

According to the results, seeds that had different irrigation treatments during water

stress exposure displayed different drought stress signals using ABA and Ca changes. ABA data showed that the selection of rain-fed and 100% FC plots for seed production was not appropriate since the seeds had high ABA concentration and showed drought signal. As Richardson *et al.* (2019) believed, high ABA stored in seeds would affect seed viability. It is evident that ABA accumulation in these seeds may induce dormancy and inhibition of germination (Papenfus *et al.*, 2013). High Ca plays an important role in adaptation to drought by improving seedling vigor and leaf water potential protection (Kaczmarek *et al.*, 2017). Thus, the existence of high Ca and low ABA in matured cotton seed makes the stress injury sign better and different

treatments have various characters (Nayyar and Kaushal, 2002). Therefore, the seeds that were cultivated in exposure to one-third and full-irrigation needs, were reproduced under rain-fed conditions. This production occurred because these seeds had more ABA and low Ca content compared to other seed treatments, which may have weak germination for the next crop season. It can be concluded that, ABA storage in seed reproduction of  $S_{32}$  in 33% FC water need exposure was not enough to prevent seed germination in the next crop season. Li *et al.* (2019) believed that ABA storage as a memory sign induced after the initial stress but was not stable for memory reactions to subsequent water stress. However, it should be noted that the most Ca content, as a secondary messenger in response to stress (Vishwakarma *et al.*, 2017), and the least ABA content of  $S_{32}$  seeds under rain-fed illustrated drought tolerant sign. Therefore, on the basis of ABA and Ca content of seed, seed production with 33% FC can be recommended for rain-fed farms.

### Antioxidant Enzyme Activity

#### SOD Activity

According to our results, the increase in SOD activity in  $S_{30}$  seeds under rain-fed conditions was representative of the water deficiency sign. Studies of SOD activity of leaves showed that with the aim of seed reproduction,  $S_{32}$  in comparison to  $S_{30}$  (control), revealed a high potential to eliminate  $O_2$  under water shortage and high irrigation conditions. Also, the most SOD activity in the seeds of  $S_{32}$  in 33% and 66% FC treatments revealed that these water deficiencies were sufficient to activate antioxidant metabolism. Moreover, the activity of SOD in seeds was more than SOD activity in leaves. Also, the seeds that were grown in this treatment would be more tolerant than rain-fed and 100% FC water stress exposure. The induction of SOD activity is recognized as plant ability to

overcome oxidative stress by up-regulation of antioxidant enzymes (Alscher *et al.*, 2002). According to this fact,  $S_{33}$  seeds under  $W_3$  and  $W_0$  and  $S_{32}$  seeds in  $W_2$  and  $W_3$  had low active oxygen species against substrate; therefore, SOD activity would decrease. Moreover, the lowest SOD activity in  $S_{33}$  seeds in exposure to  $W_1$  and  $W_2$  irrigation proved that these seeds do not show any water shortage signal. It seems that in spite of increasing SOD activity in the leaves of  $S_{32}$ , the activity of SOD was reduced in cotton seeds under  $W_1$  irrigation. This result indicated that seeds of  $S_{32}$  could activate memory of drought-tolerant stress for planting against  $W_1$ . Many reports have shown that abiotic stress could provoke activation of SOD by overproduction of ROS, but other information displayed the opposite trend. These contradictions were related to plant species and long- period or short-period exposure to stress (Szollosi, 2014).

#### APX Activity

As the results show,  $S_{32}$  would be able to decrease APX activity in seeds and increase APX activity in leaves under  $W_0$  and  $W_1$  irrigation. Higher APX activity in the seeds of  $S_{32}$  in  $W_2$  water exposure enhances tolerance to low water shortage more than in  $W_1$  treatment. APX plays an important role in the removal of  $H_2O_2$ , like its substrate, i.e., ascorbic acid. According to our data, APX activity in leaves was higher than that in seed. The activity of changes in antioxidant enzymes was also highlighted as one of the mechanisms of drought stress memory. The report of Gunes *et al.* (2008) in sunflower is in line with our result. One of the highest important signals of drought-tolerant mechanism is overexpression of SOD with increasing  $H_2O_2$  scavenging enzyme, such as APX activities (McKersie *et al.*, 1999). It is clear that  $S_{33}$  produced seeds with low APX activity in exposure to  $W_0$ ,  $W_2$ , and  $W_3$  and that it did not show any drought signal through serving memorial



stress. However,  $S_{32}$  leaves reacted to drought stress by increasing APX and SOD activation in exposure to  $W_0$  and  $W_1$ .

### Gmax and Seedling Vigor Index

The potential attributes of seed vigor index as a crucial physiological seed feature was evaluated for seedling establishment in farm. Gmax and seedling vigor traits were studied. The increase in ABA concentration in seed may be related to seed dormancy and low seed vigor (Yao *et al.*, 2019). Metabolic compound seed affects seedling vigor directly or indirectly. Therefore, seed tolerance and seed quality are assessed by seed vigor (Marcos Filho, 2015). As it was revealed in this research, seeds of  $S_{33}$  against 100% FC irrigation, had high potential seed vigor index, while in this irrigation conditions, seeds of  $S_{32}$  had fair seedling vigor index. It was found that seed reproduction, after the third 33 and 66% FC water stress exposures, were irrigated with 66% FC and had high seed vigor index. Based on Hopper and Hinton (1980) researches,  $S_{32}$  and  $S_{33}$  exposed to  $W_0$ ,  $W_1$ , and  $W_2$  were able to have potential establishment in these irrigation conditions; therefore, using a low planting rate is needed.

### Yield

The cotton yield will suffer serious irreversible damage when seeds are exposed to a range of intolerable water shortages (Niu *et al.*, 2018). Our data and Wang *et al.* (2013) report also showed that the thirty-boll weight and yield decreased under drought. Nevertheless, at the first harvesting period, thirty-boll weight and thirty-fiber weight enhanced in  $S_{32}$  when exposed to  $W_0$ ,  $W_1$ , and  $W_2$  compared to  $W_3$ . Accordingly, the severity of the compensation aspect varied in different seed treatments against different irrigations. On the other hand, being exposed to  $W_1$  and  $W_2$ ,  $S_{32}$  showed the

influence of low water use in potential yield of rain-fed fields by producing 29.7 and 150% more yield in comparison to  $S_{30}$ . It is in line with Papastylianou and Argyrokastritis (2014) studies that showed the enhancement of cotton yield and fiber weight of cotton cultivars when subjected to moderate drought stress. The greatest thirty-boll weight and the first harvest yield of  $S_{32}$ , which was exposed to  $W_2$  water stress, can be a sign of stress memory. This finding is also in line with Kim *et al.* (2020).

### CONCLUSIONS

Decrease in yield under drought stress has been a major challenge for staple crops. Therefore, the identified drought-induced changes are primarily required in intergenerational stress memory. The long-term physiological memory in response to drought stress exposure showed a decrease in ABA level of seed, Ca content of seed, and activity of SOD and APX activity. These changes might be due to the stress memory mechanism. In summary,  $S_{32}$  was able to overcome the applied stresses and produced the highest yield in the first-harvest. Memory of drought stress through activating antioxidant enzyme in leaves prevented transport and storage of free radicals in seed. Based on the basis of role regulation of memorial water stress, that is formed or forgotten in cotton during period of stress,  $S_{32}$  could balance resources allocated and improved acclimation physiological mechanisms under  $W_0$ ,  $W_1$ , and  $W_2$  irrigation. Generally, for seed propagation in *Gossypium hirsutum*, 33% FC treatment activated plant stress memory by adjusting physiological metabolism for the third generation.

### Abbreviations

Abscisic Acid (ABA), Aspartate Peroxidase (APX), Calcium (Ca), Leaf Area

(LA); Field Capacity (FC); Super Oxide Dismutase (SOD).

### ACKNOWLEDGEMENTS

We thank MS Dodangie for her helping.

### REFERENCES

- Alscher, R. G., Erturk, N. and Heath, L. S. 2002. Role of Superoxide Dismutases (SODs) in Controlling Oxidative Stress. *J. Exp. Bot.*, **53**: 1331–1341.
- Alvarenga, R. O. and Marcos-Filho, J. 2014. Vigor Evaluation of Stored Cotton Seeds, Including the Seed Vigor Imaging System (SVIS®). *J. Seed Sci.*, **36(2)**: 222-230.
- Association of Official Seed Analysts, 1993. Rules for Testing Seeds. *J. Seed Technol.*, **16(3)**: 1-126.
- Beyer, W. F. J. R. and Fridovich, I. 1987. Assaying for Superoxide Dismutase Activity: Some Large Consequences of Minor Changes in Conditions. *Annu. Biochem.*, **161**: 559-566.
- Crisp, P. A., Ganguly, D., Eichten, S. R., Borevitz, J. O. and Pogson, B. J. 2016. Reconsidering Plant Memory: Intersections between Stress Recovery, RNA Turnover, and Epigenetics. *Sci Adv.*, **2**: e1501340.
- Faghani, E., Razzaghi, M. H., Sohrabi, B., Kolahi, M. and Chekani, O.B. 2018. Investigation of Physiological Characteristics and Vigor of Cotton Seeds under Water Stress and Evaluation of Their Tolerance in Golestan Province. Final Report of Complete Project: (Code: 3-07-075195105), Cotton Research Institute.
- Foulkesa, M. J., Sylvester-Bradleyb, R., Weightmanb, R. and Snapec, J.W. 2007. Identifying Physiological Traits Associated with Improved Drought Resistance in Winter Wheat. *Field Crops Res.*, **103(1)**: 11-24.
- Gunes, A., Pilbeam, D., Inal, A. and Coban, S. 2008. Influence of Silicon on Sunflower Cultivars under Drought Stress, I: Growth, Antioxidant Mechanisms and Lipid Peroxidation. *Commun. J. Soil Sci. Plant Nutr.*, **39**: 1885–1903.
- Hopper, N. W. and Hinton, H. R. 1980. The Use of Electrical Conductivity as a Measure of Cottonseed Quality. *Agron. Abstract.*, **109**.
- Ighodaro, O. M. and Akinloye, O. A. 2018. First Line Defense Antioxidants-Superoxide Dismutase (SOD), Catalase (CAT) and Glutathione Peroxidase (GPX): Their Fundamental Role in the Entire Antioxidant Defense Grid. *Alexandria J. Med.*, **54**: 287-293.
- Kaczmarek, J. C., Kowalski, P. S. and Anderson, D. G. 2017. Advances in the delivery of RNA therapeutics: from concept to clinical reality. *Genome Med.*, **9**: 602-16.
- Kelen, M., Cubukdem-Iralay, E., Sen, S. and Ozkan, G. 2004. Separation of Abscisic Acid, Indole-3 Acetic Acid, Gibberellic Acid in 99R (*Vitis berlandieri* × *Vitis rupestris*) and Rose Oil (*Rosa damascena* Mill.) by Reversed Phase Liquid Chromatography. *Turk. J. Chem.*, **28**: 603 – 10.
- Kempa, S., Krasensky, J., Dal Santo, S., Kopka, J. and Jonak, C. 2008. A Central Role of Abscisic Acid in Stress-Regulated Carbohydrate Metabolism. *PLoS One*, **3(12)**: e3935.
- Kim, Y., Chung, Y. S., Lee, E., Tripathi, P., Heo, S., and Kim, K. H. 2020. Root response to drought stress in rice (*Oryza sativa* L.). *Int. J. Mol. Sci.* **21**:1513.
- Kolahi, M., Faghani, E., Goldson-Barnaby, A. and Sohrabi, B. 2020. Physiological Traits and Anatomic Structures of the Seed for Two Short Cotton Season Genotypes (*Gossypium hirsutum* L.) under Water Stress. *J. Integr. Agric.*, **19(1)**: 89–98.
- Kolahi, M., Faghani, E., Kazemian, M., Goldson-Barnaby, A. Dodangi, S. 2021. Changes in Secondary Metabolites and Fiber Quality of Cotton (*Gossypium hirsutum*) Seed under Consecutive Water Stress and *in Silico* Analysis of Cellulose Synthase and Xyloglucan Endotransglucosylase. *Physiol. Mol. Biol. Plants*, **27**: 1837–1857.
- Li, p., Yang, H., Wang, L., Liu, H., Huo, H., Zhang, C., Liu, A., Zhu, A., Hu, J., Lin, Y. and Liu, L. 2019. Physiological and Transcriptome Analyses Reveal Short-Term Responses and Formation of Memory under Drought Stress in Rice. *Front Genet.*, **10**: 55.



18. Marcos Filho, J. 2015. *Fisiologia de Sementes de Plantas Cultivadas*. 2nd Edition, Abrates, Londrina, 660 PP.
19. Maxwell, K. and Johnson, G. N. 2000. Chlorophyll Fluorescence: A Practical Guide. *J. Exp. Bot.*, **51(345)**: 659-68.
20. McKersie, B. D., Bowley, S. R. and Jones, K. S. 1999. Winter Survival of Transgenic Alfalfa Overexpressing Superoxide Dismutase. *Plant Physiol.*, **119(3)**: 839-48.
21. Motsara, M. R. and Roy, R. N. 2008. *Guide to Laboratory Establishment for Plant Nutrient Analysis*. FAO Fertilizer and Plant Nutrition Bulletin, Food and Agriculture Organization, Rome, 219 PP.
22. Nakano, Y. and Asada, K. 1981. Hydrogen Peroxide is Scavenged by Ascorbate-Specific Peroxidase in Spinach Chloroplasts. *Plant Cell Physiol.*, **28**: 867-880.
23. Nayyar, H. and Kaushal, S. K. 2002. Alleviation of Negative Effects of Water Stress in Two Contrasting Wheat Genotypes by Calcium and Abscisic Acid. *Biol. Plant.*, **45(1)**: 65-70.
24. Niu, J., Zhang, S., Liu, S., Ma, H., Chen, J., Shen, Q., Ge, C., Zhang, X., Pang, C. and Zhao, X. 2018. The Compensation Effects of Physiology and Yield in Cotton after Drought Stress. *J. Plant Physiol.*, **224-225**: 30-48.
25. Papenfus, H. B., Kulkarni, M. G., Stirk, W. A., Finnie, J. F. and Van Staden, J. 2013. Effect of a Commercial Seaweed Extract [Kelpak (R)] and Polyamines on Nutrient-Deprived (N, P and K) Okra Seedlings. *Sci. Hortic.*, **151**: 142-146.
26. Papastylianou P. and Argyrokastritis, I. 2014. Effect of Limited Drip Irrigation Regime on Yield, Yield Components, and Fiber Quality of Cotton under Mediterranean Conditions. *Agric. Water Management* **142**: 127-134
27. Richardson, W. C., Badrakh, T., Roundy, B. A., Aanderud, Z. T., Petersen, S.L., Allen, P. S., Whitaker, D. R. and Madsen, M. D. 2019. Influence of an Abscisic Acid (ABA) Seed Coating on Seed Germination Rate and Timing of Bluebunch Wheatgrass. *Ecol. Evol.*, **9(13)**: 7438-7447.
28. Schubert, A. M., Benedict, C. R. and Kohel, R. J. 1986. Carbohydrate distribution in bolls. In: "*Cotton Physiology*", (Eds.): Mauney, J. R. and Stewart, M. C. D. The Cotton Foundation, Memphis, TN, PP. 311-324.
29. Sun, C., Ali, K., Yan, K., Fiaz, S., Dormatey, R., Bi. Z. and Bai. J. 2021. Exploration of Epigenetics for Improvement of Drought and Other Stress Resistance in Crops: A Review. *Plants*, **10(6)**: 1226.
30. Szollosi, R. 2014. Superoxide dismutase (SOD) and abiotic stress tolerance in plants: an overview. Chapter 3. In: "*Oxidative Damage to Plants*", (Ed.): Ahmad, A., Academic Press, New York, NY, USA, PP. 89-129.
31. Tombesi, S., Frioni, T., Poni, S. and Palliotti, A. 2018. Effect of Water Stress "Memory" on Plant Behavior during Subsequent Drought Stress. *Environ. Exp. Bot.*, **150**: 106-114.
32. Vishwakarma, K., Upadhyay, N., Kumar, N., Yadav, G., Singh, J., Mishra, RK., Kumar, V., Verma, R., Upadhyay, RG., Pandey, M. and Sharma, S. 2017. Abscisic Acid Signaling and Abiotic Stress Tolerance in Plants: A Review on Current Knowledge and Future Prospects. *Front. Plant Sci.*, **8**: 161.
33. Wahid, A., Perveen, M., Gelani, S. and Basra, S. M. A. 2007. Pretreatment of Seed with H<sub>2</sub>O<sub>2</sub> Improves Salt Tolerance of Wheat Seedlings by Alleviation of Oxidative Damage and Expression of Stress Proteins. *J. Plant Physiol.*, **164**: 283-294.
34. Wang, L. R., Yang, X. N., Gao, Y. S., Zhang, X. Y., Hu, W., Zhou, Z. and Meng, Y. L. 2019. Investigating Seed Dormancy in Cotton (*Gossypium hirsutum* L.): Understanding the Physiological Changes in Embryo during after-Ripening and Germination. *Plant Biol.*, **21(5)**: 911-919.
35. Wojtyla, Ł., Paluch-Lubawa, E., Sobieszczuk-Nowicka, E. and Garnczarska, M. 2020. Drought Stress Memory and Subsequent Drought Stress Tolerance in Plants. In: "*Priming-Mediated Stress and Cross-Stress Tolerance in Crop Plants*", (Eds.): Hossain, M. A., Liu, F., Burritt, D., Fujita, M., Huang, B. Academic Press, Elsevier, Amsterdam, PP. 115-131.
36. Yao, F., Zhang, X., Ye, X., Li, J., Long, L. and Yu, C. 2019. Characterization of Molecular Diversity and Genome-Wide Association Study of Stripe Rust Resistance at the Adult Plant Stage in Northern

- Chinese Wheat Landraces. *BMC Genet.*, **20**: 38.
37. Zare M., Mohammadifard G. R., Bazrafshan F. and Zadehbagheri M. 2014. Evaluation of Cotton (*Gossypium hirsutum* L.) Genotypes to Drought Stress. *Int. J. Biosci. (IJB)*, **4(12)**: 158-166.
38. Zhang, H. J., Zhang, N., Yang, R. C., Wang, L., Sun, Q. Q., Li, D. B., Cao, Y. Y., Weeda, S., Zhao, B., Ren, S. X. and Guo, Y. D. 2014. Melatonin Promotes Seed Germination under High Salinity by Regulating Antioxidant Systems, ABA and GA4 Interaction in Cucumber (*Cucumis sativus* L.). *J. Pineal Res.*, **57**: 269-279.

## ارزیابی حافظه تنش آبی در پاسخ جبرانی گیاه پنبه (*Gossypium hirsutum* L.) در کم آبی

م. حبیبی، ا. فغانی، م. رضایی، م. ع. محمودجانلو، و م. ح. رزاقی

### چکیده

این تحقیق به منظور تامین بذر پنبه مناسب برای تکثیر بذر در مزارع دیم انجام شد، در این مطالعه، توانایی بذرهای پنبه در مواجهه با تنش به مدت سه سال متوالی جهت ارزیابی پاسخ های حافظه تنشی بررسی شدند. آزمایش در طرح اسپلیت پلات فاکتوریل با چهار سطح آبیاری،  $W_0$  (بدون آبیاری)،  $W_1$  (۳۳ درصد ظرفیت زراعی)،  $W_2$  (۶۶ درصد ظرفیت زراعی) و  $W_3$  (۱۰۰ درصد ظرفیت زراعی)، به عنوان فاکتور اصلی و پنج تیمار بذری (چهار تیمار بذر سه بار تنش دیده،  $S_{21}$  تا  $S_{24}$  و بذر گواهی شده) به عنوان فاکتور فرعی که در معرض سطوح مختلف تنش آبی برای سه سال زراعی پرورش داده شدند. نتایج نشان داد که،  $S_{32}$  پیام تنش آبی را در هر دو شرایط  $W_0$  و  $W_3$  از طریق تغییر مکانیسم های فیزیولوژیکی دریافت نمود. بذرهای  $S_{32}$ ، کمترین مقدار اسیدآبسیزیک و بیشترین مقدار کلسیم را در شرایط  $W_0$  و  $W_3$  انباشته کردند. افزایش فعالیت سوپراکسید دیسموتاز و آسپاراتات پراکسیداز در برگ های  $S_{32}$ ، در معرض  $W_0$  و  $W_3$  مکانیسم دیگری از حافظه تنشی برای خوگیری به کم آبی بود. بیشترین وزن ۳۰ غوزه، وزن الیاف ۳۰ غوزه و عملکرد چین اول از  $S_{32}$  در معرض تیمارهای  $W_0$ ،  $W_1$  و  $W_2$  بدست آمد. همچنین بذرهای  $S_{32}$  که در معرض  $W_0$ ،  $W_1$  و  $W_2$  قرار گرفتند، بیشترین بنیه بذر و درصد جوانه زنی را داشتند. می توان نتیجه گرفت که حافظه تنشی از طریق تغییر فیزیولوژی و مورفولوژی رفتار گیاه می تواند در تحمل به کم آبی به گیاهانی که به طور متوالی در معرض خشکی قرار دارند، کمک نماید.