

Allelopathic Potentials of Eight Barley Cultivars on *Brassica juncea* (L) Czern. and *Setaria viridis* (L) p. Beauv.

J. Asghari^{1*} and J. P. Tewari²

ABSTRACT

The allelopathic potential of the foliage tissues of eight barley cultivars was investigated using seedlings bioassays of Indian brown mustard (*Brassica juncea*) and green foxtail (*Setaria viridis*) as typical broadleaf and grass weeds of the croplands. The correlations of allelochemical compounds with seed germination of weeds were determined. The barley cultivars used were Jackson, Bronco, CDC Dolly, B1602, Harrington, UNA80, Stander, and TR251. Three dilutions of the extracts of each barley cultivar (20, 10 and 5 g/L) plus deionized distilled water, as a control, were assayed on the target weeds. The effects of barley cultivars and extract concentrations on seed germination and radicle and shoot lengths of the target weed seeds were highly significant ($P < 0.001$). By increasing the concentration of the allelopathic extracts, the percent germination, radicle and shoot lengths of both target plants seedlings decreased. Seed germination and seedling growth of *B. juncea* were more sensitive than those of *S. viridis*. Relative to the germination inhibition of target weed seeds in comparison to the untreated control, the selected barley cultivars were categorized as being highly allelopathic, moderately allelopathic and of low allelopathic potential. Among the phenolic compounds detected, the cumulative effect of four small-quantity-compounds (vanillic, chlorogenic, p-coumaric, and ferulic acids) on average weed germination was very high ($R^2 = 0.83$), with *B. juncea* it was high ($R^2 = 0.70$) and with *S. viridis* it was very low ($R^2 = 0.15$). The two high concentration phenolic compounds of protocatechuic and p-hydroxybenzoic acid, when accumulated with the small-quantity phenolic compounds of barley cultivars, reduced the weed seeds germination correlation to $r = -0.11$.

Keywords: Allelopathy, Aqueous extracts, Barley, *Brassica juncea*, HPLC, Phenolic acids, *Setaria viridis*.

INTRODUCTION

Allelopathy is a mechanism of plant interference in agroecosystems that offers an opportunity to manage weeds in crop sequence but could also adversely affect crop yields and influence the choice of rotation. The allelopathic potential of many crop plants has been investigated and approved (Burgos *et al.*, 1999; Baghestani, *et al.*, 1999; Wu *et al.*, 2001). Heavy use of herbicides in most integrated weed management (IWM) sys-

tems is a major concern since it causes serious threats to the environment, public health and increases costs of crop production. The degree of weed seed germination inhibition and growth suppression which can be attributed to crop allelopathy is highly important and worth while. This can be considered as a possible alternative weed management strategy (Macias, 1995).

Barley (*Hordeum vulgare* L.) is a smother crop and possesses the allelopathic potential to suppress some weeds (Liu and Lovett,

1. Department of Agronomy and Plant Breeding, Faculty of Agricultural Sciences, Guilan University, P. O. Box 41635-1314, Rasht, Islamic Republic of Iran.

2. Department of Agricultural, Food and Nutritional Science, University of Alberta, Edmonton, Alberta T6G 2P5, Canada.

* Corresponding author, e-mail: jafarasghari@yahoo.com



1993a; Overland, 1966). In a modified bioassay system with reduced environmental influences, Liu and Lovett (1993b) found that root of barley released allelopathic chemicals which delayed germination and inhibited growth of white mustard (*Sinapis alba* L.). In further studies using HPLC they found that hordenine, as a major component of barley root which released up to a maximum of 2 µg/plant/day for up to 60 days in a hydroponics system, is responsible for the growth retardation. The mustard radicle tips exposed to hordenine and gramine showed cell wall damage, an increase in both size and number of vacuoles, autophagy, and disorganization of organelles (Liu and Lovett, 1993b). A linear relationship was found between peak area and concentration of authentic standards of both hordenine and gramine using the HPLC method for barley seedlings (Hoult and Lovett, 1993). Baghestani *et al.* (1999) found that various concentrations of spring cereal cultivar root extracts inhibited *Brassica kaber* hypocotyls and root growth with no negative effects on germination. As cereal root exudate concentrations increased, *B. kaber* growth decreased. Benzoic, caffeic, o-coumaric, and vanillic acids, scopoletin and parahydroxybenzoic acid were all found in barley and other examined cereal seedlings using HPLC. Smith and Martin (1994) have found that the extract of leaf and stem tissue of Italian ryegrass (*Lolium multiflorum*), little barley (*Hordeum pusillum*) and tall fescue (*Festuca arundinacea*) suppressed seed germination and seedling growth of alfalfa (*Medicago sativa*). They have estimated that concentrations of about 5.0 g/L of aqueous extracts from foliage tissues of little barley reduced seed germination and seedling growth in alfalfa and Italian ryegrass by 50%. Production of allelopathic compounds varies within the same species (Barnes *et al.*, 1987). Understanding the allelopathic potential of crop cultivars is very important for weed management, use as cover crops in conservation tillage, breeding or engineering cultivars to enhance allelopathic production (Burgos *et al.*, 1999).

The objectives of the present experiments were to find the potential allelopathic property of eight barley cultivar aqueous extracts on the germination and seedling growth of isolates and quantify the phenolic compounds of barley cultivars to determine whether or not these allelochemicals correlated with the target weed's seed germination.

MATERIALS AND METHODS

Seedbed Preparation and Barley Cultivation

To provide an adequate seedbed for barley growth under greenhouse conditions, the silty loam provided was sterilized in an oven using 2.1 kg cm⁻² pressure under 132 °C for 1 hour, then mixed with pre-sterilized Mito-Mix 290 (a horticultural product) using machine mixture. Thirty-two 25×50×7 cm flat plastic culture trays were filled each with five kg of the prepared soil, then mixed with 30 gr of slow release 14:14:14 NPK fertilizer. The trays were placed in a greenhouse with 14 hours light per day, and a 22/16 °C day/night temperature regime. The trays were irrigated to field capacity and the soil pH was determined (pH=6.94). The seedbed was corrugated longitudinally in five rows, 4 cm apart. One hundred uniform seeds of each barley cultivar were sown in each tray (20 seeds per row). The trays were sprayed with 0.1 % of No-dump (Oxine benzoate 2.5%) fungicide to control soil-borne diseases.

Eight barley cultivar seeds were obtained from Alberta Agricultural Research Station (Canada) including Jackson, Bronco, CDC Dolly, B1602, Harrington, UNA80, Stander, and TR251. The trays were replicated four times. The trays were irrigated every other day and allowed to germinate and grow for 30 days. The foliar section of each plant was cut from the base and stored in labeled paper bags. The samples were freeze dried for three days, and ground with a Wiley mill to pass through a 40-50 mesh screen. The

ground tissues were used for allelochemical extractions.

Aqueous Extracts Preparation

Ten grams of each cultivar ground tissue were placed in a 1-L Erlenmeyer flask, and 500 ml deionized distilled water (DDI) was added to it. The flasks were covered with aluminum foil to protect them from photo-decomposition, then placed on a rotary shaker (≈ 250 revolutions per min.) at laboratory temperature ($\approx 22^\circ\text{C}$) for 10 hours. The mixtures were filtered through 4 layers of cheesecloth and two layers of Whatman No.1 using a vacuum pump. The pH and electrical conductivity of the extracts were determined using a digital pH meter and a conductivity meter. These filtrates were considered as stock solutions. A series of solutions including the stock solution (extract of 20 g dry weight per liter of water; (S_1), and concentration dilutions of 10 (S_2), and 5 g/L (S_3), were developed from the stock solutions. The extra stock solutions were kept in -20°C for later use.

Petri Dish Bioassays

The barley cultivar young plant aqueous extract bioassay was conducted on seed germination, growth of radicle and shoot appearance of *B. juncea* (Indian brown mustard) and *S. viridis* (green foxtail) as typical broadleaf and grass cropland weeds. The seeds were pre-sterilized with 1% sodium hypochlorite for 5 minutes and washed with distilled water. Three dilute series (S_1 , S_2 and S_3) of each cultivar's aqueous extract plus the control (DDI water) were used for germination tests. Fifteen *B. juncea* or *S. viridis* seeds (pre-tested seeds with $>97\%$ germinabilities) were evenly distributed on two layers of Whatman No.1 filter papers in each 9-cm disposable sterile Petri dish. Five ml of a dilute series was added to each Petri dish, covered with a lid, sealed with parafilm, then incubated at 22°C for five

days. The bioassay was replicated four times and the experiment was repeated once.

Seeds were considered germinated if the radicle had emerged 2 mm from the seed coat (An *et al.*, 1997). The number of germination and the length of the radicle and shoot of the germinated weed seedlings were compared with the control and expressed as a percentage of the control. In total, three factors including 1) Aqueous extracts of eight barley cultivars plus control, 2) Extract concentrations in three levels, and 3) The two target weed seeds (*B. juncea* and *S. viridis*), as bioassay plants, were used on a randomized complete block design with four replications.

The data were subjected to analysis of variance, and highly significant or significant differences were tested at least with 1% or 5% levels using LSMEAN, i.e. least significant mean difference comparisons. Regression between measured parameters was made if needed.

HPLC Sample Preparation

Hot water extraction was performed according to the Kajimoto *et al.*, (1999) method. Samples of barley cultivar aqueous extract stock solutions (S_1) were prepared by heating 0.10 g of each in 2.0 mL of water for one hour at 100°C . The samples were then injected onto the HPLC. The HPLC was a Varian 5000 (Varian, Mississauga, ON, Canada) equipped with a Shimadzu SIL-9A autosampler (Shimadzu Corporation, Columbia, MD) and a WATERS 486 UV detector (Waters, Milford, MA) at 280 nm. Separation was performed with a Supelcosil LC-18, $5\ \mu\text{m}$, $4.6\ \text{mm} \times 15\ \text{cm}$ (Supelco, Oakville, ON, Canada) column. The data were integrated and analyzed using Shimadzu CLASS-VP Chromatography Laboratory automated Software System (Shimadzu Corporation, Columbia, MD). The mobile phase utilized a gradient composed of a 0.01 M sodium citrate buffer (A) pH=5.4 adjusted with 50% acetic acid, and methanol (B). The best separation was obtained using the fol-



lowing gradient: 0 min, 2% B; 12 min, 4% B; 20 min, 13% B; 22 min, 13% B and 26 min, 2% B. The running time was 30 minutes. The solvent flow rate was 1.0 ml/min and separation was performed at room temperature. Each phenolic acid standard (protocatechuic, p-hydroxybenzoic, vanillic, syringic, caffeic, chlorogenic, p-coumaric, and ferulic acids) displayed a linear response ($R^2=0.95$) over 8-40 μg calibration series. The concentrations of phenolic acids in samples were calculated using syringic acid as an internal standard. Results were expressed in $\mu\text{g/g}$ dry matter of barley cultivar foliar parts.

tion were highly significant ($P<0.001$), but the three variables interaction (TWS * Cs * BC) on seed germination were also significant ($P=0.05$). The radicle lengths of target weed seedlings did not show significant differences ($P>0.05$) but the shoot did ($P=0.027$). Effects of other sources of variation on radicle and shoot lengths were the same as germination, except that the interaction of target weed seedlings with barley cultivars on radicle was not significant.

The barley cultivar aqueous extracts pH did not have a meaningful correlation with weed seed germination inhibition (Figure 1A; $R^2=0.057$). This indicates that the inhibitory property of the extracts is independ-

Table 1. Probability levels (P) of percentage germination, radicle length and shoot length of *B. juncea* and *S. viridis* seeds, treated with three concentrations of barley cultivar dry powder aqueous extracts in a Petri dish.

Source of variation	Germination		Radicle length		Shoot length	
	F value	P>F	F value	P>F	F value	P>F
Replication	1.46	0.3817	1.87	0.3105	0.88	0.5404
Target weed seeds (TWS)	376.41	0.0003	1.52	0.3054	16.44	0.0270
Rep. * TWS (Error)	-	-	-	-	-	-
Concentrations (Cs)	522.38	0.0001	504.14	0.0001	401.80	0.0001
Barley Cultivars (BC)	52.36	0.0001	118.48	0.0001	22.85	0.0001
TWS * Cs	143.36	0.0003	14.36	0.0001	7.03	0.0012
TWS * BC	5.43	0.0001	2.26	.0262	3.06	0.0031
Cs * BC	16.77	0.0001	8.46	0.0001	8.77	0.0001
TWS * Cs * BC	1.71	0.0500	3.17	0.0001	1.84	0.0300

RESULTS AND DISCUSSION

The effects of three concentrations of eight barley cultivar leaf dry powder aqueous extracts on germination, radicle and shoot lengths of two target plant species (*B. juncea* and *S. viridis*) are shown in Table 1. According to both analytical Models I and III, the effects of target weed seeds (TWS), extract concentrations (Cs) and barley cultivars (BC) on seed germinations were highly significant ($P<0.01$). Not only, the bidirectional interactions of all the variables (TWS * Cs, TWS * BC and Cs * BC) on seed germina-

ent from the pH. The correlation of aqueous extract pH and their allelopathic potential is occasionally a problem. This result is in agreement with the findings of Quayyum *et al.* (1999) and in disagreement with An *et al.* (1997). This may explain the observations that the pH range of plant aqueous extract is within the tolerance range of target seeds, even if the seeds show some inhibition (Mason-Sedun *et al.*, 1986).

The correlation between the electrical conductivity of barley cultivar aqueous extracts and seed germination was very weak (figure 1B; $R^2=0.10$). This suggests that the osmotic effects of extracts on germination is not important for observed germination inhibition.




Figure 1. Dynamics of (A) pH and (B) electrical conductivity of barley cultivar aqueous extracts on percentage germination inhibition of *B. juncea* and *S. viridis* seeds.

The negligibility of osmotic effects of aqueous extracts is supported by several investigators (An *et al.*, 1997; Mason-Sedum, 1986).

There was an indirect relationship between the concentrations of barley cultivar aqueous extracts with percentage germination, radicle, and shoot lengths of target weeds' seedlings (Table 2).

As the concentrations increased, the percentage germination, radicle and shoot lengths of the target weed seedlings significantly decreased and reached the lowest percentage compared with the control. This is in agreement with the findings of other researchers (Chaves *et al.*, 2001; An *et al.*, 2001; Wu *et al.*, 2001).

The percentage germination and shoot length of *S. viridis* seedlings were signifi-

cantly higher than the *B. juncea*, but their radicle lengths were not different (Figure 2). This indicated that barley cultivar allelochemicals were more inhibitorier on *B. juncea* germination and seedling growth than those of *S. viridis*. The sensitivity of plants to allelopathic compounds differs among species and genotypes within a species. Quayyum *et al.* (1999) have shown that lettuce seedlings were more sensitive to the aqueous extracts than wild rice. Overland (1966) has shown that the inhibitory activity of barley cultivars was selective and *Stellaria media* was more sensitive than *Cap-sula bursa-pastoris*. The importance of using adequate target weed species as a bioassay plant has been emphasized by Inderjit and Dukshini (1995).

Table 2. Effects of three levels of concentrations of barley cultivars allelochemicals on germination, root length, and shoot length of *B. juncea* and *S. viridis* seeds in Petri dishes. (Data are based on percentage of control and mean of barley cultivar aqueous extracts).

Barley extract Concentration	Germination	Radicle length	Shoot length	Germination	Radicle length	Shoot length
	<i>Setaria viridis</i>			<i>Brassica juncea</i>		
(mg ml ⁻¹)	Percent of control			Percent of control		
5	95.13 a ^a	60.36 a	101.72 a	72.93 a	73.92 a	99.67 a
10	84.02 b	22.71 b	54.94 b	24.85 b	17.05 b	33.32 b
20	10.92 c	11.24 c	11.06 c	15.18 c	8.84 c	11.20 c

^a Within columns, means followed by the same letter are not significantly different at the 0.05 levels as determined by least significant difference.




Figure 2. Comparison of *B. juncea* and *S. viridis* weeds, seed germination, radicle and shoot growth of plants treated with barley cultivars aqueous extracts.

The degree of influence of barley cultivars aqueous extracts on the seed germination of target weeds can be simplified in to three categories: 1) Cultivars with the highest allelopathic potential - Stander, CDC Dolly, 2) Cultivars with a moderate allelopathic potential - Harrington, UNA80, TR251, and 3) Cultivars with the lowest allelopathic potential - Bronco, B1602, Jackson (Figure 3). Extracts of barley cultivars reduced radicle length, seed germination and shoot length of target weeds in 76, 57 and 50% of non-treated controls, respectively (Figure 4). In other words, the seedling growth of the target weeds was more suppressed than the germination. Smith (1991), Smith and Martin (1994) and Ben-Hammouda *et al.* (1995)

found aqueous leaf tissue extracts of several species have suppressed seedling growth in target plants more than seed germination. Indeed, the radicle growth was more sensitive to allelochemicals than the coleoptile growth. This is in agreement with Ahn and Chung (2000) who found that the length and dry weight of roots of *Echinochloa crusgalli* were more affected by hull extract than the shoots. An *et al.* (2001), on evaluation of *Vulpia* (*Vulpia myuros*) allelochemicals, also found all phenolic compounds caused greater inhibition on root elongation than on shoot length.

The interaction of barley cultivars with the target weeds' seed germination, radicle and shoot length were highly significant (Table




Figure 3. Effect of barley cultivars, aqueous extract allelochemicals on germination of *B. juncea* and *S. viridis* seeds in Petri dishes (data mean of weed seeds and concentrations).

Table 3. Effect of various barley cultivar aqueous extract allelochemicals on *B. juncea* and *S. viridis* seed germination, radicle, and shoot length in a petri dish (data mean of concentrations).

Barley cultivars	<i>Setaria viridis</i>			<i>Brassica juncea</i>		
	Germination	Radicle length	Shoot length	Germination	Radicle length	Shoot length
	Percent of control			Percent of control		
Jackson	60.23 a	23.96 a	50.56 a	31.68 a	17.01 d	44.53 b
Bronco	55.56 a	20.02 a	44.81 a	36.63 a	36.02 a	64.38 a
CDC Dolly	54.97 a	19.63 a	41.44 a	22.77 bc	21.62 bcd	35.92 b
B1602	63.74 a	24.34 a	53.82 a	28.71 ab	27.07 abc	49.62 b
Harrington	57.89 a	24.70 a	51.53 a	25.74 b	30.68 ab	44.07 b
UNA80	61.40 a	19.50 a	51.43 a	22.77 bc	20.29 cd	30.55 b
Stander	54.73 a	25.92 a	56.34 a	16.83 c	15.84 d	29.17 b
TR251	59.65 a	24.84 a	53.24 a	23.76 bc	30.60 ab	34.29 b


3). The aqueous extract of cultivar Stander by reducing seed germination of *B. juncea* to 16.83%, was the most influential inhibitor and the extract of cultivar Bronco at 36.63% was the least influential on seed germination.

Significant differences were not observed in the shoot length of *B. juncea* treated with aqueous extract of various barley cultivars. The Stander extract by reducing the radicle length of *B. juncea* to 15.84% of the control was the most effective, and the extract of Bronco by reducing the radicle length of *B. juncea* to 36.02% was the least. In contrast to *B. juncea*, the germination, radicle and shoot length of *S. viridis* treated with aqueous extracts of barley cultivars were not significant. In other words, the phytotoxicity of barley cultivar aqueous extract on *S. viridis* as a grass weed is much lower than that of *B.*

juncea. Similarly, Anaya *et al.* (1999), on the allelochemicals bioassay of *Metopium brownie*, have shown that the radicle growth inhibitor of *Amaranthus hypochondriacus* as a broad leaf weed was much more severe than that of *Echinochloa crusgalli*, a grass weed.

Compared with the non-treated control, the percentage germination of target plant seeds decreased as concentrations of barley cultivars extracts increased (Figure 5). The germination suppression decreased from Stander, CDC Dolly, Harrington, UNA80, TR251, Bronco, B1602, to Jackson, respectively. By decreasing the ratio of extraction from 20 g/L to 5 g/L, the range of germination variation increased and the ability of individual cultivars' allelochemical potential in germination suppression became clearer. Ahn and Chung (2000) have found an inverse relationship between rice hull water extract concentration and target weed seed germination. This finding also agrees with the work of Chaves *et al.* (2001) who have found that by increasing the concentration of *Cistus ladanifer* leaf aqueous extracts, the percentage germination, root and cotyledon length of *Rumex crispus* decreased.

Six phenolic compounds (protocatechuic, p-hydroxybenzoic, vanillic, chlorogenic, p-coumaric, and ferulic acids) were obtained from each barley cultivar foliar aqueous extract. None of the cultivars contained syringic, caffeic, or chlorogenic acid when

**Figure 4.** Effect of barley cultivars, aqueous extract allelochemicals on the average germination, root length and shoot length of *B. juncea* and *S. viridis* seeds in comparison to a non-treated control.




Figure 5. Effects of various concentrations of barley cultivar aqueous extracts on germination, of *B. juncea* and *S. viridis* seeds in a Petri dish. (data mean of weed seeds).

compared with the standard (Table 4). The concentrations of each phenolic acid differed among cultivars. The content of protocatechuic acid was the highest and chlorogenic acid was the lowest among the detected allelochemicals. The correlation coefficient ($r = -0.11$) of total concentration of phenolic compounds with percentage germination inhibition was very low (Figure 6). This indicates that none of the phenolic compounds affect negatively or equally on germination. Both the inhibitory and stimulatory responses of the aqueous extracts are significant for assessing allelopathic properties (Rice, 1995).

In contrast to total phenolic compounds, there was a high correlation between the percentage germination reduction of *B. juncea* with the accumulation of four small-quantity-phenolic-compounds (vanillic, chlorogenic, p-coumaric, and ferulic acids) of barley aqueous extracts ($R^2 = 0.70$) as shown in Figure 7. These results indicate that allelochemicals present in large quantities possessed low activity, while those present in small quantities possessed a strong inhibitory activity. An *et al.* (2001) on the basis of evaluation of the biological activity of identified allelochemical from *Vulpia myuros* have found that individual compounds

Table 4. Barley cultivar aqueous extract allelochemicals (Protocatechuic, p-Hydroxybenzoic, vanillic, chlorogenic, p-coumaric and ferulic acid) isolated using HPLC(Micro gr/L).

Barley cultivar	Protocatechuic acid	p-hydroxy benzoic	Vanillic acid	Chlorogenic acid	p-coumaric acid	Ferulic acid
Jackson	9221.91	606.02	36.06	82.15	366.38	216.44
Bronco	2287.74	593.45	43.55	26.73	280.51	131.85
CDC Dolly	9752.91	701.60	460.84	29.64	414.75	120.27
B1602	6982.5	652.90	108.66	66.28	437.18	99.69
Harrington	11261.87	510.03	367.11	33.69	325.85	168.99
UNA80	11268.35	767.92	144.18	31.65	391.39	187.88
Stander	14466.18	680.58	472.18	29.17	368.82	201.68
TR251	16127.75	653.86	414.17	28.08	334.04	17.59




Figure 6. Effects of total concentrations of phenolic compounds (protocatechuic, p-hydroxybenzoic, vanillic, chlorogenic, p-comaric and ferulic acid) of barley cultivar aqueous extracts on germination inhibition of *B. juncea* and *S. viridis* seeds.

were not equally inhibitory to tested plants; allelochemicals present in large quantities possessed low activity, while those present in small quantities possessed a strong inhibitory activity. They have concluded that the exploration of the relative composition of a cluster of allelochemicals is more important than simply focusing on the identification of one or two compounds with strong biological activity. In contrast, the small-quantity-phenolic-compounds showed lower allelopathic effects on *S. viridis* germination ($R^2 = 0.15$), while the average two weeds germination correlation was also very high ($R^2 = 0.83$). The low effect of barley cultivar

aqueous extract on seed germination of *S. viridis* indicates the weak influence of barley phenolic compounds on grass weeds.

This research has shown that barley cultivar shoot tissue aqueous extract for a specific weed is not constant, but is related to the target species, aqueous concentrations, and the proposed cultivars. Broadleaf weeds are more sensitive than grass weeds to barley extracts. The allelochemicals present in shoot tissues may greatly contribute to plant residue in the soil. The residue will interfere with the growth of weeds in that soil and will also affect the germination and vigor of weeds in the vicinity of the crop. Under-




Figure 7. Effects of total low concentrations of barley cultivars' aqueous extract phenolic compounds (vanillic, chlorogenic, p-comaric and ferulic acid) on germination of *S. viridis*, *B. juncea* and the average of the target weed seeds.



standing the allelopathic potential of barley cultivars will help weed control management of the croplands by lowering the production costs due to herbicide application and hand-weeding. Another use of allelopathic cultivars is incorporating the genes carrying the allelopathic potential into varieties with high quantity and quality crops, which will benefit farmers and consumers as well as the environment.

ACKNOWLEDGEMENTS

The authors would like to thank the University of Guilan (Iran) and the University of Alberta (Canada) for their financial support and use of their laboratory facilities in the conduct of this study. The authors are also grateful to Mr. Gary Sedgwick of the Department of Agricultural, Food and Nutritional Sciences, University of Alberta for his kind assistance in conducting HPLC.

REFERENCES

1. Ahn, J. K. and Chung, I. M. 2000. Allelopathic Potential of Rice Hulls on Germination and Seedlings Growth of Barnyardgrass. *Agron. J.*, **92**: 1162-1167.
2. An, M., Pratley, J. E., and Haig, T. 1997. Phytotoxicity of *Vulpia* Residues: I. Investigation of Aqueous Extracts. *J. Chem. Ecol.* **23**: 1979-1994.
3. An, M., Pratley, J. E., and Haig, T. 2001. Phytotoxicity of *Vulpia* Residues: III. Biological Activity of Identified Allelochemicals from *Vulpia myuros*. *J. Chem. Ecol.* **27**: 383-394.
4. Anaya, A. L., Calera, M. R. Mata, R., and Mirandaf, R. P. 1990. Allelopathic Potential of Compounds Isolated from *Ipooea tricolor* Cav. *J. Chem. Ecol.* **16**: 2145-2152.
5. Anaya, A. L., Mata, R. Rivero-Cruz, F. Chavez-Velasco, D., and Gomez-Pompa, A. 1999. Allelochemical Potential of *Metopium brownie*. *J. Chem. Ecol.* **25**: 141-156.
6. Baghestani, A., Lemieux, C. Leroux, G. D. Baziramakenga, R. and Simard, R. R. 1999. Determination of Allelochemicals in Spring Cereal Cultivars of Different Competitiveness. *Weed Sci.* **47**: 498-504.
7. Barnes, J. P., Putnam, A. R. Burke, B. A. and Aasen, A. J. 1987. Isolation and Characterization of Allelochemicals in Rye Herbage. *Phytochem.* **26**: 1385-1390.
8. Ben-Hammouda, M., Kremer, R. J. Minor, H. C., and Sarwar, M. 1995. A Chemical Basis for Differential Allelopathic Potential of Sorghum Hybrids on Wheat. *J. Chem. Ecol.* **21**: 775-786.
9. Burgos, N. R., Talbert, R. E., and Mattice, J. D. 1999. Cultivar and Age Differences in the Production of Allelochemicals by *Secale cereale*. *Weed Sci.* **47**: 481-485.
10. Chaves, N., Sosa, T. Alias, J. C. and Escudero, J. C. 2001. Identification and Effects of Interaction Phytotoxic Compounds from Exudates of *Cistus ladanifer* Leaves. *J. Chem. Ecol.* **27**: 611-621.
11. Hoult, A. H. C., and Lovett, J. V. 1993. Biologically Active Secondary Metabolites of Barley. III. A Method for Identification and Quantification of Hordenine and Gramine in Barley by High-performance Liquid Chromatography. *J. Chem. Ecol.* **19**: 2245-2254.
12. Inderjit, and Dakshini, K. I. M. 1995. On laboratory Bioassay in Allelopathy. *Bot /Rev.* **61**: 29-44.
13. Kajimoto, G., Onitake, N. Okuda, N. and Murakami, C. 1999. Antioxidant Activity of Barley Tea and their Composition. *J. Jpn. Soc. Food Sci. Technol.* **46**(2): 67-74.
14. Liu, D. L., and Lovett, J. V. 1993a. Biologically Active Secondary Metabolites of Barley. I. Developing Techniques and Assessing Allelopathy in Barley. *J. Chem. Ecol.* **19**: 2217-2230.
15. Liu, D. L., and Lovett, J. V. 1993b. Biologically Active Secondary Metabolites of Barley. II. Phytotoxicity of Barley Allelochemicals. *J. Chem. Ecol.* **19**: 2231-2244.
16. Macias, F. A. 1995. Allelopathy in the Search for Natural Herbicide Models, pp. 310-329, In: "Allelopathy: Organisms, Processes, and Applications", (Eds.) Inderjit, K. M. M. Dakshini, and Einhellig, F. A. ACS Symposium Series 582. American Chemical Society, Washington, D.C.
17. Mason-Sedun, W., Jessop, R. S., and Lovett, J. V. 1986. Different Phytotoxicity among Species and Cultivars of the Genus Brassica to Wheat. I. Laboratory and Field Screening of Species. *Plant Soil.*, **93**: 3-16.
18. Overland, L. 1966. The Role of Allelopathic Substances in the "Smother Crop" Barley. *Am. J. Bot.* **53**: 423-432.

15. Quayyum H. A., Mallik, A. U. and Lee, P. F. 1999. Allelopathic Potential of Aquatic Plants Associated with Rice (*Zizania palustris*): Bioassay with Plant and Lake Sediment Samples. *J. Chem. Ecol.* **25**: 209-219.
16. Rice, E. L. 1995. Biological Control of Weeds and Plant Diseases: Advances in Applied Allelopathy. University of Oklahoma Press, Oklahoma.
17. Smith, A. E. 1991. The Potential Importance of Allelopathy in the Pasture Ecosystem; A Review. *Adv. Agron.* **1**: 27-37.
18. Smith, A. E. and L. D. Martin. 1994. Allelopathic Characteristics of Three Cool-season Grass Species in the Forage ecosystem. *Agron. J.* **86**: 243-246.
19. Wu, H., Haig, T. Pratley, J. Lemerle, D. and M. An. 2001. Allelochemicals in Wheat (*Triticum aestivum* L.): Variation of Phenolic Acids in Shoot Tissues. *J. Chem. Ecol.* **27**: 125-135.

توان دگرآسیبی ۸ رقم جو بر روی علفهای هرز دم روباهی سبز و خردل قهوه ای

ج. اصغری و ج. پ. تواری

چکیده

برای تعیین توان دگرآسیبی ارقام جو بر روی علفهای هرز، مطالعات زیست سنجی عصاره بافتهای هوایی ۸ رقم جو بر روی علف هرز پهن برگ (*Brassica juncea*) و باریک برگ دم روباهی (*Setaria viridis*) انجام و همبستگی ترکیبات دگرآسیب این ارقام با جوانه زنی علفهای هرز مذکور تعیین شد. بذور ارقام جکسون، برانکو، سی دی سی دالی، ب ۱۶۰۲، هرینگتون، یونا ۸۰، استاندارد، و تی آر ۲۵۱ جو برای مدت یکماه در محیط گلخانه کشت گردید، سپس بافتهای هوایی آنها پس از یخ خشکانی، پودر کرده و عصاره آبی هر یک از ارقام تهیه شد. سه غلظت از هر رقم جو (۵، ۱۰ و ۲۰ گرم پودر خشک جو در لیتر آب مقطر) همراه با آب مقطر (شاهد) بر روی جوانه زنی بذر علفهای هرز مذکور مطالعه شد. تأثیر عصاره ارقام جو و غلظتهای بکار رفته بر جوانه زنی و طول ریشه چه و ساقه چه بذور علفهای هرز بسیار معنی دار بود و با افزایش غلظت عصاره دگرآسیب درصد جوانه زنی، طول ریشه چه و ساقه چه هر دو گیاه کاهش یافت. جوانه زنی و رشد جوانه های خردل قهوه ای حساسیت بیشتری نسبت به دم روباهی سبز از خود نشان داد. با توجه به نسبت تأثیر عصاره ها در مقایسه با شاهد آب مقطر در منع جوانه زنی علفهای هرز مذکور، این ارقام به ارقام دگر آسیب شدید (استاندر و سی دی سی دالی با بیش از ۶۱٪ منع جوانه زنی)، دگر آسیب متوسط (هرینگتون، یونا ۸۰ و تی آر ۲۵۱ با ۵۸٪ منع جوانه زنی) و دگر آسیب ضعیف (جکسون، برانکو، و ب ۱۶۰۲ با کمتر از ۵۴٪) دسته بندی شدند. از میان ترکیبات فنولیک بدست آمده در مطالعه با HPLC، اثر چهار ترکیب فنولی با غلظت کم (اسیدهای وانیلیک، کروژنیک، پی کوماریک، و فرولیک) بر میانگین جوانه زنی علفهای هرز مذکور بسیار بالا ($R^2 = 0/83$) بود. در بررسی مستقل، میزان این اثر در منع جوانه زنی خردل قهوه ای نسبتاً بالا ($R^2 = 0/70$) ولی بر دم



روباهی سبز بسیار ناچیز بود ($R^2=0/15$). تجمع دو ترکیب دگرآسیب فنولی دارای غلظت زیاد (اسید پروتوکاتکوئیک و پی هیدروکسی بنزوئیک) با ترکیب‌های فنولی غلظت کم ارقام جو سبب ضریب همبستگی منفی در جوانه زنی بذور علفهای هرز مذکور شد ($r=0/11$).