

1 **ACCEPTED ARTICLE**

2 **Herbicidal and insecticidal activity of secondary metabolites from endophytic**
3 **and soil fungi**

4
5 S. A. M. Abdelgaleil^{1*}, Y. Shiono², N. E. M. Taktak³ and M. M. G. Saad¹

6 **Running title:** Bioactivity of fungal secondary metabolites

7 1. Department of Pesticide Chemistry and Technology, Faculty of Agriculture, El-Shatby,
8 Alexandria University, Alexandria 21545, Egypt

9 2. Department of Food, Life, and Environmental Science, Faculty of Agriculture, Yamagata
10 University, Tsuruoka, Yamagata, 997-8555, Japan

11 3. Department of Tropical Health, High Institute of Public Health, Alexandria University,
12 Alexandria, Egypt

13 ***Corresponding author; e-mail:** samirabdelgaleil@gmail.com

14
15 **ABSTRACT**

16 Fungi have been shown to be a good source for lead molecules in drug discovery and
17 development. However, many compounds derived from fungi were not evaluated for their
18 bioactivity against economic agricultural and public health pests. Twelve fungal secondary
19 metabolites (1-12) were evaluated for herbicidal activity against *Silybum marianum* and
20 insecticidal activity against *Culex pipiens* larvae. Among the tested metabolites, brefeldin A (6)
21 and 6-eopxy-4-hydroxy-3-methoxy-5-methyl-cyclohex-2-en-1-one (11) showed potent herbicidal
22 activity against *S. marianum* with complete inhibition of seed germination at 500 mg L⁻¹.
23 Compound 6 revealed an exceptional herbicidal activity as it caused complete inhibition of root
24 growth and strong reduction in shoot growth (I = 74.5%) and germination (10.0%) at 25 mg L⁻¹.
25 In addition, dehydroaustin (9), phomaxanthone A (4) and deacetylphomaxanthone A (5) displayed
26 a potent toxicity against fourth larval instar of *C. pipiens* with LC₅₀ values of 3.27, 57.03 and 63.50
27 mg L⁻¹, respectively. Based on the results of this study, compounds 4-6, 9 and 11 should be
28 developed as natural pesticides.

29 **Keywords:** Natural products, Fungal metabolites, Bioactivity, *Silybum marianum*, *Culex pipiens*.

30
31 **INTRODUCTION**

32 Fungi are a rich source of diverse natural compounds that have the potential for the discovery of
33 new drugs and agrochemicals (Porrás-Alfaro *et al.* 2011). It has been estimated that only 5%
34 (around 70,000) of fungi total populations (1.5 million) have been explored (Sharma *et al.*, 2016).
35 This confirms fungi as possible alternatives to plants for producing valuable natural products.

36 Although the soil fungi are well-known as a source for important antibiotics and drugs (Wasser,
37 2002; Sethi *et al.*, 2013) much effort has been devoted in last two decades to study the secondary
38 metabolites form marine and endophytic fungi as new sources for drugs and agrochemicals (Butler
39 *et al.*, 2014; Segaran *et al.*, 2019).

40 Interest in microbial metabolites as pesticides arises to overcome the problems associated with
41 the continuous use of synthetic pesticides, such as adverse effects of pesticide residues on human,
42 non-target organisms and environment, and emergence of pest resistance (Aktar *et al.*, 2009). In
43 fact **the** microbial metabolites have several advantages to be good candidates in pest control, such
44 as structure diversity, wide-spectrum of bioactivity, new modes of action and biodegradability
45 (Gerwick and Sparks, 2014). Several pesticides of microbial origin have been introduced into field
46 of pest management include: avermectin, bialaphos, milbemycin, kasugamycin, blasticidin S,
47 polyoxin, mildiomicin, validamycin and tetranactin (Tanaka and Omura, 1993). Therefore, it is
48 highly important to investigate new fungal metabolites and examine their bioactivities against
49 economic pests.

50 *Silybum marianum* (L.) Gaertn. (Asteraceae) is a dangerous weed in many parts of Middle East,
51 Africa, Australia, and North and South America (Holm *et al.*, 1997). In Egypt, it is considered as
52 an invasive weed that spread through wheat fields in both new reclaimed lands and desert areas.
53 *Culex pipiens* L. (Diptera: Culicidae) is the most widespread mosquito species in Egypt (Zahran *et*
54 *al.*, 2017; Abdelgaleil *et al.*, 2017). This insect is able to transmit several pathogens, such as West
55 Nile virus, *Bancroftian flariasis* and Rift Valley Fever virus (Meegan *et al.*, 1980).

56 In our continuous efforts to search for new natural products with possible application in pest
57 management, twelve fungal secondary metabolites (**1-12**) were examined for their effect on the
58 germination and growth of roots and shoots of *S. marianum*. In addition, the toxicity of these
59 compounds was tested against the fourth larval stage of *C. pipiens* to explore their potential for the
60 control of these economic pests.

61

62 **MATERIALS AND METHODS**

63 **Test organisms**

64 Field biotype seeds of **Holy** thistle, *Silybum marianum* (L.) Gaertn. (Asteraceae), were collected
65 from Faculty of Agriculture Farm, Alexandria (31° 12' 56.30" N, 29° 57' 18.97" E). The plant seeds
66 were identified by Prof. FathAllah Zaitoon of the Plant Pathology Department, Faculty of
67 Agriculture, Alexandria University. Voucher specimen (SM-1909) was deposited in Department

68 of Pesticide Chemistry and Technology, Faculty of Agriculture, Alexandria University. Uniform
69 and undamaged seeds were used for the germination and seedling growth tests. Seeds were
70 examined for their germination before experiments. The seed germination was 80% after 12 days
71 of sowing. *Culex pipiens* L. (Diptera: Culicidae) was reared in an insectary at 27 ± 2 °C and $75\pm 5\%$
72 **relative humidity (R.H.)** at the Department of Applied Entomology and Zoology, Alexandria
73 University, Egypt as described by Zahran *et al.* (2017).

74

75 **Fungal strains and isolation of secondary metabolites**

76 Strains of different endophytic and soil fungi were isolated from different sources (Table 1),
77 identified by BEX Co. Ltd., Japan, using a DNA analysis of the 18S rDNA regions and had been
78 deposited in the laboratory of natural products at the Faculty of Agriculture of Yamagata
79 University. After fermentation on unpolished rice media (unpolished rice 1000 g + 35 ml of 3.5%
80 NaCl at 25°C for four weeks), the fungal growth media were extracted with MeOH. The MeOH
81 extracts were concentrated and partitioned into *n*-hexane and EtOAc. The EtOAc extracts **were**
82 **subjected to silica gel column chromatography with mixtures** of *n*-hexane–EtOAc (100:0 –
83 0:100, v/v), and mixtures of EtOAc–MeOH (50:50 – 0:100, v/v). The rustling fractions were further
84 chromatographed on ODS by eluting with H₂O and an increasing ratio of MeOH (100:0 to 0:100)
85 or silica gel by eluting with a mixture of CHCl₃–MeOH (90:10, v/v). For HPLC analysis, a reversed
86 phase HPLC (Semipreparative HPLC with Shimadzu pump and UV LC-10A detector (set at 210
87 nm) on a Mitysil ODS column (150 × 6.0 mm i.d.) at a flow rate of 2.0 mL/min, solvent systems,
88 MeOH:H₂O, 60:40 – 80:20, v/v) and PTLC (EtOAc – *n*-hexane, 70:30 – 30:70, v/v) were used.
89 The purity of isolated compounds was higher than 95%.

90

91 **Germination and seedling growth inhibition bioassay**

92 The effect of fungal secondary metabolites (**1-12**) on seed germination, and root and shoot growth
93 of *S. marianum* was evaluated using a bioassay method explained by Abdelgaleil *et al.* (2009). The
94 tested compounds were dissolved first in DMSO and diluted with distilled water containing 0.02%
95 of Triton-X 100 to give final concentration of 500 mg L⁻¹. Three replicates with 10 seeds in each
96 one were prepared for each compound. The compound solutions (6 ml) were transferred to each
97 Petri dish (9 cm) lined with filter paper (Whatman No. 2). Petri dishes were then placed in
98 polyethylene bags which were expanded and closed to prevent moisture loss. In the control
99 treatment, a solution of 0.5% of DMSO and 0.02% of Triton-X 100 in distilled water was added.

100 All treatments were kept at 20±2 °C and a 12-h photoperiod for 12 days. Then, the number of
101 germinated seeds and length of root and shoot were taken. Growth inhibition (I %) of root and
102 shoot lengths was calculated from this equation:

$$103 \quad I (\%) = [1 - T/C] \times 100$$

104 Where, T and C are the root or shoot lengths (cm) in treatment and control. Moreover, two
105 compounds (**6** and **11**) were further evaluated on germination and seeding growth at a series of
106 concentrations of 25, 50, 100, 250 and 500 mg L⁻¹.

107

108 **Larvicidal bioassay**

109 The toxicity of fungal secondary metabolites (**1-12**) was carried out against *C. pipiens* larvae
110 following a recommended method of the World Health Organization (WHO, 1996) with slight
111 modifications. The tested compounds were dissolved in dimethyl sulfoxide (DMSO) and diluted
112 with distilled water containing 0.1% of Tween-20 to give three final concentrations (100, 250 and
113 500 mg L⁻¹). Water in control treatments was mixed with DMSO (0.5%) and Tween-20 (0.1%).
114 Twenty *C. pipiens* larvae were separately put into 200-ml plastic cups containing 100 ml of the
115 compound solutions. Each concentration was replicated three times. All treatments were kept under
116 the same insect rearing conditions for 24 h. Then, the number of dead larvae was counted and
117 mortality (%) was calculated. Three compounds (**4**, **5** and **9**) were further tested at concentrations
118 of 10, 25, 50, 70 and 100 mg L⁻¹. Mortality data of these compounds were subjected to probit
119 analysis to calculate LC₅₀ values for these compounds (Finney, 1971).

120

121 **Statistical analysis**

122 Significant differences among mean values of germination percentages, root and shoot lengths
123 were determined (P = 0.05) by using a one-way analysis of variance (ANOVA) followed by
124 Tukey's HSD test. The LC₅₀ values of compounds were estimated by probit analysis using the
125 SPSS 21.0 software (Statistical Package of Social Sciences Inc., USA).

126

127 **RESULTS**

128 **Isolation, structure elucidation of isolated secondary metabolites**

129 Twelve secondary metabolites (**1-12**) have been isolated from different fungal strains (Table 1)
130 using different chromatographic techniques, including silica gel and ODS columns, PTLC and
131 HPLC. Among the isolated metabolites, nine compounds (**1**, **2**, **4-7** and **10-12**) were isolated from

132 endophytic fungi and three compounds (**3**, **8**, and **9**) were isolated from soil fungi. The chemical
133 structure (Fig. 1) of the isolated compounds was elucidated on the basis of their spectroscopic data
134 of ultraviolet (UV), infrared (IR), high resolution mass spectroscopy (HRMS) and nuclear
135 magnetic resonance (NMR). The isolated compounds belong to ten different chemical groups of
136 natural products (Table 1).

137 138 **Effect of fungal metabolites on germination and seedling growth of *S. marianum***

139 Effect of fungal secondary metabolites (**1-4** and **6-12**) on germination, root and shoot growth of
140 *S. marianum* at concentration of 500 mg L⁻¹ is shown in Table 2. Among the **11** tested metabolites,
141 compounds **3**, **4**, **6**, **8**, **9**, **10** and **11** caused reduction in seed germination after 12 days of treatment.
142 Compounds **6** and **11** induced complete inhibition of seed germination at this concentration. In
143 addition, seven compounds (**2**, **3**, **4**, **6**, **8**, **11** and **12**) revealed significant inhibition in root growth
144 compared with control treatment. **Similarly**, compounds **6** and **11** which showed complete
145 inhibition of germination, compounds **3**, **4** and **12** strongly inhibited root growth with 87.5, 80.4
146 and 74.8% inhibition, respectively. Conversely, compounds **1** and **10** significantly increased the
147 root growth. Similarly, compounds **3**, **4**, **8** and **12**, **significantly reduced** in shoot growth. The
148 results revealed that the tested compounds caused higher inhibitory effects on **plant establishment**
149 **by reducing** root and shoot growth than on seed germination. Also, some of the tested compounds
150 had greater inhibitory effects on root growth than on shoot growth.

151 The previously presented results showed that compounds **6** and **11** caused complete inhibition
152 of seed germination at 500 mg L⁻¹. Therefore, these two compounds were further evaluated at
153 lower concentrations (25, 50, 100, 250 and 500 mg L⁻¹). Compound **6** caused complete inhibition
154 of root growth at all tested concentrations (Table 3). Likewise, it induced complete inhibition of
155 germination and shoot growth at all tested concentrations except 25 mg L⁻¹. This compound
156 showed strong inhibition of germination (10.0%) and shoot growth (**74.5%**) at 25 mg L⁻¹. In
157 addition, compound **11** induced significant reduction in seed germination at all of the tested
158 concentrations. Also, it caused strong inhibition of root and shoot growth at 100 and 250 mg L⁻¹.

159 160 **Larvicidal activity of fungal metabolites against *C. pipiens***

161 Toxicity of twelve fungal metabolites against the fourth larval instar of *C. pipiens* at
162 concentrations of 100, 250 and 500 mg L⁻¹ is presented in Table 4. Four compounds (**4**, **5**, **8** and
163 **9**) caused complete mortality (100%) of larvae at 500 mg L⁻¹, while three compounds (**4**, **5** and **9**)

164 caused complete mortality of larvae at 250 mg L⁻¹. At 100 mg L⁻¹, compound **9** was the only
165 compound that caused complete mortality of larvae. Also, compounds **4** and **5** induced strong larval
166 toxicity at this concentration with 95.0 and 90.0 % mortality, respectively. Comparative toxicity
167 results (Table 5) indicated that compound **9** revealed the highest larval toxicity (LC₅₀ = 3.27 mg
168 L⁻¹), followed by compound **4** (LC₅₀ = 57.03 mg L⁻¹) and compound **5** (LC₅₀ = **63.50** mg L⁻¹).

169 170 **DISCUSSION**

171 New pest management strategies are based on decrease the use of synthetic pesticides and
172 increase the use of natural alternatives. Therefore, introducing new natural products with low risk
173 on human and environment, and new modes of action is highly needed in the field pest management
174 nowadays (Schrader *et al.*, 2010). In this regard, microorganisms are a rich source for natural
175 compounds with successful use in pest management (Singh, and Yadav, 2020; Saad *et al.*, 2021).
176 **Several hundred species of fungi are known to infect and kill pests. The most important**
177 **groups are the Hyphomycetes within genera like *Beauveria*, *Metarhizium*, *Verticillium*,**
178 ***Paecilomyces* and *Hirsutella* (Keller, 1998). Moreover, fungi are also known for producing**
179 **toxic compounds against pests mainly form soil fungi of genera *Aspergillus* and *Penicillium*.**
180 **Secondary metabolites with insecticidal and antifeedant properties were also found in**
181 **endophytic and phytopathogenic fungi (Berestetskiy and Hu, 2021).** In this study, the herbicidal
182 and insecticidal activities of twelve secondary metabolites isolated from fungi were investigated to
183 explore their possible use in pest control programs.

184 The results showed that most of the tested compounds reduced seed germination and seedling
185 growth of *S. marianum* at concentration of 500 mg L⁻¹. Among the eleven tested metabolites,
186 compounds **6** and **11** revealed excellent herbicidal activity with complete inhibition of germination
187 and seedling growth at this concentration. Moreover, compound **6** showed strong herbicidal
188 activity at 25 mg L⁻¹. Based on the obtained results, this compound is among the most effective
189 natural herbicides reported so far. In our previous study, compounds **6** and **11** were also the most
190 potent inhibitors of seed germination, and growth of root and shoot of *Echinochloa crus-galli* at
191 concentration of 2mM (Saad *et al.*, 2021). Also, compound **6** was found to be more effective than
192 compound **11** against *E. crus-galli*. The herbicidal activity of compound **11** was further supported
193 by the studies of Shiono and Murayama (2005) who stated that this compound inhibited the root

194 growth of lettuce by 46% at concentration of 50 mg L⁻¹. In addition, Compound **6** has been
195 described to inhibit wheat germination and the root growth of *Allium cepa* (Betina, 1992)

196 Comparing the inhibitory effects of tested compounds on the seed germination with seedling
197 growth of *S. marianum* indicated that the tested compounds were more effective against seedling
198 growth than germination. These findings are in agreement with previous studies on many natural
199 compounds in which the seedling growth were more sensitive than germination (Leather and
200 Einhellig, 1984; Abdelgaleil and Hashinaga, 2007; Saad *et al.*, 2012). Also, the results of current
201 study revealed that the tested compounds were more potent inhibitors to root than to shoot with
202 few exceptions. The higher sensitivity of roots could be due to the roots are the first to expose and
203 absorb tested compounds from growing media (Turk *et al.*, 2002). Similar observations were
204 reported on the inhibitory effects of other natural compounds on seedling growth (Wang *et al.*,
205 2007; Gouda *et al.*, 2016).

206 The results of larvicidal bioassays against *C. pipiens* demonstrated that among the tested
207 metabolites, compounds **9**, **4** and **5** showed a promising larvicidal activity. In particular, compound
208 **9** which displayed the highest toxicity with LC₅₀ = 3.27 mg L⁻¹. Based on our results, this
209 compound is among the most toxic natural compounds reported so far against *C. pipiens* or other
210 mosquito species. This finding is supported by previous study of Geris *et al.* (2008) who found that
211 compound **9** was highly toxic against third instar larvae of *Aedes aegypti* with LC₅₀ value of 2.9
212 mg L⁻¹. Compound **9** was more toxic to *C. pipiens* larvae than thymol (LC₅₀ = 37.95 mg L⁻¹),
213 carvacrol (LC₅₀ = 44.38 mg L⁻¹), cinnamaldehyde (LC₅₀ = 58.97 mg L⁻¹), eugenol (LC₅₀ = 86.22
214 mg L⁻¹) and cuminaldehyde (LC₅₀ = 38.94 mg L⁻¹) (Radwan *et al.*, 2008; Zahran and Abdelgaleil,
215 2011). Similarly, compound **9** showed higher toxicity than curcumin (LC₅₀ = 19.07 mg L⁻¹)
216 isolated from *Curcuma longa* (Sagnou *et al.*, 2012).^[44] Moreover, this compound showed similar
217 toxicity to caulerpin (LC₅₀ = 1.99 mg L⁻¹) and caulerpinic acid (LC₅₀ = 4.89 mg L⁻¹) isolated from
218 *Caulerpa racemosa* (Alarif *et al.*, 2010).

219 The results of the current study indicate that the tested compounds showed high selectivity against
220 the examined pests. For example, compounds **6** and **11** showed promising herbicidal activity
221 against *S. marianum* while had no insecticidal activity. Likewise, compounds **5** and **9** possessed
222 potent insecticidal activity against *C. pipiens* but were not active against the tested weed. This
223 selectively is crucial for the development of these compounds as new bio-pesticides.

224 In conclusion, this study described, for the first time, the herbicidal and insecticidal activities of
225 twelve fungal metabolites against the economic agricultural pests, *S. marianum* and *C. pipiens*.
226 Among the tested metabolites, brefeldin A (**6**) and 6-epoxy-4-hydroxy-3-methoxy-5-methyl-
227 cyclohex-2-en-1-one (**11**) revealed promising herbicidal activity against *S. marianum*, in particular,
228 brefeldin A (**6**) which could be developed as a new natural herbicide. In addition, dehydroaustin
229 (**9**) exhibited remarkable larvicidal activity against *C. pipiens* indicating its possible use in
230 integrated management programs of mosquitoes.

231

232 ACKNOWLEDGEMENTS

233 The authors are grateful to Eng. Hagar Youssef for her technical assistance.

234

235 REFERENCES

- 236 1. Abdelgaleil, S. A. M., Abdel-Razeek, N. and Soliman, S. A. 2009. Herbicidal activity of
237 three sesquiterpene lactones on wild oat (*Avena fatua*) and their possible mode of action. *Weed*
238 *Sci.*, **57**: 6-9.
- 239 2. Abdelgaleil, S. A. M. and Hashinaga, F. 2007. Allelopathic potential of two sesquiterpene
240 lactones from *Magnolia grandiflora* L. *Biochem. Syst. Ecol.*, **35**: 737-742.
- 241 3. Abdelgaleil, S. A. M., Zoghroban, A. A. M., El-Bakry, A. M. and Kassem, S. M. I., 2019.
242 Insecticidal and antifungal activities of crude extracts and isolated compounds from rhizomes of
243 *Curcuma longa* (Zingiberaceae). *J. Agric. Sci. Technol.*, **21**: 1049-1061.
- 244 4. Aktar, W. and Sengupta, D. and Chowdhury, A. 2009. Impact of pesticides use in
245 agriculture: Their benefits and hazards. *Interdiscip. Toxicol.*, **2**: 1-12.
- 246 5. Alarif, W. M., Abou-Elnaga, Z. S., Ayyad, S.-E. N. and Al-lihaibi, S, S. 2010. Insecticidal
247 metabolites from the green alga *Caulerpa racemosa*. *Clean: Soil, Air, Water*, **38**: 548-557.
- 248 6. Betina, V. 1992. Biological effects of the antibiotic brefeldin A (decumbin, cyanein,
249 ascotoxin, synergisidin): a retrospective. *Folia Microbiol.*, **37**: 3-11.
- 250 7. **Berestetskiy, A., Hu, Q. 2021. The Chemical Ecology Approach to Reveal Fungal**
251 **Metabolites for Arthropod Pest Management. *Microorganisms*, 9: 1379.**
- 252 8. Butler, M. S., Robertson, A. A. and Cooper, M. A. 2014. Natural product and natural
253 product derived drugs in clinical trials. *Nat. Prod. Rep.*, **31**: 1612-1661.

- 254 9. Elsaesser, B., Krohn, K., Floerke, U., Root, N., Aust, H. J., Draeger, S., Schulz, B., Antus,
255 S. and Kurtán, T. 2005. X-ray structure determination absolute configuration and biological
256 activity of phomoxanthone. *Eur. J. Org. Chem.*, **21**: 4563–4570.
- 257 10. Finney, D. J. 1971. Probit Analysis, 3rd ed. Cambridge University Press, London, p. 318.
- 258 11. Geris, R., Rodrigues-Fo, E., da Silva, H. H. G. and da Silva, I. G. I. G. 2008. Larvicidal
259 effects of fungal meroterpenoids in the control of *Aedes aegypti* L., the main vector of dengue and
260 yellow fever. *Chem. Biodivers.*, **5**: 341-345.
- 261 12. Gerwick, B. C. and Sparks, T, C. 2014. Natural products for pest control: An analysis of
262 their role, value and future. *Pest Manag. Sci.*, **70**: 1169–1185.
- 263 13. Gouda, N. A. A., Saad, M. M. G. and Abdelgaleil, S. A. M. 2016. Pre and Post Herbicidal
264 activity of monoterpenes against barnyard grass (*Echinochloa crus-galli*). *Weed Sci.*, **64**: 191-200.
- 265 14. Hayasaka, S., Koseki, T., Murayama, T., Kwon, E. and Shiono, Y. 2011.
266 Phenylisobenzofuranones from *Fungicolous nodulisporium* sp. SH-1. *Z. Naturforsch.*, **66b**: 961–
267 964.
- 268 15. Holm, L. G., Doll, J., Holm, E., Pancho, J. and Herberger, J. 1997. World Weeds. Natural
269 Histories and Distribution. Wiley, NewYork.
- 270 16. Hutchinson, C. R., Shu-Wee, L., Mcinnes, A. G. and Walter, J. A. 1983. Comparative
271 biochemistry of fatty acid and macrolide antibiotic (Brefeldin A) formation in *Penicillium*
272 *brefeldianum*. *Tetrahedron*, **39**: 3507-3513.
- 273 17. Ikeda, M., Niwa, G.-I., Tohyama, K., Sassa, T. and Miura, Y. 1977. Structures of fasciculol
274 C and its depsipeptides, new biologically active substances from *Neamatoloma fasciculare*. *Agric.*
275 *Biol. Chem.*, **41**: 1803-1805.
- 276 **18. Keller, S. 1998. Use of Fungi for Pest Control in Sustainable Agriculture.**
277 ***Phytoprotection*, **79**: 56-60.**
- 278 19. Kim, K. H., Moon, E., Choi, S. U., Kim, S. Y. and Lee, K. R. 2013. Lanostane triterpenoids
279 from the mushroom *Naematoloma fasciculare*. *J. Nat. Prod.*, **76**: 845–851.
- 280 20. Leather, G. R. and Einhellig, A. 1985. Mechanisms of allelopathic action in bioassay, pp
281 197–205, In the Chemistry of Allelopathy. Washington, DC: American Chemical Society.
- 282 21. Lucas, E. M. F., de Castro, M. C. M. and Takahashi, J. A. 2007. Antimicrobial properties
283 of sclerotiorin, isocromophilone VI and pencolide, metabolites from Brazilian cerrado isolate
284 of *Penicillium sclerotiorum* van beyma. *Braz. J. Microbiol.*, **38**: 785-789.

- 285 22. Masters, K. S. and Bräse, S. 2002. Xanthenes from fungi, lichens, and bacteria: the natural
286 products and their synthesis. *Chem. Rev.*, 112: 3717–3776.
- 287 23. Meegan, J. M., Khalil, G. M., Hoogstraal, H. and Adham, F. K. 1980. Experimental
288 transmission and field isolation studies implicating *Culex pipiens* as a vector of Rift Valley fever
289 virus in Egypt. *Am. J. Trop. Med. Hyg.*, **29**: 1405–1410.
- 290 24. Porras-Alfaro, A. and Bayman, P. 2011. Hidden fungi, emergent properties: endophytes
291 and microbiomes. *Annu. Rev. Phytopathol.*, **49**: 291-315.
- 292 25. Radwan, M. A., El-Zemity, S. R., Mohamed, S. A. and Sherby, S. M. 2008. Larvicidal
293 activity of some essential oils, monoterpenoids and their corresponding N-methyl carbamate
294 derivatives against *Culex pipiens* (Diptera: Culicidae). *Int. J. Trop. Insect Sci.*, **28**: 61–68.
- 295 26. Ronsberg, D., Debbab, A., Mandi, A., Vasylyeva, V., Bohler, P., Stork, B., Engelke, L.,
296 Hamacher, A., Sawadogo, R., Diederich, M., Wray, V., Lin, W., Kassack, M. U., Janiak, C., Scheu,
297 S., Wesselborg, S., Kurtan, T., Aly, A. H. and Proksch, P. 2013. Pro-apoptotic and
298 immunostimulatory tetrahydroxanthone dimers from the endophytic fungus *Phomopsis longicolla*.
299 *J. Org. Chem.* **78**: 12409–12425.
- 300 27. Saad, M. M. G., Abdelgaleil, S. A. M. and Shiono, Y. 2021. Antibacterial and herbicidal
301 properties of secondary metabolites from fungi. *Nat. Prod. Res.* **35**: 5446–5451.
- 302 28. Saad, M. M. G., Abdelgaleil, S. A. M. and Suganuma, T. 2012. Herbicidal potential of
303 pseudoguaninolide sesquiterpenes on wild oat, *Avena fatua* L. *Biochem. Syst. Ecol.*, **44**: 333-337.
- 304 29. Sagnou, M., Mitsopoulou, K. P., Koliopoulos, G., Pelecanou, M., Couladouros, E. A. and
305 Michaelakis, A. 2012. Evaluation of naturally occurring curcuminoids and related compounds
306 against mosquito larvae. *Acta Trop.*, **123**: 190–195.
- 307 30. Schrader, K. K., Andolfi, A., Cantrell, C. L., Cimmino, A., Duke, S. O., Osbrink, W.,
308 Wedge, D. E. and Evidente, A. 2010. A Survey of phytotoxic microbial and plant metabolites as
309 potential natural products for pest management. *Chem. Biodivers.*, **7**: 2261-2280.
- 310 31. Schürmann, B. T. M., Sallum, W. S. T. and Takahashi, J. A. 2010. Austin, dehydroaustin
311 and other metabolites from *Penicillium brasilianum*. *Quím Nova.*, **33**: 1044–1046.
- 312 32. Segaran, G. and Sathiavelu, M. 2019. Fungal endophytes: A potent biocontrol agent and a
313 bioactive metabolite reservoir. *Biocatal. Agric. Biotechnol.*, **21**: 101284.
- 314 33. Sethi S, Kumar R, Gupta S. 2013. Antibiotic production by microbes isolated from soil. *Int.*
315 *J. Pharm. Sci. Res.*, **4**: 2967.

- 316 34. Sharma, D., Pramanik, A. and Agrawal, P. K. 2016. Evaluation of bioactive secondary
317 metabolites from endophytic fungus *Pestalotiopsis neglecta* BAB-5510 isolated from leaves of
318 *Cupressus torulosa* D.Don. *3 Biotech*, **6**: 210.
- 319 35. Shiono, Y., Kikuchi, M., Koseki, T., Murayama, T., Kwon, E., Aburai, N., Kimura, K.-
320 I. 2011. Isopimarane diterpene glycosides, isolated from endophytic fungus *Paraconiothyrium* sp.
321 MY-42. *Phytochemistry*, **72**: 1400-1405.
- 322 36. Shiono, Y. and Murayama, T. 2005. New eremophilane-type sesquiterpenoids,
323 eremoxyларins A and B from *Xylariaceous endophytic* fungus YUA-026. *Z. Naturforsch.*, **60b**:
324 885–890.
- 325 37. Shiono, Y., Murayama, T., Takahashi, K., Okada, K., Katohda, S. and Ikeda, M. 2005.
326 Three oxygenated cyclohexanone derivatives produced by an endophytic fungus. *Biosci.*
327 *Biotechnol. Biochem.*, **69**: 287–292.
- 328 38. Shiono, Y. 2006. Anthracobic acids A and B, two polyketides, produced by an endophytic
329 fungus *Anthracobia* sp. *Chem. Biodivers.*, **3**: 217-223.
- 330 39. Silvie, P. J., Martin, P., Huchard, M., Keip, P., Gutierrez, A. and Sarter, S. 2021.
331 Prototyping a knowledge-based system to identify botanical extracts for plant health in Sub-
332 Saharan Africa. *Plants*, **10**: 896.
- 333 40. Singh, J. and Yadav, A. N. 2020. Natural bioactive products in sustainable agriculture.
334 Springer, Singapore.
- 335 41. Tanaka, Y. and Omura, S. 1993. Agroactive compounds of microbial origin. *Annu. Rev.*
336 *Microbiol.*, **47**: 57-87.
- 337 42. Turk, M. A., Abdel-Rahman and Tawaha, M. 2002. Inhibitory effects of aqueous extracts
338 of black mustard on germination and growth of lentil. *Pak. J. Agron.* **1**: 28–30.
- 339 43. Wang, W., Zhu, T., Tao, H., Lu, Z., Fang, Y., Gu, Q. and Zhu, W. 2007. Two new cytotoxic
340 quinone type compounds from the Halotolerant fungus *Aspergillus varicolor*. *J. Antibiot.*, **60**:
341 603–607.
- 342 44. Wasser, S. P. 2002. Medicinal mushrooms as a source of antitumor and immunomodulating
343 polysaccharides. *Appl. Microbiol. Biotechnol.*, **60**: 258–274.
- 344 45. WHO1996. Report of the WHO informal consultation on the evaluation on the testing of
345 insecticides CTD/WHO PES/IC/96.1, p 69.

346 46. Wicklow, D. T. and Poling, S. M. 2009. Antimicrobial activity of pyrrocidines from
347 *Acremonium zeae* against endophytes and pathogens of maize. *Phytopathol.*, **99**: 109-115.

348 47. Zahran, H. A. and Abdelgaleil, S. A. M. 2011. Insecticidal and developmental inhibitory
349 properties of monoterpenes on *Culex pipiens* L. (Diptera: Culicidae) *J. Asia-Pac. Entomol.*, **14**:
350 46–51.

351 48. Zahran, H. E. D., Abou-Taleb, H. K. and Abdelgaleil, S. A. M. 2017. Adulticidal, larvicidal
352 and biochemical properties of essential oils against *Culex pipiens* L. *J. Asia-Pac. Entomol.*, **20**:
353 133–139.

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

Table 1. Names, chemical class and source of fungal secondary metabolites.

Name	Chemical group	Fungus (fungal origin)	Reference
Nodulisporone B (1)	Phenylisobenzofuranones	<i>Nodulisporium</i> sp. SH-1 (<i>Xylaria polymorpha</i>)	Hayasaka <i>et al.</i> , 2011
19-(α -D-glucopyranosyloxy)isopimarane-7,15-dien-3 β -ol (2)	Isopimarane diterpenoids	<i>Paraconiothyrium</i> sp. MY-42 (unidentified plant)	Shiono <i>et al.</i> , 2011
Pencolide (3)	Maleimides	<i>Penicillium sclerotiorum</i> (Soil)	Lucas <i>et al.</i> , 2007
Phomaxanthone A (4)	Dimeric xanthenes	<i>Phomopsis</i> sp. (<i>Rhizophora mucronata</i>)	Elsaesser <i>et al.</i> , 2005
Deacetylphomaxanthone A (5)	Dimeric xanthenes	<i>Phomopsis longicolla</i> (<i>Sonneratia caseolaris</i>)	Ronsberg <i>et al.</i> , 2013
Brefeldin A (6)	Macrolides	<i>Penicillium brefeldianum</i> (<i>Pinellia ternata</i>)	Hutchinson <i>et al.</i> , 1983
Anthracobic acid A (7)	Polyketides	<i>Anthracobia</i> sp. (unidentified plant)	Shiono, 2006
Fasciculol C (8)	Lanostane triterpenoids	<i>Neamatoloma fasciculare</i> (Mushroom)	Ikeda <i>et al.</i> , 1977; Kim <i>et al.</i> , 2013
Dehydroaustin (9)	Meroterpenoids	<i>Penicillium brasilianum</i> (Soil)	Schürmann <i>et al.</i> , 2010
Pyrrucidine A (10)	Alkaloids	<i>Acremonium zeae</i> (<i>Zea mays</i>)	Wicklow and Poling, 2009
6-Eopxy-4-hydroxy-3-methoxy-5-methyl-cyclohex-2-en-1-one (11)	Cyclohexenones	Xylariaceous endophytic fungus (YUA 026) (unidentified plant)	Shiono <i>et al.</i> , 2005
Secalonic acid A (12)	Dimeric xanthenes	<i>Claviceps purpurea</i> (<i>Secale cereale</i>)	Masters and Bräse, 2002

378

379

380

Table 2 Effect secondary metabolites isolated from endophytic fungi on germination and seedling growth of *Silybum marianum* after 12 days of sowing at 500 mg L^{-1a}.

Compound	Germination ^b (% \pm SE)	Root		Shoot	
		Length (cm) (Mean \pm SE)	I ^c (%)	Length (cm) (Mean \pm SE)	I (%)
Control	86.7 \pm 3.34ab	11.39 \pm 0.22bc	0.0	2.16 \pm 0.06abc	0.0
1	86.7 \pm 3.34ab	14.32 \pm 0.14a	-28.2	2.33 \pm 0.13ab	-7.87
2	86.7 \pm 3.34ab	5.47 \pm 0.23d	52.0	1.86 \pm 0.24bc	13.9
3	76.7 \pm 3.34b	1.42 \pm 0.06g	87.5	1.33 \pm 0.17d	38.4
4	80.0 \pm 5.78b	2.23 \pm 0.25f	80.4	1.30 \pm 0.10d	39.8
6	0.0 \pm 0.0c	0.0 \pm 0.0h	100.0	0.0 \pm 0.0e	100.0
7	96.71 \pm 3.34a	11.93 \pm 0.11b	-4.74	2.33 \pm 0.25ab	-7.87
8	83.3 \pm 3.34ab	5.05 \pm 0.42d	55.7	1.35 \pm 0.09d	37.5
9	80.0 \pm 5.78b	11.12 \pm 0.17c	2.37	2.53 \pm 0.06a	-17.1
10	83.3 \pm 6.67ab	13.82 \pm 0.17a	-21.3	1.80 \pm 0.26c	16.7
11	0.0 \pm 0.0c	0.0 \pm 0.0h	100.0	0.0 \pm 0.0e	100.0
12	90.0 \pm 5.78ab	2.87 \pm 0.11e	74.8	1.31 \pm 0.16d	39.4
df ^d	11	11		11	
F ^e	63.74	801.70		29.30	
P ^f	<0.00	<0.00		<0.00	

381

382

383

384

385

^a Values are means \pm SE of three replicates with 10 seeds in each one.^b Values followed by different letters are significantly different at 0.05 probability level.^c I = Inhibition.^d df = Degrees of freedom.^e F = F-statistic. ^f P = P-value.

386 **Table. 3** Effect of brefeldin A and 6-eopxy-4-hydroxy-3-methoxy-5-methyl-cyclohex-2-en-1-
 387 one on germination and seedling growth of *Silybum marianum* after 12 days of sowing at different
 388 concentrations^a.

Compound	Conc. (mg L ⁻¹)	Germination (% ± SE) ^b	Root		Shoot	
			Length (cm) (Mean ± SE)	I ^c (%)	Length (cm) (Mean ± SE)	I ^c (%)
Control	0.0	75.0±2.89a	6.33±0.49a	0.0	1.96±0.21a	0.0
Brefeldin A (6)	25	10.0±5.78d	0.0±0.0d	100.0	0.50±0.28c	74.5
	50	0.0±0.0d	0.0±0.0d	100.0	0.0±0.0d	100.0
	100	0.0±0.0d	0.0±0.0d	100.0	0.0±0.0d	100.0
	250	0.0±0.0d	0.0±0.0d	100.0	0.0±0.0d	100.0
	500	0.0±0.0d	0.0±0.0d	100.0	0.0±0.0d	100.0
6-Eopxy-4-hydroxy-3- methoxy-5-methyl- cyclohex-2-en-1-one (11)	25	60.0±5.78b	6.27±0.40a	0.95	1.63±0.03a	16.8
	50	56.7±3.33b	4.50±0.17b	28.9	1.87±0.20a	4.59
	100	43.3±3.33c	0.83±0.27c	86.9	1.20±0.17b	38.8
	250	6.7±3.33d	0.02±0.02d	99.7	0.17±0.17cd	91.3
	500	0.0±0.0d	0.0±0.0d	100.0	0.0±0.0d	100.0
df ^d		10	10		10	
F ^e		88.34	139.35		32.99	
P ^f		<0.00	<0.00		<0.00	

389 ^a Values are means ± SE of three replicates with 10 seeds in each one.

390 ^b Values followed by different letters are significantly different at 0.05 probability level.

391 ^c I = Inhibition.

392 ^d df = Degrees of freedom.

393 ^e F = F-statistic.

394 ^f P = P-value.

396 **Table. 4** Mortality percentages of *Culex pipiens* fourth instar larvae after 24 h of treatment with
 397 secondary metabolites isolated from endophytic fungi at different concentrations.

Compound	Mortality (%) after 24h		
	100 mg L ⁻¹	250 mg L ⁻¹	500 mg L ⁻¹
Control	0.0	0.0	0.0
Nodulisporone B (1)	0.0	0.0	0.0
19-(α -D-glucopyranosyloxy) isopimara-7,15-dien-3 β -ol (2)	0.0	- ^a	-
Pencolide (3)	0.0	0.0	0.0
Phomaxanthone A (4)	95.0	100.0	100.0
Deacetylphomaxanthone A (5)	90.0	100.0	100.0
Brefeldin A (6)	0.0	0.0	46.7
Anthracobic acid A (7)	0.0	0.0	10.0
Fasciculol C (8)	0.0	6.25	100.0
Dehydroaustin (9)	100	100.0	100.0
Pyrrocidine A (10)	0.0	0.0	15.0
6-Eopxy-4-hydroxy-3-methoxy-5- methyl-cyclohex-2-en-1-one (11)	0.0	0.0	38.9
Secalonic acid A (12)	0.0	0.0	0.0

408 a= Not tested.

410 **Table 5.** Comparative toxicity of fungal metabolites against *Culex pipiens* fourth instar larvae after
 411 24h of exposure.

Compound	LC ₅₀ ^a (mg L ⁻¹)	95% confidence limits (mg L ⁻¹)		Slope ± SE	Intercept ± SE	(χ ²) ^b	P ^c
		Lower	Upper				
Phomaxanthone A (4)	57.03	52.62	61.39	6.75±0.81	-11.85±1.47	0.001	0.978
Deacetylphomaxanthone A (5)	63.50	58.75	68.41	6.50±0.72	-11.73±1.31	0.006	0.941
Dehydroaustin (9)	3.27	0.38	6.53	1.11±0.30	-0.57±0.40	0.181	0.671

412 ^aThe concentration causing 50% mortality.

413 ^bChi square value.

414 ^cProbability value.




Figure 1. Chemical structure of fungal metabolites (1-12).

415

416

417