

Exploration And Characterization of Superior Locals Durian In Two Subdistricts of Malang East Java Indonesia

Sumeru Ashari¹, Eko Muhammad Bayu¹, Eni Fidiyawati^{2*}, Weda Makarti Mahayu³, Tri Sudaryono³, and Sudarmadi Purnama³

1. Department of Agronomy, Faculty of Agriculture, University of Brawijaya, Jl. Veteran, 65145, Indonesia.

2. Horticulture and Estate Research Centre, National Research and Inovation Agency (BRIN), Cibinong Science Center, Jl. Raya Jakarta- Bogor, 16915, Indonesia.

3. Pusat Riset Hortikultura dan Perkebunan, National Research and Inovation Agency (BRIN), Cibinong Science Center, Jl. Raya Jakarta- Bogor, 16915, Indonesia.

*Corresponding author; e-mail: kagakunoriteni@gmail.com

ABSTRACT

Durian production centers in East Java areas are quite numerous, two of which are Kasembon and Ngantang Subdistricts, Malang District. In these areas, most durian trees are propagated from seeds. This research aimed to determine the genetic diversity by exploration and characterization of superior types, it was conducted by descriptive method. Determination of samples in the study using a non probability sampling design with a purposive sampling method. The research was conducted by characterizing the qualitative and quantitative morphological characters from fruit, leaves, and trees. The characters identified are 81 characters. Characterization of the durian types was made based on the guidebook description from Bioversity International. Quantitative data was used for descriptive statistical analysis. Qualitative data analysis was carried out using the Unweighted Pair Group Method with Arithmetic Averaging (UPGMA) method using NTSYS version 2.02 software. The results of the research showed that local superior durians from Kasembon and Ngantang Subdistrict had diverse characteristics. Diversity are in the character of fruits, leaves, and trees. The genetic distance among superior durians is 0.109–0.343. Some of durian accessions observed such as Vodca, and Kunir Amad have the potential to be registered as regional assets or as local superior cultivars.

KEYWORDS: Characteristic, durian, local, morphology.

INTRODUCTION

Durian (*Durio zibethinus*) is a climacteric, seasonal tropical fruit of Southeast Asia. The ripe durian fruit owing to its unique taste and aroma is considered locally the king of fruits (Srianta et al., 2012), contain high in essential micro and macro nutrients, minerals and

vitamins (Ho & Bhat, 2015), provide phytoestrogens, anti inflammatory, and antioxidants (Zannah et al., 2022). Durian is a type of annual plant, and evergreen leaves, but it has a period of flushing, which is a period of growing new leaves and generally occurs after the fruiting period is over (Ashari, 2017).

Durian production in Indonesia reaches 1.169 million tons annually, East Java being the largest durian producer with 289,000 tons (Agriculture Ministry, 2019), provides additional income for the community (Matius et al., 2018; Prasetyo et al., 2018). In Indonesia at least there are 84 varieties have been released by the Ministry of Agriculture, based on their morphological characteristics, and originate. The wide variety of cultivars is an important source of germplasm as selection material for breeding durian and becoming superior local fruit cultivars from each region (Yulita & Nurnianjari, 2010). Efforts to identify superior types of durian from this region to various areas carried out through inventory activities, includes exploration and identification, is an out of the field activity collect data on types of durian in the area, genotype morphology characteristic expected to reveal potential seed of plant to be developed in another region, also is used as a reference to introduce all types durian in this area to a wider scope (Yuniastuti, Anggita, et al., 2018; Yuniastuti, Nandariyah, et al., 2018).

Characterization and relationship among them are very important for plant breeding program (Hariyati et al., 2013). First select a good character edible fruit, flesh color, water content, attractive flesh aroma, soft aril texture, fair flesh creaminess, sweet and sweet with bitter after-taste flesh. These two subdistricts are known as centers of local durians in Malang Regency which have wide diversity of local cultivars with delicious taste and more expensive than other types but have not been reported yet. This research was conducted to determine the genetic diversity and characterization of local superior durians in Kasembon and Ngantang Subdistricts.

MATERIALS AND METHODS

The research was conducted in peak season from January to August 2018 in Subdistricts of Kasembon 7°47'02", 112°18'32" and altitude of 500-721 m asl and Ngantang -7°51'23", 112°22'09" and altitude of 199-539 m asl.

95

Table 1. Names and subdistricts of superior durian found.

No.	Durian name	Initials	Origin subdistrict
1.	Kendi	KD	Ngantang
2.	Kunir Amad	KA	Ngantang
3.	Vodka	VK	Ngantang
4.	Manalagi Kuning	MK	Ngantang
5.	Duri Tajam	DT	Ngantang
6.	Getuk	GT	Ngantang
7.	Ranti	RT	Ngantang
8.	Selaput Udang	SU	Ngantang
9.	Toni	TN	Ngantang
10.	Aroma Kweni	AK	Ngantang
11.	Joko	JK	Ngantang
12.	Tarum	TR	Ngantang
13.	Ketan	KT	Ngantang
14.	Gipat	GP	Ngantang
15.	Klenting Kuning	KK	Kasembon
16.	Lima Ratus	LR	Kasembon
17.	Duri In	DI	Kasembon
18.	Brojo	BJ	Kasembon

96

97

98

99

100

101

102

103

104

105

106


107

108

109

Diversity exists in three levels, namely genetic, species, and ecosystem diversity. Morphological characteristics are based on the appearance of phenotypes, namely fruit shape, fruit spines, fruit flesh taste, flowers, leaf colour, and other characteristics (Rivero-Guerra, 2011; Sunaryo et al., 2015; Sundari et al., 2015; Daryono & Maryanto, 2017).

Durian fruit shape is globose (7 of 18 individuals) in general (Figure 2). Characters of shape of the fruit apex are: mammiform, convex, truncate, depressed, and pointed. Shape of the fruit base are: depressed, convex, truncate, and necked. Fruit spine shape are: convex, concave, pointed concave, and conical. Fruit spine density are: intermediate and sparse. Fruit rind color, are: yellow, yellowish green, greenish yellow, and brown. Flesh color are: yellow, orange, and white (Figure 3). Local superior durians mostly have intermediate easiness of splitting, thin-rinded durian fruit has a longer shelf life, and has a more edible portion of fruit (Fitmawati et al. 2015).



110

111

112

113

114

Figure 2. Fruit shape (a) globose; (b) oblate; (c) star; (d) obovoid; (e) oval; (f) oblong; (g) ovoid.




Figure 3. Flesh color, a. Yellow, b. Orange, and c. White.

Durian individuals with aborted and immature seeds have the potential to develop as superior durians without seeds and enlarge the edible portion. Durian had seed shape variations (Figure 4), with seed coat color intensity light and dark.




Figure 4. Seed Shape, a. Spheroid, b. Ellipsoid, c. Oblong, d. Ovoid, e. Obovoid, and f. Irregular.

The 18 durian types were varied in leaf form, generally leaf blade shape is elliptic (16 of 18 individuals). Most of the leaf attitude is drooping at 45° (17 of 18 individuals), the crown shape of the tree was 2, its growth habit variants of the durian tree were 3, most of the tree growth habit is intermediate (15 of 18 individuals). Durian tree age range from 11 until there are more than 100 years. Tree height range between 10.1-30.1 meters.

The organoleptic test showed that most of the local durians had moderate flesh aroma, soft aril texture, non-juicy aril, medium presence of fibre, fair flesh creaminess, slightly sticky flesh, sweet and sweet with bitter after-taste flesh with total sugar range between 34.96 ± 0.33 – $48.38 \pm 0.46\%$. It caused by genetic differences, environmental factors and the possibility of fruit ages at different harvest times (Hadiati et al., 2016), the sweetness level is influenced by level of fruit maturity and water content (Belgis et al. 2016)

Propagation by seeds will produce offspring that vary with the properties of the parent plant, because durian is a cross pollinating plant (Bumrungsri et al., 2009). According to (Indriyani et al. 2012), maternal parent influence the character of fruit (length, skin thickness, number of seeds, seed weight per fruit) and percentage of aborted seeds. There are many hybrids which are suitable for selection because durian is an open pollination plant (Prihatini et al. 2016), this is consistent with the results of this study. Exploration results for local durians in the two sub-districts show a high level of diversity as most durians grow from

seeds that are the result of crossing different combinations of parent trees. This increases the genetic diversity of durian plants, giving us the opportunity to obtain superior accession with high heterosis to use as mother trees and develop through vegetative propagation. However, the consequences of seed propagation will make the durian population heterogeneous, resulting individual inferior phenotypes that are less profitable for farmers.

Clustering of local durians

Cluster analysis was carried out based on qualitative data, showed that morphological similarity value of 18 durian species was 0.657–0.891 (Figure 5). Local superior durians in Kasembon and Ngantang Subdistricts based on the qualitative characters of fruit, leaves and trees, are divided into 2 Main Clusters, Ranti's accession was identified as the only individual with the lowest similarity coefficient (0.657) with the other 17 accessions, so it split from the two main cluster that formed, as it bears little morphological resemblance to other accessions. Grouping between accessions is not always related with origin, first group consists of various accessions originating from both subdistrict, Kendi and Duri In are the two accessions that show the highest morphological similarity (0.891). Grouping occurs because these accessions have similarities in morphological characters which can be caused by natural cross pollination of durian plants with the help of bats, birds and several species of beetles (Sritongchuay et al., 2016; Stewart & Dudash, 2017; Wayo et al., 2018). The highest similarity value was found between Kendi and Duri In durian (0.891), while the lowest similarity value was found between durian Ranti and 17 other durians (0.657) so it has not been identified. This is in accordance with the results of the study (Santoso et al., 2017) and (Sundari et al., 2021) that the fewer similarities, smaller similarity value or greater the genetic distance. Based on morphological observations (leaf and tree characters), genetic distance between durians is 0.109–0.343.




Figure 5. The dendrograms of Local superior durian Kasembon and Ngantang Subdistricts

	KD	KA	VK	MK	DT	GT	RT	SU	TN	AK	JK	TR	KT	GP	KK	LR	DI	BJ
KD	1.000																	
KA	0.745	1.000																
VK	0.673	0.655	1.000															
MK	0.727	0.782	0.673	1.000														
DT	0.873	0.800	0.745	0.800	1.000													
GT	0.873	0.745	0.691	0.691	0.836	1.000												
RT	0.636	0.673	0.600	0.655	0.691	0.655	1.000											
SU	0.818	0.691	0.636	0.782	0.782	0.782	0.618	1.000										
TN	0.800	0.800	0.636	0.764	0.818	0.800	0.709	0.782	1.000									
AK	0.800	0.709	0.636	0.745	0.764	0.745	0.655	0.818	0.818	1.000								
JK	0.800	0.673	0.673	0.673	0.745	0.764	0.618	0.836	0.800	0.764	1.000							
TR	0.782	0.764	0.782	0.745	0.818	0.782	0.636	0.709	0.764	0.727	0.691	1.000						
KT	0.727	0.745	0.691	0.818	0.727	0.709	0.691	0.745	0.818	0.800	0.727	0.764	1.000					
GP	0.745	0.800	0.618	0.782	0.764	0.709	0.673	0.745	0.818	0.818	0.709	0.691	0.782	1.000				
KK	0.764	0.727	0.636	0.764	0.745	0.709	0.618	0.745	0.691	0.727	0.673	0.673	0.764	0.764	1.000			
LR	0.745	0.818	0.636	0.727	0.782	0.764	0.709	0.727	0.818	0.800	0.727	0.782	0.727	0.764	0.691	1.000		
DI	0.891	0.764	0.709	0.764	0.891	0.855	0.655	0.836	0.873	0.836	0.818	0.764	0.782	0.836	0.764	0.782	1.000	
BJ	0.782	0.818	0.636	0.673	0.727	0.745	0.673	0.745	0.764	0.727	0.727	0.764	0.709	0.745	0.709	0.727	0.764	1.000

Figure 6. The resemblance matrix of local superior durian characters in Kasembon and Ngantang Subdistricts.

Differences and similarities in the emergence of outside morphology of a plant species can be used to determine the proximity of kinship. Environmental factors also influence the expression of these characteristics, even if they are only temporary (Hafizah et al., 2018).

CONCLUSIONS

The results of the research showed that local superior durians from Kasembon and Ngantang had diverse characters in the character of fruits, leaves and trees. The genetic distance between 18 durian accessions is 0.109–0.343 on the fruit, leaf and tree characters. Some of durian accessions observed such as Vodca, and Kunir Amad have the potential to be registered as regional assets or as local superior cultivars. This research was carried out observing agromorphology and organoleptics, so the results are still influenced by the environment.

ACKNOWLEDGEMENTS

Thanks are conveyed to Brawijaya University and durian farmers in the Ngantang and Kasembon subdistricts for their permission to characterize existing durian plants.

REFERENCES

- Ashari, S. (2017). *Durian: king of the fruit*. UB Press.
- Belgis, M., Wijaya, C. H., Apriyantono, A., Kusbiantoro, B., & Yuliana, N. D. (2016). Physicochemical differences and sensory profiling of six lai (*Durio kutejensis*) and four durian (*Durio zibethinus*) cultivar indigenous indonesia. *International Food Research Journal*, 23(4), 1466–1473.
- Bioversity International. (2007). *Descriptors for durian (Durio zibethinus Murr.)*. Bioversity International.

- Bumrungsri, S., Sripaoraya, E., Chongsiri, T., Sridith, K., & Racey, P. A. (2009). The pollination ecology of durian (*Durio zibethinus*, Bombacaceae) in southern Thailand. *Journal of Tropical Ecology*, 25(1), 85–92. <https://doi.org/10.1017/S0266467408005531>
- Daryono, B. S., & Maryanto, S. D. (2017). *Keanekaragaman dan Potensi Sumber Daya Genetik Melon*. Gadjah Mada University Press.
- Direktorat Budidaya Tanaman Buah. (2010). *Pedoman Standar Penilaian Durian*. Direktorat Jendral Hortikultura Kementerian Pertanian.
- Fitmawati, Aisyah, L., & Iriani, D. (2015). Comparative study of anatomical structure of durian perikarp (*Durio zibethinus* Murr.) shelf life and not shelf life from Bengkalis Island, Riau Province. In A. Soemargono, Mulyati, S. Hadiati, Martias, A. Sutanto, N. L. P. Indriyani, & Jumjunidang (Eds.), *Nusantara II. Bukittinggi 23–25 September 2015* (pp. 631–640).
- Hadiati, S., Nasution, F., & Kuswandi. (2016). Karakterisasi dan Evaluasi Koleksi Sumber Daya Genetik Durian Berdasarkan Karakter Morfologi Buah. *Bul. Plasma Nutfah*, 22(1), 1–10.
- Hafizah, R. A., Adawiyah, R., Harahap, R. M., Hannum, S., & Santoso, P. J. (2018). Aplikasi Marka SSR Pada Keanekaragaman Genetik Durian (*Durio zibethinus* Murr.) Di Kabupaten Deli Serdang, Sumatra Utara. *Al-Kauniyah: Jurnal Biologi*, 11(1), 49–56. <https://doi.org/10.15408/kauniyah.v11i1.5668>
- Handayani, F., & Rahayu, S. P. (2017). Short Communication: Assessment of genetic diversity in Lai (*Durio kutejensis*) local cultivars of Batuah (Indonesia) using ISSR marker. *Biodiversitas Journal of Biological Diversity*, 18(2), 525–529. <https://doi.org/10.13057/biodiv/d180212>
- Hariyati, T., Kusnadi, J., & Arumingtyas, E. L. (2013). Genetic diversity of hybrid durian resulted from cross breeding between *Durio kutejensis* and *Durio zibethinus* based on random amplified polymorphic DNAs (RAPDs). *American Journal of Molecular Biology*, 03(03), 153–157. <https://doi.org/10.4236/ajmb.2013.33020>
- Ho, L.-H., & Bhat, R. (2015). Exploring the potential nutraceutical values of durian (*Durio zibethinus* L.) – An exotic tropical fruit. *Food Chemistry*, 168, 80–89. <https://doi.org/10.1016/j.foodchem.2014.07.020>
- Indriyani, N. L. P., Hadiati, S., Nasution, F., Sudjijo, E., & Irawati, Y. (2012). Maternal and Paternal Effect on the Characters of Durian (*Durio Zibethinus* Murr.) Fruit from Cross-Pollination. *Journal of Fruit and Ornamental Plant Research*, 20(2), 23–33. <https://doi.org/10.2478/v10290-012-0012-x>

- 235 Kementerian Pertanian. (2019). *Produksi Durian Menurut Provinsi , Tahun 2015-2019*.
 236 Wwww. Pertanian.Go.Id.
- 237 Matius, P., Tjwa, S. J. M., Raharja, M., Sapruddin, S., Noor, S., & Ruslim, Y. (2018). Plant
 238 diversity in traditional fruit gardens (munaans) of Benuaq and Tunjung Dayaks tribes of
 239 West Kutai, East Kalimantan, Indonesia. *Biodiversitas Journal of Biological Diversity*,
 240 19(4), 1280–1288. <https://doi.org/10.13057/biodiv/d190414>
- 241 Prasetyo, B., Chikmawati, T., Walujo, eko baroto, & Amzu, E. (2018). Ethnoecology: The
 242 traditional landscape of Osing Tribe in Banyuwangi, Indonesia. *Biodiversitas Journal of*
 243 *Biological Diversity*, 19(6), 2003–2009. <https://doi.org/10.13057/biodiv/d190604>
- 244 Prihatini, R., Ihsan, F., & Indriyani, N. L. P. (2016). Genomic Profiling of F 1 Hybrids of
 245 Durian (*Durio zibethinus*) Revealed by RAPD-PCR. *Journal of Horticultural Research*,
 246 24(2), 69–76. <https://doi.org/10.1515/johr-2016-0022>
- 247 Rivero-Guerra, A. O. (2011). Morphological Variation within and Between taxa of the
 248 *Santolina rosmarinifolia* L. (Asteraceae: Anthemideae) Aggregate. *Systematic Botany*,
 249 36(1), 171–190. <https://doi.org/10.1600/036364411X553261>
- 250 Santoso, P. J., Granitia, A., Indriyani, N. L. P., & Pancoro, A. (2016). Analisis Lokus dan
 251 Keragaman Sumber Daya Genetik Durian (*Durio sp.*) Berdasarkan Marka Mikrosatelit.
 252 *Jurnal Hortikultura*, 26(1), 9. <https://doi.org/10.21082/jhort.v26n1.2016.p9-20>
- 253 Santoso, P. J., Pancoro, A., Suhandono, S., & Aryantha, I. N. P. (2017). Development of
 254 Simple-Sequence Repeats Markers from Durian (*Durio zibethinus* Murr. cultv. Matahari)
 255 Genomic Library. *AGRIVITA Journal of Agricultural Science*, 39(3), 257–265.
 256 <https://doi.org/10.17503/agrivita.v39i3.1171>
- 257 Srianta, I., Hendrawan, B., Kusumawati, N., & Blanc, P. J. (2012). Study on durian seed as a
 258 new substrate for Angkak production. *International Food Research Journal*, 19(3), 941–
 259 945.
- 260 Sritongchuay, T., Kremen, C., & Bumrungsri, S. (2016). Effects of forest and cave proximity
 261 on fruit set of tree crops in tropical orchards in Southern Thailand. *Journal of Tropical*
 262 *Ecology*, 32(4), 269–279. <https://doi.org/10.1017/S0266467416000353>
- 263 Stewart, A. B., & Dudash, M. R. (2017). Flower- visiting bat species contribute unequally
 264 toward agricultural pollination ecosystem services in southern Thailand. *Biotropica*,
 265 49(2), 239–248. <https://doi.org/10.1111/btp.12401>
- 266 Sunaryo, W., Hendra, M., Rudarmono, Suprpto, H., Pratama, A. N., & Rahman. (2015).
 267 Exploration and identification of Lai Durian, new highly economic potential cultivars
 268 derived from natural crossing between *Durio zibethinus* and *Durio kutejensis* in East

- Kalimantan. *Asian Journal of Microbiology, Biotechnology and Environmental Sciences*, 17(2), 365–371.
- Sundari, Arumingtyas, E. L., Hakim, L., & Azrianingsih, R. (2015). Exploration and Morphological Character Identification of Local Durian (*Durio zibethinus* Murr.) from Tidore Island, North Maluku. *Proceeding of 6th ICGRC*, 1–4.
- Sundari, Mas'Ud, A., Wahyudi, D., Arumingtyas, E. L., Hakim, L., & Azrianingsih, R. (2021). Genetic diversity of local durian from Tidore Island based on morphological and molecular data for tropical fruit conservation in North Maluku. *IOP Conference Series: Earth and Environmental Science*, 739(1). <https://doi.org/10.1088/1755-1315/739/1/012073>
- Wayo, K., Phankaew, C., Stewart, A. B., & Bumrungsri, S. (2018). Bees are supplementary pollinators of self-compatible chiropterophilous durian. *Journal of Tropical Ecology*, 34(1), 41–52. <https://doi.org/10.1017/S0266467418000019>
- Yulita, K. S., & Nurnianjari, M. (2010). Keragaman Genetik Beberapa Klon Durian (*Durio zibethinus* Murray) asal Jawa Barat berdasarkan Sidik Random Amplified Polimorphic DNA. *Jurnal Berita Biologi*, 10(3), 269–275.
- Yuniastuti, E., Anggita, A., Nandariyah, & Sukaya. (2018). Local durian (*Durio zibethinus* murr.) exploration for potentially superior tree as parents in Ngrambe District, Ngawi. *IOP Conference Series: Earth and Environmental Science*, 142(1). <https://doi.org/10.1088/1755-1315/142/1/012029>
- Yuniastuti, E., Nandariyah, N., & Bukka, S. R. (2018). Karakterisasi Durian (*Durio zibethinus*) Ngrambe di Jawa Timur, Indonesia. *Caraka Tani: Journal of Sustainable Agriculture*, 33(2), 136. <https://doi.org/10.20961/carakatani.v33i2.19610>
- Zannah, F., Kamaliah, Pramudiyanti, Ayatusaadah, & Hidayati, N. (2022). Exploration of the Potential of Local Plants of *Melastoma malabathricum* Fruit for Food Fortification. *Journal of Tropical Life Science*, 12(3), 333–338. <https://doi.org/10.11594/jtls.12.03.06>