

In Press, Pre-Proof Version

Sample storage and fasting times affect serum and plasma concentrations of metabolites in fasted and non-fasted broiler chickens

Lucas Wachholz¹, Cleison de Souza¹, Clauber Polese¹, Jomara Broch¹, André Sanches de Avila¹, Vaneila Daniele Lenhart Savaris¹, Nilton Rohloff Junior¹, Tânia Luiza Köhler¹, Jansller Luiz Genova^{2*}, and Ricardo Vianna Nunes

ABSTRACT

Sample storage and fasting times leads to some changes of blood metabolite in broilers. Therefore, a study was conducted with the aim to assess the influence of storage and fasting times in serum and plasma fractions on glucose, total cholesterol, triacylglycerols, aspartate aminotransferase (AST), alanine aminotransferase (ALT), and gamma-glutamyl transferase (GGT) concentrations in broilers. A total of 70 male broiler chickens fasted at 7 times (0, 2, 4, 6, 8, 10, and 12 h) to collect blood fractions (serum and plasma) stored at -20 °C for 0, 30, and 60 days. Glucose and GGT were affected by fasting times×blood fraction. Serum glucose concentration decreased linearly ($\cong 2.48 \text{ mg dL}^{-1}$), whereas total cholesterol and plasma GGT increased linearly ($\cong 0.92 \text{ mg dL}^{-1}$ and $\cong 0.19 \text{ IU L}^{-1}$, respectively) with fasting time. There was a quadratic effect on plasma glucose and serum GGT (maximum at 3.95 h and minimum at 5.22 h of fasting, respectively), and triacylglycerol (minimum at 8.75 h of fasting) and ALT concentrations (maximum at 8.45 h of fasting). Glucose, total cholesterol, AST, ALT, GGT concentrations were higher in serum, while triacylglycerols were higher in plasma. Glucose concentration had the lowest values at 30 days, while ALT was higher on day 0. However, GGT concentrations were lower on days 0 and 30. **Samples of plasma for glucose**, ALT, and GGT stored at -20 °C for long periods should be avoided. In addition, serum samples and 6 h fasting are recommended for the assessment of blood biochemical metabolites in broilers.

Keywords: Blood fractions, Broiler, Fasting, Liver enzyme, Sample storage.

INTRODUCTION

The assessment of biochemical parameters in poultry research can be better targeted, allowing the measurement of the concentration of several blood biochemical constituents that can be used as indicators of metabolic disorders, nutritional status, and in the diagnosis of

¹Department of Animal Science, Postgraduate Program in Animal Science, Universidade Estadual do Oeste do Paraná, Marechal Cândido Rondon, PR 85960-000, Brazil.

²Department of Animal Science, Postgraduate Program in Animal Science, Universidade Federal de Viçosa, MG 85960-000, Brazil.

*Corresponding author; e-mail: jansller.genova@ufv.br

33 diseases. Experiments involving broiler chickens commonly aim to assess bird performance,
34 promoting satisfactory feed conversion without compromising the metabolism and health
35 integrity of the birds. The changes that occur in the metabolism of birds can be caused by the
36 inclusion of ingredients, additives, chemotherapeutics, or other components of the diet (Hagan
37 *et al.*, 2022), in addition to environmental effects, which can alter the nutritional status and
38 health of the animals. Therefore, blood biochemical assessments are an important tool for the
39 diagnosis of diseases and metabolic disorders, providing efficient, fast, and safe diagnoses.

40 According to Gattani *et al.* (2016), the measurement of blood concentrations of glucose,
41 total cholesterol, triacylglycerol, aspartate aminotransferase (AST), alanine aminotransferase
42 (ALT), and gamma-glutamyl transferase (GGT) activities can be used to aid in the diagnosis of
43 numerous metabolic disorders in broiler chickens. However, there is a wide divergence in the
44 conditions under which blood samples are taken for analysis, as well as no standardization of
45 the postprandial fasting period performed before blood collection. Córdova-Noboa *et al.* (2018)
46 did not fast birds to collect blood samples, but Sadeghi *et al.* (2014), Behboudi *et al.* (2016),
47 Zakaria *et al.* (2017), and Swarna *et al.* (2018) performed fasting between 2 to 12 h prior sample
48 collection. Taking into account these variations in fasting times, it is not possible to state that
49 the results obtained in biochemical blood analyses will not be influenced by a long postprandial
50 fasting period, as recently demonstrated in a study conducted by Wachholz *et al.* (2023).

51 Biochemical parameters can be measured in serum samples, or plasma obtained with the
52 use of anticoagulants. The difference between these two processes is that in plasma, a larger
53 volume of supernatant is obtained compared to serum (Lumeij, 2008). Some divergence in the
54 use of serum or plasma in research results in a lack of standardization of results. Some authors
55 used samples of serum to measure blood biochemical parameters (Chand *et al.*, 2018, Gilani *et al.*
56 *et al.*, 2018, Rehman *et al.*, 2018, Subhani *et al.*, 2018), while others assessed in plasma (Sharideh
57 *et al.*, 2016, Yang *et al.*, 2017, Zhang *et al.*, 2017, Kim *et al.*, 2019).

58 Regarding the storage times that each sample can be subjected to before the biochemical
59 analyses, it is important to consider that the samples will usually be collected at different times
60 from the day of analysis because factors such as distance between the laboratory and the
61 experimental facility, time to obtain sufficient volume of samples, and transportation period,
62 can delay the date of the measurements of the biochemical variables in blood (Livesey *et al.*,
63 2008). This is supported by the study conducted by Wachholz *et al.* (2023), who assessed the
64 effect of time and storage condition on biochemical metabolites in serum or plasma samples of
65 broiler chickens and observed significant changes in the concentrations of glucose, total
66 cholesterol, triglycerides, AST, ALT and GGT.

67 Here, a study was conducted based on the hypothesis that both independent variables tested
68 can significantly alter blood metabolite concentrations and the results in broiler chicken
69 experiments. Therefore, this study aimed to assess the influence of storage and fasting times in
70 serum and plasma fractions on the concentrations of glucose, total cholesterol, triacylglycerols,
71 AST, ALT and GGT in broiler chickens.

72

73 **MATERIALS AND METHODS**

74 **Place of study, birds, handling, housing, and diets**

75 The present study was conducted at Universidade Estadual do Oeste do Paraná (Unioeste),
76 Marechal Cândido Rondon, PR, Brazil. The University Animal Ethics Committee approved the
77 experiment under number 23/20. A total of 70 45-day-old Cobb 500[®] male broiler chickens
78 were used, with an average body weight of 3,072±859 g. The birds were raised from 1 to 42
79 days of age, receiving water and diet *ad libitum*, and the same care to management, lighting and
80 ambient temperature recommended by the lineage manual. The diets for each phase (starter,
81 grower, and finisher) were corn-soybean based supplemented with industrial amino acids,
82 isonutritional and isoenergetic, and according to the nutritional requirements proposed by
83 Rostagno *et al.* (2017).

84 At 42 days of age, the birds were assigned in a completely randomized design in 7 pens
85 replicates (1.76 m²), with a masonry floor covered with 10 cm of pine wood shavings. Each pen
86 was composed of 10 birds and a density of 5.7 birds per m². The pens were equipped with a
87 tubular feeder and nipple drinkers. The facility was equipped with electrical elements, hoods,
88 evaporative pads to assist cooling and air exchange.

89

90 **Blood sampling, preparation and analysis**

91 After 3 days of adaptation at 45 days-old, the birds were fasted for 1 h, then fed for 30 min.
92 This procedure was adopted so that all birds had the same postprandial feeding condition. After
93 this period of feeding, the first blood collection was performed. At every 2-h interval, within a
94 total period of 12 h (0, 2, 4, 6, 8, 10, and 12 h of fasting), blood was collected by puncture of
95 the ulnar vein from 1 bird per pen ($n = 7$ birds per fasting time randomly selected). Every 7
96 birds were used only for a single sampling at a given fasting time. Blood collection was
97 performed with the birds in the decubitus position lateral, using specific vacuum collection
98 tubes (Vacutainer[®], Curitiba, PR, Brazil) with a capacity of 10 mL, specific adapter and 25×0.8
99 mm needles (21G 1”) (Labor Import brand, Maringá, PR, Brazil).

100 Two tubes of approximately 4 mL each were collected within the allowable for species and
101 body weight (Kelly and Alworth, 2013). The first blood sample was collected to obtain the
102 serum in a tube (BD Vacutainer[®], Curitiba, PR, Brazil) with clot activator (silica powder)
103 blasted on the tube wall to accelerate the process of coagulation. The second sample collected
104 was to obtain plasma in a tube (BD Vacutainer[®], Curitiba, PR, Brazil) with 5 mg of sodium
105 fluoride as a glycolysis inhibitor and 4 mL EDTA-K₃ anticoagulant. These two draws collected
106 per bird were consecutive. After collection, the samples remained for 15 min at room
107 temperature in a horizontal position and then were centrifuged (Centrifuge Kasvi K14-4000,
108 Kasvi, São Paulo, SP, Brazil) at 2,500 g for 10 min at room temperature.

109 After centrifugation and separation of serum and plasma, the samples were identified and
110 divided into three aliquots as technical triplicates, which were placed in 2 mL microtubes
111 (Eppendorf[®] brand, Minispin[®], Hamburg, Germany). An aliquot was immediately sent to the
112 laboratory for analysis. The other aliquots (two microtubes as technical duplicates) were stored
113 at -20 °C for times of 30 and 60 days for further analysis (Wachholz *et al.*, 2023).

114 Within each storage times, the samples were thawed under refrigeration (4 °C) and kept in
115 a refrigerator for 24 h. Before performing the analysis, the samples were centrifuged in an
116 Eppendorf microcentrifuge (Eppendorf[®] brand, Minispin[®], Hamburg, Germany) to remove
117 fibrin formation. Biochemical analyzes were performed using commercial kits and calibrators
118 (Elical II multiparametric Calibrator, ref. CALI-0550), and measurement standards for birds
119 (Elitrol I normal multiparametric control, ref. CONT-0060) (Elitech Clinical Systems, ELITech
120 Group, Paris, France) in automatic calibration spectrophotometer (Elitech[®] brand, Flexor
121 EL200 model, Puteaux, France).

122 The determination of glucose concentration (Glucose PAP) was performed by Trinder's
123 method, enzymatic colorimetric kinetic (Trinder, 1969), total cholesterol was performed by
124 Trinder's method, enzymatic colorimetric endpoint (Allain *et al.*, 1974), triacylglycerols
125 (Triglycerides SL New), enzymatic colorimetric endpoint (Fossati and Prencipe, 1982), AST
126 and ALT were performed according to International Federation of Clinical Chemistry method
127 without pyridoxal phosphate, kinetic, UV (Schuman *et al.*, 2002a, Shuman *et al.*, 2002b), and
128 GGT (GAMMA GT plus) was performed according the procedure Glupa-C substrate method,
129 kinetic (Schuman *et al.*, 2002c).

130

131 **Statistical procedures**

132 Data were analyzed considering blood fraction (serum and plasma), fasting (0, 2, 4, 6, 8,
133 10, and 12 h) and storage times of the sample (0, 30, and 60 days) as fixed effects. Residual

134 error was considered as a random factor. Each bird belonging to the same pen was considered
135 an experimental unit. Data were subjected to normality analysis using the Shapiro-Wilk test.
136 Afterward, the two-way analysis of variance (ANOVA) was performed, considering the
137 isolated effects and interactions between the studied factors: fasting or storage times×blood
138 fractions (serum and plasma) as fixed factors. All tests were performed at 5% significance level.
139 In case of significance, F-test was performed for the blood fraction, regression analysis for
140 fasting times (linear or quadratic models), and Tukey's post hoc test for storage times. The triple
141 interaction was not tested. All statistical procedures were performed using the GLM procedure
142 of SAS University Edition.

143

144 RESULTS AND DISCUSSION

145 Glucose concentrations were influenced by the interaction between fasting times×blood
146 fractions (Table 1). Serum glucose concentration decreased linearly with fasting time, reducing
147 $\cong 2.48 \text{ mg dL}^{-1}$ every 2 h of fasting. Total cholesterol concentrations increased linearly with
148 prolonged fasting time, increasing $\cong 0.92 \text{ mg dL}^{-1}$ (Figure 1). There was a quadratic effect on
149 the concentrations of plasma glucose, with maximum concentrations at 3.95 h of fasting, and
150 on triacylglycerol concentrations, with minimum concentration at 8.75 h of fasting. Glucose
151 and total cholesterol concentrations were higher in serum than in plasma, while triacylglycerol
152 concentrations were higher in plasma compared to serum. Glucose concentrations were
153 influenced by storage times, with the lowest values at 30 days of storage compared to 0 and 60
154 days.

155 The linear decreasing effect shown in serum glucose and quadratic for plasma glucose is
156 explained by the decrease of glycogen in the liver not being enough to maintain blood glucose
157 concentrations, causing its gradual reduction (Rodrigues *et al.*, 2017). The increase in total
158 cholesterol concentration and the decrease in triacylglycerols is due to lipid mobilization, as
159 lipids enter the bloodstream and are consequently transported by very-low-density lipoproteins,
160 causing a greater circulation of total cholesterol in the blood and, hence, the use of
161 triacylglycerols by the tissues (Lumeij, 2008).

162 In addition, blood concentrations of triacylglycerols are also influenced by fasting times
163 because when a negative energy balance occurs, bird metabolism is stimulated to mobilize
164 circulating triacylglycerols to provide fatty acids and glycerol via β -oxidation and glycolysis,
165 respectively, which are used for energy production, lowering blood triacylglycerols
166 concentrations (Coelho *et al.*, 2013). These results are in agreement with Vosmerova *et al.*

167 (2010), who observed that birds exposed to prolonged fasting times showed reduced glucose
168 and triacylglycerol concentrations.

169 The storage times of serum and plasma can lead to a reduction in glucose concentrations.
170 The present study showed variation in glucose concentrations caused by storage even at -20 °C.
171 This suggested that the temperature used during storage was not sufficient to keep glucose
172 concentrations stable for 60 days. The results of this study corroborate with the findings of
173 Cuhadar *et al.* (2013) and Clark *et al.* (1990), who stored serum samples from humans at -20
174 °C for 30 and 60 days, and observed a decrease in glucose concentrations.

175 The results observed for total cholesterol and triacylglycerols demonstrate that these
176 analytes can remain stable for 60 days at a temperature of -20 °C without any change. According
177 to Stokes *et al.* (1986), and Tiedink and Katan (1989), cholesterol and triacylglycerols present
178 in plasma and serum do not change their concentrations stored at -20 °C for 19 to 27 weeks.
179 The stability of triacylglycerols for long storage periods was also observed by Shimizu and
180 Ichihara (2019) storing serum samples for 2 months at -20 °C.

181 Although the differences found between the concentrations of biochemical analytes in the
182 blood fraction are statistically significant, Picheth *et al.* (2001) reported that these differences
183 between fractions for glucose do not represent chemically significant differences, pointing out
184 that both fractions can be used analytically for glucose.

185 Gamma-glutamyl transferase concentrations were affected by the interaction between
186 fasting times×blood fractions (Table 2). Plasma GGT concentrations increased linearly with
187 prolonged fasting time, increasing $\cong 0.19$ IU L⁻¹ (Figure 1). There was a quadratic effect on the
188 concentrations of serum GGT, with a minimum concentration at 5.22 h of fasting. In addition,
189 there was a quadratic effect on ALT concentrations, with a maximum concentration at 8.45 h
190 of fasting. Aspartate aminotransferase, ALT and GGT concentrations were higher in serum than
191 in plasma. Alanine aminotransferase and GGT concentrations were influenced by storage times.
192 It was observed that ALT concentration was higher on day 0 compared to the other storage
193 times. However, GGT concentrations were lower on days 0 and 30 than on day 60.

194 The quadratic response for ALT enzyme activity corroborates the results described of Veiga
195 *et al.* (1978), who reported that fasting time increases ALT concentrations in birds due to the
196 greater hepatic influx of gluconeogenic substrates, but less correlated with hepatic
197 gluconeogenesis than AST activity. The occurrence of a quadratic effect with the increase in
198 serum GGT concentration, and the linear increase in plasma GGT is because the metabolism of
199 the animals in prolonged fasting triggers the protein catabolism and, consequently, increases
200 the transport of amino acids. This highlights the main function of GGT, which according to Yu

201 and Long (2016), has the function of acting in the extracellular environment by catalyzing the
202 conversion of glutathione into glutamate dipeptides or cysteine.

203 Another metabolite stable to storage is AST, which did not change its concentration during
204 the 2 months of storage at -20 °C. According to Kaneko *et al.* (2008), this enzyme is stable to
205 storage during freezing. This response was also observed by Thoresen *et al.* (1995) and Oliveira
206 *et al.* (2011), who studied the same enzyme, but in blood samples from dogs and lambs,
207 respectively. However, ALT concentrations tend to decrease with storage times (Kaneko *et al.*,
208 2008). This decrease in blood ALT concentration is the result of enzyme denaturation as
209 reported by Ikeda *et al.* (2015) in a study conducted with human blood samples, who observed
210 enzyme denaturation and a decrease in its concentrations at -20 °C due to its instability at this
211 temperature.

212 Storage times can negatively interfere with the results of blood analyses in experiments
213 because researchers can often make conclusions based on results influenced by storage times.
214 The concentrations of glucose, total cholesterol, AST, ALT, and GGT evaluated in the current
215 study were higher in the serum sample compared to those observed in the plasma. This higher
216 concentration in serum can be attributed to the action that low molecular weight anticoagulants,
217 such as sodium fluoride, can exert on the samples. The osmotic effect of these anticoagulants
218 tends to remove an amount of water from inside the erythrocytes and transfer it to the plasma,
219 which would cause a dilution of the plasma concentrations of some metabolites (Alper *et al.*,
220 1974). According to Grande *et al.* (1964), this can cause dilution of the plasma constituents,
221 resulting in lower values in this blood fraction and, therefore, an anticoagulant that does not
222 interfere with the analysis results should be chosen. In addition, another important point to note
223 is that an osmotic redistribution between cells and plasma can occur, and this can interfere with
224 the analyte results.

225 The use of sodium fluoride as an anticoagulant can interfere analytically in
226 spectrophotometric assessments because this anticoagulant can cause fibrin formation in
227 samples during their collection (Fernandez *et al.*, 2013, Al-Kharusi *et al.*, 2014, Bonetti *et al.*,
228 2016). Thus, after plasma separation, the formation of hemolysis must be observed and, if
229 possible, to perform a new collection or make a higher centrifugal force.

230 Gamma-glutamyl transferase concentrations showed different effects on blood fraction
231 during the storage times. The difference in GGT concentrations between serum and plasma, as
232 well as a decrease in metabolite values in plasma analysis, are due to the use of sodium fluoride
233 anticoagulant, which reduces GGT activity (Burtis *et al.*, 2012).

234 **Du et al. (2023)** examined serum blood metabolites in broilers at 49-day-old on restricted
235 feeding and 16-h fasting and found lower average values than our findings for total cholesterol
236 (67.14 vs. 136 mg dL⁻¹), triacylglycerols (7.92 vs. 42 mg dL⁻¹), and glucose (151 vs. 233 mg
237 dL⁻¹) and ALT (2.06 vs. 9.51 IU L⁻¹) compared to the 12-h fasting in our study. However, the
238 values for AST (587 vs. 412 IU L⁻¹) and GGT (24.78 vs. 22.59 IU L⁻¹) were higher than those
239 of the present study. Corroborating the results of the present study, Hagan *et al.* (2022) reported
240 serum reference values of 197 to 299 mg dL⁻¹ for glucose, 129 to 297 mg dL⁻¹ for total
241 cholesterol, and average values of 96.15 mg dL⁻¹ for triacylglycerol concentrations in 56-day-
242 old Cobb and Ross broiler chickens fasted for 12 h.

243 Based on the criteria assessed in the current study, the results indicated that the
244 determination of AST is not influenced by the sample storage and fasting times. Total
245 cholesterol and triacylglycerol measurements can be performed on samples stored for up to 60
246 days at -20 °C. Plasma glucose, ALT, and GGT analyses stored at -20 °C for long periods should
247 be avoided. In addition, serum samples and 6 h fasting are recommended for the assessment of
248 blood biochemical metabolites in broilers.

249

250 **ACKNOWLEDGEMENTS**

251 The authors gratefully acknowledge the Coordenação de Aperfeiçoamento de Pessoal de Nível
252 Superior (CAPES), Fundação Araucária and the Conselho Nacional de Desenvolvimento
253 Científico e Tecnológico (CNPq).

254

255 **REFERENCES**

256 Al-Kharusi, A., Al-Lawati, N., Al-Kindi, M. and Mula-Abed, W. A. 2014. Are tubes containing
257 sodium fluoride still needed for the measurement of blood glucose in hospital laboratory
258 practice?. *Oman Med. J.*, **29**: 404-409.

259 Allain, C. C., Poon, L. S., Chan, C. S., Richmond, W. F. P. C. and FU, P. C. 1974. Enzymatic
260 determination of total serum cholesterol. *Clin. Chem.*, **20**: 470-475.

261 Alper, C. 1974. Specimen collection and preservation. In: *Clinical Chemistry, Principles and*
262 *Techniques*, (Eds.): Henry, R. J., Cannon, D. C. and Winkelman, J. W. Harpper and Row
263 Publishers, New York, PP. 373-388.

264 Behboudi, H., Esmailipour, O., Mirmahmoudi, R. and Mazhari, M. 2016. The Influence of
265 drinking water containing lemon juice and thyme supplemented diet on performance and
266 some blood parameters of broilers under heat stress. *Iran. J. Appl. Anim. Sci.*, **6**: 169-174.

- 267 Burtis, C. A., Ashwood, E.R., Bruns, D. E., Sawyer, B. G. 2008. Tietz fundamentals of clinical
268 chemistry. 6th Edition. Saunders: St Louis, Missouri.
- 269 Bonetti, G. Cancelli, V., Coccoli, G., Piccinelli, G., Brugnani, D., Caimi, L. and Carta, M. 2016.
270 Which sample tube should be used for routine glucose determination?. *Prim. Care Diabetes*,
271 **10**: 227-232.
- 272 Chand, N., Naz, S., Rehman, Z. and Khan, R. U. 2018. Blood biochemical profile of four fast-
273 growing broiler strains under high ambient temperature. *Appl. Biol. Chem.*, **61**: 273–279.
- 274 Clark, M. L., Humphreys, S. M. and Frayn, K. N. 1990. Stability of plasma glucose during
275 storage. *Ann. Clin. Biochem.*, **27**: 373-377.
- 276 Coelho, M., Oliveira, T. and Fernandes, R. 2013. Biochemistry of adipose tissue: an endocrine
277 organ. *Arch. Med. Sci.*, **9**: 191-200.
- 278 Cuhadar, S., Koseoglu, M., Atay, A. and Dirican, A. 2013. The effect of storage time and
279 freeze-thaw cycles on the stability of serum samples. *Biochem. Med.*, **23**: 70-77.
- 280 Córdova-Noboa, H. A., Oviedo-Rondón, E. O., Sarsour, A. H., Barnes, J., Sapkota, D., López,
281 D., Gross, L., Rademacher-Heilshorn, M. and Braun, U. 2018. Effect of guanidinoacetic
282 acid supplementation on live performance, meat quality, pectoral myopathies and blood
283 parameters of male broilers fed corn-based diets with or without poultry by-products. *Poult.*
284 *Sci.*, **97**: 2494-2505.
- 285 Du, P., Wang, H., Shi, X., Zhang, X., Zhu, Y., Chen, W. and Huang, Y. 2023. A comparative
286 study to determine the effects of breed and feed restriction on glucose metabolism of
287 chickens. *Anim. Nutr.*, **13**: 261-269.
- 288 Fernandez, L., Jee, P., Klein, M. J., Fischer, P., Perkins, S. L. and Brooks, S. P. 2013. A
289 comparison of glucose concentration in paired specimens collected in serum separator and
290 fluoride/potassium oxalate blood collection tubes under survey 'field' conditions. *Clin.*
291 *Biochem.*, **46**: 285-288.
- 292 Fossati, P. and Prencipe, L. 1982. Serum triglycerides determined colorimetrically with an
293 enzyme that produces hydrogen peroxide. *Clin. Chem.*, **28**: 2077-2080.
- 294 Gattani, A., Pathak, A., Kumar, A., Mishra, V. and Bhatia, J. S. 2016. Influence of season and
295 sex on hemato-biochemical traits in adult turkeys under arid tropical environment. *Vet.*
296 *World*, **9**: 530-534.
- 297 Gilani, S. M. H., Zehra, S., Galani, S. and Ashraf, A. 2018. Effect of natural growth promoters
298 on immunity, and biochemical and haematological parameters of broiler chickens. *Trop. J.*
299 *Pharm. Res.*, **17**: 627-633.

- 300 Grande, F., Amatuzio, D. S. and Wada. S. 1964. Cholesterol measurement in serum and in
301 plasma. *Clin. Chem.*, **10**: 619-626.
- 302 Hagan, J. K., Hagan, B. A., Ofori, S. A. and Etim, N. N. 2022. Haematological and serum
303 biochemical profiles of two broiler strains fed rations with varying levels of palm kernel oil
304 residue. *Ghana. J. Anim. Sci.*, **13**: 30-41.
- 305 Ikeda, K., Ichihara, K., Hashiguchi, T., Hidaka, Y., Kang, D., Maekawa, M., Matsumoto, H.,
306 Matsushita, K., Okubo, S., Tsuchiya, T. and Furuta, K. 2015. Evaluation of the short-term
307 stability of specimens for clinical laboratory testing. *Biopreserv. Biobank*, **13**: 135-143.
- 308 Kaneko, J. J., Harvey, J. W., Bruss, M. L. 2008. Clinical biochemistry of domestic animals. 6th
309 Edition. Academic Press, San Diego.
- 310 Kelly, L. M. and Alworth, L. C. 2013. Techniques for collecting blood from the domestic
311 chicken. *Lab. Animal*, **42**: 359-61.
- 312 Kim, J. H., Choi, H. S., Goo, D., Park, G. H., Han, G. P., Delos Reyes, J. and Kil, D. Y. 2019.
313 Effect of dietary melamine concentrations on growth performance, excreta characteristics,
314 plasma measurements, and melamine residue in the tissue of male and female broiler
315 chickens. *Poult. Sci.*, **98**: 3204-3211.
- 316 Livesey, J. H., Ellis, M. J. and Evans, M. J. 2008. Pre-analytical requirements. *Clin. Biochem.*
317 *Rev.*, **29**: 11-15.
- 318 Lumeij, J. T. 2008. Avian Clinical Biochemistry. In: *Clinical Biochemistry of Domestic*
319 *Animals*, (Eds.): Kaneko, J. J., Harvey, J.W. and Bruss, M. L. Academic Press, San Diego,
320 PP. 839-872.
- 321 Oliveira, F. S., Falbo, M. K., Sandini, I. E. and Ishiy, L. E. 2011. Effect of freezing and storage
322 time of blood serum of lambs in the determination of biochemical parameters. *Semin. Cienc.*
323 *Agrar.*, **32**: 717-722.
- 324 Picheth, G., Jaworski, M. C. G., Pinto, A. P., Kikuti, M. Y., Scartezini, M., Alcântara, V. M.
325 and Fadel-Picheth, C. M. T. 2001. Fluoride-plasma compared with serum in blood glucose
326 determination. *Rev. Bras. Anál. Clín.*, **33**: 167-170.
- 327 Rehman, Z., Chand, N., Khan, R. U., Naz, S. and Alhidary, I. A. 2018. Serum biochemical
328 profile of two broilers strains supplemented with vitamin E, raw ginger (*Zingiber officinale*)
329 and L-carnitine under high ambient temperatures. *S. Afr. J. Anim. Sci.*, **48**: 935-942.
- 330 Rodrigues, D. R., Café, M. B., Jardim Filho, R. M., Oliveira, E., Trentin, T. C., Martins, D. B.
331 and Minafra, C. S. 2017. Metabolism of broilers subjected to different lairage times at the
332 abattoir and its relationship with broiler meat quality. *Arq. Bras. Med. Vet. Zootec.*, **69**: 733-
333 741.

- 334 Rostagno H. S., Albino L. F. T., Hannas M. I., Donzele J. L., Sakomura N. K., Perazzo F. G.,
335 Saraiva A., Teixeira M. L., Rodrigues P. B., Oliveira R. F. D., Barreto S. L. D. T. and Brito
336 A. O. 2017. Brazilian tables for poultry and pigs: food composition and nutritional
337 requirements. 4th Edition. UFV, Viçosa.
- 338 Sadeghi, A. A., Mohamadi-Saei, M. and Ahmadvand, H. 2014. The efficacy of dietary savory
339 essential oil on reducing the toxicity of aflatoxin b1 in broiler chicks. *Kafkas Univ. Vet. Fak.*
340 *Derg.*, **20**: 481-486.
- 341 Schumann, G., Bonora, R., Ceriotti, F., Férard, G., Ferrero, C. A., Franck, P. F., Gella, F. J.,
342 Hoelze, W., Jorgensen, P. J., Kanno, T., Kessner, A., Klauke, R., Kristiansen, N., Lessinger,
343 J. M., Linsinger, T. P. J., Misaki, H., Panteghini, M., Pauwels, J., Schiele, F., Schimmel, H.
344 G., Weidemann, G. and Siekmann, L. 2002a. IFCC primary reference procedures for the
345 measurement of catalytic activity concentrations of enzymes at 37°C. Part 5. Reference
346 procedure for the measurement of catalytic activity concentration of aspartate-
347 aminotransferase [L-aspartate: 2-oxoglutarate-aminotransferase (AST), EC 2.6.1.1]. *Clin.*
348 *Chem. Lab. Med.*, **40**: 725-733.
- 349 Schumann, G., Bonora, R., Ceriotti, F., Férard, G., Ferrero, C. A., Franck, P. F., Gella, F. J.,
350 Hoelze, W., Jorgensen, P. J., Kanno, T., Kessner, A., Klauke, R., Kristiansen, N., Lessinger,
351 J. M., Linsinger, T. P. J., Misaki, H., Panteghini, M., Pauwels, J., Schiele, F., Schimmel, H.
352 G., Weidemann, G. and Siekmann, L. 2002b. IFCC primary reference procedures for the
353 measurement of catalytic activity concentrations of enzymes at 37°C. Part 4. Reference
354 procedure for the measurement of catalytic activity concentration of alanine
355 aminotransferase [L-alanine: 2-oxoglutarate aminotransferase (ALT), EC 2.6.1.2]. *Clin.*
356 *Chem. Lab. Med.*, **40**: 718-724.
- 357 Schumann, G., Bonora, R., Ceriotti, F., Férard, G., Ferrero, C. A., Franck, P. F., Gella, F. J.,
358 Hoelze, W., Jorgensen, P. J., Kanno, T., Kessner, A., Klauke, R., Kristiansen, N., Lessinger,
359 J. M., Linsinger, T. P. J., Misaki, H., Panteghini, M., Pauwels, J., Schiele, F., Schimmel, H.
360 G., Weidemann, G. and Siekmann, L. 2002c. IFCC primary reference procedures for the
361 measurement of catalytic activity concentrations of enzymes at 37°C. Part 6. Reference
362 procedure for the measurement of catalytic activity concentration of γ -glutamyltransferase
363 [(γ -glutamyl)-peptide: amino acid γ -glutamyltransferase (GGT), EC 2.3.2.2]. *Clin. Chem.*
364 *Lab. Med.*, **40**: 734-738.
- 365 Sharideh, H., Zhandi, M., Zaghari, M. and Akhlaghi, A. 2016. Effect of dietary zinc oxide and
366 phytase on the plasma metabolites and enzyme activities in aged broiler breeder hens. *Iran.*
367 *J. Vet. Res.*, **9**: 263-270.

- 368 Shimizu, Y. and Ichihara, K. 2019. Elucidation of stability profiles of common chemistry
369 analytes in serum stored at six graded temperatures. *Clin. Chem. Lab. Med.*, **57**: 1388-1396.
- 370 Subhani, Z., Shahid, M., Hussain, F. and Khan, J. A. 2018. Efficacy of *Chlorella pyrenoidosa*
371 to ameliorate the hepatotoxic effects of aflatoxin b1 in broiler chickens. *Pak. Vet. J.*, **38**: 13-
372 18.
- 373 Stokes, Y. M., Salmond, C. E., Carpenter, L. M. and Welby, T. J. 1986. Stability of total
374 cholesterol, high-density-lipoprotein cholesterol, and triglycerides in frozen sera. *Clin.*
375 *Chem.*, **32**: 995-999.
- 376 Swarna, C. L., Srinivas, G. and Rao, S. V. 2018. Effect of unsaturated to saturated fatty acids
377 ratio of supplemental fat in the diet with or without l-carnitine on performance of broiler
378 chicken. *Indian J. Poult. Sci.*, **53**: 150-155.
- 379 Thoresen, S. I., Tverdal, A., Havre, G. and Morberg, H. 1995. Effects of storage time and
380 freezing temperature on clinical chemical parameters from canine serum and heparinized
381 plasma. *Vet. Clin. Pathol.*, **24**: 129-133.
- 382 Tiedink, H. G. and Katan, M. B. 1989. Variability in lipoprotein concentrations in serum after
383 prolonged storage at -20° C. *Clin. Chim. Acta*, **180**: 147-155.
- 384 Trinder, P. 1969. Determination of glucose in blood using glucose oxidase with an alternative
385 oxygen acceptor. *Ann. Clin. Biochem.*, **6**: 24-27.
- 386 Veiga, J. A., Roselino, E. S. and Migliorini, R. H. 1978. Fasting, adrenalectomy, and
387 gluconeogenesis in the chicken and a carnivorous bird. *Am. J. Physiol. Regul. Integr. Comp.*
388 *Physiol.*, **234**: 115-121.
- 389 Vosmerova, P., Chloupek, J., Bedanova, I., Chloupek, P., Kruzikova, K., Blahova, J. and
390 Vecerek, V. 2010. Changes in selected biochemical indices related to transport of broilers
391 to slaughterhouse under different ambient temperatures. *Poult. Sci.*, **89**: 2719-2725.
- 392 Wachholz, L., Genova, J. L., Polese, C., Broch, J., Savaris, V. D. L., Köhler, T. L. and Nunes,
393 R. V. 2023. Light intensity, blood fraction, fasting and storage time affect blood biochemical
394 metabolites in broiler chickens. *Semin. Cienc. Agrar.*, **44**: 2095-2112.
- 395 Yang, J. Y., Zhang, H. J., Wang, J., Wu, S. G., Yue, H. Y., Jiang, X. R. and Qi, G. H. 2017.
396 Effects of dietary grape proanthocyanidins on the growth performance, jejunum
397 morphology and plasma biochemical indices of broiler chicks. *Anim.*, **11**: 762-770.
- 398 Yu, X. and Long, Y. C. 2016. Crosstalk between cystine and glutathione is critical for the
399 regulation of amino acid signaling pathways and ferroptosis. *Sci. Rep.*, **6**: 1-10.

- 400 Zakaria, H. A., Jalal, M., Al-Titi, H. H. and Souad, A. 2017. Effect of sources and levels of
401 dietary zinc on the performance, carcass traits and blood parameters of broilers. *Rev. Bras.*
402 *Cienc. Avic.*, **19**: 519-526.
- 403 Zhang, Y. N., Wang, J., Qi, B., Wu, S. G., Chen, H. R., Luo, H. Y., Yin, D. J., Lu, F. J., Zhang,
404 H. J. and Qi, G. H. 2017. Evaluation of mango saponin in broilers: effects on growth
405 performance, carcass characteristics, meat quality and plasma biochemical indices. *Asian-*
406 *Australas J. Anim. Sci.*, **30**: 1143-1149.

407 **Table 1.** Concentrations (mg dL⁻¹) of glucose, total cholesterol and triacylglycerols in serum and
 408 plasma stored at different times in 45-day-old broiler chickens subjected to different fasting times.


Fasting times (h)	Glucose		Total cholesterol		Triacylglycerols	
	Serum	Plasma	Serum	Plasma	Serum	Plasma
0	260±12	233±15	130±13	114±10	76±19	78±19
2	260±18	236±13	134±13	118±13	48±10	52±12
4	258±21	255±17	141±14	136±13	47±12	53±13
6	247±21	239±12	138±12	123±12	41±14	42±11
8	241±11	232±10	132±11	121±12	38±6	41±08
10	231±03	219±10	146±14	127±11	32±7	33±06
12	238±21	228±17	145±14	128±13	42±5	43±06
Storage times (days)						
0	251±20	238±16 ^A	137±14	124±14	44±19	49±20
30	243±17	230±15 ^B	138±14	125±12	47±17	50±71
60	249±23	235±19 ^A	138±14	122±15	47±16	48±18
Average	248±20 ^a	234±17 ^b	138±14 ^a	124±14 ^b	46±17 ^b	49±18 ^a
SEM	0.961		0.765		0.847	
Fasting times	<0.001		<0.001 (L)		<0.001 (Q)	
Blood fraction	<0.001		<0.001		0.017	
Storage times	<0.001		0.501		0.364	
Fasting times×blood fraction	<0.001		0.061		0.899	
Linear	<0.001	<0.001				
Quadratic	<0.001	<0.001				
Storage times×blood fraction		0.848		0.594		0.355

409 ^{a-b} Lowercase letters in the same row differ by F test ($P < 0.05$); ^{A-B} Capital letters in the same column differ by
 410 Tukey's post hoc test ($P < 0.05$); SEM: Pooled standard error of the mean; L: Linear effect; Q: Quadratic effect.

411
 412 **Table 2.** Concentrations (IU L⁻¹) of aspartate aminotransferase (AST), alanine aminotransferase
 413 (ALT) and gamma-glutamyl transferase (GGT) in serum and plasma stored at different times
 414 in 45-day-old broiler chickens subjected to different fasting times.

Fasting times (h)	AST		ALT		GGT	
	Serum	Plasma	Serum	Plasma	Serum	Plasma
0	353±60	322±56	9.17±2.43	7.18±2.13	23.95±6.97	14.57±2.62
2	408±87	407±81	9.34±2.29	8.28±2.83	21.10±6.78	16.36±4.86
4	450±76	440±70	9.95±2.84	9.59±2.85	21.32±5.49	18.05±4.01
6	436±63	403±71	10.25±2.45	9.12±2.15	19.55±6.01	15.08±2.67
8	413±66	380±73	9.93±2.00	8.80±2.30	20.96±5.10	18.57±4.02
10	460±67	417±70	10.65±2.33	8.85±2.91	23.25±6.07	16.14±4.22
12	436±71	387±65	10.65±3.01	8.38±2.75	27.04±5.63	18.15±2.99
Storage times (days)						
0	409±71	388±65	11.14±2.40	9.76±2.26 ^A	16.80±3.85	16.66±4.08 ^B
30	431±74	397±71	9.40±2.42	8.02±2.64 ^B	16.48±3.69	16.03±4.43 ^B
60	426±85	396±94	9.44±2.37	8.04±2.64 ^B	24.22±6.45	16.82±4.32 ^A
Average	422±77 ^a	394±77 ^b	9.99±2.52 ^a	8.60±2.64 ^b	22.46±6.41 ^a	16.70±3.94 ^b
SEM	3.826		0.130		0.296	
Fasting times	0.140		<0.001(Q)		<0.001	
Blood fraction	<0.001		<0.001		<0.001	
Storage times	0.135		<0.001		0.009	
Fasting times×blood fraction	0.483		0.352		0.002	
Linear					0.032	0.004
Quadratic					<0.001	0.010
Storage times×blood fraction	0.703		0.998		0.005	
Unfolding					<0.001	0.851

415 ^{a-b} Lowercase letters in the same row differ by F test ($P < 0.05$); ^{A-B} Capital letters in the same column differ by
 416 Tukey's post hoc test ($P < 0.05$); SEM: Pooled standard error of the mean; L: Linear effect; Q: Quadratic effect.



417

418 **Figure 1.** Serum and plasma concentrations of glucose, total cholesterol, triacylglycerols,
 419 alanine aminotransferase and gamma-glutamyl transferase in broiler chickens at 45-day-old
 420 subjected to different fasting times (R²: coefficient of determination).
 421

421