

***Juglans regia* kernel powder supplementation in Broiler Chickens fed Aflatoxin-contaminated diets: Effect on growth, serum chemistry indices, immunoglobulin and pro-inflammatory cytokines**

Running title: Walnut kernel in chickens fed an aflatoxin-infected feed

O. D. Oloruntola^{1*}, A. B. Falowo¹, S. O. Ayodele², O. J. Olarotimi¹, S. A. Adeyeye³, C. O. Osowe⁴, D. A. Oloruntola⁵, F. A. Gbore¹ and F. S. Oladebeye¹

1. Department of Animal Science, Adekunle Ajasin University, Akungba Akoko, Nigeria.

2. Department of Agricultural Technology, The Federal Polytechnic, Ado Ekiti, Nigeria.

3. Department of Animal Health and Production Technology, Federal College of Agriculture, Akure, Nigeria.

4. Department of Animal Production and Health, The Federal University of Technology, Akure, Nigeria.

5. Department of Medical Laboratory Science, University of Medical Sciences, Ondo City, Nigeria.

***Corresponding author: olugbenga.oloruntola@aaua.edu.ng or oloruntoladavid@gmail.com**

ABSTRACT

The study aimed to investigate the impact of *Juglans regia* kernel powder (JKP) on broiler chickens subjected to Aflatoxin (AF)-contaminated diets during a 42-day feeding trial conducted in February and March 2022. A total of 240 one-day-old broiler chickens were divided into four dietary groups: Diet 1 (Control), Diet 2 (0.5 mg/kg AF), Diet 3 (0.5 mg/kg AF +250 mg/kg JKP), and Diet 4 (0.5 mg/kg AF+500 mg/kg JKP). Birds on Diet 2 exhibited a significantly lower ($P=0.01$) relative growth rate compared to other diets. JKP supplementation at 250 mg/kg (Diet 3) and 500 mg/kg (Diet 4) mitigated the negative impact of AF on growth. Birds on Diet 2 showed significantly lower ($P=0.01$) serum concentrations of total protein, albumin, and globulin compared to those on Diets 1, 3, and 4. Elevated levels of aspartate aminotransferase (AST) and creatinine in Diet 2 indicated liver and kidney damage. Alanine transaminase (ALT) concentrations in Diet 2 were higher ($P=0.01$) than Diets 1 and 4. Birds fed diet 2 had lower glucose levels ($P=0.01$) than diets 1 and 4. IgA levels in birds fed Diet 2 were lower ($P=0.03$) than those in the birds fed Diet 4. Birds fed diet 2 had considerably ($P<0.05$) lower IgE and IgG levels than birds fed diets 1 and 4. Nuclear Factor Kappa B (NFκB) was higher ($P=0.01$) in birds fed Diet 2 compared to other diets. Interleukin 6 (IL 6) concentration was significantly ($P=0.01$) higher in the birds fed Diet 2 than in the rest

40 diets. A recommended dietary supplementation of 250 mg/kg JKP is suggested based on
41 observed ameliorative effects.

42 **Keywords:** aflatoxicosis, botanicals, broiler chickens, immunity, inflammation.

43

44 INTRODUCTION

45 According to reports, the recent surge in animal output in tropical and subtropical nations has
46 been attributed to an increase in the human population (Godfray *et al.*, 2010). Additionally,
47 broiler chicken production has been identified as a potential solution to the issue of animal
48 protein deficiency in these regions (Hatab *et al.*, 2019). However, broiler chickens face
49 challenges in reaching their genetic potential in tropical and subtropical climates due to factors
50 such as the scarcity of feedstuffs, heat stress, and contaminated feed (Kpomasse *et al.*, 2019).

51 Aflatoxin (AF), the most common mycotoxin, is produced by *Aspergillus flavus*, *A. nomius*,
52 and *A. parasiticus* (Morrison *et al.*, 2017). *A. flavus* produces four toxins (AFB1, AFB2, AFG1,
53 and AFG2) with similar chemical structures, but AFB1 is the most potent hepatotoxin and a
54 recognized hepatocarcinogen (Quezada *et al.*, 2000). The prevalence of warm and humid
55 conditions favorable for aflatoxin growth, combined with the limited effectiveness of feed
56 processing techniques in removing aflatoxins from contaminated diets due to their thermal
57 resistance, contributes to the prevalence of aflatoxicosis in the tropics and subtropics (Medina
58 *et al.*, 2017; Mahato *et al.*, 2019).

59 Aflatoxicosis in poultry induces anorexia, lethargy, stunted growth, reduced fertility, and
60 microbial stress, leading to impaired gut health, economic losses, and toxicity (Sarma *et al.*,
61 2017). Furthermore, aflatoxin damage extends to the liver and kidneys, resulting in impaired
62 immune function and an upregulation of proinflammatory gene expression (Quezada *et al.*,
63 2000; Li *et al.*, 2022). It was reported that AF induces the generation of intracellular Reactive
64 Oxygen Species (ROS) such as hydroxyl radicals, superoxide anions, and hydrogen peroxide
65 in mammalian cells (Sohn *et al.*, 2003; An *et al.*, 2017). This suggests the potential of
66 antioxidants to ameliorate the negative effects of AF toxicity in animals when fed AF-
67 contaminated diets.

68 The potential of medicinal plant-derived antioxidant dietary intake in ameliorating the
69 damage caused by oxidative stress through inhibiting the initiation or propagation of oxidative
70 chain reactions and quenching singlet oxygen and reducing agents has been reported (Cai *et*
71 *al.*, 2004; Baiano and del Nobile, 2015; Adegbeye *et al.*, 2020). Broadly speaking, these natural
72 antioxidants have anti-inflammatory, antiviral, antibacterial, and anticancer activities (Xu *et*
73 *al.*, 2017).

74 *Juglans regia* Linn is a potential nutraceutical and medicinal plant used traditionally to
75 address various maladies, including diarrhea, stomachaches, arthritis, asthma, and endocrine
76 problems like diabetes mellitus, thyroid dysfunctions, and cancer (Taha and Al-wadaan, 2021).
77 As documented by Oloruntola (2022), *Juglans regia* kernel powder contains saponins (43.49
78 mg/g), alkaloids (120.80 mg/g), flavonoids (14.72 mg/g), tannins (1.69 mg/g), phenol (35.93
79 mg/g), and steroids (4.84 mg/g), contributing to its nutraceutical properties. Recent study
80 (Oloruntola, 2022) have highlighted *Juglans regia* kernel powder's anti-inflammatory,
81 antioxidant, and anti-diabetic effects, encouraging its usage as a nutritional supplement for
82 feed.

83 Including green husk walnut powder (Mousavi Razi *et al.*, 2017) and walnut leaves (Popescu
84 *et al.*, 2020) in the diet has been reported to enhance the function of the broiler immune system
85 and promote gastrointestinal tract health. However, research on the dietary supplementation of
86 *Juglans regia* kernel powder in broiler nutrition is relatively scarce. Therefore, this study aims
87 to investigate the effects of *Juglans regia* kernel powder dietary supplementation on the
88 growth, serum chemistry indices, immunoglobulin, and pro-inflammatory cytokines of broiler
89 chickens fed aflatoxin-contaminated diets.

90

91 **MATERIALS AND METHODS**

92 **Animal Ethics, *Juglans regia* kernel powder, Aflatoxin B1, Experimental diets and Birds**

93 The broiler care and use procedures have obtained approval from the Department of Animal
94 Science's Animal Care and Use Committee at Adekunle Ajasin University, Akungba Akoko,
95 Nigeria. *Juglans regia* kernel powder (JKP) was produced, as previously detailed by
96 Oloruntola (2022). The *Juglans regia* fruits were sourced from villages in Akungba Akoko,
97 Nigeria. Raw kernels were carefully extracted, finely chopped, sparingly scattered, and air-
98 dried in the shade for 14 days. Subsequently, the dried kernels were milled to form *Juglans*
99 *regia* kernel powder (JKP), which was then stored for subsequent laboratory analysis.

100 The *Aspergillus flavus* (NRRL 3251) pure culture, maintained on potato dextrose agar, served
101 as the source of the aflatoxin. Autoclavable polypropylene bags containing 500 grams of maize
102 grits were heated to 121 degrees Celsius and exposed to a pressure of 120 kPa for 60 minutes.
103 Following inoculation with an *A. flavus* spore suspension, the autoclaved grit maize was
104 cultivated for seven days at a temperature of 28°C. After the fungus developed, the grit maize
105 was dried in a 70°C oven and ground into powder.

106 In formulating experimental diets with 0.5mg/kg AFB1 contamination, 100g of AFB1
107 cultured maize was carefully blended with 1kg of broiler feed and subsequently analyzed for

108 AFB1 concentration. The analysis indicated an AFB1 concentration of 17mg/kg.
109 Consequently, these findings were utilized to calculate the necessary amount of cultured maize
110 required for 1kg of broiler feed to achieve the targeted 0.5mg/kg AFB1 concentration. The
111 amount of aflatoxin (AF) in the blend of maize and broiler feed was measured in triplicate
112 using thin-layer chromatography (AOAC, 2010).

113 A baseline diet (Table 1) for the starter and finisher stages was produced following the
114 recommendations of the National Research Council (NRC, 1994). Subsequently, thin-layer
115 chromatography was employed to check the baseline diet for any AF that may have been
116 present (AF was not present in any significant amount). The proximate composition of the
117 baseline diets was investigated (AOAC, 2010), and the diets were split into four equal parts.
118 Each part was sufficiently contaminated with AF-maize powder, added JKP, and labeled as
119 necessary: Diet 1: Control; Diet 2: 0.5 mg/kg AF; Diet 3: 0.5 mg/kg AF +250 mg/kg JKP; Diet
120 4: 0.5 mg/kg AF+500 mg/kg JKP. The 0.02 mg/kg limit allowed by NAFDAC, the EU, the
121 USFDA, the CFIA, and ANAC was 25 times lower than the 0.5 mg AF/kg feed concentration
122 in the chicken diet used in this study (Burel *et al.*, 2009).

123 A total of 240 Cobb 500 broiler chickens that were 1 day old were randomly assigned to 4
124 diets, each having 6 replicates of 10 chickens. The experiment consisted of two phases: 1-21
125 days and 22-42 days. For the entire six-week testing period, both feed and water were freely
126 available.

127 **Determination of Relative Growth Rate, Serum Chemistry Indices, Immunoglobulin,** 128 **and Pro-inflammatory Cytokines**

129 **Measurement of Relative Growth Rate (RGR)**

130 At the onset of the feeding experiment (day 1) and upon its conclusion (day 42), the weights
131 of the broiler chicks were meticulously measured. The Relative Growth Rate (RGR) was
132 estimated using the formula published by Adebayo *et al.* (2020):

$$133 \text{RGR} = [(w_2 - w_1) / ((w_1 + w_2) / 2)] * 100.$$

134 Where:

135 w represents the initial weight of the broiler chickens before the experiment,

136 w represents the weight of the broiler chicks on the final day of the experiment.

137 Three randomly selected birds per replication were tagged, and approximately 10 ml of blood
138 samples were obtained using a syringe and needle from the brachial vein. The blood was drawn
139 into plain bottles, allowed to stand at room temperature for around 30 minutes, centrifuged at
140 3,000 rpm for 10 minutes, and the serum was then decanted into new plain bottles. The labeled

141 serum samples were stored at -20 °C until required for analysis of chemistry indices,
142 immunoglobulins, and pro-inflammatory cytokines.

143

144 **Serum Chemistry Indices Analysis**

145 A Reflectron® Plus 8C79 (Roche Diagnostic, GmbH Mannheim, Germany) with commercial
146 kits was employed to measure total protein, albumin, aspartate aminotransferase (AST), alanine
147 transferase (ALT), cholesterol, creatinine, and glucose (Oloruntola *et al.*, 2018). The
148 discrepancies between total protein and albumin were utilized to determine globulin.

149

150 **Immunoglobulin and Pro-inflammatory Cytokines Analysis**

151 Immunoglobulins A (IgA), E (IgE), G (IgG), and M (IgM) were determined using ELISA kits
152 from Fortress Diagnostics Limited, United Kingdom. Nuclear Factor Kappa B (NFK B) was
153 determined using a Rat NFKB-p65 ELISA kit from Elabscience Biotechnology Inc. USA.
154 Tumor Necrosis Factor Alpha (TNF α) was determined with an ELISA kit, also from
155 Elabscience Biotechnology Inc. USA, while Interleukin 6 (IL 6) was determined using a Rat
156 IL-6 ELISA kit from the same manufacturer.

157

158 **Statistical Data Analysis**

159 The obtained data were subjected to analysis of variance (ANOVA) using SPSS version 20. To
160 identify differences in treatment means, the Duncan multiple range test from the same
161 statistical program was employed (Oloruntola *et al.*, 2018).

162

163 **RESULTS**

164 Figure 1 illustrates the impact of *Juglans regia* kernel powder (JKP) supplementation on the
165 relative development of broiler chickens fed diets contaminated with aflatoxin (AF). Broiler
166 chickens on diet 2 (AF-contaminated) exhibited a significantly lower ($P<0.05$) relative growth
167 rate compared to those on control (diet 1) and other diets. Birds on diet 4 (AF + 500 mg/kg
168 JKP) displayed a relative growth rate comparable ($P>0.05$) to diet 3 (AF + 250 mg/kg JKP) but
169 significantly higher ($P<0.05$) than diet 1.

170 The results of JKP supplementation on the serum chemistry indices are presented in Table 2.
171 Birds on diet 2 showed significantly lower ($P<0.05$) serum concentrations of total protein,
172 albumin, and globulin compared to diets 1, 3, and 4. Additionally, diet 2 resulted in
173 significantly higher ($P<0.05$) levels of aspartate aminotransferase (AST) and creatinine
174 compared to the control and other diets. Alanine transaminase (ALT) concentrations in diet 2
175 were comparable ($P>0.05$) to diet 3 but significantly higher ($P<0.05$) than diets 1 and 4. Broiler

176 chickens on diet 2 had glucose levels comparable ($P>0.05$) to diet 3 but significantly lower
177 ($P<0.05$) than diets 1 and 4.

178 Table 3 shows the impact of JKP supplementation on the immunoglobulin levels of broiler
179 chickens fed diets contaminated with AF. Birds on diet 2 had significantly lower ($P<0.05$) IgA,
180 IgE, and IgG levels compared to birds on diets 1 and 4. IgA levels in diet 2 were comparable
181 to diets 1 and 3 but significantly lower than diet 4.

182 The effects of JKP supplementation on pro-inflammatory cytokines are presented in Table 4.
183 Nuclear Factor Kappa B (NFK B) and Interleukin 6 (IL 6) concentrations were significantly
184 higher ($P<0.05$) in birds on diet 2 compared to diets 1, 3, and 4. IL 6 concentrations were
185 similar in birds fed the control (diet 1) and diet 4.

186

187 **DISCUSSION**

188 *Juglans regia* kernels are rich in antioxidants, including polyphenols, omega-3 fatty acids, and
189 melatonin, providing potential benefits for anti-inflammatory responses and cardiovascular
190 health (Bhat *et al.*, 2023). Constituents such as ellagic acid, gallic acid, Alpha-Linolenic Acid
191 (ALA), and melatonin contribute to the nutraceutical profile of *Juglans regia* kernels, making
192 them a promising dietary supplement (Shah *et al.*, 2018).

193 Aflatoxin feed contamination (0.5 mg/kg) significantly reduced the relative growth rate of
194 broiler chickens, aligning with previous studies (Denli *et al.*, 2004; Denli *et al.*, 2009). Due to
195 the degradation of the digestive and metabolic efficiency of the birds exposed to AF dietary
196 contamination, the retarded growth rate was connected to decreased energy and poor protein
197 utilisation (Verma *et al.*, 2002; Denli *et al.*, 2009). However, supplementation with JKP,
198 especially at 250 and 500 mg/kg, mitigated the negative effects of AF contamination on growth,
199 suggesting a protective role for JKP in the digestive system and physiological processes. The
200 use of medicinal or herbal plants parts in controlling or preventing cases of toxicity has been
201 reported (Khafaga and Bayad, 2016; Aboelhassan *et al.*, 2018). Certain components present in
202 *Juglans regia* kernel, including bioactive compounds like saponins, alkaloids, flavonoids,
203 tannins, phenols, and steroids, may contribute to aflatoxin-binding properties (Oloruntola,
204 2022). These compounds may interact with aflatoxins, potentially reducing their absorption
205 and mitigating their adverse effects on the gastrointestinal tract (Pathaw *et al.*, 2022).
206 Nevertheless, this assertion is contingent upon further and more comprehensive research.

207 The blood total protein test determines the quantity of all proteins, specifically blood globulin
208 and albumin (Tothova *et al.*, 2016) and is one of the sensitive early biomarkers of poultry
209 exposure to aflatoxin B1 (Quezada *et al.*, 2000). Also, significant clinical problems such as

210 inflammatory illnesses, liver disorders, kidney disorders, malnutrition, and others were linked
211 to low total protein levels (Tothova *et al.*, 2016).

212 The reduction in blood concentrations of total protein, albumin, and globulin observed in this
213 investigation is consistent with the finding of Safameher (2008) that broiler chickens fed diets
214 containing 0.5 to 2.0 ppm AFB1/kg indicate a decrease in total serum protein concentration.
215 The blocking of RNA synthesis, followed by the inhibition of protein synthesis in the liver, and
216 ultimately the reduction in plasma protein concentration could be the reason for the decreased
217 serum protein concentration brought on by aflatoxin exposure (Del Bianchi *et al.*, 2005).
218 Furthermore, complications with the liver or kidneys (Tothova *et al.*, 2016) could be
219 responsible for the reported decrease in serum total protein concentration in this study. This is
220 supported by concurrent elevated serum AST, ALT, and creatinine recorded in the same group
221 of birds (diet 2). The elevated serum AST and ALT concentrations recorded in birds fed diet 2
222 indicate liver damage. This finding aligns with the observations of Tessari *et al.* (2010), who
223 documented elevated AST levels in birds fed 50 and 200 µg AFB1/kg. Additionally, Valchev
224 *et al.* (2014) reported increased ALT activity in broiler chickens fed 0.5 mg/kg AFB1.

225 In addition, records on the toxic effects of aflatoxin on blood parameters exhibited through
226 increased creatinine and uric acid were reported (Valchev *et al.*, 2014). Hence, the observed
227 elevation in serum creatinine concentration among birds fed aflatoxin-contaminated feed (diet
228 2) in this study underscores the potential peril of aflatoxin dietary contamination on the normal
229 physiological and anatomical functions of the kidney (Valchev *et al.*, 2014).

230 In a nutshell, the production of a reactive metabolite called AFB1- 8,9-epoxide, which is
231 formed quickly by the action of at least five members of the mixed-function oxidase family, is
232 the cause of aflatoxin's renal toxicity. AFB1-8,9-epoxide reacts with DNA to yield the 8,9-
233 dihydro-8-(N7-guanyl)-9- hydroxy aflatoxin B1 adduct (AFB1-N7-Gua), which has been
234 positively correlated with DNA strand breaks, hepatic tumor development, and the
235 development of renal lesions (O'Brien and Dietrich, 2004).

236 According to this study, broiler chickens fed AF-contaminated diets had lower blood glucose
237 levels, possibly attributed to aflatoxin's hepatotoxic effects, which cause problems with lipid
238 and carbohydrate metabolism (Rosa *et al.*, 2001; Basmacioglu *et al.*, 2005). This outcome was
239 consistent with data from Basmacioglu *et al.* (2005), who noted hypoglycemia in broiler
240 chickens fed a diet contaminated with 2 mg AF/kg feed.

241 Free radicals and reactive oxygen species (ROS) produced by mycotoxins harm cells (Marin
242 and Taranu, 2012). Aflatoxin-induced ROS generation can harm the cells of target organs like
243 the liver and kidney. In addition to an increase in lipid peroxidation metabolites in the liver and

244 kidney (Alpsoy and Yalvac, 2011) and a decrease in the cellular total antioxidant in birds, there
245 is a considerable shift in blood biochemical indices after this increase (Sirajudeen *et al.*, 2011).
246 Therefore, the observed ameliorative activities of JKP in the birds fed diets contaminated with
247 AF (0.5mg/kg) and supplemented with JKP (250 and 500 mg/kg) diets in this study about the
248 serum total protein, albumin, globulin, AST, ALT, creatinine, and glucose could be an outcome
249 of the nutraceutical and antioxidant activity of JKP. It has been claimed that JKP has
250 antioxidant, anti-inflammatory, and anti-diabetic characteristics and is a helpful nutraceutical
251 feed additive (Oloruntola, 2022).

252 JKP's ameliorative effects on broiler chickens fed diets contaminated with aflatoxin in this
253 study were consistent with those of curcumin (Damiano *et al.*, 2022) and aloe vera powder
254 (Seifi *et al.*, 2022) on poultry/birds fed diets containing aflatoxin.

255 It was discovered that aflatoxins impair the innate and acquired responses of the immune
256 system (Weaver *et al.*, 2013). The decreased concentrations of IgA, IgE, and IgA observed in
257 broiler chickens given a diet contaminated with aflatoxin in this study may be caused by the
258 dysregulation of dendritic cells' ability to present antigens and impaired cell-mediated
259 immunity as a result of aflatoxin exposure (Mehrzaad *et al.*, 2014). However, the improved
260 levels of IgA, IgE, and IgA in birds fed a supplemented diet in this study unveil the
261 immunomodulatory properties of phytochemicals or bioactive compounds in the JKP.
262 Inferentially, JKP supplementation stops the mechanisms leading to immune system
263 dysfunction typically linked to aflatoxin dietary contamination. As previously explained,
264 several dietary phytochemicals interact with immunological signal transduction pathways
265 connected to inflammation to exhibit immune modulatory actions (Zhao *et al.*, 2021).

266 The triggered Nuclear Factor Kappa B (NF- κ B) and interleukin 6 (IL-6) observed in the birds
267 fed the aflatoxin-contaminated diet in this study could be associated with the typical expression
268 of aflatoxicosis because exposure to aflatoxin frequently results in elevated reactive oxygen
269 species (ROS), oxidative stress, lipid peroxidation, apoptosis, mitochondrial dysfunction,
270 necrosis, and inflammatory response (Dai *et al.*, 2022). For instance, NF- κ B is one of the
271 several pathways that have been shown to support AFB1-mediated toxicity in mammalian cells
272 (Dai *et al.*, 2022), and according to Karunaweera *et al.* (2015), the activation of NF- κ B requires
273 the degradation of the inhibitor kappa B alpha and mediates the production of more than 500
274 genes, including tumor necrosis factor-alpha (TNF-alpha) and IL-6 (Yamashita *et al.*, 2014).

275 As recently reported, the administration of low doses of aflatoxin may also upregulate the
276 expression of NF- κ B, TNF- α , and IL-6, causing a significant inflammatory response in the
277 liver tissues (Guo *et al.*, 2022; Dai *et al.*, 2022). However, the identical or similar NF- κ B and

278 IL-6 gene expression observed in the birds fed aflatoxin-contaminated diets supplemented with
279 JKP and the control diet in this study further demonstrates the nutraceutical properties and the
280 activities of bioactive components of JKP. JKP achieves this by inhibiting the activation of
281 TLR4/MyD88, which is followed by the activation of NF-κB and its downstream IL-6, and
282 TNF-α genes' expression (Li *et al.*, 2022; Guo *et al.*, 2022). This outcome agrees with Li *et al.*
283 (2022) findings, which showed that curcumin supplementation slowed the expression of the
284 NF-κB and IL-6 genes.

285

286 CONCLUSIONS

287 Dietary supplementation with JKP at 250 and 500 mg/kg demonstrated ameliorative effects on
288 broiler chickens exposed to aflatoxin B1. The improvements in growth rate, serum chemistry
289 indices, immunoglobulins, and pro-inflammatory cytokines suggest the potential of JKP as a
290 nutraceutical feed supplement. A recommended dietary supplementation of 250 mg/kg JKP is
291 suggested for optimal broiler chicken production.

292

293 REFERENCES

- 294 1. Aboelhasan, D.M., Hafiz, N.A., Darwish, H.R., Shabana, M.E., Eshak, M.G.,
295 Hassanane, M.M., Farag, I.M., and Abdalla, A.M. (2018). Enhancing effects of *Moringa*
296 *oleifera* leaf extract on carcinogenic Aflatoxin B1-induced genetic alterations, haematotoxicity
297 and histological changes in liver and kidney of rats. *Bioscience Research*, 15(2), 814-833.
- 298 2. Adebayo, F.B., Adu, O.A., Chineke, C.A., Oloruntola, O.D., Omoleye, O.S., Adeyeye,
299 S.A., and Ayodele, S.O. (2020). The performance and hematological indices of broiler
300 chickens fed Chromium Picolinate and vitamin C supplemented diets. *Asian Journal of*
301 *Research in Animal and Veterinary Science*, 6(4), 54-61.
- 302 3. Adegbeye, M.J., Oloruntola, O.D., Asaniyan, E.K., Agunbiade, B., Oisagah, E.A. and
303 Ayodele, S.O. (2020). Pawpaw, Black Cumin, and Mustard Seed Meals Dietary
304 Supplementation in Broiler Chickens: Effect on Performance, Gut Microflora, and Gut
305 Morphology, *Journal of Agricultural Science and Technology*, 22(5): 1235-1246.
- 306 4. Alpsy, L. and Yalvac, M.E. (2011). Key roles of vitamins A, C, and E in aflatoxin B1-
307 induced oxidative stress. *Vitamins and Hormones*, 86, 287-305.
- 308 5. An, Y., Shi, X., Tang, X., Wang, Y., Shen, F., Zhang, Q., Wang, C., Jiang, M., Liu, M.,
309 and Yu, L (2017). Aflatoxin B1 Induces Reactive Oxygen Species-Mediated Autophagy and
310 Extracellular Trap Formation in Macrophages. *Frontiers of Cell and Infectious Microbiology*,
311 7, 53. doi: 10.3389/fcimb.2017.00053

- 312 6. AOAC. (2010). Official method of analysis, 18th edition. Association of Official
313 Analytical Chemists, Washington D.C.
- 314 7. Baiano, A., and del Nobile, M.A. (2015). Antioxidant compounds from vegetable
315 matrices: Biosynthesis, occurrence, and extraction systems. *Critical Reviews in Food Science*
316 *and Nutrition*, 56, 2053–2068. doi: 10.1080/10408398.2013.812059.
- 317 8. Basmacioglu, H., Oguz, H., Ergul, M., Col, R., and Birdane, Y.O. (2005). Effect of
318 dietary esterified glucomannan on performance, serum biochemistry and haematology in
319 broilers exposed to aflatoxin. *Czech Journal of Animal Science*, 50, 2005 (1): 31–39.
- 320 9. Bhat, A. A., Shakeel, A., Rafiq, S., Farooq, I., Malik, A. Q., Alghuthami, M. E.,
321 Alharthi, S., Qanash, H., and Alharthy, S. A. (2023). *Juglans regia* Linn.: A Natural Repository
322 of Vital Phytochemical and Pharmacological Compounds. *Life* (Basel, Switzerland), 13(2),
323 380. <https://doi.org/10.3390/life13020380>
- 324 10. Burel, S.D., Favrot, M.C., Fremy, J.M., Massimi, C., Prigent, P., Debongnie, L.P. and
325 Morgavi, D. (2009). Review of mycotoxin-detoxifying agents used as feed additives: mode of
326 action, efficacy and feed/food safety. EFSA-Q- pp.2009-00839, EFSA J, Dec 82012;
327 2012:564367. doi: 10.1100/2012/564367.
- 328 11. Cai, Y.Z., Luo, Q., Sun, M., and Corke, H. (2004). Antioxidant activity and phenolic
329 compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life*
330 *Science*, 74, 2157–2184. doi: 10.1016/j.lfs.2003.09.047.
- 331 12. Dai, C., Tian, E., Hao, Z., Tang, S., Wang, Z., Sharma, G., Jiang, H., and Shen, J.
332 (2022). Aflatoxin B1 Toxicity and Protective Effects of Curcumin: Molecular Mechanisms and
333 Clinical Implications. *Antioxidants* (Basel, Switzerland), 11(10), 2031.
334 <https://doi.org/10.3390/antiox11102031>
- 335 13. Damiano, S., Jarriyawattanachaikul, W., Girolami, F., Longobardi, C., Nebbia, C.,
336 Andretta, E., Lauritano, C., Dabbou, S., Avantaggiato, G., Schiavone, A., Badino, P., and
337 Ciarcia, R. (2022). Curcumin Supplementation Protects Broiler Chickens Against the Renal
338 Oxidative Stress Induced by the Dietary Exposure to Low Levels of Aflatoxin B1. *Frontiers of*
339 *Veterinary Science*, 8, 822227. doi: 10.3389/fvets.2021.822227
- 340 14. Del Bianchi, M., Oliveira, C.A.F., Albuquerque, R., Guerra, J.L., and Correa, B. (2005).
341 Effects of Prolonged Oral Administration of Aflatoxin B1 and Fumonisin B1 in Broiler
342 Chickens. *Poultry Science*, 84:1835–1840.
- 343 15. Denli, M., Blandon, J.C., Guynot, M.E., Salado, S., and Perez, J.F. (2009). Effects of
344 dietary AflaDetox on performance, serum biochemistry, histopathological changes, and

- 345 aflatoxin residues in broilers exposed to aflatoxin B1, *Poultry Science*, 88 (7), 1444-1451.
346 <https://doi.org/10.3382/ps.2008-00341>.
- 347 16. Denli, M., Okan, F., and Doran, F. (2004). Effect of conjugated linoleic acid (CLA) on
348 the performance and serum variables of broiler chickens intoxicated with aflatoxin B1. *South*
349 *African Journal of Animal Science*, 34 (2004), 97-103.
- 350 17. Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J.
351 F., Pretty, J., Robinson, S., Thomas, S. M., and Toulmin, C. (2010). Food security: The
352 challenge of feeding 9 billion people. *Science*, 3275(967), 812–818. 10.1126/science.118538
- 353 18. Guo, J., Yan, W.R., Tang, J.K., Jin, X., Xue, H.H., Wang, T., Zhang, L.W., Sun, Q.Y.,
354 and Liang, Z.X. (2022). Dietary phillygenin supplementation ameliorates aflatoxin B(1)-
355 induced oxidative stress, inflammation, and apoptosis in chicken liver. *Ecotoxicology and*
356 *Environmental Safety*, 236:113481. doi: 10.1016/j.ecoenv.2022.113481.
- 357 19. Hatab, A. A. , Cavinato, M. E. R. , and Lagerkvist, C. J. (2019). Urbanization, livestock
358 systems and food security in developing countries: A systematic review of the literature. *Food*
359 *Security*, 11, 279–299. 10.1007/s12571-019-00906-1
- 360 20. Karunaweera, N., Raju, R., Gyengesi, E., and Munch, G. (2015). Plant polyphenols as
361 inhibitors of NF-kappa B induced cytokine production a potential anti-inflammatory treatment
362 for Alzheimer's disease? *Frontiers in Molecular Neuroscience*, 8:24. doi:
363 10.3389/fnmol.2015.00024.
- 364 21. Khafaga, A.F., and Bayad, A.E. (2016). Impact of Ginkgo biloba extracts on
365 reproductive toxicity induced by single or repeated injection of cisplatin in adult male rats.
366 *International Journal Pharmacology*, 12: 340-350
- 367 22. Kpomasse, C.C., Oke, O.E., Houndonougbo, F.M., and Tona, K. (2021). Broiler
368 production challenges in the tropics: A review. *Veterinary Medicine and Science*, 7(3), 831–
369 842. <https://doi.org/10.1002/vms3.435>
- 370 23. Li, S., Liu R., Xia, S., Wei, G., Ishfaq, M., Zhang, Y., and Zhang, X. (2022). Protective
371 role of curcumin on aflatoxin B1-induced TLR4/RIPK pathway mediated-necroptosis and
372 inflammation in chicken liver. *Ecotoxicology and Environmental Safety*, 233, 113319.
373 <https://doi: 10.1016/j.ecoenv.2022.113319>.
- 374 24. Mahato, D.K., Lee, K.E., Kamle, M., Devi, S., Dewangan, K.N., Kumar, P. and Kang,
375 S.G. (2019). Aflatoxins in Food and Feed: An Overview on Prevalence, Detection and Control
376 Strategies. *Frontiers in Microbiology*, 10, 2266. <https://doi: 10.3389/fmicb.2019.02266>
- 377 25. Marin, D.E. and Taranu, I. (2012). Overview on aflatoxins and oxidative stress. *Toxin*
378 *Reviews*, 31(3), 32-43.

- 379 26. Medina, A., Gilbert, M. K., Mack, B. M., OBrian, G. R., Rodríguez, A., Bhatnagar, D.,
380 Payne, G., and Magan, N. (2017). Interactions between water activity and temperature on the
381 *Aspergillus flavus* transcriptome and aflatoxin B1 production. *International Journal of Food*
382 *Microbiology*, 256, 36–44. <https://doi.org/10.1016/j.ijfoodmicro.2017.05.020>
- 383 27. Mehrzad, J., Devriendt, B., Baert, K., and Cox, E. (2014). Aflatoxin B(1) interferes
384 with the antigen-presenting capacity of porcine dendritic cells. *Toxicology in Vitro*, 28:531–
385 537.
- 386 28. Mousavi Razi, B., Roostaei Ali Mehr, M., and Mohiti Asli, M. (2017). Effect of walnut
387 green husk (*Juglans regia*) powder on immune responses of broiler chickens. *Iranian*
388 *Veterinary Journal*, 13(2), 86-95. doi: 10.22055/ivj.2017.50827.1712
- 389 29. Morrison, D.M., Ledoux, D.R., Chester, L.F.B., and Samuels, C.A.N. (2017). A limited
390 survey of aflatoxins in poultry feed and feed ingredients in Guyana. *Veterinary Sciences*, 4 (4),
391 60. <https://DOI: 10.3390%2Fvetsci4040060>.
- 392 30. NRC (1994). Nutrient Requirements of Poultry. Ninth Revised Edition. The National
393 Academies Press. Pp. 29. <http://nap.edu/2114>
- 394 31. O'Brien, E., and Dietrich, D.R. (2004). Mycotoxins affecting the kidney. First
395 published in: Toxicology of Kidney. Ed. J.B Hook *et al.* Boca Raton: CRC Press. Pp. 895-936.
396 <http://www.crcpress.com/>
- 397 32. Oloruntola, O.D., Agbede, J.O., Onibi, G.E., Igbasan, F.A., Ayodele, S.O. Arogunjo,
398 M.A. and Ogunjo, S.T. (2018). Rabbits fed fermented cassava starch residue I: Effect on
399 performance and health status. *Archivos de Zootecnia*, 67(260), 578-586.
- 400 33. Oloruntola, O. D. (2022). *Juglans regia* kernel meal; A prospective nutraceutical feed
401 supplement. *Biotech Studies*, 31(2), 87-94. <http://doi.org/10.38042/biotechstudies.1222785>
- 402 34. Pathaw, N., Devi, K. S., Sapam, R., Sanasam, J., Monteshori, S., Phurailatpam, S.,
403 Devi, H. C., Chanu, W. T., Wangkhem, B., and Mangang, N. L. (2022). A comparative review
404 on the anti-nutritional factors of herbal tea concoctions and their reduction strategies. *Frontiers*
405 *in Nutrition*, 9, 988964. <https://doi.org/10.3389/fnut.2022.988964>
- 406 35. Popescu, R. G., Voicu, S. N., Pircalabioru, G. G., Ciceu, A., Gharbia, S., Hermenean,
407 A., Georgescu, S. E., Panaite, T. D., and Dinischiotu, A. (2020). Effects of Dietary Inclusion
408 of Bilberry and Walnut Leaves Powder on the Digestive Performances and Health of Tetra SL
409 Laying Hens. *Animals*, 10(5), 823. <https://doi.org/10.3390/ani10050823>
- 410 36. Quezada, T., Cuéllar, H., Jaramillo-Juárez, F., Valdivia, A.G. and Reyes, J.L (2000).
411 Effects of aflatoxin B1 on the liver and kidney of broiler chickens during development,

- 412 *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and*
413 *Endocrinology*, 125 (3), 265-272. [https://doi.org/10.1016/S0742-8413\(99\)00107-3](https://doi.org/10.1016/S0742-8413(99)00107-3).
- 414 37. Quezada, T., Cuellar, H., Jaramillo-Juarez, F., Valdivia, A.G. and Reyes, J.L. (2000).
415 Effects of aflatoxin B1 on the liver and kidney of broiler chickens during development.
416 *Comparative Biochemistry and Physiology Part C*, 125 (2000), 265 – 272.
- 417 38. Rosa, C.A., Miazzo, R., Magnoli, C., Salvano, M., Chiac, S.M., Ferrero, S., Saenz, M.,
418 Carvalho, E.C. and Dalcerro, A. (2001): Evaluation of the efficacy of bentonite from the south
419 of Argentina to ameliorate the toxic effects of aflatoxin in broilers. *Poultry Science*, 80, 139–
420 144.
- 421 39. Safameher, A. (2008). Effects of clinoptilolite on performance, biochemical parameters
422 and hepatic lesions in broiler chickens during aflatoxicosis. *Journal of Animal Veterinary*
423 *Advances*, 7, 381-388.
- 424 40. Sarma, U.P., Bhetaria, P.J., Devi, P. and Varma, A. (2017). Aflatoxins: implications on
425 health. *Indian Journal of Clinical Biochemistry*, 32 (2), 124-133. DOI: 10.1007%2Fs12291-
426 017-0649-2.
- 427 41. Seifi, S., Sadighara, P. and Mohajer, A. (2022). Protective effects of Aloe vera powder
428 supplementation on some quantitative and qualitative characteristics of egg, histopathological
429 changes and serum biochemistry of laying hens fed by Aflatoxin B1. *Veterinary Research*
430 *Forum*, 13 (4) 507 – 512. doi: 10.30466/vrf.2021.530920.3186.
- 431 42. Shah, U. N., Mir, J. I., Ahmed, N., Jan, S., and Fazili, K. M. (2018). Bioefficacy
432 potential of different genotypes of walnut *Juglans regia* L. *Journal of food science and*
433 *technology*, 55(2), 605–618. <https://doi.org/10.1007/s13197-017-2970-4>
- 434 43. Sirajudeen, M., Gopi, K., Tyagi, J.S., Moudgal, R.P. Mohan, J. and Singh, R. (2011).
435 Protective effects of melatonin in reduction of oxidative damage and immunosuppression
436 induced by aflatoxin B1-contaminated diets in young chicks. *Environmental Toxicology*, 26(
437 2), 153 -160.
- 438 44. Sohn, D.H., Kim, Y.C., Oh, S.H., Park, E.J., Li, X. and Lee, B.H. (2003).
439 Hepatoprotective and free radical scavenging effects of *Nelumbo nucifera*. *Phytomedicine* 10,
440 165–169. doi: 10.1078/094471103321659889
- 441 45. Taha, N.A. and Al-wadaan, M.A. (2021). Significance and use of walnut, *Juglans regia*
442 Linn: A review. *Advanced Journal of Microbiology Research*, 15 (1), 001-010.
- 443 46. Tessari, E.N., Kobashigawa, E., Cardoso, A.L., Ledoux, D.R., Rottinghaus, G.E. and
444 Oliveira, C. A. (2010). Effects of aflatoxin B(1) and fumonisin B(1) on blood biochemical
445 parameters in broilers. *Toxins*, 2(4), 453–460. <https://doi.org/10.3390/toxins2040453>

- 446 47. Tothova, C., Nagy, O. and Kovac, G. (2016). Serum proteins and their diagnostic utility
447 in veterinary medicine: a review. *Veterinarni Medicina*, 61, 2016 (9), 475–49.
- 448 48. Valchev, I., Kanakov, D., Hristov, T. and Lavarov, L. (2014). Investigations on the liver
449 function of broiler chickens with experimental aflatoxicosis. *Bulgarian Journal of Veterinary*
450 *Medicine*, 17(4), 302-313.
- 451 49. Valchev, I., Kanakov, D., Ts. Hristov, Ts., Lazarov, L., Binev, R., Grozeva, N. and
452 Nikolov, Y. (2014). Effects of experimental aflatoxicosis on renal function in broiler chickens.
453 *Bulgarian Journal of Veterinary Medicine*, 17(4), 314-324.
- 454 50. Verma, J., Swain, B.K. and Johri, T.S. (2002). Effect of various levels of aflatoxin and
455 ochratoxin A and combinations thereof on protein and energy utilisation in broilers. *Journal of*
456 *Science of Food and Agriculture*, 82, 1412-1417.
- 457 51. Weaver, A.C., See, M.T., Hansen, J.A., Kim, Y.B., De Souza, A.L. and Middleton, T.F.
458 (2013). The use of feed additives to reduce the effects of aflatoxin and deoxynivalenol on pig
459 growth, organ health and immune status during chronic exposure. *Toxins* (Basel), 5, 1261–
460 1281.
- 461 52. Xu, D.P., Li, Y., Meng, X., Zhou, T., Zhou, Y., Zheng, J., Zhang, J.J. and Li, H.B.
462 (2017). Natural Antioxidants in Foods and Medicinal Plants: Extraction, Assessment and
463 Resources. *International Journal of Molecular Sciences*, 18(1), 96.
464 <https://doi.org/10.3390/ijms18010096>
- 465 53. Yamashita, Y., Ueyama, T., Nishi, T., Yamamoto, Y., Kawakoshi, A., Sunami, S.,
466 Iguchi, M., Tamai, H., Ueda, K., Ito, T., Tsuruo, Y. and Ichinose, M. (2014). Nrf2-inducing
467 anti-oxidation stress response in the rat liver--new beneficial effect of lansoprazole. *PLoS one*,
468 9(5), e97419. <https://doi.org/10.1371/journal.pone.0097419>
- 469 54. Zhao, F., Ci, X., Man, X., Li, J., Wei, Z. and Zhang, S. (2021). Food-Derived
470 Pharmacological Modulators of the Nrf2/ARE Pathway: Their Role in the Treatment of
471 Diseases. *Molecules*, 26(4):1016. doi: 10.3390/molecules26041016.

472
473
474
475
476
477
478
479
480
481
482

483
484
485
486
487
488
489
490
491
492
493
494
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515


Table 1. Composition of the experimental diets.

Ingredients (%)	Starter phase	Finisher phase
Rice bran	0.00	3.02
Maize	50.36	58.36
Maize bran	3.00	0.00
Soy oil	1.00	1.00
Fish meal	3.00	3.00
Soybean meal	38.00	30.00
Bone meal	3.00	3.00
**Premix	0.31	0.31
Limestone	0.49	0.47
Salt	0.31	0.31
Methionine	0.29	0.29
Lysine	0.24	0.24
Nutrient composition (%)		
Metabolizable energy (Kcal/kg)	3018.10	3108.20
Available phosphorus	0.48	0.43
Calcium	1.03	1.04
*Crude fibre	3.52	3.58
*Crude fat	4.23	2.38
*Crude protein	22.17	20.04

495

*Analyzed composition

**1kg of vitamin-mineral premix contains Vitamin D3 - 2,000,000IU, Vitamin K - 2,250mg, Vitamin A - 10,000,000IU, Vitamin E - 20,000IU, Thiamine B1 - 1,750mg, Niacin - 27,500mg, Pantothenic acid - 7,500mg, Biotin - 50mg, Choline chloride - 400g, Riboflavin B2 - 5,000mg, Pyridoxine B6 - 2,750mg, Antioxidant - 125g, Magnesium - 80g, Iodine - 1.2g, Selenium - 200mg, Cobalt - 200mg, Zinc - 50mg, Iron - 20g, Copper - 5g.



516

517 **Figure 1.** Effects of *Juglans regia* kernel powder supplementation on growth of broiler
 518 chickens fed Aflatoxin B1 contaminated diets. AF: Aflatoxin; Diet 1: Control; Diet 2: 0.5
 519 mg/kg AF; Diet 3: 0.5 mg/kg AF+250 mg/kg JKP; Diet 4: 0.5 mg/kg AF+500 mg/kg JKP.

520

521 **Table 2.** Effects of *Juglans regia* kernel powder supplementation on serum chemistry of
 522 broiler chickens fed Aflatoxin-contaminated diets

Parameters	Diet 1	Diet 2	Diet 3	Diet 4	SEM	P value
Total protein (mmol/l)	39.80 ^a	26.40 ^b	37.00 ^a	39.47 ^a	1.76	0.01
Albumin (mmol/l)	21.35 ^a	13.75 ^b	20.45 ^a	20.52 ^a	1.17	0.04
Globulin (mmol/l)	18.45 ^a	12.65 ^b	16.55 ^a	18.95 ^a	0.87	0.01
Aspartate aminotransferase (IU/L)	87.05 ^b	111.20 ^a	94.30 ^b	85.05 ^b	3.45	0.01
Alanine transaminase (IU/L)	46.20 ^c	52.65 ^a	50.65 ^{ab}	49.45 ^b	0.75	0.01
Cholesterol (mmol/l)	5.05	5.40	5.65	5.30	0.08	0.07
Creatinine (mmol/l)	45.24 ^b	53.22 ^a	35.21 ^c	36.86 ^c	2.30	0.01
Glucose (mmol/l)	17.64 ^a	14.10 ^b	15.51 ^{ab}	17.16 ^a	0.46	0.01

523 ^{a-c}Means within a row with different letters are significantly different (P<0.05); AF: Aflatoxin; Diet 1: Control;
 524 Diet 2: 0.5 mg/kg AF; Diet 3: 0.5 mg/kg AF +250 mg/kg JKP; Diet 4: 0.5 mg/kg AF+500 mg/kg JKP; SEM:
 525 Standard error of means.

526

527 **Table 3.** Effects of *Juglans regia* kernel powder supplementation on immunoglobulins of
 528 broiler chickens fed aflatoxin-contaminated diets.

Parameters	Diet 1	Diet 2	Diet 3	Diet 4	SEM	P value
Immunoglobulin A (mg/dl)	218.80 ^{ab}	170.69 ^b	221.64 ^{ab}	266.34 ^a	12.49	0.03
Immunoglobulin E (mg/dl)	1071.50 ^a	931.52 ^b	1047.93 ^a	1089.23 ^a	20.81	0.01
Immunoglobulin G (mg/dl)	315.65 ^a	212.06 ^b	297.68 ^a	336.67 ^a	15.64	0.02
Immunoglobulin M (mg/dl)	371.41	330.88	353.21	343.51	7.44	0.28

529 ^{a-b}Means within a row with different letters are significantly different (P<0.05); AF: Aflatoxin; Diet 1: Control;
 530 Diet 2: 0.5 mg/kg AF; Diet 3: 0.5 mg/kg AF +250 mg/kg JKP; Diet 4: 0.5 mg/kg AF+500 mg/kg JKP; SEM:
 531 Standard error of means.

532

533

534

535 **Table 4.** Effects of *Juglans regia* kernel powder supplementation on pro-inflammatory
 536 cytokines of broiler chickens fed aflatoxin-contaminated diets.

Parameters	Diet 1	Diet 2	Diet 3	Diet 4	SE M	P value
Nuclear Factor Kappa B (pg/ml)	26.93 ^b	38.37 ^a	27.92 ^b	28.06 ^b	1.59	0.01
Tumour necrosis factor alpha (pg/ml)	34.82	66.13	43.82	40.58	4.91	0.09
Interleukin 6 (pg/ml)	14.43 ^c	39.82 ^a	27.31 ^b	18.11 ^c	3.18	0.01

537 ^{a-c}Means within a row with different letters are significantly different (P<0.05); AF: Aflatoxin; Diet 1: Control;
 538 Diet 2: 0.5 mg/kg AF; Diet 3: 0.5 mg/kg AF +250 mg/kg JKP; Diet 4: 0.5 mg/kg AF+500 mg/kg JKP; SEM:
 539 Standard error of means.

540

541

542