

1 **ACCEPTED ARTICLE**

2 **Alleviation of drought stress in German Chamomile (*Matricaria chamomilla* L.) in**
3 **response to suppressive oxidative stress and water deficit-induced stomatal closure by**
4 **exogenous polyamines**

5
6 Mohammad Javad Ahmadi-Lahijani^{1*}, Jafar Nabati², Saeed Moori³, Mohammad Kafi¹
7

8 ¹ Department of Agrotechnology, Ferdowsi University of Mashhad, Iran.

9 ² Research Center for Plant Sciences, Ferdowsi University of Mashhad, Iran.

10 ³ Department of Agronomy and Plant Breeding, Lorestan University, Iran.
11

12 * Corresponding Author: Department of Agrotechnology, Faculty of Agriculture, Ferdowsi
13 University of Mashhad, Mashhad, Islamic Republic of Iran. ORCID: 0000-0001-7356-7276.

14 E-mail: mjahmadi@um.ac.ir
15

16 **Running head:** Polyamines alleviates drought stress in German Chamomile

17 **ABSTRACT**

18 Polyamines (PAs) are signaling molecules that exhibit promising roles in improving stress
19 tolerance in plants. Limited information is available concerning the effects of the exogenous
20 PAs on medicinal plants including chamomile. This experiment was carried out to study the
21 effects of foliar application of PAs [putrescine (Put), spermidine (Spd), and spermine (Spm)]
22 on physiological and biochemical processes to understand the possible mechanisms concerning
23 the water deficit stress [soil field capacity (FC) as control, 80% of FC (FC₈₀), and 60% of FC
24 (FC₆₀)] alleviation in German Chamomile. We found that PAs partially inhibited water deficit-
25 induced stomatal closure and induced antioxidant enzymes to eliminate the increased H₂O₂.
26 Spd increased stomatal conductance (g_s) by 66, 65, and 35% at FC, FC₈₀, and FC₆₀, respectively,
27 compared with the control. The increased g_s enhanced leaf net photosynthesis (A_N) by 52 and
28 86% at FC₈₀ and FC₆₀, respectively, compared with the control. The role of PAs in oxidative
29 damage alleviation was approved by the negative correlation of leaf antioxidant activities and
30 malondialdehyde (MDA) and H₂O₂ content. According to the results, PAs function as stress-
31 protecting compounds to instigate the antioxidative enzymes to scavenge stress-induced H₂O₂,
32 improve membrane stability, and enhance water deficit tolerance. Generally, our results
33 suggested that PAs could be potential growth regulators to alleviate mild to severe water deficit
34 stress.

35 **Keywords:** Chlorophyll fluorescence; enzymatic antioxidant; hydrogen peroxide; gas
36 exchange variables; non-photochemical quenching.
37

38 **INTRODUCTION**

39 Drought is considered the most crucial worldwide factor in plant production systems (Hafezi
40 Ghehestani *et al.*, 2021). Water deficit stress affects the metabolism and growth of plants,
41 agricultural ecosystems, and human societies (Tezara *et al.*, 1999). Various physiological and
42 metabolic responses such as stomatal closure and decline in photosynthesis and growth rate are
43 induced in plants under water deficiency (Flexas and Medrano, 2002). Plant response to
44 stressful conditions is initiated when the stress is recognized at the cellular level. In both
45 unstressful and stressful environments, plants produce reactive oxygen species (ROS) that react
46 with proteins, lipids, and DNA and impair normal cellular functions (Apel and Hirt, 2004).
47 Water deficit stress disturbs the balance between ROS production and scavenging in plants,
48 leading to the accumulation of ROS and accelerating cell membrane damage and lipid
49 peroxidation (Farooq *et al.*, 2009).

50 Polyamines (PAs) are classified as a group of phytohormone-like aliphatic amine natural
51 compounds with aliphatic nitrogen structure and are considered secondary messengers in
52 signaling pathways (Liu *et al.*, 2023). Generally, naturally occurring PAs in the higher plants
53 including putrescine (Put), spermidine (Spd), and spermine (Spm) are not only involved in
54 numerous cellular and molecular processes in plants but also have been shown to improve plant
55 tolerance to abiotic stresses (Baghalian *et al.*, 2011). PAs trigger several molecular,
56 biochemical, and physiological responses of plants including increasing membrane stability and
57 osmolyte accumulation, protection of photosynthetic apparatus, activation of antioxidant
58 machinery, regulation of redox homeostasis, upregulation of stress-related genes, and
59 promotion of plant stress tolerance (Alcázar *et al.*, 2020).

60 The interaction of PAs with membrane phospholipids induces membrane stability under
61 stressful conditions. PAs play a vital role as signaling molecules that regulate several metabolic
62 pathways. The abiotic stress adaptations are enhanced by the PAs' functions as stress signaling
63 molecules (Pál *et al.*, 2015). Exogenously applied PAs increased antioxidant enzyme activities
64 under various stressful conditions; which could reduce cell damage and enhance the stress
65 tolerance of plants (Hassan *et al.*, 2018; Alcázar *et al.*, 2020). The accumulation of PAs under
66 adverse conditions can directly act as an antioxidant in eliminating ROS or may activate the
67 ROS-scavenging enzyme system (Liu *et al.*, 2023). Previous studies showed that exogenous
68 PAs significantly increased the activity of antioxidants such as SOD, POD, and CAT and
69 decreased ROS synthesis in *Vicia faba*, *Citrus reticulata*, *Arabidopsis thaliana*, and *Rosa*
70 *damascene* (Hasan *et al.*, 2021; Liu *et al.*, 2023).

71 German chamomile (*Matricaria chamomilla*) is one of the most valuable medicinal plants of
72 the Asteraceae (Compositae) family with many applications in the pharmaceutical, nutritional,
73 and cosmetic industries. Chamomile is relatively adaptable to a wide range of climates
74 including arid and semi-arid regions (Das *et al.*, 1998). However, drought negatively affects
75 chamomile performance and productivity. Although studies of PAs have been performed on
76 various crops (Alcázar *et al.*, 2006; Farooq *et al.*, 2009; Liu *et al.*, 2023), the available
77 information concerning the effects of PAs on medicinal plants is still limited. The effect of
78 foliar application of polyamines on leaf gas exchanges, chlorophyll fluorescence, and
79 physiological and biochemical processes was studied to understand the possible mechanisms
80 concerning water stress alleviation in German chamomile.

81 MATERIALS AND METHODS

82 Experimental site and procedure

83 The experiment was carried out at the Greenhouse of the Research Center for Plant Sciences,
84 Ferdowsi University of Mashhad, in 2021. Chamomile seeds (*cv.* Presov, obtained from Isfahan
85 Natural Resources Research Center) were surface sterilized with 0.2% sodium hypochlorite
86 solution for 5 min and rinsed three times with tap water. Ten chamomile seeds were sown in
87 each 10 kg pot (20 and 25 cm in diameter and depth, respectively) filled with clay loam soil at
88 a depth of 1 cm and thinned to 5 plants per pot after establishment. Plants were grown under
89 greenhouse conditions with day/night temperatures of 20/15±2 °C, natural light (~800 μmol.m⁻².
90 s⁻¹ PPFD), photoperiod of 13/11 h day/night, and relative humidity of 50±10%.

91 Soil preparation and treatments

93 Irrigation was applied until the full establishment (4-leaf), and then, water deficit was applied
94 at three levels of control (FC₁₀₀), moderate stress (80% of FC; FC₈₀), and severe stress (60% of
95 FC; FC₆₀) according to the method of Topp and Davis (1985). Plants were fertilized with the
96 Hoagland nutrient solution once a week along with irrigation water. Polyamines were foliar
97 applied as (a) spermine [Spm], (b) spermidine [Spd], (c) putrescine [Put], and (d) control, at a
98 concentration of 10 μM (Farooq *et al.*, 2009; Ali *et al.*, 2009). 10 mL of the solution was applied
99 to each plant using a handheld sprinkler. The first spray was made at the 5-leaf stage and
100 repeated at 15-day intervals until the flowering onset. The control plants were sprayed with
101 distilled water. The experiments were carried out in three replicates. All physiological and
102 biochemical data were taken from fully expanded leaves at the middle of the flowering stage.

103 Gas exchange parameters

105 Net photosynthetic rate (A_N), intercellular CO_2 concentration (C_i), transpiration rate (T_r), and
106 stomatal conductance (g_s) were measured between 9:00–11:00 h using a portable
107 photosynthetic system (ADC Bio Scientific Ltd, UK). Photosynthetically active radiations
108 (PAR), air temperature, relative humidity, and CO_2 concentration inside the sensor head were
109 set at $800 \mu mol.m^{-2}.s^{-1}$, 25 ± 2 °C, $50 \pm 5\%$, and 400 ± 20 ppm, respectively. Instantaneous (WUE_i)
110 and intrinsic (A_N/g_s) water use efficiency were calculated by dividing A_N by T_r and g_s ,
111 respectively. Mesophyll conductance (g_m) was also calculated as A_N/C_i (Fischer *et al.*, 1998).

112
113
114
115

116 Chlorophyll fluorescence (Chf)

117 A portable fluorometer (PAM-2500, Walz, Effeltrich, Germany) was used to measure the dark-
118 and light-adapted leaf chlorophyll fluorescence between 10:00–12:00 h. After 30 min of dark
119 adaptation, F_v/F_m was calculated as $(F_m - F_o)/F_m$, where F_m and F_o were the maximum
120 fluorescence elicited by a saturating light pulse and steady-state chlorophyll fluorescence,
121 respectively (Genty *et al.*, 1989). The maximum (F_m') and the steady-state (F_s) fluorescence
122 signals were measured from the light-adapted leaves after 4 min of illumination with
123 continuous red wavelength, non-saturating actinic light, and saturating pulses every 25 sec
124 (Murchie and Lawson, 2013). To measure the minimal fluorescence after the PSI excitation
125 (F_o'), the actinic light was then turned off, and far-red pulses were applied. Photochemical
126 quenching (qP) was calculated as $(F_m' - F_s)/(F_m' - F_o')$. Non-photochemical quenching, NPQ,
127 which is a proportion of the rate of the thermal energy dissipation, was estimated as $(F_m -$
128 $F_m')/F_m'$ (Van Kooten and Snel, 1990).

129
130

130 Electrolyte leakage (EL)

131 Leaf EL was measured to determine leaf membrane damage using an electrical conductivity
132 (EC) meter (Jenway Model 4510) according to Eq. 1 (Lutts *et al.*, 2016):

$$133 \quad EL (\%) = \frac{EC_1}{EC_2} \times 100 \quad (1)$$

134 Here, EC_1 and EC_2 are the EC of the solution after 24 h and the autoclaved (120 °C for 20 min)
135 samples, respectively.

136
137

137 Relative water content (RWC)

138 Leaf RWC was estimated using Eq. 2 (Smart and Bingham, 1974):

139
$$RWC (\%) = \left[\frac{W_f - W_d}{W_t - W_d} \right] \times 100 \quad (2)$$

140 Here, W_f , W_t , and W_d are fresh weight, turgid weight, and oven-dried weight (at 70 °C until
141 constant mass), respectively.

142

143 **Leaf osmotic potential (ψ_o)**

144 Leaf ψ_o was determined according to the freezing point depression method using an osmometer
145 (Wogel, model OM802.D). The leaf osmolytes content was calculated based on the van't Hoff
146 equation, and the leaf water content was measured by Eq. (3):

147
$$\frac{mMol}{g} = \left[\left(-\frac{Op}{RT} \right) \times \left(\frac{WC}{1 - WC} \right) \right] \quad (3)$$

148 Where the osmolytes content is based on mM g⁻¹ dry weight, R is the universal gas constant
149 (0.00831-liter MPa mol⁻¹ °K⁻¹), T is the absolute temperature (273 °K), Op is the leaf osmotic
150 potential (MPa), and WC is the leaf water content". The solute potential was determined at
151 room temperature (25 °C).

152

153 **Photosynthetic pigments**

154 Fresh leaves (100 mg) were homogenized in ethanol 70% and kept at 4 °C for 24 h. Leaf
155 pigments content (Chlorophylls a, b, and carotenoids) were determined spectrophotometrically
156 (U-2000, Hitachi Instruments, Tokyo, Japan) according to Lichtenthaler and Wellburn (1983).

157

158 **Leaf antioxidant enzymes**

159 100 mg leaf fresh weight was ground in liquid nitrogen, and 1 ml potassium phosphate (0.1 M,
160 pH = 7.8) containing 1 mM EDTA was added. The insoluble solids were removed by
161 centrifuging the mixture in 12,000 g at 4 °C (Sigma, model K18-3). The supernatant was kept
162 at -80°C to assay the enzymatic antioxidant activities (Yamaguchi *et al.*, 1995). Leaf enzymatic
163 antioxidants activity, including Ascorbate peroxidase (APX, EC 1.11.1.11), Superoxide
164 dismutase (SOD, EC 1.15.1.1), Catalase (CAT, EC 1.11.1.6), and Peroxidase (POD, EC
165 1.11.1.7), were assayed by the methods described by Nakano and Asada (1981), Giannopolitis
166 and Ries (1977), Cakmak and Horst (1991), and Ghanati *et al.* (2002), respectively.

167

168 **Malondialdehyde (MDA) and H₂O₂ content**

169 One hundred mg of leaf fresh weight was used to measure leaf MDA by the methods described
170 by Jiang and Zhang (2001). Leaf MDA was measured by homogenizing leaf fresh weight in 5
171 ml of trichloroacetic acid (100 g⁻¹) containing 250 g l⁻¹ thiobarbituric acid. The supernatant

172 absorbance was read at 532 nm spectrophotometrically (Jenway UV-Visible, Model 6305) and
173 was corrected at A600. For H₂O₂ content measurement, leaf tissues (500 mg) were
174 homogenized in an ice bath with 5 ml 0.1% (w:v) TCA. The homogenate was centrifuged at
175 12000×g for 15 min and the supernatant absorbance was read at 390 nm. The content of H₂O₂
176 was given on a standard curve (Sergiev *et al.*, 1997).

177 178 **Statistical analysis**

179 The experiment was carried out as a factorial (3 levels of water deficit × 4 levels of PAs)
180 arrangement in a randomized complete block design with three replications. The experiment
181 was carried out twice and the pooled data were analyzed. Data were subjected to a two-way
182 analysis of variance, and the LSD $p \leq 0.05$ was the test criterion for assessing differences
183 between the means of the main and/or interaction effects using SAS v.9.4 software. Data was
184 presented as \pm SE.

185 186 **RESULTS**

187 **Photosynthetic parameters**

188 Although the gas exchange parameters were reduced by the water deficit, they were
189 significantly improved by the application of PAs. Water deficit at FC₆₀ diminished the untreated
190 plant A_N by 40% compared with FC₁₀₀ (Fig. 1A). PAs application increased A_N by ~60%
191 compared with the untreated plants under FC₁₀₀ (Fig. 2A). However, under FC₈₀ and FC₆₀, Spd
192 showed the greatest improving effect on A_N; Spd application enhanced leaf A_N by 52 and 86%
193 at FC₈₀ and FC₆₀, respectively, compared with the untreated plants (Fig. 1A). Chamomile leaf
194 T_r reduced by 1.1 and 1.6 times, respectively, at FC₈₀ and FC₆₀ compared with FC₁₀₀ (Fig. 1B).
195 The highest leaf T_r was observed in Spd-treated plants at FC₁₀₀; 41, 17, and 21%, respectively,
196 higher than the untreated, Put, and Spm-treated plants (Fig. 1B).

197 Foliar application of PAs reduced the diminishing effects of water deficit on g_s. Spd increased
198 leaf g_s by 66, 65, and 35% at FC₁₀₀, FC₈₀, and FC₆₀, respectively, compared with the untreated
199 plants (Fig. 1C). The lowest leaf g_m was observed when Spd and Spm were applied,
200 respectively, at FC₁₀₀ and FC₈₀ (Fig. 1D). Although water deficit decreased C_i and C_i:C_a, PAs
201 significantly increased C_i and C_i:C_a compared with the untreated plants (Fig. 3A and B). At
202 FC₁₀₀ and FC₈₀, C_i was the highest in the Spd-treated plants by 1.4 and 1.1 times higher than
203 the untreated plants, respectively. The highest WUE_i was observed in the untreated plants at
204 FC₈₀ (Fig. 3C). Spm-treated plants showed 43% higher WUE_i at FC₁₀₀ compared with the
205 untreated plants; however, WUE_i reduced when PAs were applied under water deficit

206 conditions. A_N/g_s decreased by reducing the soil moisture. The highest A_N/g_s was observed in
207 Spm and Spd-treated plants at FC₁₀₀ and FC₆₀, respectively; 24 and 38% higher than the
208 untreated plants (Fig. 3D).

209

210 **Leaf chlorophyll fluorescence (Chf)**

211 Water deficit at the level of FC₆₀ decreased F_v/F_m by 22% compared with FC₁₀₀ (Fig. 4A). At
212 FC₈₀, Put-treated plants showed the highest F_v/F_m , which was 13% higher than the untreated
213 plants. Non-photochemical quenching (NPQ) increased by increasing the water deficit intensity
214 (Fig. 4B). Spd-treated plants showed 85 and 65% lower leaf NPQ than the untreated plants,
215 respectively, at FC₈₀ and FC₆₀. Photochemical quenching (qP) reduced by 26 and 33% at FC₈₀
216 and FC₆₀, respectively, compared with FC₁₀₀ (Fig. 4C). The highest qP was observed in the Put-
217 treated plants; 36, 69, and 34% higher than the untreated plants, respectively, at FC₁₀₀, FC₈₀,
218 and FC₆₀. The linear electron transport rate, ETR, decreased by 24 and 48%, respectively, at
219 FC₈₀ and FC₆₀ compared with FC₁₀₀ (Fig. 4D). The greatest ETR was recorded in the Spm-
220 followed by Put-treated plants at FC₈₀ by on average ~25% higher than the untreated plants
221 (Fig. 4D and 2A). However, at FC₆₀, Spd increased ETR by 39% compared with the untreated
222 plants.

223

224 **Leaf RWC, ψ_o , and EL**

225 Water deficit at FC₈₀ and FC₆₀ reduced leaf RWC by 13 and 22%, respectively, compared with
226 FC₁₀₀ (Fig. 5A). Put, Spm, and Spd enhanced leaf RWC by an average of ~25% compared with
227 the untreated plants at FC₁₀₀ (Fig. 5A and 2A). Water deficit at the level of FC₆₀ reduced ψ_o by
228 45% compared with FC₁₀₀ (Fig. 5B). In contrast, leaf ψ_o was enhanced in the PAs-treated
229 plants. Put application increased leaf ψ_o 35% compared with the untreated plants at FC₆₀ (Fig.
230 5B). Leaf EL was increased by 1.6 and 2.8 times at FC₈₀ and FC₆₀, respectively, compared with
231 FC₁₀₀ (Fig. 5C). However, Spm and Spd decreased leaf EL by 17 and 28%, respectively,
232 compared with untreated plants at FC₆₀ (Fig. 5C).

233

234 **Leaf photosynthetic pigment**

235 A significant decrease was observed in leaf photosynthetic pigments content exposed to water
236 deficit (Table 1). However, Spd increased leaf Chlt by 51, 60, and 79%, respectively, compared
237 with the untreated plants at FC₁₀₀, FC₈₀, and FC₆₀ (Table 1). The highest Chl a:b was observed
238 in the put-treated plants at FC₈₀; 32, 13, and 18% higher than the untreated, Spm-, and Spd-
239 treated plants, respectively. Water deficit increased leaf carotenoid content on average by ~18%

240 compared with FC₁₀₀ (Fig. 2B). The highest leaf carotenoid content was observed in Spd-treated
241 plants at all water-deficit levels (Table 1).

242 243 **Enzymatic antioxidant**

244 Leaf enzymatic antioxidant activity was significantly influenced by the water deficit, foliar
245 application of PAs, and their interaction (Fig. 6A). Water deficit and PAs increased leaf
246 antioxidant activity. Spd-treated plants showed the highest CAT at FC₈₀ and FC₆₀ than the
247 untreated plants. Spd increased CAT by 82 and 100% compared with the untreated plants at
248 FC₈₀ and FC₁₀₀, respectively (Fig. 6A). Leaf POD activity showed an increasing trend by
249 increasing the water deficit intensity and PAs application. At FC₈₀ and FC₆₀, Spd- and Spm-
250 treated plants showed the greatest POD, respectively, which were nearly double the untreated
251 plants at the respective water deficit level. At FC₈₀, the highest APX activity was recorded in
252 Spm- followed by Spd-treated plants by an average of ~35% over the untreated plants (Fig. 6C
253 and 2B). Leaf SOD activity showed a similar trend as POD. The highest SOD activity was
254 recorded in Spd- and Spm-treated plants at FC₈₀ and FC₆₀ by 42 and 36%, respectively, over
255 the untreated plants at their respective water deficit levels (Fig. 6D).

256 257 **Leaf MDA and H₂O₂ content**

258 Water deficit increased leaf MDA content; water deficit at FC₈₀ and FC₆₀ increased leaf MDA
259 content by 85% and 1.6 times, respectively, compared with FC₁₀₀ (Fig. 6E). Although PAs-
260 treated plant MDA also increased at FC₈₀, it remained almost unaltered at FC₆₀ compared with
261 FC₈₀. Leaf MDA content of Put, Spm, and Spd-treated plants were 38, 34, and 31%,
262 respectively, lower than the untreated plants (Fig. 6E). Leaf H₂O₂ content was almost doubled
263 at FC₆₀ compared with FC₁₀₀. However, PAs treatments reduced leaf H₂O₂ content compared
264 with the untreated plants (Fig. 6F). Spd application reduced leaf H₂O₂ by 18 and 10% compared
265 with the untreated plants at FC₈₀ and FC₆₀, respectively.

266 267 **DISCUSSION**

268 Abiotic stresses such as drought, cold, and K deficiency stresses simultaneously stimulate
269 abscisic acid (ABA) and PAs biosynthesis (Li *et al.*, 2021; Réthoré *et al.*, 2021; Zhu *et al.*,
270 2020). It has been supposed that PAs and ABA either alone or synergically induce stomatal
271 closure to increase plant tolerance during stressful conditions (Gong *et al.*, 2021). However,
272 most recent findings revealed that the ABA-induced stomatal closure is directly inhibited by
273 PAs in *Vicia faba*, and exogenous applications of PAs could reopen stomata even if they were

274 partially closed by ABA treatment (Liu *et al.*, 2023). We observed that exogenously sprayed
275 PAs stimulated leaf g_s of the water deficit-stressed chamomile plants, resulting in the improved
276 leaf A_N , T_r , F_v/F_m , ETR, and qP , and decreased NPQ, which may indicate the alleviating role
277 of PAs on drought-induced stomatal closure. Spd or Spm increased g_s and photosynthesis of
278 Chinese dwarf cherry but did not affect the F_v/F_m under drought stress (Yin *et al.*, 2014). Those
279 observations indicate that PAs can enhance photosynthesis by inhibiting stomatal closure
280 without affecting the stability of the photosynthetic system.

281 PAs are involved in plant protection against different environmental stresses (Baghalian *et al.*,
282 2011; Farooq *et al.*, 2009). PAs with acid-neutralizing, antioxidative, and membrane-stabilizing
283 properties positively influence photosynthetic efficiency under stressful conditions (Mapelli *et*
284 *al.*, 2008). Exogenously application of Put increased the net photosynthetic rate of basil
285 (*Ocimum basilicum* L.) plants under drought stress, while electrolyte leakage was reduced
286 (Darabi *et al.*, 2020). PAs with high net positive charges can stabilize PSII proteins such as D_1
287 and D_2 and by binding to membrane proteins can stabilize the structure of the proteins during
288 stress (Hamdani *et al.*, 2011).

289 The reductions in F_v/F_m and qP were correlated with an increase in NPQ (Fig. 7). A decline in
290 F_v/F_m indicates photoinhibition damage resulting from the incident PPFD when plants are
291 exposed to environmental stresses (Wang *et al.*, 2018). We found that water-deficit-induced
292 NPQ was ameliorated by the PAs, meanwhile, leaf qP of PAs-treated plants was improved
293 under water deficit conditions. Sang *et al.* (2016) found that the NPQ of water-stressed
294 cucumber (*Cucumis sativus*) leaves treated with Spd was enhanced, indicating that Spd can
295 accelerate the dissipation of absorbed light under drought conditions. Water deficit leads to a
296 decrease in the ETR and the generation of excess excitation energy (Tezara *et al.*, 2005). The
297 ETR of PAs-treated plants was higher than the untreated plants under water deficit, indicating
298 that the PSII reaction center of PAs-treated leaves maintained high activity. The negative
299 charges of LHCII can be neutralized with the PA's positive charges, resulting in the LHCII
300 complexes quenching by minimizing the exertion between the complexes (Hamdani *et al.*,
301 2011).

302 Exogenously applied PAs significantly increased leaf enzymatic antioxidant activities.
303 Enzymatic antioxidant activity and higher ROS scavenging ability were enhanced by
304 exogenously applied Spd in cucumber roots under stressful conditions (Wu *et al.*, 2018). In a
305 study on tomato plants under heat stress, exogenous application of Spd regulated various signal
306 transduction factors; mainly associated with genes related to stress signaling pathways such as

307 hormonal and sugar metabolisms (Cheng *et al.*, 2012). To decrease the ROS content during
308 stressful conditions, the concentration of several nonenzymatic antioxidants, such as ASA and
309 lycopene, as well as enzymatic antioxidants including SOD, POD, APX, and CAT activity can
310 be enhanced by PAs (Hasan *et al.*, 2021). ABA or Spd treatment alone decreased the activities
311 of both POD and CAT, which might be due to their contribution to the ROS accumulation
312 induced by ABA or Spd. However, Spd could activate the antioxidant enzymes to scavenge
313 H_2O_2 induced by ABA in the presence of ABA (Liu *et al.*, 2023). Our results also showed that
314 the increase in the antioxidant enzymes activities by PAs, when plants were fully irrigated, was
315 not as high as the water-stressed plants, which might be due to their lower ABA content.
316 However, under stressful conditions, higher ABA content probably activates the antioxidant
317 enzymes to scavenge higher H_2O_2 . Alcázar *et al.* (2006) also found that Put accumulation was
318 mainly an ABA-dependent metabolic response during drought.

319 PAs are regulators of redox homeostasis that play a dual role in plant oxidative stress (Saha *et*
320 *al.*, 2015). Although PAs might be responsible for cellular breakdown due to generating the
321 strong oxidizers H_2O_2 under stressful conditions, H_2O_2 can act as a signaling molecule involved
322 in stress signal transduction (Groppa and Benavides, 2008). The H_2O_2 -mediated signaling
323 pathway, which is involved in salt stress-responsive genes (*SIWRKY1*, *SIHKT1*, *SIDREB2*, and
324 *SIMYB102*), was induced by exogenous Spd in tomato plants and significantly reduced the
325 adverse effects of salt stress (Raziq *et al.*, 2022). Working on *Salvia officinalis* L. under drought
326 stress revealed that the enzymatic activities of APX and CAT were strongly coordinated
327 (Mohammadi-Cheraghabadi *et al.*, 2021).

328 PAs foliar application alleviated the ROS-induced membrane damage and reduced leaf MDA,
329 EL, and H_2O_2 . MDA and H_2O_2 accumulation can be indicators of cell damage. A lower leaf
330 MDA and H_2O_2 content led to lower membrane damage and leaf EL. PAs can accelerate the
331 antioxidant enzyme activities to protect plants against the oxidative damages and membrane
332 injury or may enhance the biosynthesis of protective substances under stressful conditions.

333 Tomato (*Solanum lycopersicum*) leaf MDA content and EL were decreased by exogenous Spd
334 under drought stress (Sang *et al.*, 2016). Besides their properties as free radical scavengers, PAs
335 also stabilize biological membranes by binding to membrane phospholipids under stressful
336 conditions (Pál *et al.*, 2015). Moradi Peynevandi *et al.* (2018) also observed that exogenously
337 applied PAs significantly decreased leaf H_2O_2 and MDA contents and enhanced the membrane
338 stability of cold-stressed stevia (*Stevia rebaudiana* Bertoni) plants.

339 Leaf *Chla* was decreased greater than *Chlb* content under water deficit conditions. This might
340 be due to a greater susceptibility of *Chla* than *Chlb* to stress or the generation of *Chlb* through
341 the degradation of *Chla* products (Sen *et al.*, 2014; Shahba *et al.*, 2010). This eventually led to
342 a decrease in *Chla:b*. It has been reported that PAs could protect the functional and structural
343 integrity of chloroplasts and slow down the rate of photosynthetic pigment degradation (Li *et*
344 *al.*, 2014; Nahar *et al.*, 2015). In our experiment, Spd foliar application increased the carotenoid
345 content of leaves. Carotenoids are also among the essential compounds playing a role in
346 protecting photosynthesis and stress-signaling pathways. Leaf carotenoid content was
347 positively correlated with leaf A_N , chlorophyll fluorescence, and antioxidative enzyme
348 activities (Fig. 7). Due to an increase in leaf carotenoid content under PAs application, it seems
349 that carotenoids are likely to prevent chlorophyll degradation due to their protective role (Dhar
350 *et al.*, 2020).

351 CONCLUSIONS

353 Our results revealed that exogenously applied PAs improved the drought tolerance of
354 chamomile plants in multiple ways. PAs improved leaf water status and alleviated oxidative
355 damage on the biological membranes by instigating leaf antioxidant content and reducing
356 membrane damage. The results here indicated that exogenously applied PAs act as regulators
357 to prevent chlorophyll degradation, protect the photosynthetic antenna and PSII structure, and
358 improve the photosynthetic efficiency of chamomile plants. Leaf antioxidant activities were
359 found to be determining and effective mechanisms to alleviate water deficit effects on
360 chamomile plants, induced by PAs. PAs inhibit water deficit-induced stomatal closure through
361 antioxidant enzyme-dependent H_2O_2 elimination. PAs can potentially improve water deficit
362 tolerance in chamomile. Furthermore, Spd showed the most effective impact in alleviating the
363 water deficit effects.

364 ACKNOWLEDGMENT

366 The research leading to these results received funding from the Vice President for Research and
367 Technology of Ferdowsi University of Mashhad under Grant Agreement No N.1/53684.

368 REFERENCES

- 370 1. Alcázar, R., Bueno, M. and Tiburcio, A. F. 2020. Polyamines: Small amines with large
371 effects on plant abiotic stress tolerance. *Cell*, **9(11)**: 2373.


- 372 2. Alcázar, R., Cuevas, J. C., Patron, M., Altabella, T. and Tiburcio, A. F. 2006. Abscisic acid
373 modulates polyamine metabolism under water stress in *Arabidopsis thaliana*. *Physiol.*
374 *Plant.*, **128(3)**: 448-455.
- 375 3. Ali, R., Abbas, H. and Kamal, R. 2009. The effects of treatment with polyamines on dry
376 matter and some metabolites in salinity–stressed chamomile and sweet majoram
377 seedlings. *Plant Soil Environ.*, **55(11)**: 477-483.
- 378 4. Apel, K. and Hirt, H. 2004. Reactive oxygen species: metabolism, oxidative stress, and
379 signal transduction. *Ann. Rev. Plant. Biol.*, **55**: 373-399.
- 380 5. Baghalian, K., Abdoshah, S., Khalighi-Sigaroodi, F. and Paknejad, F. 2011. Physiological
381 and phytochemical response to drought stress of German chamomile (*Matricaria*
382 *recutita* L.). *Plant Physiol. Biochem.*, **49(2)**: 201-207.
- 383 6. Cakmak, I. and Horst, W. J. 1991. Effect of aluminium on lipid peroxidation, superoxide
384 dismutase, catalase, and peroxidase activities in root tips of soybean (*Glycine max*).
385 *Physiol. Plant.*, **83(3)**: 463-468.
- 386 7. Cheng, L., Sun, R.-r., Wang, F.-y., Peng, Z., Kong, F.-l., Wu, J., Cao, J.-s. and Lu, G. 2012.
387 Spermidine affects the transcriptome responses to high temperature stress in ripening
388 tomato fruit. *J. Zhejiang Univ. Sci. B.*, **13(4)**: 283-297.
- 389 8. Darabi, F., Abbasi, N. and Zarea, M. J. 2020. Evaluation of the effect of putrescine and
390 brassinosteroid on induction of drought tolerance and physiological changes in basil
391 plant (*Ocimum basilicum* L.). *Environ. Str. Crop Sci.*, **13(4)**: 1183-1202.
392 doi:10.22077/escs.2020.2449.1644
- 393 9. Das, M., Mallavarapu, G. and Kumar, S. 1998. Chamomile (*Chamomilla recutita*):
394 Economic botany, biology, chemistry, domestication and cultivation. *J. Med. Arom.*
395 *Plant. Sci.*, **20**: 1074-1109.
- 396 10. Dhar, M. K., Mishra, S., Bhat, A., Chib, S. and Kaul, S. 2020. Plant carotenoid cleavage
397 oxygenases: structure–function relationships and role in development and metabolism.
398 *Brief. Func. Gen.*, **19(1)**: 1-9.
- 399 11. Farooq, M., Wahid, A. and Lee, D.-J. 2009. Exogenously applied polyamines increase
400 drought tolerance of rice by improving leaf water status, photosynthesis and
401 membrane properties. *Acta Physiol. Plant.*, **31(5)**: 937-945.
- 402 12. Fischer, R., Rees, D., Sayre, K., Lu, Z. M., Condon, A. and Saavedra, A. L. 1998. Wheat
403 yield progress associated with higher stomatal conductance and photosynthetic rate,
404 and cooler canopies. *Crop Sci.*, **38(6)**: 1467-1475.

- 405 13. Flexas, J. and Medrano, H. 2002. Drought-inhibition of photosynthesis in C₃ plants:
406 stomatal and non-stomatal limitations revisited. *Ann. Bot.*, **89(2)**: 183-189.
- 407 14. Genty, B., Briantais, J.-M. and Baker, N. R. 1989. The relationship between the quantum
408 yield of photosynthetic electron transport and quenching of chlorophyll fluorescence.
409 *Biochim. Biophysic. Acta (BBA)-Gen. Sub.*, **990(1)**: 87-92.
- 410 15. Ghanati, F., Morita, A. and Yokota, H. 2002. Induction of suberin and increase of lignin
411 content by excess boron in tobacco cells. *Soil Sci. Plant Nut.*, **48(3)**: 357-364.
- 412 16. Giannopolitis, C. N. and Ries, S. K. 1977. Superoxide dismutases: I. Occurrence in higher
413 plants. *Plant Physiol.*, **59(2)**: 309-314.
- 414 17. Gong, L., Liu, X.-D., Zeng, Y.-Y., Tian, X.-Q., Li, Y.-L., Turner, N. C. and Fang, X.-W.
415 2021. Stomatal morphology and physiology explain varied sensitivity to abscisic acid
416 across vascular plant lineages. *Plant Physiol.*, **186(1)**: 782-797.
- 417 18. Groppa, M. and Benavides, M. 2008. Polyamines and abiotic stress: recent advances.
418 *Amino acid.*, **34(1)**: 35-45.
- 419 19. Hafezi Ghehestani, M. M., Azari, A., Rahimi, A., Maddah-Hosseini, S. and Ahmadi-
420 Lahijani, M. J. 2021. Bacterial siderophore improves nutrient uptake, leaf
421 physiochemical characteristics, and grain yield of cumin (*Cuminum cyminum* L.)
422 ecotypes. *J. Plant Nut.*: 1-13.
- 423 20. Hamdani, S., Yaakoubi, H. and Carpentier, R. 2011. Polyamines interaction with
424 thylakoid proteins during stress. *J. Photochem. Photobiol. B: Biol.*, **104(1-2)**: 314-319.
- 425 21. Hasan, M. M., Alharbi, B. M., Alhathloul, H. A. S., Abdulmajeed, A. M., Alghanem, S.
426 M., Al-Mushhin, A. A., Jahan, M. S., Corpas, F. J., Fang, X.-W. and Soliman, M. H.
427 2021. Spermine-mediated tolerance to selenium toxicity in wheat (*Triticum aestivum*
428 L.) depends on endogenous nitric oxide synthesis. *Antioxidant.*, **10(11)**: 1835.
- 429 22. Hassan, F., Ali, E. and Alamer, K. 2018. Exogenous application of polyamines alleviates
430 water stress-induced oxidative stress of *Rosa damascena* Miller var. trigintipetala
431 Dieck. *South Afr. J. Bot.*, **116**: 96-102.
- 432 23. Jiang, M. and Zhang, J. 2001. Effect of abscisic acid on active oxygen species,
433 antioxidative defence system and oxidative damage in leaves of maize seedlings. *Plant*
434 *Cell Physiol.*, **42(11)**: 1265-1273.
- 435 24. Li, M., Wang, C., Shi, J., Zhang, Y., Liu, T. and Qi, H. 2021. Abscisic acid and putrescine
436 synergistically regulate the cold tolerance of melon seedlings. *Plant Physiol.*
437 *Biochem.*, **166**: 1054-1064.

- 438 25. Li, X., Gong, B. and Xu, K. 2014. Interaction of nitric oxide and polyamines involves
439 antioxidants and physiological strategies against chilling-induced oxidative damage in
440 *Zingiber officinale* Roscoe. *Sci Hort*, **170**: 237-248.
- 441 26. Lichtenthaler, H. K. and Wellburn, A. R. (1983). Determinations of total carotenoids and
442 chlorophylls a and b of leaf extracts in different solvents: Portland Press Ltd.
- 443 27. Liu, X. D., Zeng, Y. Y., Zhang, X. Y., Tian, X. Q., Hasan, M. M., Yao, G. Q. and Fang,
444 X. W. 2023. Polyamines inhibit abscisic acid-induced stomatal closure by scavenging
445 hydrogen peroxide. *Physiol. Plant.*: e13903.
- 446 28. Lutts, S., Benincasa, P., Wojtyla, L., Kubala, S., Pace, R., Lechowska, K., Quinet, M. and
447 Garnczarska, M. 2016. *Seed priming: new comprehensive approaches for an old*
448 *empirical technique*, PP.
- 449 29. Mapelli, S., Brambilla, I., Radyukina, N., Ivanov, Y. V., Kartashov, A., Reggiani, R. and
450 Kuznetsov, V. V. 2008. Free and bound polyamines changes in different plants as a
451 consequence of UV-B light irradiation. *Gen. App. Plant Physiol.*, **34(1-2)**: 55-66.
- 452 30. Mohammadi-Cheraghabadi, M., Modarres-Sanavy, S. A. M., Sefidkon, F., Rashidi-
453 Monfared, S. and Mokhtassi-Bidgoli, A. 2021. Improving water deficit tolerance of
454 *Salvia officinalis* L. using putrescine. *Sci. Rep.*, **11(1)**: 1-15.
- 455 31. Moradi Peynevandi, K., Razavi, S. M. and Zahri, S. 2018. The ameliorating effects of
456 polyamine supplement on physiological and biochemical parameters of *Stevia*
457 *rebaudiana* Bertoni under cold stress. *Plant Prod. Sci.*, **21(2)**: 123-131.
- 458 32. Murchie, E. H. and Lawson, T. 2013. Chlorophyll fluorescence analysis: a guide to good
459 practice and understanding some new applications. *J. Exp. Bot.*, **64(13)**: 3983-3998.
- 460 33. Nahar, K., Hasanuzzaman, M., Alam, M. and Fujita, M. 2015. Exogenous spermidine
461 alleviates low temperature injury in mung bean (*Vigna radiata* L.) seedlings by
462 modulating ascorbate-glutathione and glyoxalase pathway. *Int. J. Mol. Sci.*, **16(12)**:
463 30117-30132.
- 464 34. Nakano, Y. and Asada, K. 1981. Hydrogen peroxide is scavenged by ascorbate-specific
465 peroxidase in spinach chloroplasts. *Plant Cell Physiol.*, **22(5)**: 867-880.
- 466 35. Pál, M., Szalai, G. and Janda, T. 2015. Speculation: polyamines are important in abiotic
467 stress signaling. *Plant Sci.*, **237**: 16-23.
- 468 36. Raziq, A., Din, A. M. U., Anwar, S., Wang, Y., Jahan, M. S., He, M., Ling, C. G., Sun, J.,
469 Shu, S. and Guo, S. 2022. Exogenous spermidine modulates polyamine metabolism

- 470 and improves stress responsive mechanisms to protect tomato seedlings against salt
471 stress. *Plant Physiol. Biochem.*, **187**: 1-10.
- 472 37. Réthoré, E., Jing, L., Ali, N., Yvin, J.-C., Pluchon, S. and Hosseini, S. A. 2021. K
473 deprivation modulates the primary metabolites and increases putrescine concentration
474 in *Brassica napus*. *Front. Plant Sci.*, **12**: 681895.
- 475 38. Saha, J., Brauer, E. K., Sengupta, A., Popescu, S. C., Gupta, K. and Gupta, B. 2015.
476 Polyamines as redox homeostasis regulators during salt stress in plants. *Front.*
477 *Environ. Sci.*, **3**: 21.
- 478 39. Sang, Q., Shu, S., Shan, X., Guo, S. and Sun, J. 2016. Effects of exogenous spermidine on
479 antioxidant system of tomato seedlings exposed to high temperature stress. *Russ J*
480 *Plant Physiol*, **63(5)**: 645-655.
- 481 40. Sang, T., Shan, X., Li, B., Shu, S., Sun, J. and Guo, S. 2016. Comparative proteomic
482 analysis reveals the positive effect of exogenous spermidine on photosynthesis and
483 salinity tolerance in cucumber seedlings. *Plant cell rep*, **35(8)**: 1769-1782.
- 484 41. Sen, G., Eryilmaz, I. E. and Ozakca, D. 2014. The effect of aluminium-stress and
485 exogenous spermidine on chlorophyll degradation, glutathione reductase activity and
486 the photosystem II D₁ protein gene (psbA) transcript level in lichen *Xanthoria*
487 *parietina*. *Phytochem*, **98**: 54-59.
- 488 42. Sergiev, I., Alexieva, V. and Karanov, E. 1997. Effect of spermine, atrazine and
489 combination between them on some endogenous protective systems and stress markers
490 in plants. *Compt Rend Acad Bulg Sci*, **51(3)**: 121-124.
- 491 43. Shahba, Z., Baghizadeh, A. and Yosefi, M. 2010. The salicylic acid effect on the tomato
492 (*Lycopersicon esculentum* Mill.) germination, growth and photosynthetic pigment
493 under salinity stress (NaCl). *J Str Physiol Biochem*, **6(3)**.
- 494 44. Smart, R. E. and Bingham, G. E. 1974. Rapid estimates of relative water content. *Plant*
495 *Physiol.*, **53(2)**: 258-260.
- 496 45. Tezara, W., Marín, O., Rengifo, E., Martínez, D. and Herrera, A. 2005. Photosynthesis
497 and photoinhibition in two xerophytic shrubs during drought. *Photosynthetica*, **43(1)**:
498 37-45.
- 499 46. Tezara, W., Mitchell, V., Driscoll, S. and Lawlor, D. 1999. Water stress inhibits plant
500 photosynthesis by decreasing coupling factor and ATP. *Nat.*, **401(6756)**: 914-917.
- 501 47. Topp, G. and Davis, J. 1985. Time-domain reflectometry (TDR) and its application to
502 irrigation scheduling. In: "Adv. Irr.", (Ed.): Elsevier, PP. 107-127.


- 503 48. Van Kooten, O. and Snel, J. F. 1990. The use of chlorophyll fluorescence nomenclature in
504 plant stress physiology. *Photosynth. Res.*, **25(3)**: 147-150.
- 505 49. Wang, Z., Li, G., Sun, H., Ma, L., Guo, Y., Zhao, Z., Gao, H. and Mei, L. 2018. Effects of
506 drought stress on photosynthesis and photosynthetic electron transport chain in young
507 apple tree leaves. *Biol. Open*, **7(11)**.
- 508 50. Wu, J., Shu, S., Li, C., Sun, J. and Guo, S. 2018. Spermidine-mediated hydrogen peroxide
509 signaling enhances the antioxidant capacity of salt-stressed cucumber roots. *Plant*
510 *Physiol. Biochem.*, **128**: 152-162.
- 511 51. Yamaguchi, K., Mori, H. and Nishimura, M. 1995. A novel isoenzyme of ascorbate
512 peroxidase localized on glyoxysomal and leaf peroxisomal membranes in pumpkin.
513 *Plant and Cell Physiology*, **36(6)**: 1157-1162.
- 514 52. Yin, Z. P., Li, S., Ren, J. and Song, X. S. 2014. Role of spermidine and spermine in
515 alleviation of drought-induced oxidative stress and photosynthetic inhibition in
516 Chinese dwarf cherry (*Cerasus humilis*) seedlings. *Plant Growth Regul.*, **74**: 209-218.
- 517 53. Zhu, M.-D., Zhang, M., Gao, D.-J., Zhou, K., Tang, S.-J., Zhou, B. and Lv, Y.-M. 2020.
518 Rice OsHSFA3 gene improves drought tolerance by modulating polyamine
519 biosynthesis depending on abscisic acid and ROS levels. *Int. J. Molecul. Sci.*, **21(5)**:
520 1857.
- 521
- 522



523


524 **Fig. 1.** Changes in (A) Net photosynthetic rate, (B) Transpiration rate, (C) Stomatal
 525 conductance, and (D) Mesophyll conductance of German chamomile leaves exposed to water
 526 deficit and foliar application of polyamines. **WD: water deficit, PA_s: polyamines.** FC: field
 527 capacity, FC₈₀: 80% of FC, FC₆₀: 60% of FC. *, **, ***, and ns: significant at $p \leq 0.05$, $p \leq$
 528 0.01 , $p \leq 0.001$, and non-significant. Means with the same letters are not significantly different.
 529 **LSD $p \leq 0.05$. $n=9$.**

530




531

532 **Fig. 2.** Percent changes of (A) physiological and growth parameters, and (B) biochemical traits
 533 of German chamomile leaves affected by polyamines foliar application under water deficit
 534 levels relative to the untreated plants. AN: Net photosynthetic rate, gs: Stomatal conductance,
 535 Tr: Transpiration rate, Ci: Intercellular CO₂ concentration, AN/g: Intrinsic water use efficiency,
 536 WUEi: instantaneous water use efficiency, Fv/Fm: Maximum photochemical quantum yield of
 537 PSII, qP: Photochemical quenching, NPQ: Nonphotochemical quenching, ETR: Electron
 538 transport rate, EL: electrolyte leakage, RWC: Relative water contents, OP: Leaf osmotic
 539 potential, MDA: Malondialdehyde, H₂O₂: Hydrogen peroxide, EO: Essential oil, CAT:
 540 Catalase, POX: Peroxidase, APX: Ascorbate peroxidase, SOD: Superoxide dismutase, Chla:
 541 Chlorophyll a, Chlb: Chlorophyll b, Chlt: Total chlorophyll, Car: Carotenoids.
 542




543

544 **Fig. 3.** Changes in (A) Intercellular CO_2 concentration, (B) Intercellular to ambient CO_2
 545 concentration (C) Instantaneous water use efficiency, and (D) Intrinsic water use efficiency of
 546 German chamomile leaves exposed to water deficit and foliar application of polyamines. WD:
 547 water deficit, PAs: polyamines. FC: field capacity, FC₈₀: 80% of FC, FC₆₀: 60% of FC. *, **,
 548 ***, and ns: significant at $p \leq 0.05$, $p \leq 0.01$, $p \leq 0.001$, and non-significant. Means with the
 549 same letters are not significantly different. LSD $p \leq 0.05$. $n=9$.



550

551 **Fig. 4.** Changes in (A) Maximum photochemical quantum yield of PSII, (B) Nonphotochemical
 552 quenching, (C) Photochemical quenching, and (D) Linear electron transport rate of German
 553 chamomile leaves exposed to water deficit and foliar application of polyamines. FC: field
 554 capacity, FC₈₀: 80% of FC, FC₆₀: 60% of FC. WD: water deficit, PAs: polyamines. *, **, ***,
 555 and ns: significant at $p \leq 0.05$, $p \leq 0.01$, $p \leq 0.001$, and non-significant. Error bars indicate the
 556 differences between the water deficit levels, \pm SE. n=9.




557

558 **Fig. 5.** Changes in (A) Relative water content, (B) Osmotic potential, and (C) Electrolyte
 559 leakage of German chamomile leaves exposed to water deficit and foliar application of
 560 polyamines. WD: water deficit, PAs: polyamines. FC: field capacity, FC₈₀: 80% of FC, FC₆₀:
 561 60% of FC. *, **, ***, and ns: significant at $p \leq 0.05$, $p \leq 0.01$, $p \leq 0.001$, and non-significant.
 562 Means with the same letters are not significantly different. LSD $p \leq 0.05$. n=9.

563

564




565

566 **Fig. 6.** Changes in the activity of (A) Catalase, (B) Peroxidase, (C) Ascorbate peroxidase, (D)
 567 Superoxide dismutase, the content of (E) Malondialdehyde, and (F) Hydrogen peroxide German
 568 chamomile leaves exposed to water deficit and foliar application of polyamines. FC: Field
 569 capacity, FC₈₀: 80% of FC, FC₆₀: 60% of FC. WD: water deficit, PAs: polyamines. *, **, ***,
 570 and ns: significant at $p \leq 0.05$, $p \leq 0.01$, $p \leq 0.001$, and non-significant. Error bars indicate the
 571 differences between the water deficit levels, \pm SE. n=9.
 572

573

574



575

576 **Fig. 7.** Corplot analysis of OP: Leaf osmotic potential, NPQ: Nonphotochemical quenching,
 577 MDA: Malondialdehyde, EL: electrolyte leakage, H2O2: Hydrogen peroxide, gs: Stomatal
 578 conductance, AN: Net photosynthetic rate, Chlt: Total chlorophyll, EO: Essential oil, AO:
 579 Antioxidant enzymes, Car: Carotenoids, qP: Photochemical quenching, RWC: Relative water
 580 contents, F_v/F_m: Photochemical quantum yield of PSII german chamomile plants under water
 581 deficit and polyamines foliar application.

582

583

Table 1. Effect of water deficit and foliar application of polyamines on photosynthetic pigments of German chamomile leaves.

Water deficit	PAs [‡] (10 μM)	mg.gFW ⁻¹				Chlorophyll a:b
		Chlorophyll a	Chlorophyll b	Total chlorophyll	Carotenoids	
FC [†]	Control	0.71	0.31	1.03	0.64	2.41
	Put□	0.93	0.36	1.29	0.75	2.58
	Spm	0.98	0.39	1.38	0.78	2.50
	Spd	1.09	0.46	1.55	0.82	2.39
LSD		0.46	0.23	0.46	0.23	1.32
FC ₈₀	Control	0.53	0.26	0.79	0.75	2.02
	Put	0.83	0.27	1.11	0.86	3.38
	Spm	0.72	0.29	1.01	0.89	2.58
	Spd	0.90	0.37	1.27	0.96	2.42
LSD		0.41	0.21	0.54	0.46	2.21
FC ₆₀	Control	0.33	0.20	0.54	0.76	1.64
	Put	0.54	0.25	0.78	0.80	2.07
	Spm	0.58	0.27	0.84	0.88	2.14
	Spd	0.66	0.30	0.96	0.99	2.25
LSD		0.35	0.15	0.43	0.51	1.29
Water deficit		***	***	***	ns	ns
Polyamines		***	**	***	*	ns
WD*PAs		ns	ns	ns	ns	ns

[†] FC: field capacity, FC₈₀: 80% of FC, FC₆₀: 60% of FC, [‡] PAs: polyamines. □ Put: putrescine, Spm: spermine, Spd: spermidine. WD: water deficit, PAs: polyamines. *, **, ***, and ns: significant at $p \leq 0.05$, $p \leq 0.01$, $p \leq 0.001$, and non-significant. Means with the same letters are not significantly different. LSD $p \leq 0.05$.

584

585

586 کاهش تنش خشکی در بابونه آلمانی (*Matricaria chamomilla* L.) در پاسخ به تنش اکسیداتیو و بسته شدن روزنه
587 ناشی از کمبود آب توسط کاربرد خارجی پلی آمین ها

588 محمدجواد احمدی لاهیجانی^{1*}، جعفر نباتی¹، سعید موری²، محمد کافی¹

589

590

چکیده

591 پلی آمین ها مولکول های پیام رسانی هستند که نقش های امیدوارکننده ای در بهبود تحمل به تنش در گیاهان نشان داده اند. اطلاعات
592 محدودی در مورد اثرات کاربرد خارجی پلی آمین ها بر روی گیاهان دارویی از جمله بابونه در دسترس است. این آزمایش به منظور
593 بررسی اثرات محلول پاشی پلی آمین ها [پوترسین (Put)، اسپرمیدین (Spd) و اسپرمین (Spm)] بر فرآیندهای فیزیولوژیکی و
594 بیوشیمیایی برای درک مکانیسم های احتمالی مربوط به کاهش اثرات تنش کمبود آب [ظرفیت مزرعه (FC) به عنوان شاهد، 80
595 درصد ظرفیت مزرعه (FC80) و 60 درصد ظرفیت مزرعه (FC60)] در بابونه آلمانی انجام شد. نتایج نشان داد که پلی آمین ها تا حدی
596 بسته شدن روزنه ناشی از کمبود آب را مهار می کند و آنزیم های آنتی اکسیدانی را برای از بین بردن افزایش پراکسید هیدروژن القا
597 می کند. اسپرمیدین هدایت روزنه ای را در FC، FC80 و FC60 به ترتیب 66، 65 و 35 درصد در مقایسه با شاهد افزایش داد. افزایش
598 هدایت روزنه ای فتوسنتز خالص برگ را در FC80 و FC60 به ترتیب 52 و 86 درصد در مقایسه با شاهد افزایش داد. نقش پلی آمین ها

599 در کاهش تنش اکسیداتیو با همبستگی منفی فعالیت‌های آنتی‌اکسیدانی برگ و محتوای مالون‌دی‌آلدئید و پراکسید هیدروژن تایید
600 شد. با توجه به نتایج، پلی‌آمین‌ها به عنوان ترکیبات محافظ تنش عمل می‌کنند تا آنزیم‌های آنتی‌اکسیدانی را برای حذف
601 پراکسید هیدروژن ناشی از تنش، بهبود پایداری غشاء و افزایش تحمل کمبود آب تحریک کنند. به‌طور کلی، نتایج نشان داد که
602 پلی‌آمین‌ها می‌توانند تنظیم‌کننده‌های بالقوه رشد برای کاهش تنش کم‌آبی خفیف تا شدید باشند.

603 کلیدواژگان: فلورسانس کلروفیل؛ آنتی‌اکسیدان آنزیمی؛ پراکسید هیدروژن؛ متغیرهای تبادل گاز؛ خاموشی غیر فوتوشیمیایی

604

605