

ACCEPTED ARTICLE

Population Changes and Spatial Distribution Pattern of the Date Palm Spider Mite *Oligonychus afrasiaticus* (McGregor) in Natural Conditions of Ramshir, Khuzestan Province

Manuchehr Jahangiri¹, Leila Ramezani^{2*}, and Ebrahim Tamoli Torfi³

¹ Department of Plant Protection, Faculty of Agriculture, Agricultural Science and Natural Resources University of Khuzestan.

² Department of Plant Protection, Faculty of Agriculture, Agricultural Sciences and Natural Resources University of Khuzestan, Mollasani, Khuzestan, Islamic Republic of Iran.

³ Department of Plant Protection, Faculty of Agriculture, Agricultural Science and Natural Resources University of Khuzestan.

*Corresponding author; e-mail: ramezani@asnrukh.ac.ir or danaus.lpp@gmail.com

ABSTRACT

The date palm dust mite *Oligonychus afrasiaticus* (McGregor) is one of the most important pests of the date palms *Phoenix dactylifera* L. around the world. This pest causes reduction in the economic value of the product, and sometimes its damage reaches up to 100% in the southern provinces of Iran. The population fluctuations and spatial distribution of this pest on two more common varieties (Sayer and Barhi) were investigated in Ramshir city, Khuzestan province, southwestern Iran. For this purpose, two date palm plantations of 2 ha planted with mixed those two varieties of date palm trees were selected and 10 trees of each variety in each orchard were selected randomly. Sampling was done from four directions (North, South, East, and West) of each date palm every four days from mid-May to Early-November during 2020 and 2021. The results showed that the spider mite appeared on the clusters of two varieties from the beginning of June and the peak population of mites on both varieties was recorded in September in 2020 and 2021. Significant difference was observed between different sampling times during those two years and two varieties: Sayer ($F_{1.916, 9.580}=183.695, P<0.0001$) and ($F_{2.344, 11.720}=58.104, P<0.0001$) and Barhi ($F_{1.541, 4.622}=89.010, P<0.0001$) and ($F_{1.688, 5.065}=31.137, P<0.002$) in 2020 and 2021 respectively. The spatial distribution pattern of mites on both varieties in two years were assessed through the regression models of Taylor's power law and Iwao's patchiness regression method. The results of dispersion indicated an aggregated pattern of this pest in both varieties. The information obtained from the results can be used to prepare a successful integrated pest management program to reduce the consumption of chemical poisons for control of this pest.

38 **Keywords:** Barhi, Date spider mite, Sayer, Iran, Spatial distribution.

39
40 **INTRODUCTION**

41 Dates palm (*Phoenix dactylifera* L.) with more than four thousand years old are one of the most
42 important and strategic products in Iran. The countries of Iran, Iraq, Pakistan, Tunisia, Egypt and
43 Saudi Arabia, are the most important exporters of date around the world. With more than 250
44 thousand hectares of date orchards, Iran is one of the largest producers and exporters of dates in
45 the world. Currently, dates are cultivated and produced in 13 provinces of the country, of which
46 five provinces produce more than 99% of the country's dates. In terms of the amount of production,
47 Sistan and Baluchistan province ranks first and followed by Kerman, Fars, Bushehr, Khuzestan
48 and Hormozgan provinces (Ahmadi *et al.*, 2021).

49 One of the most important pests on date palm trees is the date spider mite *Oligonychus*
50 *afrasiaticus* (McGregor) (El-Shafie, 2022). This pest has a global distribution and can be found in
51 the Afrotropical, Nearctic, and Palearctic regions. It has been reported in several countries,
52 including Iran, Iraq, Saudi Arabia, United Arab Emirates, Bahrain, Oman, Kuwait, Yemen, Jordan,
53 Israel, Libya, Tunisia, Algeria, Morocco, Egypt, Sudan, Chad, Mauritania, Mali, and Nigeria (Al-
54 Zadjali, 2006; Alatawi and Kamran, 2018; Sanad *et al.*, 2017 Migeon and Dorkeld, 2021). In Iran,
55 this mite is distributed in the palm cultivation areas including Khuzestan, Bushehr, Fars, Isfahan
56 (Khour and Biabanak) and it have also been collected from ornamental date palm, corn, sugarcane,
57 sorghum and weeds (Farahbakhsh, 1961; 32.Modarres Awal, 1994; Gharib, 1991; Mossadegh and
58 Kocheili, 2002). The damage of this pest is high in Khuzestan province and Bushehr and the
59 infestation may lead high economic losses (Arbabi *et al.*, 2010; Latifian *et al.*, 2021).

60 One of the important features of phytophagous mite's population, is their spatial distribution
61 and population dynamics which can help determining the time of infestations, time of pest
62 outbreak, and the period when they frequently occur in the field (Ferraz *et al.*, 2020). Different
63 developmental stages of a species can create different distribution patterns according to their
64 activity level. The distribution pattern is the result of the interaction between a population and its
65 surrounding environment. Being aware of this pattern in the environment allows us to have more
66 information and the ability to describe and recognize the studied population (Pool, 1974). The
67 spatial distribution pattern can be used in modeling, determining the appropriate sampling method
68 and designing sampling programs (Taylor, 1984; Franco *et al.*, 2008; Ramezani *et al.*, 2016).
69 Therefore, the better the spatial distribution of a species is known, the more and better the

70 population dimensions of that species in natural and agricultural ecosystems can be measurable
71 (Pool, 1974).

72 Population members can be described in three types: random, regular, and aggregated
73 dispersion. Random distribution uses all points equally, while regular distribution is suitable for
74 small areas and rarely describes large areas. (Pool, 1974). Aggregative distribution occurs when
75 one species' presence increases the likelihood of another species' presence near the same point,
76 often due to irregularly distributed environmental factors or species' tendency to aggregate. (Pool,
77 1974, Latifian *et al.*, 2012, 2018, 2021; Ramezani and Zandi sohani, 2013; Ramezani *et al.*, 2016;
78 Ghaedi *et al.*, 2020).

79 Although a number of researchers have investigated the population fluctuations of *O.*
80 *afrasiaticus* in different parts of the world, for example, Elhalawany *et al.* (2020), investigated the
81 population dynamics of *O. afrasiaticus* on dates fruits in Qalyubia and New Valley governorates
82 of Egypt, while Ben-Chaban *et al.* (2011) assessed the seasonal occurrence of *O. afrasiaticus* and
83 phytoseiid predators on date palm in Tunisian oases and again Ben-Chaban *et al.* (2017) conducted
84 another study on the seasonal occurrence of the date palm spider mite *O. afrasiaticus* on the cultivar
85 Deglet Nour in a pesticide-free Tunisian date palm oasis, however, such studies are rare in Iran and
86 researchers have worked more in the field of studying the biology and control of this pest (Latifian
87 *et al.*, 2006; Latifian, 2017; Arbabi *et al.*, 2017). Aims of this research were studying population
88 fluctuations and spatial distribution of date spider mite in order to better understand the behavioral
89 characteristics and help the effective control methods.

90

91 **MATERIALS AND METHODS**

92 **Study site**

93 In order to investigate the population changes of the date spider mite *O. afrasiaticus* in Ramshir
94 city (E 30°53'28.0" N49°24'26.3"), Khuzestan Province, Iran, two separate date palm orchards,
95 with two varieties (one with Barhi and the other with Sayer) were selected and 10 date palm trees
96 (two date palm trees from the center of date palm plantation and two from each geographical
97 direction: East, West, North and South) from each variety with relatively uniform infestation were
98 chosen randomly. Sampling palm trees were between 13-15 years old and their height was about
99 two meters and they were not treated with any insecticides or acaricides during the study.

100 **Sampling**

101 Mite populations were sampled every four days from June to November of two years (2020 and
102 2021). In each sampling date eight fruits were chosen randomly from the four directions of the
103 north, south, east and west (two fruits from each direction and about 80 fruits per sampling date in
104 each date palm plantation for each variety) and mites were extracted by washing the fruits in 70%
105 ethanol and then counted under a stereomicroscope. Also the minimum and maximum temperature
106 values and the average temperature were recorded on each sampling date. To analyze the effect of
107 sampling date (during 2020 and 2021) and date varieties (Barhi and Sayer) on mite population,
108 one-way analysis of variance were used. Also, to determine the significant difference between the
109 average populations of mites in each of the geographical directions and compare it with the average
110 population in all directions (all four directions) *Student's t* method at the level of 0.05 was used.

111 **Spatial distribution**

112 In order to determine the spatial distribution of mites on each date palm tree, linear regression
113 method (Taylor's power law and Iwao's patchiness regression method) was used. In this method,
114 the data related to each date was considered separately and the variance and average of each
115 sampling date were calculated.

116

117 **Taylor's power law b index**

118 Taylor (1961) found a function between mean and variance as:

$$119 \quad S^2 = a\bar{x}^b \Rightarrow \log S^2 = \log a + b \log(\bar{x})$$

120 In this relationship, "a" and "b" are two constant parameters, where "a" depends on the sample size
121 and "b" indicates the amount of accumulation. To calculate Taylor's *b* index, after calculating the
122 mean and variance of the samples, by establishing a regression relationship between the logarithm
123 of the variance as the dependent variable and the logarithm of the mean as the independent variable,
124 the slope of the regression line or Taylor's *b* coefficient is calculated. If *b* is greater, equal or smaller
125 than one, it indicates a negative binomial (aggregated), a Poisson (random), or a positive binomial
126 (regular) distribution, respectively (Taylor, 1984).

127

128 **Iwao's patchiness regression method**

129 The average Lloyd's crowding index is denoted by m^* . This index was calculated by the following
130 relationship.

$$131 \quad m^* = m + \frac{s^2}{m - 1}$$

132 In this relationship, m and s are the mean and variance of the investigated samples, respectively.
133 Iowa established the following regression relationship between Lloyd's crowding index and
134 average samples: $m^* = \alpha + \beta m$
135 α : indicates the tendency of individuals to crowding and is one of the inherent properties of the
136 species and β reflects the population distribution in space and is interpreted in the same manner as
137 the b of Taylor's power law (Iwao, 1968). If β is greater than or equal to one, the distribution of
138 samples corresponds to negative binomial or Poisson, respectively, while β less than one is not
139 biologically defined (Southwood, 2000). Then the calculated t was compared with the Student's t -
140 table and based on that the type of mite spatial distribution was determined (Southwood, 2000).
141 Test $b = 1$ $t = (b - 1) / SE_b$ and Test $\beta = 1$ $t = (b - 1) / SE_\beta$
142 where SE_b and SE_β are the standard errors of the slope for the mean crowding regression.
143 Calculated values are compared with Two-tailed t -test at $n-2$ degrees of freedom. If the calculated
144 $t (t_c) < t$ -table (t_t), the null hypothesis ($b= 1$) would be accepted and spatial distribution would be
145 random and if it is the opposite, then the null hypothesis would be rejected and if $b > 1$ and < 1 , the
146 spatial distribution would be aggregated and uniform, respectively. SPSS software (version 16.0)
147 was used to perform all analyzes and Excell (2010) software was used to draw graphs.

148 149 **RESULTS**

150 Based on the sampling results, the emergence and activity of the date palm dust mite was from
151 the beginning of June on the fruits of both Sayer and Barhi varieties (Figure 1 and 2). Beginning
152 of *O. afrasiaticus* infestation was recorded from the early of Jun on Sayer and late of May on Barhi
153 in 2020 and 2021. On the other hand, the peak population (98.31 and 97.38 mites per fruit) of spider
154 mite was observed on Sayer on 10th September 2020 and 17th August 2021, respectively and (111.8
155 and 158.09 mites per fruit) on Barhi on 13th September 2020 and 24th August 2021, respectively.
156 The mean temperature at those times was between 37-39°C for both years. After the peak of mite
157 population, the trend of population changes decreased and the lowest population was recorded in
158 mid-October for both varieties (mean temperature was about 14.5°C) and after the end of October,
159 due to the entry of fruit into the Rutab and mature stage and crop harvest, no mite were observed
160 (Figure 1 and 2). Significant differences were observed between different sampling times in 2020
161 and 2021 on Sayer ($F_{1.916, 9.580}=183.695, P<0.0001$) and ($F_{2.344, 11.720}=58.104, P<0.0001$) and Barhi
162 ($F_{1.541, 4.622}=89.010, P<0.0001$) and ($F_{1.688, 5.065}=31.137, P <0.002$), respectively. In addition, a


163 significant difference was observed in population dynamics of the spider mite on the two varieties
 164 during the sampling dates in 2020 ($t_{1,342}=8.630, P=0.004$). However, no significant difference was
 165 observed in 2021 ($t_{1,438}= 0.283, P=0.595$). The analysis of variance revealed no significant
 166 difference in the changes of mite population on the Sayer variety between the two years of sampling
 167 ($t_{1,434}=1.823, P=0.178$). However, a significant difference was observed in the mite population on
 168 the Barhi variety between the two sampling years ($t_{1,345}=16.856, P=0.0001$), with the population
 169 density on Barhi higher in 2021 compared to 2020. According to **Table 1** the population dynamics
 170 of the spider mite on four directions of each tree during two years of sampling on both Barhi and
 171 Sayer varieties was similar and the statistical analysis showed that the changes in the mite
 172 population in the north, south, east and west directions of date palm were not significantly different
 173 for both varieties (Sayer and Barhi) during 2020 ($F_{3,168}=0.010, P=0.999; F_{3,168}=0.029, P=0.993$)
 174 and 2021 ($F_{3,172}=0.514, P=0.673; F_{3,172}=0.420, P=0.739$), respectively. These findings suggest that
 175 the presence and distribution of date spider mite on the Sayer and Barhi varieties in all directions
 176 of the tree in the studied date palm plantations was relatively uniform.

177


178 **Table 1. Multiple Comparisons of the spider mite population in four geographical directions of**
 179 **sampling date palm trees.**

(I) Directi on	(J) Directi on	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
North	South	-0.0233	8.17677	0.998	-16.0778	16.0313
	East	3.151	8.17677	0.700	-12.9034	19.2057
	West	-1.384	8.17677	0.866	-17.4383	14.6709
South	North	0.0233	8.17677	0.998	-16.0313	16.0778
	East	3.174	8.17677	0.698	-12.8802	19.2290
	West	-1.361	8.17677	0.868	-17.4151	14.6941
East	North	-3.151	8.17677	0.700	-19.2057	12.9034
	South	-3.174	8.17677	0.698	-19.2290	12.8802
	West	-4.535	8.17677	0.579	-20.5895	11.5197
West	North	1.384	8.17677	0.866	-14.6709	17.4383
	South	1.361	8.17677	0.868	-14.6941	17.4151
	East	4.535	8.17677	0.579	-11.5197	20.5895


180



181
 182 **Figure 1.** Population fluctuation of *Oligonychus afrasiaticus* on Sayer and Barhi varieties of date
 183 palm orchards in 2020.
 184



185
 186 **Figure 2.** Population fluctuation of *Oligonychus afrasiaticus* on Sayer and Barhi varieties of date
 187 palm orchards in 2021.
 188



189
190 **Figure 3.** The time duration of different developmental stages of date palm fruit in Ramshir.

191
192 The spatial distribution patterns and parameters of *O. afrasiaticus* in general (Table 2) and in
193 different geographical directions (Figure 3 and 4) was assessed according to Taylor's power law
194 and Iwao's patchiness regression on both varieties and the slope of the regression line was
195 calculated during 2020 and 2021. The results showed that both methods provided a significant
196 relationship between variance and mean density (in Taylor's power law), and between mean
197 crowding and mean density (in Iwao's patchiness regression). Also in both methods the calculated
198 t (t_c) was greater than the t in the table, thus indicating a significant difference in the slope of the
199 line with the number one ($b > 1$) (Table 2). In other words, the spatial distribution of spider mite was
200 aggregative (variance is greater than the average). Although the Iwao's patchiness method
201 compared to Taylor's power law better demonstrated fitness to this spider mite data, the results of
202 two years sampling showed that in both methods the explanatory coefficients (R^2) have a good
203 correlation with the data (Table 2) and both methods showed that the spatial distribution pattern of
204 this spider mite was aggregated in general and in four geographical directions (Figure 3 and 4).

205
206 **Table 2.** Spatial distribution statistics of *Oligonychus afrasiaticus* on two varieties of date palm
207 (Sayer and Barhi) using Taylor's power law and Iwao's patchiness regression analysis.

Spatial distribution method	Year of sampling	Variety	N	Regression slope \pm SE	Intercept \pm SE	R^2	Spatial distribution	t_c	$F_{(df)}$	$P_{\text{regression}}$
-----------------------------	------------------	---------	---	---------------------------	--------------------	-------	----------------------	-------	------------	-------------------------

Taylor	2020	Sayer	44	1.584±0.073	0.517±0.147	0.921	Clumped	8	475.034(1,41)	<0.0001
		Barhi	44	1.589±0.078	0.366±0.161	0.910	Clumped	7.551	412.467(1,43)	<0.0001
	2021	Sayer	44	1.369±0.052	0.182±0.122	0.941	Clumped	7.096	691.405(1,43)	<0.0001
		Barhi	44	1.391±0.081	0.390±0.201	0.875	Clumped	4.827	293.543(1,43)	<0.0001
Iwao	2020	Sayer	44	1.043±0.012	1.483±1.028	0.995	Clumped	3.583	7.983(1,43)	<0.0001
		Barhi	44	1.050±0.020	1.243±1.814	0.985	Clumped	2.5	2.784(1,43)	<0.0001
	2021	Sayer	44	1.032±0.005	1.779±2.527	0.999	Clumped	6.4	4.391(1,43)	<0.0001
		Barhi	44	1.028±0.009	17.684±5.670	0.996	Clumped	3.111	1.182(1,43)	<0.0001

209 Taylor: Taylor's power law and Iwao.: Iwao's patchiness regression.

210
211 **Table 3.** Linear regression between temperature and the mean population densities of *Oligonychus*
212 *afraziaticus* on varieties of Barhi and Sayer during 2020 and 2021 in Ramshir.

Year of sampling	Variety	N	Regression		R ²	F _(df)	P _{regression}
			slope ± SE	Intercept ± SE			
2020	Sayer	44	1.441±0.422	18.682±13.746	0.453	11.634(1,43)	<0.0001
	Barhi	44	1.867±0.417	29.46±13.753	0.556	20.091(1,43)	<0.0001
2021	Sayer	44	2.819±0.215	48.526±6.987	0.793	172.441(1,43)	<0.0001
	Barhi	44	3.597±0.402	53.647±13.075	0.640	80.138(1,43)	<0.0001

213
214 **DISCUSSION**

215 According to the current research, the appearance of the date spider mite on the Sayer and Barhi
216 during two consecutive years was at the beginning of June. With the increase in temperature, the
217 mite's population increased and the peak population was recorded in the beginning and middle of
218 September 2020 and in the middle to the end of August 2021 in Sayer and Barhi, respectively. The
219 effect of temperature on the mite population is shown in Table 3. As it shown, temperature has a
220 positive effect on increasing the mite population. Faez *et al.* (2018) assessed the population density
221 of *Panonychus citri* on Thomson navel orange in Ghaemshahr, Iran. They showed that the peak of
222 mite population occurred in the summer during July to September and they conclude that the
223 temperature is one of the reasons for the increase of mite population. Elhalawany *et al.* (2020)
224 indicated that the population dynamics of *O. afraziaticus* started with attacks fruits at second week
225 of April during Kimri stages, and reached its peak on June on Sayer date palm variety. After that
226 the mites migrate from fruits to fronds during Khalal stage.


227 Other researchers have shown that climatic factors such as temperature and dust affect the spider
228 mite population (Ikegami *et al.*, 2000). Latifian (2014) assessed the effect of dust phenomenon on

229 date palm important pests and diseases and showed that the dust phenomenon had the highest
230 effects on the incidence of the date spider mite. According to the observations of different
231 researchers, the population of date spider mites increases until the beginning of the Khalal stage
232 then decreases. At this stage, when the fruits color change from green to yellow or red, the increase
233 in fruit weight is slow, but the sucrose content increases. In the moist stage, the weight of the fruit
234 decreases due to the decrease in humidity, the color of the skin becomes light brown and the texture
235 of the fruit becomes soft. In these stages, the reduction of the mite population continues. As it can
236 be seen in Figures 1, 2 and 3, this decrease in our research started from the middle of September
237 and continued until the middle of October, and it is completely correlated with the ripening stages
238 (Rutab and Tamr) of date fruits. Also at the full ripening stage (Tamr), mites are rarely found on
239 the fruit and this time coincides with mid-October in Khuzestan province (Figure 3), which the
240 mite population extremely reduces and rarely be seen on the fruits (Figure 1 and 2). These
241 observations are completely consistent with the observations of other researchers (Paloski *et al.*,
242 2005; Aldosari and Ali, 2007; Ben Chaaban *et al.*, 2011). Based on the results of Latifian (2014),
243 the activity and damage of *O. afrasiaticus* starts from May and gradually increases with the
244 warming of the air and the increase of relative humidity. Also, this researcher reported the peak
245 activity of the mite in the months of July and August. On the other hand, the decrease in population
246 and mite damage was reported from September and when the fruit entered the moist stage (Latifian,
247 2014). In another study, Latifian and Kajbaf Vala (2016) investigated the effect of releasing
248 *Stethorus gilvifrons* (Mulsant) ladybug inoculum to control the date spider mite *O. afrasiaticus* in
249 Shadegan, Khuzestan. According to their results, the emergence and activity of date mite started
250 from the beginning of June and the peak population was also announced in July which is slightly
251 different from our result in the present study. As can be seen in Figure 3, in the present study, the
252 rate of increase in the spider mite population was observed in the Kimri stage and the peak
253 population was seen in the Khalal stage, which is completely consistent with the observations of
254 Alatawi *et al.* (2019). In another study, the density of date spider mite *O. afrasiaticus* was
255 investigated in Tunisia from 2006-2007. The results showed peak populations in 2006 on Kentichi,
256 Alig, Deglent Noor, Besser recorded in July and August; and in July 2007 on Deglent Noor,
257 Kentichi, Alig, and Besser varieties. (Ben Chaaban *et al.*, 2011). Ben Chaaban *et al.* (2017) and
258 Hussain (1969) found that *O. afrasiaticus* infestation on date palms starts in mid-May to June,
259 peaking in July and August, and declining at Khalal stage. Resistance to mite infestation varies


260 among different date palm cultivars, with some cultivars more susceptible to infestation than others
261 (Palevsky *et al.* 2005; Aldosari and Ali 2007; Ben Chaaban and Chermiti 2009). **Therefore, the**
262 **results of our study are consistent with the results of other researchers in terms of the appearance**
263 **and activity of the date spider mite. But in terms of the time of the peak population of spider mites,**
264 **it does not match with some of the reported results. This inconsistency can be caused by different**
265 **studied varieties and different geographical regions with different weather conditions.**

266 The results of analyzing the spatial distribution of the spider mite *O. afrasiaticus* using Taylor's
267 power law and Iwao's patchiness regression (Table 2) during 2020 and 2021 showed that there was
268 a significant relationship between mean log and variance of population density. In Both method,
269 the coefficient (b or β) was greater than 1 and the distribution was aggregated and follows the
270 negative binomial pattern. Aggregative spatial distribution of other species of spider mites
271 (*Panonychus ulmi* in apple orchards (Rahmani *et al.*, 2010), *Panonychus citri* on Thomson navel
272 orange (Faez *et al.*, 2018), and *Eotetranychus frosti* on apple (Darbemamieh *et al.* 2012)) has been
273 reported by several researchers. Rahmani *et al.* (2010) studied the spatial distribution and seasonal
274 activity of *P. ulmi* and its predator *Zetzellia mali* in Zanzan apple orchards using Taylor's power
275 law and Iwao's patchiness regression methods. They found higher accuracy in Iwao's and Tylor's
276 methods, and Taylor's power law provided more accurate population estimations. In the oresent
277 study, Iwao's patchiness regression showed more accurate estimations of populations than Taylor's
278 power law, however, both of them had a high correlation with the data (Table 2). Gharib (1996)
279 during the study on date palm sucking pests declared that the relative density of this group of pests
280 is high and their distribution in date palm plantations is mostly uniform but varies on clusters and
281 leaves. Also, according to different researches, aggregative distribution is the most common pattern
282 of spatial distribution among insects, especially pest insects (Ramezani & Zandi Sohani, 2017;
283 Ramezani *et al.*, 2016; Zarei Sarchogha *et al.*, 2018). This distribution may be seen in all
284 developmental stages of the pest or in some developmental stages.

285



286 **Figure 4.** Spatial distribution of *Oligonychus afrasiaticus* in **four** directions of Barhi cultivar in
 287 2020: left and 2021: right, using Taylor's Power law. B: Barhi, N: North, S: South, W: West, E:
 288 East



290 **Figure 5.** Spatial distribution of *Oligonychus afrasiaticus* in four directions of Sayer cultivar in
291 2020: left and 2021: right, using Taylor's Power law. Es: Sayer, N: North, S: South, W: West, E:
292 East.

293

294 CONCLUSIONS

295 The results of the current study provided valuable information about the population dynamics,
296 appearance time, peak population, and spatial distribution of the date spider mite *O. afrasiaticus*
297 on two important date palm varieties (Sayer and Barhi) in the climatic conditions of Ramshir city,
298 Khuzestan Province, Iran. These findings are essential in the development of effective management
299 strategies and the preparation of successful integrated pest management programs for the date
300 spider mite. By reducing the consumption of chemical poisons and protecting the environment,
301 these programs can help to minimize the economic losses caused by this pest and maintain the
302 productivity of date palm crops. Overall, this research provides a foundation for further studies on
303 the behavior and management of the date spider mite, which can help to improve the sustainability
304 and profitability of date palm cultivation.

305

306 ACKNOWLEDGEMENTS

307 This study was funded by the Agricultural Science and Natural Resources University of
308 Khuzestan, Iran.

309

310 REFERENCES

- 311 1. Ahmadi, K., Hatami, F., Hoseinpoor, R., Abdeshah, H. and Ebadzadeh, H. R. 2021.
312 *Agricultural Statistics*. Ministry of Jihad-Agriculture. Deputy of Planning and Economics.
313 Tehran, 392pp.
- 314 2. Alatawi, F. J. and Kamran, M. 2018. Spider mites (Acari: Tetranychidae) of Saudi Arabia:
315 two new species, new records and a key to all known species. *J. Nat. His.*, 52(7–8): 429–
316 55.
- 317 3. Alatawi, F. J., Mirza, J. H., Alsahwan, K. A. and Kamran, M. 2019. Field population sex
318 ratio of the date date palm mite, *Oligonychus afrasiaticus* (McGregor). *Afr. Entomol.*,
319 27(2):336–43.
- 320 4. Aldosari, S. and Ali, A. G. 2007. Susceptibility of date palm fruit cultivars to the natural
321 infestation by *Oligonychus afrasiaticus* (Mcg.) (Acari: Tetranychidae) in relation to their
322 chemical composition. *Ass. Univ. Bull. Environ. Res.*, 10(2):1–7.

- 323 5. Al-Zadjali, T. S., Abd-Allah, F. F. and El-Haidari, H. S. 2006. Insect pests attacking date
324 palms and dates in Sultanate of Oman. *Egypt. J. Agric. Res.*, 84:51–9.
- 325 6. Arbabi, M., Latifian, M., Askari, M., Fasihi, M. T., Damghani, M. R.,
326 Golmohammadzadeh-Khiaban, N. and Rezai, H. 2017. Evaluation of different treatments
327 in control of *Oligonychus afrasiaticus* in date date palm orchards of Iran. *Persian J. Acarol.*,
328 6(2):125–35.
- 329 7. Arbabi, M., Asgari, M., Fasihi, M. T., Golmohammadzadeh-Khiaban, N., Damghani, M.
330 R., Latifian, M. and Babai, M. 2010. Evaluation of water spray application for organic
331 control of date date palm spider mite *Oligonychus afrasiaticus* (McGregor)
332 (Acari:Tetranychidae) of date date palm orchards in southern parts of Iran. *J. Entomol. Res.*,
333 1: 269–77.
- 334 8. Ben Chaaban, S. and Chermiti, B. 2009. Characteristics of date fruit and its influence on
335 population dynamics of *Oligonychus afrasiaticus* in the southern of Tunisia. *Acarologia*,
336 49(1–2): 29–37.
- 337 9. Ben Chaaban, S., Chermiti, B. and Kreiter, S. 2011. *Oligonychus afrasiaticus* and
338 phytoseiid predators' seasonal occurrence on date palm *Phoenix dactylifera* (Deglet Noor
339 cultivar) in Tunisian oases. *Bull. Insectology*, 64 (1): 15–21.
- 340 10. Ben Chaaban, S., Chermiti, B. and Kreiter, S. 2017. The spatio-temporal distribution
341 patterns of the spider mite, *Oligonychus afrasiaticus*, on date date palm (Deglet Nour
342 cultivar) in a pesticide free Tunisian oasis. *Tunis. J. Plant Prot.*, 12(2):159–72.
- 343 11. Darbemamieh, M., Fathipour, Y., and Kamali, K. 2012. Seasonal activity and spatial
344 distribution pattern of *Eotetranychus frosti* (Acari: Tetranychidae) in an unsprayed apple
345 orchard of Kermanshah, Western Iran. *Persian J. Acarol.*, 1(2): 137-146.
- 346 12. Elhalawany, A. S., Sayed, A. A. and Khalil, A. E. 2020. Biodiversity and population
347 dynamics of mites inhabiting date palm trees in Qalyubia and New Valley Governorates,
348 *Egypt. J. Plant Prot. Res. Inst.*, 3(1):346–364.
- 349 13. El-Shafie, H. A. 2022. The Old World date palm mite *Oligonychus afrasiaticus* (McGregor
350 1939) (Acari: Tetranychidae), a major fruit pest: biology, ecology, and management. *CABI*
351 *Reviews*, 17(020), 1-15.

- 352 14. Faez, R., Shojaii, M., Fathipour, Y., and Ahadiyat, A. 2018. Effect of initial infestation on
353 population fluctuation and spatial distribution of *Panonychus citri* (Acari: Tetranychidae)
354 on Thomson navel orange in Ghaemshahr, Iran. *Persian J. Acarol.*, 7(3):256-278
- 355 15. Farahbakhsh, Q. 1961. *List of important pests of plants and agricultural products of Iran.*
356 Publications of the Plant Protection Organization. Tehran, 153 pp.
- 357 16. Ferraz, J. C. B., Da Silva, S. M., De França, S. M., Silva, P. R. R., Melo, J. W. D. S., & De
358 Lima, D. B. 2020. Host preference, population dynamics, distribution, and injuries of
359 *Oligonychus punicae* (Acari: Tetranychidae) in an eucalyptus clonal minigarden. *Syst.*
360 *Appl. Acarol.*, 25(9), 1649-1660.
- 361 17. Franco, R. A., Reis, P. R., Zacarias, M. S., Altoe, B. F., & Pedro, N. 2008. Population
362 dynamics of *Oligonychus ilicis* (McGREGOR, 1917) (Acari: Tetranychidae) in coffee
363 plants and of their associated phytoseiids. *Coffee Sci.*, 3(1), 38-46.
- 364 18. Ghaedi, H., Kocheili, F., Latifian, M., & Nejad, R. F. 2020. Spatial and temporal
365 distribution of rhinoceros beetles *Oryctes Hellwig* (Col.: Scarabaeidae) in date palm
366 plantations of Khuzestan province. *Plant Pest Res.*, 10(2).
- 367 19. Gharib, A. R. 1996. *Fauna of insects and animal pests, diseases and weeds in date orchards*
368 *production of Iran.* Plant Pests and Diseases Research Institute Publication, Ministry of
369 Agriculture and Rural Development, Tehran, 24 pp.
- 370 20. Gharib, A. R. 1991. *Important date palm pests in Iran.* Agricultural Extension Organization
371 Publication, 41 pp.
- 372 21. Hussain, A. A. 1969. Biology of *Paratetranychus afrasiaticus* McG. infesting date date
373 palm in Iraq. *Egypt. Acad. J. Biolog. Sci.*, 33: 221–5.
- 374 22. Ikegami, Y., Yano, S., Takabasyi, J. and Takafuji, A. 2000. Function of quiescence of
375 *Tetranychus kanzawai* (Acari: Tetranychidae), as a defense mechanism against rain. *Appl.*
376 *Entomol. Zool.*, 35: 339–343.
- 377 23. Iwao, S. 1968. A new regression method for analyzing the aggregation pattern of animal
378 populations. *Res. Popul. Ecol.*, 10: 1–20.
- 379 24. Latifian, M., Rahnama, A. A., & Sharifnezhad, H. 2012. Effects of planting pattern on
380 major date palm pests and diseases injury severity. *Intl. J. Agri. Crop. Sci.*, 4(19), 1443-
381 1451.

- 382 25. Latifian, M. 2014. Date palm spider mite (*Oligonychus afrasiaticus* McGregor) forecasting
383 and monitoring system. *WALIA journal*, 30: 79-85.
- 384 26. Latifian M. 2017. Integrated pest management of date palm fruit pests: A review. *J.*
385 *Entomol.*, 14(3):112–121.
- 386 27. Latifian, M., Rad, B., & Habibpour, B. 2018. Termites of Iranian date palm orchards and
387 their spatial and temporal distribution. *Sociobiology*, 65(1), 24-30.
- 388 28. Latifian, M., Ahmadi, A., and Pezhman, H. A. 2006. A study of the extension factors
389 affecting suitable control of date date palm pests and diseases. *Iran. J. Agric. Sci.*,
390 37(2):155–63.
- 391 29. Latifian, M., Assari, M. J., Modarresi-Najafabadi, S. S., Amani, M., Basavand, F., Fasihi,
392 M. T., Zohdi, H. Bagheri, A. (2021). Economic injury level of date spider mite,
393 *Oligonychus afrasiaticus* (Acari: Tetranychidae) on six commercial date cultivars. *Persian*
394 *J. Acarol*, 10(4), 451-466.
- 395 30. Latifian, M. and Kajbafevala, G. 2016. Studying the efficiency of *Stethorus gilvifrons*
396 inoculate release for date palm spider mite (*Oligonychus afrasiaticus*) biological control in
397 field condition. Tehran, Iran: Ministry of Jihad-E-Agriculture. Research and education
398 organization. Date palm and Tropical Fruits Research Institute of Iran, 75 pp.
- 399 31. Migeon, A, Dorkeld, F. 2021. Spider Mites Web: a comprehensive database for the
400 Tetranychidae. Available from: <http://www1.montpellier.inra.fr/CBGP/spmweb>
401 [Accessed: February 24, 2022].
- 402 32. Modarres *Awal*, M. 1994. *List of agricultural pests of Iran and their natural enemies.*
403 Publications of Ferdowsi University of Mashhad, 364 pages.
- 404 33. Mossadegh, M. S. and Kocheili, F. 2002. *Semi-descriptive list of identified identified*
405 *agricultural arthropods, health and other agricultural pests of Khuzestan province.*
406 Publications of Shahid Chamran University of Ahvaz, 475 pages.
- 407 34. Palevsky, E., Borochoy-Neori, H. & Gerson, U. 2005. Population dynamics of *Oligonychus*
408 *afrasiaticus* in the southern Arava Valley of Isreal in relation to date fruit characteristics
409 and climatic conditions. *Agric. and Forest Entomol.*, 7: 283–290.
- 410 35. Poole, R.W. 1974. *An introduction to quantitative ecology.* McGraw – Hill. NewYork, 532
411 pp.

- 412 36. Rahmani, H., Fathipour, Y., & Kamali, K. 2010. Spatial distribution and seasonal activity
413 of *Panonychus ulmi* (Acari: Tetranychidae) and its predator *Zetzellia mali* (Acari:
414 Stigmaeidae) in apple orchards of Zanjan, Iran. *J. Agric. Sci. Technol.*, 12: 155-165.
- 415 37. Ramezani, L., and Zandi Sohani, N. 2013. Population dynamics and spatial distribution of
416 important Thysanoptera species on wheat. *Iran J. Plant Prot. Sci.*, 44(2):283-290.
- 417 38. Ramezani, L., Rajabpour, A., Zandi Sohani, N. and Yarahamdi. F. 2016. Sequential
418 sampling with constant accuracy of aphids in wheat fields in Ahvaz. *Iran Plant Prot.*
419 *Research*, 29(4): 582-588.
- 420 39. Sanad, A. S., Elhalawany, A. S., Abou-Setta, M. M., and El-Khateeb, H. M. 2017. Partial
421 survey of date palm dust mite, *Oligonychus afrasiaticus* (McGregor) in Egypt including
422 historical trait. *Acarines: J. Egyptian Soc. Acarol*, 11(1): 53-55.
- 423
- 424 40. Southwood, T. R. E. 2000. *Ecological methods, with the particular reference to the study*
425 *of insect populations*. 3rd edition. Chapman & Hall Pub. London, 524 pp.
- 426 41. Taylor, L. R. 1984. Assessing and interpreting the spatial distribution of insect populations.
427 *Annu. Rev. Entomol.*, 29: 321–357.
- 428 42. Taylor, L. R. 1961. Aggregation, variance to the mean. *Nature*, 189: 732–735.
- 429 43. Zarei Sarchogha, R., Zandi-Sohani, N. and Ramezani, L. 2018. Spatial dispersion and
430 sequential sampling of *Sitobion avenae* (Fabricius) in wheat fields of SarPol-e Zahab,
431 Kermanshah province. *Plant Protection (Sci. Journal Agri.)*, 41(2): 49-60.

432
433
434 تغییرات جمعیت و الگوی پراکنش فضایی کنه تارتن خرما *Oligonychus afrasiaticus* McGregor در شرایط طبیعی
435 رامشیر، استان خوزستان

436 کنه تارتن خرما *Oligonychus afrasiaticus* (McGregor) یکی از آفات مهم خرما در مناطق خرما خیز دنیا می باشد. خسارت
437 این کنه باعث کاهش ارزش اقتصادی محصول می شود، به طوری که خسارت آن در سال های طغیانی در استان های جنوبی ایران
438 تا 100 درصد نیز می رسد. بررسی تغییرات جمعیت، تعیین زمان اوج جمعیت و توزیع فضایی جهت مدیریت این آفت در نخلستان
439 ها اهمیت فراوانی دارد. در پژوهش حاضر، تغییرات جمعیت و توزیع فضایی این آفت روی دو واریته استعمران و برحی طی سال های
440 1399 و 1400 در شهرستان رامشیر مورد بررسی قرار گرفت. بدین منظور پنج تکرار از هر واریته خرما با آلودگی نسبتاً یکسان جهت
441 نمونه برداری انتخاب شد. نمونه برداری از چهار جهت نخل خرما و به صورت دو روز در هفته از اواسط بهار تا اواخر تابستان انجام
442 شد. نتایج نشان داد که ظهور کنه تارتن خرما روی خوشه های دو واریته استعمران و برحی از اوایل خرداد ماه و به صورت پراکنده
443 می باشد. همچنین اوج جمعیت کنه روی هر دو واریته در سال 1399 و 1400 به ترتیب در شهریور و مرداد ماه ثبت گردید. بین
444 زمانهای مختلف نمونه برداری طی سال های 1399 و 1400 به ترتیب روی دو واریته استعمران ($F=1.916, 9.580=183.695, P<0$).

445 (F_{2, 688, 5. 065}=31.137, و $P<0. 0001$) (F_{1. 541, 4. 622}=89. 010 و برحی (F=2. 344, 11. 720=58.104, $P<0. 0001$) و 0001)
446 ($P<0. 002$) اختلاف معنی دار مشاهده شد. همچنین اختلاف معنی داری بین تغییرات جمعیت کنه تارتن خرما روی دو وارپته طی
447 سال 1399 مشاهده شد. از طرفی توزیع فضایی کنه تارتن خرما روی هر دو وارپته خرما طی دو سال مورد بررسی تجمعی بود. از
448 اطلاعات نتایج حاصل از این پژوهش جهت تهیه یک برنامه موفق مدیریت تلفیقی آفت جهت کاهش مصرف سموم شیمیایی در
449 شهرستان رامشیر می توان استفاده نمود.

450