

The Relations between Some Phytochemical Properties and Fatty Acid Content of Pumpkin (*Cucurbita pepo* L.) Seeds

Haluk Çağlar Kaymak^{a*}, Selen Akan^b, Faika Yarali Karakan^c, and Serpil Tıraşçı^a

^aDepartment of Horticulture Faculty of Agriculture, Atatürk University, Erzurum, Türkiye .

^bDepartment of Horticulture Faculty of Agriculture, Ankara University, Ankara, Türkiye .

^cDepartment of Horticulture Faculty of Agriculture, Kilis 7 Aralık University, Kilis, Türkiye .

*Corresponding author, e-mail: hckaymak@atauni.edu.tr

Running title: Some Phytochemical Properties in different pumpkin seeds

ABSTRACT

Pumpkin (*Cucurbita pepo* L.) seeds are popular for their dietary and health benefits. However, there were limited data on the pathway between phytochemical and nutritional values of pumpkin seeds. For this purpose, the seeds of some Turkish pumpkin genotypes (NVS-1, NVS-2, KNY, KYS-1, KYS-2, BRS, EDR and KRK) were analysed for their amino acids, organic acids, fatty acids, and mineral content. The wide variation between seeds in organic acids (KYS-2, 8.105 ng μl^{-1} ; KRK, 1.939 ng μl^{-1}) and amino acids (KYS-2, 32.99 nmol μl^{-1} ; KNY, 15.65 nmol μl^{-1}) content was observed. C18:2n6 and C18:1n9 were the most predominant fatty acids in the seeds, whereas C16:1n7 was the least abundant. Considering the mineral contents, seeds were relatively rich in potassium (2560.3-6697.5 mg kg^{-1}), phosphorus (529.8-1120,9 mg kg^{-1}), and magnesium (426- 1124,5 mg kg^{-1}). Moreover, the path diagram of phytochemical properties, nutritional value, and fatty acids of pumpkin seeds was determined. Consequently, the seeds of pumpkin cultivars were examined to find the best potential for a high nutritional value and contribution to the food industry.

Keywords: Pumpkin seeds, organic acid, amino acid, fatty acid, minerals.

INTRODUCTION

Pumpkin (*Cucurbita pepo* L.) is commercially grown in many regions of the world. The seeds of pumpkin have economic significance and are consumed as snack food in the Mediterranean region, particularly in Turkey and other Middle Eastern countries (Al-Khalifa, 1996). Recently, pumpkin silage is also used in cow feed owing to its valuable source of bioactive compounds (Halik *et al.*, 2018). Moreover, many growers and breeders have focused on pumpkins and their seeds, mainly aiming for drought resistance and high nutritional values, including proteins as well as high oil content (Idouraine *et al.*, 1996). For example, pumpkin seed is an abundant source of fatty acids and 98% of them contain oleic, linoleic, stearic and palmitic acids (Younis *et al.*, 2000; Murkovic *et al.*, 2004). One great source of polyunsaturated fatty acids is pumpkin seed oil,

38 representing about 84% of the total fatty acids of seeds (Procida *et al.*, 2013). Owing to these core
39 phytochemical components, pumpkin seeds were valued as an oil-rich source for nutritional
40 purposes (Stevenson *et al.*, 2007; Gohari *et al.*, 2011). The oil of pumpkin seed is currently used
41 for its medicinal properties, i.e., it is involved in the regulation of hypertension as well as the
42 mitigation of hypercholesterolemia and prevents benign prostatic hyperplasia proliferation
43 (Zuhair *et al.*, 2000; Fu *et al.*, 2006; Gossell-Williams *et al.*, 2006).

44 More than 2 billion people worldwide suffer from mineral deficiencies in many developing
45 countries. Pumpkin seeds have been indicated as considerable potential sources of potassium and,
46 to a lesser extent, sources of sodium, magnesium, calcium, phosphorus, zinc, iron, manganese,
47 and copper (Rezig *et al.*, 2012). Pumpkin seeds have also biochemical importance as a source of
48 phenylalanine, tyrosine and tryptophan (Tinoco *et al.*, 2012). As for amino acid content, it has
49 been reported that arginine and aspartic, glutamic acids were higher than threonine, methionine,
50 tyrosine, and histidine in seeds of previously studied pumpkin lines (Amin *et al.*, 2019).
51 Antioxidant activity of organic acids provide protection against a range of ailments in humans
52 (Nawirska-Olszanska *et al.*, 2014). To the best of our knowledge, data on organic acids in
53 pumpkin seeds have not been reported up-to-date. To qualify pumpkin seeds as a functional food,
54 it is important to ascertain the levels of organic acids in them. On the other hand, there is limited
55 knowledge regarding the interrelationships among the phytochemicals present in pumpkin seeds,
56 despite their rich nutritional content. Thus, understanding the relationship between the fatty acids
57 and phytochemical components in pumpkin seeds is of utmost importance. In addition to being an
58 excellent source of fatty acids, pumpkin seeds also serve as a significant phytochemical reserve.
59 However, a more detailed understanding of how phytochemical compounds are related to the fatty
60 acid composition of pumpkin seeds is necessary. Furthermore, uncovering these relationships will
61 aid in emphasizing the potential health benefits and nutritional value of pumpkin seeds, serving
62 as a valuable food source for the food industry, nutritionists, and consumers. Therefore, the
63 primary goal of the research was to determine the pathway between phytochemical and nutritional
64 values of pumpkin seeds. Another major objective was to contribute to their potential industrial
65 applications and to make recommendations as a prospective material for plant breeders to develop
66 enhanced functional crops.

67 68 MATERIAL AND METHODS

69 Material

70 The seeds of pumpkin (*Cucurbita pepo* L.) genotypes were collected from the following
71 prominent provinces of commercial pumpkin seed production in Turkey in 2020; Nevşehir (NVS-
72 1 and NVS-2), Konya (KNY), Kayseri (KYS-1 and KYS-2), Bursa (BRS), Edirne (EDR), and

73 Kırklareli (KRK). In these provinces, pumpkin seeds are produced from commercial populations
74 of national importance, but they have not been registered as cultivars. Moreover, these provinces
75 account for 86% of Turkey's overall pumpkin seed production. Seeds are sown in late April to
76 early May in these production areas, depending on weather and soil conditions, with harvesting
77 occurring after about four months. The minimum, maximum, and average temperatures of these
78 provinces ranged between 11°C and 17°C, 27°C and 30°C, and 20°C and 24°C, respectively.

79

80 **Methods**

81 **Proximate analysis**

82 The pumpkin seeds were dried to a constant weight prior to subsequent analyses. Moisture, ash
83 contents and crude protein (protein factor 6.25) were determined following the AOAC procedure
84 (1995). Total carbohydrate content and energy levels were calculated according to Idouraine *et al.*
85 (1996).

86

87 **Amino acid analysis**

88 Following the procedures described by Aristoy and Toldra (1991). The amino acid derivatives
89 were analyzed by HPLC (Agilent 1200, USA) on a Zorbax Eclipse-AAA 4.6 x 150 mm, 3.5 µm
90 columns at a temperature 40 °C with detection at 254 nm. The amino acids were expressed in
91 nmol µl⁻¹.

92

93 **Organic acid analysis**

94 A method of Gunes *et al.* (2014) was used to identify organic acids.. In summary, ultra turraks
95 were used to homogenize 1.0 g of the seed sample in 10 mL of deionized water. The mixture was
96 centrifuged at 1200 rpm for 50 min. Subsequently, supernatants were filtered through 0.22 µm
97 filters (Millex Millipore). The supernatants were transferred to glass vials and injected into HPLC
98 (Agilent 1200, USA) for separation using columns (Zorbax Eclipse-AAA 4.6 x 250 mm, 5 µm)
99 with UV detector at 220 nm absorbance. The column temperature was at 25 °C and the flow rate
100 was 1 mL min⁻¹. Organic acids were determined with 25 mM potassium phosphate (pH 2.5) as the
101 mobile phase.

102

103 **Fatty acid analysis**

104 Five sets of samples of each genotype were analysed described by Folch *et al.* (1957). The samples
105 (c. 1 g) were homogenized in a solution of chloroform/methanol (2:1, v/v) containing 0.01% (w/v)
106 of butylated hydroxytoluene (Sigma, Gas Chromatography (GC), B1378) as antioxidant 20 vol.
107 (w/v) for 1 min. The homogenized was carried out at 20-22°C on ice, filtration and incubation etc.
108 After the organic solvent was evaporated under a nitrogen stream, the amount of lipid was

109 measured gravimetrically. Fatty acid methyl esters (FAMES) were prepared as described by
110 Metcalfe and Schmitz (1961). The crude lipid extract was saponified with NaOH in methanol and
111 FAMES were prepared by transmethylation with boron trifluoride (BF₃) in methanol (Kaymak,
112 2014; Kaymak *et al.*, 2022). The samples and reference solution were analysed by a GC (Hewlett–
113 Packard 6890, USA) equipped with a flame ionization detector and a 7673A injector tower.
114 Methyl esters were separated on a DB 23 capillary column (Agilent, 60 m, 0.25 mm i.d. and 0.25
115 μm). The temperature program was set for 35 min at 190°C, after which it increased by 30°C per
116 min until 220°C, where it remained for 5 min. Hydrogen gas (2mL min⁻¹ and split ratio was 30:1)
117 was used as a carrier. By comparing their retention times and peak with a standard mix of fatty
118 acids (FAs) (“Supelco 37” component FAME mix, Cat No. 47885-U) the characteristic FAs were
119 identified and quantified (David *et al.*, 2003).

120

121 Mineral content

122 The mineral contents were determined according to AOAC official procedures 922.02 and 975.03
123 (AOAC, 1995). Moreover, tissue P, K, Ca, Mg, S, Na B, Fe, Mn, Zn, Cu, Cd, Mo, Ni and Pb were
124 determined by using an Inductively Couple Plasma Spectrophotometer (Perkin-Elmer, Optima
125 2100 DV, ICP/OES, Shelton, CT, USA). Vapodest 10 Rapid Kjeldahl Distillation Unit (Gerhardt,
126 Konigswinter, Germany) was used for determined the total N by Kjeldahl method (Bremner,
127 1996).

128

129 Statistical analyses

130 All analyses performed in this study were replicated five times. Statistical analyses were
131 performed using SPSS 20.0 (SPSS Inc., Chicago, IL, USA) software. The data were presented as
132 mean and assessed by variance analysis procedures (PROC GLM). The significant differences
133 between the mean values were compared by Duncan’s multiple range tests at $P \leq 0.05$ level.
134 Principal component analysis (PCA) was carried out to evaluate the relationships among
135 phytochemical content of seeds. In addition, AMOS 20 software (IBM Corporation, Chicago, IL)
136 was used to develop a structural equation model to show complex relationships among studied
137 variables.

138

139 RESULT AND DISCUSSION

140 Table 1 shows the chemical analysis results of the seeds in eight pumpkin genotypes. The moisture
141 content significantly ($P \leq 0.05$) differed among genotypes and varied from 4.45% (KRK) to 4.93%
142 (NVS-1 and 2). The protein contents revealed that genotype KRK (36.12%) indicated the highest
143 protein content while genotype KYS-2 (28.63%) had the lowest. The observed differences are

144 most likely caused by the results of genetic background. However, environmental factors
145 particularly location may have played a role in the differences (Idouraine *et al.*, 1996; Charaya *et*
146 *al.*, 2023). Accordingly, the differences in the results of the protein content among genotypes
147 could also arise from climatic conditions ranged between 11°C and 30°C during growing period.

148 In the study, the protein content of *Cucurbita pepo* seeds results is in line with Idouraine *et al.*
149 (1996), Younis *et al.* (2000), Rezig *et al.* (2012), and Nawirska-Olszanska *et al.* (2014). Crude oil
150 content was over 40% in genotypes BRS, NVS-1 and NVS-2 and no lower than 37% in the
151 remaining genotypes. Idouraine *et al.* (1996) reported that the observed variations might be related
152 to growing conditions, crop practices and harvest time. Additionally, the oil contents of the
153 genotypes are in the high range (13-33%) cited by Stevenson *et al.* (2007) and Rezig *et al.* (2012).
154 These outcomes are also equivalent to those reported by Idouraine *et al.* (1996) and Kaymak
155 (2012) but lower than those reported by Nawirska-Olszanska *et al.* (2014). As elucidated in Table
156 1, the seeds from genotype KYS-2 showed the highest level of total carbohydrates (21.54%)
157 whereas genotype KRK represented the lowest level (14.76%). No significant variation in the
158 carbohydrates was observed between seed samples, but these values were higher than that
159 recorded by Idouraine *et al.* (1996) and lower than those recorded by Younis *et al.* (2000). This
160 may vary concerning some environmental factors as indicated by Idouraine *et al.* (1996). Ash
161 content ranged from 9.40% (NVS-2) to 11.60% (NVS-1) and varied significantly among
162 genotypes. Calculated energy was high and varied significantly among genotypes. The energy
163 levels of present genotypes are similar to those reported by Idouraine *et al.* (1996). Moreover,
164 genotype KYS-1 (545.30 kcal 100⁻¹) showed the lowest energy, while genotype NVS-2 (578.99
165 kcal 100⁻¹) had the highest energy. It can be clearly said that the chemical analysis of seed from
166 the eight pumpkin genotypes exhibited higher oil and protein content compared to previous
167 studies.

168 The total yield of amino acids in the seeds of pumpkin genotypes ranged from 15.65 nmol µl⁻¹
169 (KNY) to 32.99 nmol µl⁻¹ (KYS-2). The genotype KYS-2 had the highest of all the amino acids
170 (Table 2). The lowest amino acid values changed according to the genotypes. Namely, glutamate,
171 glycine, tyrosine, tryptophan, leucine, hydroxyproline and proline were recorded at lower contents
172 in the KYS-1 genotype; isoleucine in EDR; phenylalanine and methionine in KRK. Thus, these
173 variations may result from the effect of different factors, including ripe stage, plant age, and crop
174 conditions (Song *et al.*, 2013). It is found that 22 amino acids in the seeds of pumpkin genotypes.
175 All of the essential amino acids (EAA) including threonine, valine, methionine, isoleucine,
176 leucine, phenylalanine, tryptophan, lysine, histidine, and arginine were detected in tested pumpkin
177 genotypes. Firstly, arginine and threonine turned out to be the predominant EAA, accounting for

178 1.00–2.13 nmol μl^{-1} and 1.05–2.12 nmol μl^{-1} , respectively. Secondly, histidine, methionine,
179 tryptophan and leucine were in a small amount at 0.52–1.10 nmol μl^{-1} , 0.40–1.02 nmol μl^{-1} , 0.45–
180 1.14 nmol μl^{-1} and 0.49–1.20 nmol μl^{-1} , respectively. Finally, valine, phenylalanine, isoleucine
181 and lysine were in a trace amount at 0.16–0.32 nmol μl^{-1} , 0.38–0.94 nmol μl^{-1} , 0.35–0.85 nmol μl^{-1}
182 ¹ and 0.32–0.72 nmol μl^{-1} , respectively. The composition of EAA in the eight pumpkin genotypes
183 is quite similar to that reported by previous studies (Mansour et al. 1993; Idouraine *et al.*, 1996;
184 Glew *et al.*, 2006), but the concentrations differed.

185 As explained in Table 3, 11 organic acids were determined. The results showed, the highest
186 organic acid content, except for oxalic, malonic, malic and citric acids were determined in the
187 KYS-2 genotype. Lactic and maleic acids were the predominant organic acids in KYS-2,
188 accounting for 3106.1 ng μl^{-1} and 2644.8 ng μl^{-1} , respectively. The lowest amino acid values were
189 changed based on the pumpkin genotypes. Low amounts of oxalic acid in tested pumpkin
190 genotypes were observed. Lactic, propionic, and tartaric acids differed in the tested pumpkin
191 seeds. In addition, organic acids, especially lactic and propionic acid, are known to exhibit good
192 antibacterial activity. Due to these properties, E.C., FAO/WHO and US-FDA has been approved
193 as a food additive (Surekha and Reddy, 2000). Tartaric acid is also one of the most important
194 organic acids, which has been widely applied in fields such as; food industry, pharmaceutical
195 industry, chemical industry (Zhang *et al.*, 2011).

196 Malic and citric acid, the main organic acids of horticultural crops, are responsible for the flavour,
197 taste and microbial stability of the products (Saavedra and Barbas, 2003). Moreover, citric acid
198 retard enzymatic activity by the way playing an important role in conjunction with antioxidants
199 to chelate trace metals (McCluskey *et al.*, 2004; Bellion *et al.*, 2006). Additionally, malic acid is
200 used as a common parameter to food control points in the food process and evaluate the quality
201 of agricultural products (Kim, 2006). The tested pumpkin genotype seeds are relatively rich in
202 organic acids, and the richness of organic acids in pumpkin seeds enables us to classify them as a
203 functional food.

204 In seeds of pumpkin genotypes, palmitic (C16:0), oleic (C18:1n-9) and linoleic (C18:2n-6) acids
205 were the highest in concentration, followed by stearic acid (C18:0) at less than 10%, and the other
206 fatty acids at an even lower content (<1%) (Table 4). Similar results have been documented by
207 Rezig *et al.* (2012), which found the major fatty acids of pumpkin seeds were oleic (44.11%),
208 linoleic (34.77%) and palmitic (15.97%) acids. Some researchers made the same observations as
209 four dominant fatty acids linoleic, oleic, palmitic, and stearic acids were determined in the seed
210 oil of pumpkin cultivars (Younis *et al.*, 2000; Stevenson *et al.*, 2007). The total saturated fatty
211 acids (SFA), monounsaturated fatty acids (MUFA), n-6 and n-3 polyunsaturated fatty acids

212 (PUFA) were also different. Similarly, Procida *et al.* (2013) reported that the content of these four
213 predominant fatty acids ranged from 97.5 to 98.7% of the total fatty acid content of tested pumpkin
214 seed oils of various origins. The results of our study are in agreement with different crop studies
215 made by Safdari-Monfared *et al.* (2019) and Akçalı (2022), who recorded that total fatty acid
216 content decreased due to the increasing temperature and sowing dates depending on different
217 ecologies.

218 The results propound that pumpkin seeds are a good source of the essential fatty acid, linoleic
219 acid. Concerning genotypes, wide variations were found related to the major fatty acids of tested
220 seeds (Table 4). SFA, MUFA, and n-6 and n-3 PUFA also differed among genotypes. Both n-6
221 and n-3 PUFA ranged from 26.29% (KYS-2) to 46.83% (KNY), and 0.16% (BRS) to 0.23%
222 (KNY), respectively; MUFA ranged from 35.60% (KNY) to 43.92% (KYS-2). In addition, all
223 seed oils contain low amounts (16-19%) of SFA. Such variations may be the result of variations
224 in cultivar, soil, and seasonal variation, stage of maturity, harvest time, drying conditions, and
225 storage (Al-Khalifa 1996; De Mello *et al.*, 2000). Furthermore, it is known that variations in oil
226 and fatty acid content are governed more by the genotypes than the growing location (Bhardwaj
227 and Hamama, 2009). Nawirska-Olszanska *et al.* (2014) also declared that the composition of fatty
228 acids differed depending on the variety and the species of the pumpkin seeds.

229 From Table 5, it is clear that the tested pumpkin seeds are excellent sources of potassium
230 magnesium (Mg) and phosphor (P), respectively. However, sodium (Na), sulfur (S), calcium (Ca),
231 iron (Fe), copper (Cu), boron (B), manganese (Mn) and zinc (Zn) levels were observed to be low.
232 In addition, cadmium (Cd) contents were the lowest, and lead (Pb) was lower in the genotypes but
233 did not prove to be statistically significant. The nitrogen (N) content of seeds ranged from 2.28%
234 (KYS-2) to 2.88% (KRK). However, environmental factors might be responsible for a small
235 portion of the differences observed (Idouraine *et al.*, 1996). The human body needs a variety of
236 minerals for almost all aspects of body function, such as potassium known for decreasing blood
237 pressure, magnesium known for its role in the structure, iron known for its essential component
238 of many of proteins and enzymes, copper known its crucial role in redox reactions and the
239 scavenging of free radicals, zinc known for its role in the structure of proteins and cell membranes
240 of the human body. Overall, the seeds of the Turkish pumpkin genotypes appeared to be a good
241 source of minerals. Although reporting different levels, Idouraine *et al.* (1996), Juranovic *et al.*
242 (2003), Glew *et al.* (2006) and Rezig *et al.* (2012) indicated similar trends for K, Mg, P, Na, and
243 the remaining minerals.

244 To assess the presence of any relationships among the 64 variables identified in the study, a
245 correlation analysis was carried out, which indicated the presence of multicollinearity among these

246 variables. To mitigate this multicollinearity issue, Principal Component Analysis (PCA) was
247 conducted. The PCA analysis revealed that the 9 parameters with the highest factor loadings
248 explained 94.93% of the total variance (Fig.1). A path analysis was also conducted to assess the
249 direct and indirect effects of the obtained 9 parameters on the dependent variables (C18:1n9 and
250 C18:2n6) (Fig.2). The path analysis showed that the Root Mean Square Approximation (RMSA)
251 value was significant at the 1% level.

252 The goodness-of-fit index (GFI), which measures the extent to which the covariance matrix is
253 captured by the model, was calculated as 0.865 (Excellent fit). Additionally, the comparative fit
254 index (CFI), which assesses whether there is any relationship between the variables and aims to
255 demonstrate the difference between the constructed model and the zero model, was computed as
256 0.915 (Excellent fit). Considering the path coefficients, it is seen in Fig.2 that the indirect effect
257 of lactic acid on the dependent variables (C:18 1n9 and C:18 2n6) has the highest coefficient
258 (801695.05) in terms of contribution to the model. Serine (52011.58), Alanine (277365.62),
259 Leucine (52106.96), Tryptophan (46267.06), Glycine (42350.87), Glutamate (31918.84),
260 Isoleucine (21414.17) and Malic acid (34.38) followed lactic acid, respectively.

261 262 CONCLUSIONS

263 The seeds of the Turkish pumpkin genotypes examined differed in most of the chemical
264 parameters. Some genotypes were distinguished from others by having higher protein, total
265 carbohydrates, and crude oil content. The seeds of the eight Turkish pumpkin genotypes are also
266 discovered to be a good source of K, Mg, P, and Na minerals. It is found that the studied
267 genotypes-dependent fatty acid content. Based on the results, the genotypes NVS-1, NVS-2, KYS-
268 1, and KYS-2 appeared to have the best potential in biochemical content for human health.
269 Through significant variances in the biochemical composition of Turkish pumpkin genotypes,
270 these genotypes can be selected, developed commercially, and registered as nutrient-rich cultivars.

271 272 REFERENCES

- 273 1. Akçalı, C. T. 2022. Increased Temperature and Shortened Grain Filling Duration Due to
274 Sowing Dates Significantly Affect Fatty Acids Composition of Corn (*Zea mays indentata*
275 Sturt.). *J. Agr. Sci. Tech*, **24**: 1143-1153.
- 276 2. Al-Khalifa, A.S. 1996. Physicochemical characteristics, fatty acid composition, and
277 lipoxygenase activity of crude pumpkin and melon seed oils. *J. Agric. Food Chem.*, **44**:
278 964–966.
- 279 3. Amin, M.Z., Islam, T., Uddin, M.R., Uddin, M.J., Rahman, M.M., Satter, M.A. 2019.
280 Comparative study on nutrient contents in the different parts of indigenous and hybrid

- 281 varieties of pumpkin (*Cucurbita maxima* Linn.). *Heliyon*, 5: e02462.
282 <https://doi.org/10.1016/j.heliyon.2019.e02462>
- 283 4. AOAC. 1995. Official methods of analysis, (15th ed.). Washington DC: Association of
284 Official Analytical Chemists
- 285 5. Aristoy, M.C., Toldra, F. 1991. Deproteinization techniques for HPLC amino acid analysis
286 in fresh pork muscle and dry-cured ham. *J. Agric. Food Chem.*, **39**: 1792-1795.
- 287 6. Bellion, M., Courbot, M., Jacob, C., Blaudez, D., Chalot, M. 2006. Extracellular and
288 cellular mechanisms sustaining metal tolerance in ectomycorrhizal fungi. *FEMS Microbiol.*
289 *Lett*, **254**: 173-181.
- 290 7. Bhardwaj, H.L., Hamama, A.A. 2009. Cultivar and growing location effects on oil content
291 and fatty acids in canola sprouts. *HortSci*, **44**: 1628-1631.
- 292 8. Bremner, J.M. 1996. Nitrogen total. Sparks D. L. (Ed.), *Methods of Soil Analysis. Part III.*
293 *Chemical Methods* 2nd ed (pp. 1085–1122). Madison, WI, USA
- 294 9. Charaya, A., Chawla, N., Dhatt, A. S., Sharma, M., Sharma, S., & Kaur, I. 2023. Evaluation
295 of biochemical composition of hulled and hull-less genotypes of pumpkin seeds grown in
296 subtropical India. *Heliyon*, e12995.
- 297 10. David, F., Sandra, P., Wylie, P.L. 2003. Improving the analysis of fatty acid methyl esters
298 using retention time locked methods and retention time databases. URL
299 <http://www.chem.agilent.com/Library/applications/5988-5871EN.pdf>. Accessed 11.03.14.
- 300 11. De Mello, M.L.S., Narain, N., Bora, P.S. 2000. Characterisation of some nutritional
301 constituents of melon (*Cucumis melo* hybrid AF-522) seeds. *Food Chem*, **68**: 411–414.
- 302 12. Folch, J., Less, M., Stanley, G.H.S. 1957. A simple method for the isolation and purification
303 of total lipids from animal tissues. *J. Biol. Chem*, **226**: 497–509.
- 304 13. Fu, C., Shi, H., Li, Q. 2006. A review on pharmacological activities and utilization
305 technologies of pumpkin. *Plant Foods Hum. Nutr*, **61**: 73-80.
- 306 14. Glew, R.H., Glew, R.S., Chuang, L.T., Huang, Y.S., Millson, M., Constans, D., Vanderjagt,
307 D.J. 2006. Amino acid, mineral and fatty acid content of pumpkin seeds (*Cucurbita* spp)
308 and *Cyperus esculentus* nuts in the Republic of Niger. *Plant Foods Hum. Nutr*, **61**: 51-56.
- 309 15. Gohari, A. A., Farhoosh, R., Haddad, K. M. 2011. Chemical composition and
310 physicochemical properties of pumpkin seeds (*Cucurbita pepo* Subsp. *pepo* Var. *Styriaka*)
311 grown in Iran. *J. Agr. Sci. Tech*, **13**: 1053-1063.
- 312 16. Gunes, A., Turan, M., Gulluce, M., Sahin, F. 2014. Nutritional content analysis of plant
313 growth-promoting rhizobacteria species. *Eur. J. Soil Biol*, **60**: 88-97.

- 314 17. Gossell-Williams, M., Davis, A., O'Connor, N. 2006. Inhibition of testosterone-induced
315 hyperplasia of the prostate of Sprague-Dawley rats by pumpkin seed oil. *J Med Food*, **9**:
316 284-286.
- 317 18. Halik, G., Lozicki, A., Wilczak, J., Arkuszewska, E., Makarski, M. 2018. Pumpkin
318 (*Cucurbita maxima* D.) Silage as a Feed that Improves Nutritional Properties of Cow's
319 Milk. *J. Agr. Sci. Tech*, **20**: 1383-1394.
- 320 19. Idouraine, A., Kohlhepp, E.A., Weber, C.W. 1996. Nutrient constituents from eight lines
321 of naked seed squash (*Cucurbita pepo* L.). *J. Agric. Food Chem.*, **44**: 721-724.
- 322 20. Juranovic, I., Breinhoelder, P., Steffan, I. 2003. Determination of trace elements in
323 pumpkin seed oils and pumpkin seeds by ICP-AES. *J Anal At Spectrom*, **18**: 54-58.
- 324 21. Kaymak, H.C. 2012. The relationships between seed fatty acids profile and seed
325 germination in cucurbit species. *Žemdirbystė-Agriculture*, **99**: 299-304.
- 326 22. Kaymak, H.C. 2014. Seed fatty acid profiles: Potential relations between seed germination
327 under temperature stress in selected vegetable species. *Acta Sci. Pol., Hortorum Cultus*, **13**:
328 119-133.
- 329 23. Kaymak H.C., Akan S., Yarali Karakan F. 2022. Pathway among Fatty Acid Profile, Seed
330 Germination, and Vigor of Watermelon Cultivars. *Emir J Food Agric*, **34**: 494-501.
- 331 24. Kim, M. 2006. Determining citrate in fruit juices using a biosensor with citrate lyase and
332 oxaloacetate decarboxylase in a flow injection analysis system. *Food Chem*, **99**: 851-857.
- 333 25. Mansour, E.H., Dworschak, E., Lugasi, A., Barna, E.Ä., Gergely, A. 1993. Nutritive value
334 of pumpkin (*Cucurbita pepo* Kakai 35) seed products. *J. Sci. Food Agric*, **61**: 73-78.
- 335 26. McCluskey, J., Herdman, L., Skene, K.R. 2004. Iron deficiency induces changes in
336 metabolism of citrate in lateral roots and cluster roots of *Lupinus albus*. *Physiologia*
337 *Plantarum* **121**: 586-594.
- 338 27. Metcalfe, L.D., Schmitz, A.A. 1961. The rapid preparation of fatty acid esters for gas
339 chromatographic analysis. *Anal Chem*, **33**: 363-364.
- 340 28. Murkovic, M., Piironen, V., Lampi, A.M., Kraushofer, T., Sontag, G. 2004. Changes in
341 chemical composition of pumpkin seeds during the roasting process for production of
342 pumpkin seed oil (Part 1: Nonvolatile compounds). *Food Chem* **84**: 359-365.
- 343 29. Nawirska-Olszanska, A., Biesiada, A., Sokol-Letowska, A., Kucharska, A.Z. 2014.
344 Characteristics of organic acids in the fruit of different pumpkin species. *Food Chem*, **148**:
345 415-419.

- 346 30. Procida, G., Stancher, B., Cateni, F., Zacchigna, M. 2013. Chemical composition and
347 functional characterisation of commercial pumpkin seed oil. *J. Sci. Food Agric*, **93**: 1035-
348 1041. <https://doi.org/10.1002/jsfa.5843>
- 349 31. Rezig, L., Chouaibi, M., Msaada, K., Hamdi, S. 2012. Chemical composition and profile
350 characterisation of pumpkin (*Cucurbita maxima*) seed oil. *Ind Crops Prod*, **37**: 82-87.
- 351 32. Saavedra, L., Barbas, C. 2003. Validated capillary electrophoresis method for small-anions
352 measurement in wines. *Electrophoresis* **24**: 2235-2243.
- 353 33. Safdari-Monfared, N., Noor-Mohammadi, G., Shirani Rad, A. H., Majidi Hervan, E. 2019.
354 Effect of Sowing Date and Glycinebetaine on Seed Yield, Oil Content, and Fatty Acids in
355 Rapeseed Cultivars. *J. Agr. Sci. Tech*, **21**: 1495-1506.
- 356 34. Song, J.F., Liu, C.Q., Li, D.J., Gu, Z.X. 2013. Evaluation of sugar, free amino acid, and
357 organic acid compositions of different varieties of vegetable soybean (*Glycine max* [L.]
358 Merr). *Ind Crops Prod*, **50**: 743-749.
- 359 35. Stevenson, D.G., Eller, F.J., Wang, L.P., Jane, J.L., Wang, T., Inglett, G.E. 2007. Oil and
360 tocopherol content and composition of pumpkin seed oil in 12 cultivars. *J. Agric. Food*
361 *Chem.* **55**: 4005-4013.
- 362 36. Surekha, M., Reddy, S.M. 2000. Preservatives. Classification and properties. In R. K.
363 Robinson CA, Batt Patel (Eds.), *Encyclopedia of food microbiology* (pp. 1710-1717). New
364 York: Academic Press.
- 365 37. Tinoco, L.P., Do, N., Porte, A., Porte, L.H.M., Godoy, R.L.O., Pacheco, S. 2012. Amino
366 acid profile of pumpkin seed flour. *UNOPAR Científica Ciências Biológicas e da Saúde*
367 *Journal* **14**: 149-153.
- 368 38. Younis, M.H., Ghirmay, S., Al-Shihry, S.S. 2000. African *Cucurbita pepo* L.: properties
369 of seed and variability in fatty acid composition of seed oil. *Phytochem*, **54**: 71-75.
- 370 39. Zhang, K., Wang, M., Gao, C. 2011. Tartaric acid production by ion exchange resin-filling
371 electrometathesis and its process economics. *J. Membr. Sci.* **366**: 266-271.
- 372 40. Zuhair, H.A., Abd El-Fattah, A.A., El-Sayed, M.I. 2000. Pumpkin seed oil modulates the
373 effect of feloipine and captopril in spontaneously hypersensitive rats. *Pharmacol. Res*, **41**:
374 555-563.

381

Table 1. Proximate composition of seeds of Turkish pumpkin genotypes ^a.

Genotypes	Moisture (%)	Protein (%)	Crude oil (%)	CHO ^b (%)	Ash (%)	Energy ^c (100 kcal ⁻¹)
KYS-1	4.49 ± 0.001 f	35.35 ± 1.20 ab	37.30 ± 1.34 b	17.05 ± 2.26 ^{NS}	10.30 ± 0.28 ab	545.30 ± 7.81 b 557.39 ± 1.65
KYS-2	4.53 ± 0.001 d	28.63 ± 0.43 c 31.29 ± 0.27	39.64 ± 0.24 ab	21.54 ± 1.38	10.20 ± 0.71 ab	ab 563.39 ± 3.47
NVS-1	4.93 ± 0.001 a	abc	41.96 ± 1.03 ab	15.15 ± 1.19	11.60 ± 0.42 a	ab 578.99 ± 16.15
NVS-2	4.93 ± 0.001 a	30.43 ± 1.14 bc	43.32 ± 3.12 a	16.86 ± 1.84	9.40 ± 0.14 b	a 557.85 ± 18.70
KRK	4.45 ± 0.003 g	36.12 ± 1.63 a 33.50 ± 3.80	39.37 ± 3.34 ab	14.76 ± 1.22	9.75 ± 0.50 b	ab 554.81 ± 1.56
EDR	4.51 ± 0.001 e	abc	39.28 ± 0.03 ab	16.82 ± 4.25	10.40 ± 0.42 ab	ab 560.10 ± 6.27
BRS	4.65 ± 0.001 c	30.31 ± 1.92 bc	40.38 ± 0.29 ab	18.86 ± 2.83	10.45 ± 1.20 ab	ab 560.57 ± 24.38
KNY	4.84 ± 0.001 b	30.13 ± 3.83 bc	39.95 ± 4.42 ab	20.12 ± 7.68	9.80 ± 0.57 b	ab

382 ^a Expressed on dry weight basis (mean ± SD, *n* = 5). Mean values with the different letter within columns are
 383 significantly different ($P \leq 0.05$). ^b CHO = carbohydrates, calculated by difference: 100 - (protein + crude fat + ash).

384 ^c Energy determined by multiplying fat by 9 and CHO and protein by 4.
 385

386

Table 2. Amino acid composition of seeds of Turkish pumpkin genotypes (nmol μl⁻¹).

Amino acids	Genotypes (n=5)							
	KYS-1	KYS-2	NVS-1	NVS-2	KRK	EDR	BRS	KNY
Aspartate	0.95 c	1.71 a	0.90 c	1.20 ab	0.89 c	0.92 c	1.27 ab	0.84 c
Glutamate	0.48 d	1.06 a	0.59 cd	0.71 b	0.52 d	0.53 d	0.69 bc	0.56 d
Asparagine	2.54 ab	2.77 a	1.41 d	2.08 bc	2.18 abc	1.91 cd	2.07 bc	1.44 d
Serine	1.97 c	4.17 a	1.92 c	2.73 b	2.07 c	2.33 bc	2.82 b	2.07 c
Glutamine	0.90 bcd	1.44 a	0.70 d	0.92 bcd	0.77 cd	0.93 bc	1.07 b	0.70 d
Histidine	0.67 b	1.10 a	0.55 c	0.70 b	0.64 b	0.72 b	0.71 b	0.52 c
Glycine	0.49 d	1.17 a	0.69 bc	0.71 bc	0.58 cd	0.61 cd	0.76 b	0.52 d
Threonine	1.37 b	2.12 a	1.32 bc	1.47 b	1.38 b	1.22 bc	1.39 b	1.05 c
Arginine	1.21 bc	2.13 a	1.00 d	1.50 b	1.22 bc	1.18 bc	1.32 ab	1.04 d
Alanine	1.60 c	3.09 a	1.38 d	1.90 b	1.61 c	1.58 c	1.90 b	1.33 d
Tyrosine	0.50 e	1.47 a	0.69 d	0.90 bc	0.77 cd	0.71 d	0.97 b	0.67 d
Cystine	0.61 bc	1.02 a	0.68 b	0.62 bc	0.70 b	0.64 bc	0.69 b	0.50 c
Valine	0.20 bc	0.32 a	0.16 c	0.20 bc	0.22 bc	0.22 bc	0.22 b	0.22 bc
Methionine	0.41 c	1.02 a	0.58 bc	0.53 bc	0.40 c	0.53 bc	0.61 b	0.46 bc
Tryptophan	0.45 d	1.14 a	0.58 cd	0.70 bc	0.49 d	0.54 d	0.79 b	0.52 d
Phenylalanine	0.46 c	0.94 a	0.47 c	0.67 b	0.38 c	0.44 c	0.68 b	0.47 c
Isoleucine	0.56 b	0.85 a	0.51 b	0.50 b	0.50 b	0.35 c	0.51 b	0.37 c
Leucine	0.49 d	1.20 a	0.54 cd	0.75 b	0.51 cd	0.62 c	0.80 b	0.52 cd
Lysine	0.45 bc	0.72 a	0.39 cd	0.48 bc	0.40 bcd	0.40 bcd	0.50 b	0.32 d
Hydroxyproline	0.39 c	0.87 a	0.41 c	0.56 b	0.40 c	0.43 c	0.59 b	0.43 c
Sarcosine	1.44 c	2.59 a	1.21 d	1.71 b	1.49 c	1.48 c	1.69 b	1.06 d
Proline	0.04 d	0.09 a	0.07 b	0.05 c	0.05 c	0.05 d	0.06 bc	0.05 c
Total	18.18	32.99	16.74	21.57	18.17	18.34	22.13	15.65

387 The number in parenthesis indicates the number of specimens of each genotype sampled. The different lowercase
 388 letters within rows indicate a significant difference ($P \leq 0.05$) between the genotypes.
 389

390

391

392

393

394

395

Table 3. Organic acid composition of seeds of Turkish pumpkin genotypes (ng μl^{-1})

Organic acids	Genotypes (n=5)							
	KYS-1	KYS-2	NVS-1	NVS-2	KRK	EDR	BRS	KNY
Oxalic acid	27.4 d	15.6 e	21.2 de	39.3 c	39.4 c	52.6 b	45.2 c	59.8 a
Propionic acid	342.9 b	461.7 a	350.6 b	237.1 d	290.9 c	356.5 b	251.1 cd	464.7 a
Tartaric acid	17.3 bc	14.3 c	24.7 a	24.9 a	18.7 bc	13.9 c	20.2 ab	17.3 bc
Butyric acid	168.9 e	489.4 a	263.0 c	196.1 de	178.5 de	216.9 d	214.6 de	321.5 b
Malonic acid	57.9 d	71.9 ab	61.8 bcd	80.3 a	58.9 cd	70.9 abc	61.8 bcd	56.8 d
Malic acid	14.6 c	23.3 abc	27.1 ab	24.0 abc	24.9 ab	30.3 ab	20.9 bc	31.8 a
Lactic acid	277.8 c	3106.1 a	384.8 e	878.7 c	310.3 e	321.9 e	1276.7 b	671.7 d
Citric acid	471.2 bc	100.4 g	406.4 de	518.0 ab	428.5 cd	547.4 a	280.7 f	351.5 e
Maleic acid	47.4 e	2644.8 a	648.7 c	250.7 d	49.1 e	53.7 e	1063.4 b	78.8 e
Fumaric acid	74.3 c	146.4 a	117.7 ab	73.5 c	67.9 c	99.4 bc	111.7 b	85.1 bc
Succinic acid	498.0 e	1033.1 a	835.1 b	716.1 c	472.4 e	623.6 d	716.8 c	537.8 e

396 The number in parenthesis indicates the number of specimens of each genotype sampled. The different lowercase
 397 letters within rows indicate a significant difference ($P \leq 0.05$) between the genotypes.
 398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

Table 4. Fatty acid composition of seeds of Turkish pumpkin genotypes (%).

Fatty acids	Genotypes (n=5)							
	KYS-1	KYS-2	NVS-1	NVS-2	KRK	EDR	BRS	KNY
C16:0	10.45 ± 0.10 bc	11.38 ± 0.10 abc	10.73 ± 0.33 abc	10.91 ± 0.83 abc	12.31 ± 0.11 a	11.18 ± 0.51 abc	11.95 ± 0.58 ab	10.29 ± 1.34 b
C16:1n7	0.19 ± 0.01 ^{ns}	0.12 ± 0.02	0.12 ± 0.00	0.14 ± 0.02	0.27 ± 0.17	0.20 ± 0.04	0.14 ± 0.00	0.12 ± 0.03
C18:0	6.95 ± 0.33 ab	7.70 ± 0.14 a	6.51 ± 0.20 abc	5.99 ± 1.41 a- d	5.45 ± 0.14 bcd	4.72 ± 0.64 d	4.90 ± 0.25 cd	6.58 ± 1.21 abc
C18:1n9	37.71 ± 0.30 ab	43.81 ± 3.55 a	37.25 ± 1.03 ab	38.19 ± 5.30 ab	37.69 ± 1.33 ab	42.23 ± 4.73 ab	42.37 ± 2.08 ab	35.49 ± 2.41 b
C18:2n6	44.02 ± 0.55 ab	36.29 ± 3.53 b	44.78 ± 1.61 ab	44.13 ± 7.65 ab	43.65 ± 1.15 ab	41.13 ± 5.92 ab	40.08 ± 1.24 ab	46.83 ± 0.29 a
C18:3n3	0.21 ± 0.00 ab	0.17 ± 0.01 ab	0.17 ± 0.01 ab	0.20 ± 0.01 ab	0.20 ± 0.04 ab	0.19 ± 0.02 ab	0.16 ± 0.00 b	0.23 ± 0.04 a
C20:0	0.47 ± 0.03 ab	0.54 ± 0.01 a	0.44 ± 0.06 ab	0.45 ± 0.14 ab	0.43 ± 0.00 ab	0.36 ± 0.10 b	0.39 ± 0.01 ab	0.46 ± 0.07 ab
SFA	17.87 ± 0.26 ^{ns}	19.61 ± 0.03	17.68 ± 0.58	17.35 ± 2.38	18.19 ± 0.03	16.25 ± 1.25	17.24 ± 0.84	17.34 ± 2.62
MUFA	37.91 ± 0.29 ab	43.92 ± 3.57 a	37.37 ± 1.03 ab	38.32 ± 5.28 ab	37.96 ± 1.16 ab	42.44 ± 4.69 ab	42.52 ± 2.08 ab	35.60 ± 2.38 b
n-6 PUFA	44.02 ± 0.55 ab	36.29 ± 3.53 b	44.78 ± 1.61 ab	44.13 ± 7.65 ab	43.65 ± 1.15 ab	41.13 ± 5.92 ab	40.08 ± 1.24 ab	46.83 ± 0.29 a
n-3 PUFA	0.21 ± 0.00 ab	0.17 ± 0.01 ab	0.17 ± 0.01 ab	0.20 ± 0.01 ab	0.20 ± 0.04 ab	0.19 ± 0.02 ab	0.16 ± 0.00 b	0.23 ± 0.04 a

421 C16:0 – palmitic acid, C16:1n-7, palmitoleic acid, C18:0 – stearic acid, C18:1n-9 – oleic acid C18:2n-6 – linoleic acid, C18:3n-3 – linolenic acid, C20:0 – arachidic acid, SFA
422 – saturated fatty acids, PUFA – polyunsaturated fatty acids, MUFA – monounsaturated fatty acids; means followed by different small letters in rows are significantly different
423 at $P \leq 0.05$., ns – not significant. The number in parenthesis indicates the number of specimens of each genotype sampled.
424


Table 5. Mineral contents of seeds of Turkish pumpkin genotypes (mg kg⁻¹).

Minerals	Genotypes (n=5)							
	KYS-1	KYS-2	NVS-1	NVS-2	KRK	EDR	BRS	KNY
B	5.84 a	4.68 a	1.14 b	5.07 a	5.36 a	5.70 a	6.20 a	2.21 b
Ca	68.66 b	63.53 b	13.49 c	32.67 c	78.87 b	107.01 a	63.96 b	34.11 c
Cd	0.03 b	0.03 b	0.09 a	0.03 b	0.03 b	0.02 b	0.03 b	0.08 a
Cu	6.77 a	4.07 c	0.56 e	3.98 c	5.28 b	5.66 b	5.16 b	2.59 d
Fe	7.76 a	4.41 c	1.51 e	3.23 d	6.69 b	7.22 ab	6.26 b	2.54 d
K	5064.50 abc	4353.50 bc	6697.50 a	5277.00 abc	4535.40 bc	5789.50 ab	2560.35 d	3542.90 cd
Mg	1124.55 a	775.65 c	426.00 d	735.30 c	924.45 b	1109.10 a	919.65 b	479.93 d
Mn	4.82 a	2.70 b	0.77 c	2.63 b	5.12 a	5.40 a	3.06 b	2.85 b
Mo	0.51 bcd	0.66 abc	0.71 ab	0.27 d	0.34 cd	0.62 a-d	0.87a	0.50 bcd
Na	19.59 c	81.21 b	157.50 a	132.80 a	13.34 c	17.66 c	22.35 c	22.82 c
Ni	0.50 bc	1.29 a	0.63 b	0.41 bc	0.27 c	0.69 b	0.69 b	0.41 bc
P	1014.60 ab	651.00 cd	859.46 bc	529.80 d	921.00 ab	1120.95 a	890.55 ab	889.55 ab
Pb	0.11 ^{ns}	0.04	0.08	0.05	0.11	0.05	0.09	0.11
S	128.70 a	46.67 c	85.45 b	58.51 c	63.66 bc	59.28 c	72.63 bc	58.28 c
Zn	4.71 a	2.51 bc	1.66 dc	1.88 cd	4.32 a	5.21 a	3.09 b	0.88 d
N (%)	2.82 ab	2.28 c	2.42 bc	2.49 abc	2.88 a	2.67 abc	2.41 bc	2.40 bc

426

427


The number in parenthesis indicates the number of specimens of each genotype sampled. The different lowercase letters within rows indicate a significant difference ($P \leq 0.05$) between the genotypes, ns – not significant.



428

429

Figure 1. Principle component analyses of phytochemical content of pumpkin seeds.



430
431
432

Figure 2. Pathway among C18:1n9, C18:2n6 and phytochemical properties of pumpkin seeds.