

Effects of Diverse Microclimates and Soil Water Contents on Water-Use Efficiency and Carbon Isotope Discrimination for Bush Bean

M. Raeini-Sarjaz^{1*} and V. Chalavi²

ABSTRACT

Environmental variables, including soil water content (SWC), act as constraints on crop growth and productivity. Therefore, open air (E_0), perforated (E_1) and non-perforated (E_2) plastic housings were used with well-watered (W_0), moderately-watered (W_1) and water-stressed (W_2) bush bean plants to explore the relationships between water-use efficiency (WUE), carbon isotope discrimination (Δ) and isotopic composition (δ_p), leaf assimilation rate (A) and leaf Kjeldahl nitrogen (N) under diverse environments. The CO_2 concentration and air carbon isotopic composition (δ_a) varied with the environment. The δ_a values were reduced by about 0.8×10^{-3} and 3.8×10^{-3} in E_1 and E_2 , respectively, compared with that in E_0 . SWC significantly affected WUE, Δ , δ_p in both E_0 and E_1 but not in E_2 . The decoupling of plants from the outside atmosphere might have contributed in maintaining the above quantities almost constant in E_2 . The Δ -value increased by about 2.2×10^{-3} in E_0 and 1.7×10^{-3} in E_1 compared with E_2 . Water stress reduced the Δ -value by about 1.1×10^{-3} in both E_0 and E_1 . WUE and Δ were significantly correlated in E_0 and E_1 ($r = -0.72$, and -0.75 , respectively) whereas there was no definite relationship between WUE and Δ in E_2 indicating that stomatal conductance was almost independent of SWC. The N-content had little effect on Δ . Leaf N significantly increased in water-stressed plants depending upon the time of harvest and the environment. The mean leaf assimilation rate was significantly higher in E_0 than in either E_1 or E_2 .

Keywords: Bush bean, Carbon isotope discrimination, Plastic tunnel, Water-use efficiency.

INTRODUCTION

Two stable isotopes of carbon occur naturally in the atmosphere as $^{12}CO_2$ and $^{13}CO_2$ with their respective quantities of 98.9% and 1.1% (Fritz and Fountner, 1980; Farquhar, *et al.*, 1989; Yeh and Wang, 2001). Although the isotope effect due to mass difference is usually a nuisance in radiotracer methodology, the same effect can be turned around and used as a tool especially in studying chemical reactions that proceed in tandem. Isotope effects occur in plant tissues due to

differences in the diffusivities of $^{13}CO_2$ and $^{12}CO_2$ in the ambient air (Farquhar *et al.*, 1989) and also in biochemical reactions involved in photosynthesis (Melander and Saunders, 1979; O'Leary *et al.*, 1981; Farquhar *et al.*, 1989; Yeh and Wang, 2001). Since crops encounter different environmental conditions during their growth, the ^{13}C to ^{12}C ratio varies significantly in the tissues of C_3 plants (Farquhar *et al.*, 1982; Yeh and Wang, 2001). This variation in the ratio of isotopes can be used to evaluate the effects of genetic and environmental factors

1. Department of Irrigation, Faculty of Agricultural Engineering, University of Mazandaran, Sari, Islamic Republic of Iran.

2. Department of Horticulture, Faculty of Agricultural Sciences, University of Mazandaran, Sari, Islamic Republic of Iran.

* Corresponding author, e-mail: m.raeini@umz.ac.ir



on the yield performance of cultivars. Any environmental factor that affects stomatal conductance and enzymatic activity may result in changes in water-use efficiency (WUE) and ^{13}C discrimination (Δ) (Farquhar *et al.*, 1982) as defined in Eq. (2).

The theory of isotope effect has been established through a linear relationship between Δ and the ratio C_i/C_a of the internal CO_2 concentration in the plant tissue, C_i , to that of the ambient air, C_a (Farquhar and Richards, 1984). Ambient CO_2 concentration (C_a) is almost constant in a wide range of environmental conditions, while internal CO_2 concentration (C_i) can vary as a photosynthetic response to environmental variables. Eq. (1) describes the relationship between the WUE_i of a single leaf defined as the ratio of the instantaneous photosynthetic and transpiration rates ($\text{mol CO}_2/\text{mol H}_2\text{O}$) and C_i/C_a (Farquhar *et al.*, 1989),

$$\text{WUE}_i = \frac{C_a(1 - C_i/C_a)}{1.6v} \quad (1)$$

where, v = water vapor pressure difference between the intercellular spaces and the ambient air. The factor 1.6 is the ratio of the diffusivity of water vapor to that of CO_2 in the air. A negative correlation was found between Δ and both the long-term transpiration efficiency, WUE_t , defined as the total dry matter per kg of water transpired (Farquhar and Richards, 1984; Ehdaie *et al.*, 1991; Ismail and Hall, 1992; Raeini-Sarjaz *et al.*, 1998; Ebdon *et al.*, 1998) and WUE_i (Wright *et al.*, 1988 and 1994) for several crops. For legumes, Meinzer *et al.* (1990) reported simultaneous reductions in stomatal conductance and in Δ with increased WUE_t for water-stressed cowpea. Ehleringer *et al.* (1991) found a high correlation between C_i/C_a and Δ for common bean. Wright *et al.* (1994) demonstrated a significant effect of various water regimes on Δ for peanut. Ehdaie *et al.* (1991) obtained higher Δ -values for greenhouse-grown wheat than those grown in the field. Hall *et al.* (1990) observed an association between gas exchange and Δ when the roots of the same cowpea genotypes were subjected to varying environments. Rao and Wright (1994) showed a

significant effect of location and water regime on Δ for cowpea. Johnson *et al.* (1995) observed a significant correlation between Δ and WUE_t for lentil. Hubick (1990) reported that low-N peanut plants accumulated less dry matter and used less water than the high-N plants.

Water use efficiency can be increased if the major factors that influence water loss are evaluated. For example, the rate of transpiration from a canopy is mainly a function of stomatal and boundary layer conductances, water vapor pressure deficit (VPD), net radiation, wind speed, and temperature. The effect of each of these factors may depend on the canopy structure and the surrounding growing environment. The canopy boundary layer conductance may have a crucial impact on the relation between water use and isotope discrimination of plants inside plastic housing. In the open air, wind increases canopy conductance and enhances mass and heat transfer, while in enclosed environments, the boundary layer conductance is relatively small, which lowers the heat and mass transfer processes (Jarvis and McNaughton, 1986; Jones, 1992). The open air canopy is well-coupled to the atmosphere and transpiration is mostly controlled by stomatal conductance, while inside a plastic housing the canopy is decoupled from the outside air and energy input becomes the governing factor for transpiration (Jones, 1992).

The CO_2 concentrations of a modified atmosphere and under enclosed environments, such as plastic tunnels and greenhouses, may be higher than that of open air (Yeh and Wang, 2001). Long-term elevated CO_2 concentrations will lead to a stomatal conductance reduction (Jones and Jongen, 1996; Pospisilova and Catsky, 1999), and therefore, will tend to reduce leaf transpiration rate and increase WUE (Jones and Jongen, 1996; Saralabai *et al.*, 1997). Hence, the increase of water use efficiency might be the positive effect of environmental elevated CO_2 (Pospisilova and Catsky, 1999). Although bulk air δ_a value is around -8×10^{-3} (Freidli *et al.*, 1986; Goodman and Francey,

1988), but elevated CO₂ concentrations under enclosed environments, due to C₃ organic matter decomposition, may reduce ¹³CO₂ enrichment (Clark and Fritz, 1997), therefore it affects air source CO₂ composition.

Although literature on ¹³C discrimination is extensive in plant breeding, plant physiology, eco-physiology and other fields (Ehleringer *et al.*, 1993), the interaction of crop growth and soil moisture on Δ needs to be examined for diverse environments so that a suitable growing environment can be determined for a particular crop. The physical environment can be modified to create a favorable microclimate for optimum plant growth. For example, plastic culture is being increasingly used, especially in temperate climates, to promote early-season vegetable production and to reduce the detrimental effects of low air and soil temperatures. Plant physiological responses, especially long-term stomatal conductance, to these artificial microclimates are not well known. ¹³C discrimination, as a long term indicator of C_i/C_a, stomatal conductance, and a probe of long-term WUE (Farquhar *et al.*, 1989; Condon *et al.*, 1990; Hubick, 1990; Brugnoli and Farquhar, 2000) may provide a tool to assess responses of plants to changes in the growing environments. Thus, the objectives of this study were to examine whether decoupling plants from the outside atmosphere (closed plastic housings), or providing facilities for re-coupling through holes (perforated plastic enclosures) in combination with soil moisture availability may influence the WUE_t, leaf N, and Δ of the bush bean.

MATERIALS AND METHODS

Well-watered (W₀), moderately-watered (W₁), and water-stressed (W₂) plants were used in combination with the experimental environments of the open air (E₀), a plastic housing perforated uniformly with 400 holes m⁻² each of 0.5 cm diameter (E₁), and a closed plastic cover (E₂). The housings were made of clear polyethylene plastic sheets,

which were transformed into tunnels of 1 m in width, 10 m in length, and 0.8 m in height. The plastic sheets were 87% transparent (Raeini-Sarjaz and Barthakur, 1997) to photosynthetic photon flux density (PPFD). Bush bean seeds (*Phaseolus vulgaris* L. cv. Provider) were germinated in the greenhouse as previously described (Raeini-Sarjaz and Barthakur, 1997). The one-week old seedlings were transferred from the greenhouse to the sites E₀, E₁, and E₂ when the ambient temperature was not harmful for the open air plants, on May 15, Macdonald Campus, Experimental Farm, McGill University (45°25' 45"N and 73°56'00"W). All plants were initially watered to 100% field capacity (FC). The soil water content (SWC) of W₀ plants was kept at 100% FC, while for W₁ and W₂ plants water was supplied to 100% FC only when SWC reached 50% and 30% FC, respectively. Two control pots with an identical amount of soil but without plants were watered and weighed similarly to monitor non-plant evaporative losses. The transpiration rate was calculated from the difference between water added and lost.

Water was replenished every day and the loss was measured with a balance. CO₂ concentrations were measured instantaneously at each site using a photosynthesis system (LI-6200, LI-COR Inc., Lincoln, NE, USA). During growth and development, plants were fertilized uniformly each week with 20-20-20 NPK. To reduce leaf sunburn during sunny and hot days, the tunnel ends were opened at noon for three hours of ventilation. At mid-June with the increasing air temperature during the pre-flowering stage (Day 35), the plastic covers were removed. Shelter was provided whenever there was rain or a risk of rain. In order to evaluate the effects of crop growth conditions on yield and WUE, half of the plants were harvested after the removal of the plastic cover (HT1= 35 d), and the rest were kept in ambient air until pod harvest time (HT2= 50 d).

For each harvest, five pots from each treatment were selected randomly, and the plant parts were put into paper bags. The



contents were dried in an oven at 60°C for 72 h to determine the total dry matter (TDM) with an electronic balance of ± 0.001 g precision, and WUE_t was calculated. Leaf N at both harvest times was measured on 0.2 ± 0.005 g of subsamples of ground leaf tissue using Automated Ion Analyzer (Lachat Instruments, model Quikchem AE, USA).

To evaluate the effect of previous soil moisture conditions on leaf gas exchange (LGE) in each experiment, plants were transferred to the greenhouse before measurements. The LGE measurements were made using a steady-state LI-6200 photosynthesis system (LI-COR, Inc., Lincoln, NE, USA). The measurements were made on newly developed trifoliate leaves of recently watered (100% FC) plants under going all water regime treatments between 11:00 and 14:00 local time on bright sunny days. The mean air temperature (37°C) and relative humidity (56%) were constant during the LGE measurements.

Carbon Isotope Discrimination

Trifoliate leaves at harvest times HT1 and HT2 were dried at 60°C for 72 h and then ground to a fine powder using a Wiley mill. From each sample, 4 to 5 mg subsamples of leaf tissues were combusted under vacuum using Vycor tubes containing silver wire and cupric oxide. Combustion took place at 820°C for 5 h to release CO₂. Combusted samples were left at room temperature for 12 h prior to the liquid N cryogenic purification of CO₂. Eq. (2) was used to calculate carbon isotope discrimination (Δ) from the measurements of $\delta_p = (R_p/R_s - 1)$, where R_p is the ¹³C/¹²C ratio in the plant and R_s that of Pee Dee Belemnite (PDB) standard (Ehleringer and Osmond, 1989).

$$\Delta = \frac{(\delta_a - \delta_p)}{(1 + \delta_p)} \quad (2)$$

where, δ_a and δ_p are isotopic composition or enrichment of air and plant samples, respectively.

To measure carbon isotopic composition of

the air, δ_a , at each site, air was collected in special aluminum bags (Mil-B-131H, Ludlow Corp.) and air CO₂ was immediately purified cryogenically. The isotopic composition measurements for duplicated plant samples were made by an isotope ratio mass spectrometer (VG T50 GAS 903D Device, Middlewich, UK). The mean of the duplicated samples of δ_p is reported along with Δ .

Statistical Analysis

Each of the three experiments of E₀, E₁, and E₂ was conducted on a 10-replicate completely random design model. Variances of the experiments were found to be homogeneous using Bartlett's test. A combined analysis of variance (ANOVA) was employed for the entire data of three experiments using SAS software (SAS Institute Inc., Cary, NC, 1990), where sources of variations were environment (E), soil water content (SWC), harvest time (HT), and their interactions. An ANOVA was performed on the data from each experiment separately, where SWC and HT were the sources of variation. For LGE analysis, leaf temperature and PPFD were employed as covariates in the model. Student-Neuman-Keuls' post hoc tests were used to identify significant effects of treatments. For determining correlations, the Pearson correlation procedure was employed, and an unpaired Cochran's t-test was used to compare harvest time results within each experiment.

RESULTS

The mean CO₂ concentrations were 453, 732 and 1478 $\mu\text{mol mol}^{-1}$ within E₀, E₁ and E₂ environments, respectively. The measured δ_a values were: -8×10^{-3} in E₀, -8.8×10^{-3} in E₁, and -11.8×10^{-3} in E₂. The mean day-time air temperature and relative humidity of the three sites, E₀, E₁ and E₂, during the last two weeks of May were 24.6, 32.4, 34.5°C and 65, 75, 95%, respectively.

The result of the combined ANOVA statis-

tics showed that environmental diversity had significant ($p < 0.001$) effects on N, Δ , δ_p , WUE_t , and the assimilation rate (A). Therefore, each of the environmental data was analyzed separately by ANOVA. Leaf N of W_2 plants were significantly higher than those of W_0 in E_0 at HT1; and in E_2 environment regardless of harvesting times (Table 1). The previous SWC history had little effect on A, and there was no significant interaction between SWC and environment, and so the pooled data of A for different SWC were run to test the effect of various environments. E_0 significantly increased the assimilation rate compared with E_1 and E_2 . The mean A-values increased in the order of $E_0 > E_1 > E_2$ environments (Table 1). A

definite relationship did not emerge between N and Δ except showing a correlation at the HT1 harvest in E_0 ($r = -0.81$; $p < 0.01$).

There were significant differences in both ^{13}C enrichment (δ) and ^{13}C discrimination (Δ) values at both HT1 and HT2 in E_0 and E_1 for different water regimes, while such differences almost disappeared in E_2 . WUE_t generally increased with reduction of soil water contents in both E_0 and E_1 at both HT1 and HT2 stages, while WUE_t remained almost constant along different SWC in E_2 (Table 1). WUE_t and Δ relationships were linear and significant in both E_0 ($r = -0.76$, $p < 0.02$) and E_1 environments ($r = -0.75$, $p < 0.02$), while no significant relationship was found ($r = 0.18$, $p > 0.6$) with E_2 (Figure 1).

Table 1. Mean leaf N ($\mu\text{g}/\text{mg}$), Δ ($\times 10^{-3}$), δ ($\times 10^{-3}$), WUE_t (g DM/ kg H_2O) and leaf assimilation rate, A ($\mu\text{mol m}^{-2} \text{s}^{-1}$) with harvest time (HT), environment (E) and soil water content (SWC) for bush bean.

Environment	HT	SWC	N	Δ	δ	WUE_t	A
E_0	1	W_0	36.44 ^b	19.55 ^a	-27.03 ^b	5.99 ^b	15.75
		W_1	37.93 ^b	19.29 ^a	-26.78 ^b	6.09 ^b	16.10
		W_2	48.05 ^a	18.42 ^b	-25.94 ^a	6.65 ^a	16.61
	Mean		40.81	19.08	-26.57	6.24	16.15 ^x
E_1	1	W_0	33.20	19.01 ^a	-27.29 ^b	5.68 ^b	14.13
		W_1	32.21	18.95 ^a	-27.24 ^b	6.52 ^a	12.75
		W_2	34.55	17.99 ^b	-26.32 ^a	6.98 ^a	15.75
	Mean		33.32	18.65	-26.95	6.72	14.21 ^y
E_2	1	W_0	30.80 ^b	17.13	-28.51	5.99	12.58
		W_1	34.61 ^b	16.90	-28.28	5.80	15.48
		W_2	41.76 ^a	16.69	-28.09	5.98	13.50
	Mean		35.72	16.91	-28.29	5.92	13.85 ^y
E_0	2	W_0	50.53	19.56 ^a	-27.03 ^b	3.55 ^b	
		W_1	55.58	18.59 ^b	-26.09 ^a	3.63 ^b	
		W_2	52.33	18.58 ^b	-26.11 ^a	4.64 ^a	
	Mean		52.81	18.91	-26.41	3.94	
E_1	2	W_0	48.38	20.34 ^a	-27.78 ^b	3.67 ^b	
		W_1	46.52	20.01 ^{ab}	-27.47 ^{ab}	3.64 ^b	
		W_2	43.78	19.42 ^b	-26.90 ^a	4.12 ^a	
	Mean		46.22	19.92	-27.38	3.81	
E_2	2	W_0	43.85 ^b	19.95	-27.41	3.29 ^b	
		W_1	50.59 ^a	19.61	-27.04	3.51 ^b	
		W_2	53.85 ^a	19.57	-27.09	3.77 ^a	
	Mean		49.43	19.71	-27.18	3.52	

Different letters of ^a and ^b show significant differences across different SWC ($p < 0.05$), while ^x and ^y show significant differences between different environments.




Figure 1. Relations between carbon isotope discrimination (Δ) and water-use efficiency (WUE_t) in open (E_0), perforated (E_1) and closed (E_2) growth environments. Legends: filled circles hallow circles and filled triangles represent well-watered (W_0), moderate-watered (W_1) and water-stressed (W_2) plants, respectively.

The t-test showed significant differences ($p \leq 0.04$) in Δ -values between HT1 and HT2 for E_1 and E_2 , but not for E_0 . The δ_p values showed significant differences along soil

water contents in both E_0 and E_1 , decreasing with an increase in SWC (Table 1).

DISCUSSION

Although WUE_t enhancements for water-stressed bean plants in E_0 and E_1 environments may seem to be unlikely at first glance, yet several authors (Nobel, 1991; Ismail and Hall, 1992; Raeini-Sarjaz and Barthakur, 1997; Raeini-Sarjaz *et al.*, 1998) have concurred with the present finding. An explanation for this apparent anomaly can be found in decreased biomass production and a drastic reduction in the rate of transpiration for the plants growing in these environments. The plants were decoupled from the outside atmosphere and a high relative humidity and temperature prevailed in E_2 (Raeini-Sarjaz and Barthakur, 1997), which might have prevented them from obtaining a similar enhancement of WUE_t in E_2 . For example, the mean daytime air temperature and relative humidity during the last two weeks of May were 24.6, 32.4, 34.5°C and 65, 78, 95% in E_0 , E_1 and E_2 environments, respectively.

The less negative δ or lower Δ value in plant tissue corresponds to the lesser discrimination against ^{13}C and is an indicator of environmental stresses imposed on stomatal conductance. To compare these values, the source (air) CO_2 should have the same enrichment in ^{13}C , δ_a . The less negative mean δ values in E_0 and E_1 environments (-26.57×10^{-3} and -26.95×10^{-3} , respectively) compared with that of E_2 environment (-28.29×10^{-3}) can be speculated to be the result of a greater stress condition imposed on E_0 and E_1 plants, while the lower Δ values on plants of the E_2 environment compared with those of other environments contradicts the δ values. The Δ values increased in E_0 and E_1 environments by about 2.2×10^{-3} and 1.7×10^{-3} , respectively, compared with E_2 at HT1. As Δ values were adjusted against the source isotopic composition, therefore, the reduction in Δ values of E_2 must be due to other environmental stresses imposed on stomatal behavior other than water stress alone. Ehdaie *et al.* (1991) reported higher Δ -values for greenhouse-grown plants at higher humidity

than those of field-grown crops in less humid air. Our results were not in agreement with the above finding perhaps due to the variation in δ_a values inside the growing environments. In microclimates, source CO_2 enrichment in ^{13}C might be different from ambient air, which mostly is influenced by soil respiration and spatial variation in δ_a (Jones, 1992; Clark and Fritz, 1997). The above authors assumed the δ_a value to remain constant in both open air and in the greenhouse. If we make the same assumption, our findings agree with their data. The validity of this assumption needs to be tested in the future. Within each growth environment, water stress reduced the Δ value by about 1.1×10^{-3} , except in E_2 , where Δ changes indicated minimal. The reduction in Δ values due to water stress in our experiments (E_0 and E_1) agreed fairly well with the results of previous authors on other plant species (Meinzer *et al.*, 1990; Ismail and Hall, 1992; Wright *et al.*, 1994). Thus, stomatal conductance and carbon isotope discrimination decreased with water stress in a similar fashion, except when plants were not coupled with the ambient air, as was the case for E_2 environment. Well-watered plants growing in an atmosphere of high relative humidity had a high stomatal conductance (Comstock and Ehleringer, 1993). The essentially constant value of Δ in E_2 environment across different soil water contents might be attributed to the imposition of a lower restriction expected for stomatal conductance and enzymatical activity of water-stressed plants compared with those of W_0 plants. Soil water depletion is expected to be less for plants in an environment of higher relative humidity. Photosynthesizing plants in enclosures are exposed to low-velocity air movements. This might contribute to a depletion of air CO_2 due to photosynthesis and also release of CO_2 from the decaying organic matters in the soil. The elevated CO_2 concentrations observed in the E_1 and E_2 environments indicates the latter process to be dominant. The δ_a values varied with the growing environment, which was due to decomposing organic matters of C_3



plant residues (Boutton, 1991). Photosynthesizing C_3 plants discriminate against $^{13}CO_2$ which increases δ_a in the canopy. However, CO_2 originated from decaying C_3 plant organic matters is not enriched in $^{13}CO_2$ (Farquhar *et al.*, 1989) and will reduce δ_a . Our measurements of lower δ_a values inside the enclosures compared with the open air indicated the contribution of CO_2 from decaying organic matters. The intermediate δ_a value in the perforated housing showed that air movement through the holes facilitated the exchange between the enriched $^{13}CO_2$ of the open air with the low enriched $^{13}CO_2$ inside the enclosure.

Harvest time had a significant effect on mean Δ values. After harvesting, HT1 plants of all environments were kept in the open air, which experienced increasingly higher air temperatures as the summer season approached, while the relative humidity was reduced. The increased Δ values in E_1 and E_2 environments at HT2 could result from a higher availability of $^{13}CO_2$ in the ambient air ($\delta_a \approx -8 \times 10^{-3}$) compared with previous condition under plastic covers ($\delta_a < -8 \times 10^{-3}$) and also the coupling of stomatal movement with open air conditions. The negative relationships and significant correlations between WUE_t and Δ in E_0 and E_1 environments of our experiment agreed not only with the theory of the isotope effect but also with the experimental works of previous authors (Wright *et al.*, 1994; Johnson *et al.*, 1995; Raeini-Sarjaz *et al.*, 1998).

Although the relationship between Δ and WUE_t in E_2 was not significant, the decoupling of plants and their surroundings from outside air (Jarvis and McNaughton, 1986; Jones, 1992) may have contributed to practically no change in WUE_t across soil water contents. Therefore, the poor mixing condition of E_2 might have reduced plant water use, resulting in an almost no variation in WUE_t and Δ . Under closed environments such as E_2 , canopy transpiration is mainly controlled by energy input (radiation) rather than stomatal conductance (Jarvis and McNaughton, 1986; Slavich *et al.*, 1998). The driving force for transpiration from

stomatal cavity to boundary layer is the gradient of saturation vapor pressure deficit (VPD). In an isothermal condition under an enclosed environment, transpiration from a canopy increases the relative humidity toward saturation point, which in turn reduces the driving force of VPD gradient towards zero. Under such environments, energy input increases air temperature which, in turn, builds up a VPD gradient and therefore enhances transpiration. The overall leaf-to-air VPD in E_2 was lower than those of the other environments, and no air movement was registered inside E_2 (Raeini-Sarjaz and Barthakur, 1997). Therefore, the absence of an air mixing mechanism combined with a low leaf-to-air VPD in E_2 resulted in a relatively small leaf boundary layer conductance and relatively low transpiration. This is expected to reduce the driving force for mass transfer. The likelihood of any increase in WUE or a reduction of Δ from low soil water content might be due to the stomatal activity. Although decoupling of plants inside the E_2 caused little change in WUE in plants across different soil water contents, the modification of plastic housing through perforations (E_1) increased the WUE and reduced Δ of water-stressed plants. Mass and heat transfer were facilitated by the presence of holes, so that the plants were coupled actively with the outside environment. Linear regression analyses showed different slopes and y-intercepts for Δ versus WUE_t relationships for different environments. This indicated that each growth environment affected the WUE_t and Δ relationship somewhat differently. The absence of a relationship between Δ and WUE_t in E_2 indicated that stomatal conductance in the closed tunnel was almost independent of soil water contents.

Since all plants received the same amount of fertilizer, N content was expected to be dependent on the rate of growth. Plants with a low growth rate contained more leaf N. The results on Δ and N were in fair agreement with those reported by Ehleringer (1990). The higher average assimilation rates for plants in E_0 than those of E_1 and E_2 supported the findings of Comstock and

Ehleringer (1993) who reported that leaf N increases photosynthetic rates. The increased assimilation rates of W_2 compared with those of W_0 plants, although not significant, indicated the effect of leaf N on photosynthesis. In general, our data showed that perforated plastic tunnels (E_1) were superior in terms of soil plant water relations, water use, WUE and plant growth rate.

In conclusion, the present results showed that carbon isotope discrimination is a valuable tool in exploring the relationships of transpiration efficiency and growing environment in C_3 plants. This type of research is expected to be increasingly important in view of the expected environmental changes and their influence on plant growth.

ACKNOWLEDGEMENTS

The authors thank Dr. D. Smith, Department of Plant Science, McGill University, for providing the photosynthesis system, and Dr. Peter Jones, School of Dietetics and Human Nutrition, McGill University, for stable isotopes facilities.

REFERENCES

1. Boutton, T. W. 1991. Stable Carbon Isotope Ratios of Natural Materials: II. Atmospheric, Terrestrial, Marine, and Freshwater Environments. In: "Carbon Isotope Techniques." (Eds.) Coleman, D. C. and Fry, B., Academic Press Inc., New York. pp. 173-185.
2. Brugnoli, E. and Farquhar, G. D. 2000. Photosynthetic Fractionation of Carbon Isotopes. In: "Photosynthesis: Physiology and Metabolism (Advances in Photosynthesis Volume 9)", (Eds.): Leegood, R. C., Sharkey, T. D. and von Caemmerer, S., Kluwer Academic Publishers, pp. 399-434.
3. Clark, I., and Fritz, P. 1997. Environmental Isotopes in Hydrogeology. CRC Press LLC: Boca Raton, F. L., U.S.A., pp.111-136.
4. Comstock, J., and Ehleringer, J. R. 1993. Stomatal Response to Humidity in Common Bean (*Phaseolus vulgaris*): Implications for Maximum Transpiration Rate, Water-use Efficiency and Productivity. *Aust. J. Plant Physiol.*, **20**: 669-691.
5. Condon, A. G., Farquhar, G. D. and Richards, R. A. 1990. Genotypic Variation in Carbon Isotope Discrimination and Transpiration Efficiency in Wheat. Leaf Gas Exchange and Whole Plant Studies. *Aust. J. Plant Physiol.*, **17**: 9-22.
6. Ebdon, J. S., Petrovic, A. M. and Dawson, T. E. 1998. Relationship between Carbon Isotope Discrimination, Water Use Efficiency, and Evapotranspiration in Kentucky Bluegrass. *Crop Sci.*, **38**: 157-162.
7. Ehdaie, B., Hall, A. E., Farquhar, G. D., Nguyen, H. T. and Waines, J. G. 1991. Water-use Efficiency and Carbon Isotope Discrimination in Wheat. *Crop Sci.*, **31**: 1282-1288.
8. Ehleringer, J. R. 1990. Correlation between Carbon Isotope Discrimination and Leaf Conductance to Water Vapor in Common Beans. *Plant Physiol.*, **93**: 1422-1425.
9. Ehleringer, J. R. and Osmond, C. B. 1989. Stable Isotopes. In: "Plant Physiological Ecology, Field Methods and Instrumentation", (Eds.) R. W. Pearcy, J. R. Ehleringer, H. A. Moodney, and P. W. Rundel. Chapman and Hall, London, pp. 280-300.
10. Ehleringer, J. R., Klassen, S., Clauton, C., Sherrill, D., Fuller-Holbrook, M., Fu, Q. and Cooper, T. A. 1991. Carbon Isotope Discrimination and Transpiration Efficiency in Common Bean. *Crop Sci.*, **31**: 1611-1615.
11. Ehleringer, J. R., Hall, A. E. and Farquhar, G. D. 1993. *Stable Isotopes and Plant Carbon-Water Relations*. Academic Press, San Diego.
12. Evans, J. R. and von Caemmerer, S. 1996. Carbon Dioxide Diffusion inside Leaves. *Plant Physiol.*, **110**: 339-346.
13. Farquhar, G. D., O'Leary, M. H. and Berry, J. A. 1982. On the Relationship between Carbon Isotope Discrimination and the Inter-cellular Carbon Dioxide Concentration in Leaves. *Aust. J. Plant Physiol.*, **9**: 121-137.
14. Farquhar, G. D. and Richards, R. A. 1984. Isotope Composition of Plant Carbon Correlates Water-use Efficiency of Wheat Genotypes. *Aust. J. Plant Physiol.*, **11**: 539-552.
15. Farquhar, G. D., Ehleringer, J. R. and Hubick, K. T. 1989. Carbon Isotope Discrimination and Photosynthesis. *Ann. Rev. Plant Physiol. Plant Biol.*, **40**: 503-537.
16. Friedli, H., Lotscher, H., Oeschger, H., Siegenthaler U. and Stauffer, B. 1986. Ice



- Core Record of the $^{13}\text{C}/^{12}\text{C}$ Ratio of Atmospheric CO_2 in the Past Two Centuries. *Nature*, **324**: 237-238.
17. Fritz, P., and Fontes, J. C. (Eds.), 1980. *Handbook of Environmental Isotope Geochemistry*, Vol. I. *The Terrestrial Environment*, Elsevier, Amsterdam.
 18. Goodman, H. S. and Francey, R. F. 1988. $^{13}\text{C}/^{12}\text{C}$ and $^{18}\text{O}/^{16}\text{O}$ in Baseline CO_2 . In: "Baseline Atmospheric Program (Australia)". (Eds.) Forgan, B. W. and Fraser, P. J., CSIRO: Canberra, Australia. pp 54-58.
 19. Hall, A. E., Mutters, R. G., Hubick, K. T. and Farquhar, G. D. 1990. Genotypic Differences in Carbon Isotope Discrimination by Cowpea under Wet and Dry Field Conditions. *Crop Sci.*, **30**: 300-305.
 20. Hubick, K. T. 1990. Effects of Nitrogen Source and Water Limitation on Growth, Transpiration Efficiency and Carbon-isotope Discrimination in Peanut Cultivars. *Aust. J. Plant Physiol.*, **17**: 413-430.
 21. Ismail, A. I. and Hall, A. E. 1992. Correlation between Water-use Efficiency and Carbon Discrimination in Diverse Cowpea Genotypes and Isogenic Lines. *Crop Sci.*, **32**: 7-12.
 22. Jarvis, P. G. and McNaughton, K. G. 1986. Stomatal Control of Transpiration: Scaling up from Leaf to Region. *Adv. Ecol. Res.*, **15**: 1-49.
 23. Jones, H. G. 1992. *Plants and Microclimate: a Quantitative Approach to Environmental Plant Physiology*. Cambridge University Press. Cambridge, UK, pp. 131-162.
 24. Jones, M. B. and Jongen, M. 1996. Sensitivity of Temperate Grassland Species to Elevated Atmospheric CO_2 and the Interaction with Temperature and Water Stress, *Agr. Food Sci. Finland*, **5**: 271-283.
 25. Johnson, R. C., Muehlbauer, F. J. and Simon, C. J. 1995. Genetic Variation in Water-use Efficiency and its Relation to Photosynthesis and Productivity in Lentil Germplasm. *Crop Sci.*, **35**: 457- 463.
 26. Meinzer, F. C., Goldstein, G. and Grantz, D. A. 1990. Carbon Isotope Discrimination in Coffee Genotypes Grown under Limited Water supply. *Plant Physiol.*, **92**: 130-135.
 27. Melander, L., and Saunders, W. H. 1979. *Reaction Rates of Isotopic Molecules*. John Wiley and Sons, New York.
 28. Nobel, P. S. 1991. *Physicochemical and Environmental Plant Physiology*, Academic Press, New York, pp. 425-453.
 29. O'Leary, M. H. 1981. Carbon Isotope Fractionation in Plants. *Phytochemistry*, **20**: 553-567.
 30. Pospisilova, J. and Èatsky, J. 1999. Development of Water Stress under Increased Atmospheric CO_2 Concentration. *Biol. Plant.*, **42**: 1-24.
 31. Raeini-Sarjaz, M. and Barthakur, N. N. 1997. Water-use Efficiency and Total Dry Matter Production of Bush Bean under Plastic Covers. *Agr. Forest Meteorol.*, **87**: 75-84.
 32. Raeini-Sarjaz, M., Barthakur, N. N., Arnold, N. P. and Jones, R. J. H. 1998. Water Stress, Water-use Efficiency, Carbon Isotope Discrimination and Leaf Gas Exchange Relationships of Bush Bean. *J. Agron. Crop Sci.*, **180**: 173-179.
 33. Rao, N. R. C. and Wright, G. C. 1994. Stability of the Relationship between Specific Leaf Area and Carbon Isotope Discrimination across Environments in Peanut. *Crop Sci.*, **34**: 98-103.
 34. Saralabai, V. C., Vivekandan, M. and Suresh Babu, R. 1997. Plant Responses to High CO_2 Concentration in the Atmosphere. *Photosynthetica*, **33**: 7-37.
 35. Slavich, P. G., Hatton, T. J. and Dawes, W. R. 1998. *The Canopy Growth and Transpiration Model of WAVES: Technical Description and Evaluation*. CSIRO Land and Water, Australia.
 36. SAS. 1994. *SAS Procedure Guide. Release 6.12 Edition*. SAS Institute Inc., Cary, NC.
 37. Yeh, H. W. and Wang, M. W. 2001. Factors Affecting the Isotopic Composition of Organic Matter. (1) Carbon Isotopic Composition of Terrestrial Plant Materials. *Proc. Nat. Sci. Counc. ROC (B)*, **25**:137-147
 38. Wright, G. C., Hubick, K. T. and Farquhar, G. D. 1988. Discrimination in Carbon Isotopes of Leaves Correlates with Water-use Efficiency of Field Grown Peanut Cultivars. *Aust. J. Plant Physiol.*, **15**: 815-825.
 39. Wright, G. C., Rao, N. R. C. and Farquhar, G. D. 1994. Water-use Efficiency and Carbon Isotope Discrimination in Peanut under Water Stress Conditions. *Crop Sci.*, **34**: 92-97.

اثر خُرداقلیم‌های گوناگون و ظرفیت آبی خاک روی کارایی مصرف آب و فرق‌گذاری ایزوتوپی کربن در لوبیا سبز

م. رائینی سرجاز و و. چالوی

چکیده

متغیرهای محیطی همچون گنجایش آبی خاک (SWC) گاهی همانند بازدارندگان رشد و حاصلخیزی گیاهان عمل می‌کنند. بنابراین، برای کنکاش در روابط میان کارایی مصرف آب (WUE)، فرق‌گذاری ایزوتوپی (Δ) و ترکیب ایزوتوپی (δ_p) کربن، بهره‌فستتری برگ (A) و محتوی ازت برگ (N) در لوبیا سبز، آزمایشی در سه خُرداقلیم هوای باز (E_0)، تونل پلاستیکی سوراخ دار (E_1) و تونل پلاستیکی بسته (E_2)، همراه با سه تیمار آبیاری خوش آبی (W_0)، آبیاری متوسط (W_1) و تنش آبی (W_2) انجام شد. غلظت CO_2 و ترکیب ایزوتوپی کربن هوا (δ_a) در محیط‌های آزمایشی متفاوت بود. مقدار δ_a در تونل‌های پلاستیکی سوراخ دار و بسته در مقایسه با هوای باز به ترتیب به اندازه $10-3 \times 0/8$ و $10-3 \times 3/8$ کاهش یافت. آب خاک به طور معنی‌داری روی کارایی مصرف آب، δ_a و Δ در دو خُرداقلیم باز و تونل سوراخ دار اثر گذاشت، ولی در خُرداقلیم تونل بسته اثر معنی‌داری نداشت. جداسازی (decoupling) گیاهان از هوای بیرون، تونل پلاستیکی بسته، ممکن است عامل ثابت ماندن متغیرهای بالا در این محیط بوده باشد. مقدار Δ در محیط‌های باز و تونل پلاستیکی سوراخ دار در مقایسه با تونل پلاستیکی بسته به ترتیب به اندازه $10-3 \times 2/2$ و $10-3 \times 1/1$ افزایش یافت. همبستگی معنی‌داری میان کارایی مصرف آب و Δ در دو محیط باز و تونل سوراخ دار (به ترتیب، $r = -0/72$ و $r = -0/75$) دیده شد، در حالی که چنین رابطه‌ای در محیط بسته یافت نشد. این مسئله می‌تواند گویای این باشد که رسانایی روزنه‌ای در این محیط تقریباً مستقل از مقدار آب خاک بوده است. مقدار ازت برگ اثر چندانی روی Δ نداشت. ازت برگ، بسته به زمان برداشت و محیط، در گیاهان زیر تنش آبی به طور معنی‌داری افزایش یافت. میانگین بهره‌فستتری برگ در محیط باز، در مقایسه با دو محیط دیگر، به طور معنی‌داری بیشتر بود.