

ACCEPTED ARTICLE

Potassium foliar application to enhance high temperature tolerance and productivity of canola under late sown conditions

Saba Iqbal^{1*}, Asmat Ullah¹, Arbab Jahangeer¹, and Naveed Akhtar¹

Agronomic Research Station Khanewal, Ayub Agricultural Research Institute (AARI), Faisalabad, Pakistan.

*Corresponding author: E-mail address: sabaiqbal87@gmail.com

Abstract

Potassium (K) mitigates the effect of high temperature on canola, especially during its later growth stages when sowing of canola is delayed. To explore the effect of K on high temperature tolerance and canola production a field experiment (2019 – 2021) was conducted. The experiment had two sowing dates (October 15 and November 1) and four K treatments, i.e. control, soil application (50 kg K ha⁻¹), soil application + 0.25% K foliar spray at pre- and post-flowering stages and soil application + 0.50% K foliar spray at pre and post-flowering stages. Application of 50 kg K ha⁻¹ as soil application along with 0.50% K-foliar sprays resulted in higher production of biochemicals (superoxide dismutase, peroxidase, catalase, and total soluble sugars) besides net photosynthetic rate and stomatal conductance along with less malondialdehyde production and relative cell injury in crops sown on November 1st than untreated plants. Moreover, it also enhanced chlorophyll fluorescence and chlorophyll (a and b) contents of late- sown crops. Plants sown on November 1st and received 50 kg K ha⁻¹ as soil application along with 0.50% K foliar spray also gave higher yield components, yield and economic returns than control. Therefore, it is suggested to supply 50 kg K ha⁻¹ at sowing and a foliar spray of 0.50% K at the pre- and post-flowering stages to canola crops sown late in the season to achieve optimal and economical yield levels.

Keywords: high temperature; reactive oxygen species; antioxidant activities; foliar spray; lipid per oxidation.

INTRODUCTION

Climate change is contributing to global warming, which has implications for the regional distribution and cultivation conditions of crops (Nesar et al., 2022). From last many years, the air temperature of earth is increasing steadily and it is expected that the rise in temperature will continue which would result in significant rise in average temperature of earth (IPCC, 2018). High

36 temperature can have a severe impact on crop productivity (Mostofa et al., 2022). Canola is
37 vulnerable to high temperature especially during its reproductive stage (Annisa et al., 2013). In
38 canola the highly heat sensitive stage is flowering stage and considerable yield reduction occurs
39 when temperature goes higher than 28-30°C at flowering stage (Chen et al., 2020). High
40 temperature can cause abnormal flower development, leading to fewer pods and seeds (Chen et al.,
41 2021). The problem is further exacerbated when canola planting is delayed due to intensive
42 cultivation practices. Farmers often struggle to plant canola on time, especially after cultivating
43 kharif crops like cotton, rice, and potatoes (Yousaf et al., 2002). Pakistan is among the countries
44 which are facing the severity of climate change. Based on records from the International Disasters
45 Database (EM-DAT), Pakistan has experienced a significant rise in the frequency and intensity of
46 extreme meteorological as well as hydrological events, including droughts, storms, floods and
47 extreme temperatures, over the past three decades. In addition to that the annual average
48 temperature of Pakistan has also increased up to 1.68°C during 1901 to 2021 whereas the global
49 rise of average annual temperature during the same period is 1.1°C; hence the increase in Pakistan's
50 average annual temperature is also higher than the global annual average temperature rise.

51 Fertilization strategies significantly reduce abiotic stress impacts by promoting plant
52 development, structure, biochemical processes, and nutrients stores, enabling plants to tolerate
53 hostile ecological circumstances (Xu et al., 2021). Within vital group of macronutrients, K assumes
54 a predominantly pivotal part in affecting growth and development of plant, whether in typical or
55 challenging environmental conditions. Its primary function in plants revolves around boosting
56 stress tolerance. Through its ability to decrease transpiration rates and enhance water absorption,
57 K contributes to increased agricultural productivity (Aslam et al., 2021). Furthermore, it assists in
58 preserving turgidity of cell and counteracting detrimental impacts of reactive oxygen species (ROS)
59 (Jan et al., 2019). Enzymatic activities and metabolic processes are enhanced by K (Zaman et al.,
60 2019), thus enhancing physiological systems and building up antioxidant defense mechanisms
61 (Hasanuzzaman et al., 2020). Importantly, K performs a pivotal function in stress mitigation by
62 accelerating the metabolism of plant proteins, which regulate numerous plant processes during
63 adverse environmental conditions. Furthermore, it supports proline synthesis, contributes to
64 osmotic regulation, and bolsters plant resilience to stress (Zamani et al., 2020).

65 Implementing proper K fertilization strategies can, therefore, perform a crucial part in boosting
66 productivity and health of plant, even in harsh conditions. Li et al. (2023) found that the application

67 of potassium dihydrogen phosphate foliar spray at flag leaf stage in wheat resulted in increase in
 68 net photosynthetic rate (Pn), chlorophyll content, antioxidant enzymes activity, dry matter
 69 accumulation, yield related traits and overall yield of crop as well as quality of produce facing high
 70 temperature. Likewise, Sarwar *et al.* (2022) reported that high temperature considerably damaged
 71 physiology of leaf as well as grain yield of wheat. Nonetheless, existing data are scarce concerning
 72 the effectiveness of K foliar spray for mitigation of high temperature effects in canola plants.
 73 Consequently, current research project was formulated with the following aims: (i) to monitor the
 74 physiological and developmental responses of canola under high temperature conditions; (ii) to
 75 appraise the influence of K foliar spray on physiology of canola which enables the plant to cope
 76 with high temperature; and (iii) to evaluate the influence of K foliar spray on canola yield.

77 MATERIAL AND METHODS

78 Experimental Site

79 Experiment was carried out at the Agronomic Research Station Khanewal, Pakistan (Fig. 1). This
 80 field trial was conducted for two growing seasons, from 2019 to 2021. The experimental soil was
 81 sandy loam with an 8.6 pH, 4 ds cm⁻¹ electrical conductivity (EC), 0.06% nitrogen (N), 6.9 ppm
 82 phosphorus (P), and 206.7 ppm K.
 83



Fig. 1: Geographic location of the experimental site.

84 Experimental Treatments and Design

85 The experiment was laid out in RCBD with split-plot arrangement with two sowing dates (main
 86 plots) and four K application treatments (sub-plots) having net plot size on 7.0 m × 2.7 m and three
 87 replications. Sowing dates were: October 15th; November 1st, and K treatments were: T₁: control
 88

89 (no application of K), T₂: 50 kg K ha⁻¹ (soil applied), T₃: T₂ + foliar spray of 0.25% K at pre- and
90 post-flowering stages, T₄: T₂ + foliar spray of 0.50% K at pre- and post-flowering stages. The
91 canola variety Faisal Canola was used in the study. Potassium nitrate (KNO₃) was used as K source,
92 and K solutions were prepared distilled water as solvent and sprayed manually using a hand
93 sprayer.

94 **Crop husbandry**

96 Canola was planted in rows 45 cm apart with a rabi drill using 4 kg ha⁻¹ seed rate. At four- leaf
97 stage, the crop was thinned to maintain 10 cm plant to plant distance. Weeds were removed
98 manually when required. Overall 85 kg N and 74 kg P ha⁻¹ were given to crop; with all P and K
99 fertilizers applied at sowing time and N fertilizer was applied in two equivalent splits at sowing
100 and flowering. Three irrigations were provided, at 30 days after sowing, flowering, and pod
101 formation. Harvesting for treatments sown on October 15th and November 1st took place on March
102 8th and March 24th, respectively, in both years of the study.

103 **Observations**

105 **Leaf Biochemical Analysis**

106 Leaf samples (from middle portion of main branch) were collected 10 days after K-foliar sprays
107 application to recorded superoxide dismutase (SOD), peroxidase (POD), catalase (CAT), total
108 soluble sugars (TSS) and malondialdehyde (MDA). The SOD (U/mg protein) activity was
109 determined using procedure described by Winterbourun et al. (1993). The (POD) (U/mg protein)
110 activity was determined by employing the procedure described by Ogawa et al. (1985). The amount
111 of CAT (U/mg protein) was determined by following the procedure described by Sinha (1972).
112 The TSS (mg/g dry weight) was determined using the procedure described by Dubois et al. (1956).
113 The MDA (U/mg) was determined by employing the procedure described by Velikova et al. (2000).

114 **Relative Cell Injury**

116 Ten days following the K-foliar spray treatment, fresh leaf samples (from the middle region of
117 the main branch) were taken to record relative cell injury. The following formula was used (Lutts
118 et al., 1999). Electrical conductivity (EC) was measured using EC meter (Model, Jenway 4510
119 Japan).

120 **Relative Cell Injury (RCI %)** = (EC of the sample - EC of the control) / (EC of the maximum
121 leakage) × 100

122

123 **Photosynthetic Parameters**

124 Ten days following the application of K-foliar sprays, the net photosynthetic rate (Pn, $\mu\text{mol m}^{-2}$
125 s^{-1}) and stomatal conductance (Gs, $\text{mol m}^{-2} \text{s}^{-1}$) were measured using infrared gas analyzer (LC
126 Analyzer with Broad Head, Part Number LC-002/B with Serial Number 32455) (Sarwar et al.,
127 2022). Chlorophyll fluorescence (Fv/Fm) was recorded to measure thylakoid membrane stability.
128 Chlorophyll was extracted from leaf samples (from middle portion of main branch) using a standard
129 procedure and the fluorescence was measured using a chlorophyll fluorimeter (OptiScience, UK
130 with Serial Number 0729501) (Murchie and Lawson, 2013). Fresh leaf samples (from middle
131 portion of main branch) were collected 10 days after K-foliar sprays to record chlorophyll a/b using
132 the following equations (Lichtenthaler and Wellburn, 1983):

133 Chlorophyll a = 12.7 (A663) - 2.69 (A647)

134 Chlorophyll b = 22.9 (A647) - 4.68 (A663)

135 **Yield and Yield Related Traits**

136 When the plants reached maturity, ten were chosen at random from each plot. Measurements
137 included plant height, number of branches and pods per plant, and length of pod. The quantity of
138 seeds per pod was calculated by counting and threshing fifteen pods that were taken from the
139 chosen plants. After that, the complete plots were gathered and threshed. A total of 1000 seeds
140 were weighed and counted in order to record the weight. The yield per plot was calculated by taking
141 the total weight of seeds in each plot and converting it to yield per hectare.

142

143 **Economic Analysis**

144 The gross income per hectare (Rs.) was calculated by multiplying canola seed yield (kg ha^{-1}) by
145 the market rate (Rs. /kg) of canola seed (Byerlee, 1988). Fixed and variable costs per hectare (Rs.
146 /ha) were calculated by combining the costs associated with standard field operations and
147 treatment-specific expenses, respectively. The total cost of production (Rs. /ha) was obtained by
148 combining the fixed and variable costs. Net income (Rs. /ha) was calculated by deducting the total
149 cost of production from the gross income. The benefit-cost ratio (BCR) for each treatment was
150 determined by dividing the net income by the total cost of production.

151

152 **Weather**

153 During 2019-20, average maximum temperature was 24.50°C, while average minimum
 154 temperature was 11.36°C and total rainfall was 232 mm from October to April. However, from
 155 February to March of 2020, when the crop was in the reproductive stage, the highest temperature
 156 range was 25.50–36.50°C. During 2020-21, average maximum temperature was 27.27°C, while
 157 average minimum temperature was 10.87°C and total rainfall was 64.20 mm from October to April.
 158 However, in February and March of that year the highest temperature range was 22.50–35.50°C.







Fig. 2: Climate conditions at the experimental location throughout the study duration.

159 **RESULTS**
 160 **Leaf Physiology**
 161 Late sowing reduced SOD, POD, CAT, and TSS contents in leaves, and increased MDA
 162 production. Foliar sprays of K enhanced these contents and reduced MDA production, especially
 163 for plants sown on November 1st (Fig. 3, 4). Late sowing resulted in high RCI, which was reduced
 164 by foliar sprays of K, especially at 0.50% concentration (Fig. 4). Delayed sowing reduced Pn and
 165 Gs, which were preserved by foliar spray of K, especially at 0.50% concentration (Fig. 4). Late
 166 sowing also reduced chlorophyll fluorescence and leaf chlorophyll contents (a and b), which were
 167 maintained by foliar sprays of K, especially at 0.50% concentration (Fig. 5). Overall, foliar sprays
 168 of K mitigated the negative effects of late sowing on canola plants.
 169
 170

Potassium mitigates effects of high temperature in canola








Fig. 3: Influence of **two sowing dates** and potassium application on biochemical of canola. Means sharing same case letter do not differ significantly at **P<0.05**.

171



Potassium mitigates effects of high temperature in canola





Fig. 4: Influence of **two sowing dates** and potassium application on physiology of canola.






Fig. 5: Influence of two sowing dates and potassium application on chlorophyll fluorescence (Fv/Fm) and chlorophyll contents of canola.

172

173 Yield and Yield Related Traits

174 Data regarding yield and related traits has been presented in table 1. During 2019-20, the canola
 175 crop sown on 15th of October resulted in 9% higher plant height as compared to canola crop sown
 176 on November 1st. Moreover regarding number of pods per plant, 52% higher were recorded in crop
 177 sown on October 15th and sprayed with 0.50% K-foliar spray than crop sown on same date but no
 178 K-foliar spray was applied. Additionally, regarding pod length and number of seed per pod, 10%
 179 and 21% higher pod length and number of seed per pod, respectively, were documented in plants
 180 sprayed with 0.50% K-foliar spray as compared with control. Regarding 1000-seed weight, 14%
 181 higher was recorded when canola was sown on October 15th than canola sown on November 1st,
 182 whereas among K treatments, the 0.25 and 0.50% K-foliar sprays resulted in 29 higher 1000-seed
 183 weight than control. Regarding yield, canola crop sown on October 15th and sprayed with 0.50%
 184 K-foliar spray resulted in 7% higher yield as compared with crop sown on same date but not
 185 sprayed with K. Whereas, crop sown on November 1st and sprayed with 0.50% K-foliar spray
 186 resulted in 43% higher yield as compared with crop sown on similar date but not sprayed with K
 187 at pre- and post-flowering stages.

188 During 2020-21, the canola crop sown on 15th of October resulted in 4% and 17% higher plant
 189 height and number of branches per plant, respectively, than canola crop sown on November 1st.
 190 Moreover regarding number of pods per plant, 39% higher were recorded in crop sown on
 191 November 1st and sprayed with 0.50% K-foliar spray than crop sown on same date but no K foliar
 192 spray was applied. Additionally, regarding pod length, 16% higher pod length was documented in
 193 plants sown on October 15th than plants sown on November 1st whereas among K treatments, crop
 194 sprayed with 0.50% K-foliar sprays gave 7% higher pod length as compared with control.

195 Furthermore, regarding number of seed per pod, 48% higher number of seed per pod was recorded
 196 in plants sown on October 15th as compared to plants sown on November 1st whereas among K
 197 treatments, crop sprayed with 0.50% K-foliar sprays gave 48% higher number of seed per pod as
 198 compared with control. Regarding 1000-seed weight, 14% higher was recorded when canola was
 199 sown on October 15th than canola sown on November 1st, whereas among K treatments, the 27%
 200 higher 1000-seed weight as compared with control. In case of yield, canola crop sown on October
 201 15th and sprayed with 0.50% K-foliar spray resulted in 7%, higher yield as compared with crop
 202 sown on same date but not sprayed with K. Whereas, crop sown on November 1st and sprayed with
 203 0.50% K-foliar spray resulted in 43% higher yield as compared with crop sown on similar date but
 204 not sprayed with K at pre- and post-flowering stages.

205
 206 **Table 1:** Influence of two sowing dates and potassium application on yield and related traits of
 207 canola.

Treatments	Plant height (cm)		Number of branches per plant		Number of pods per Plant		Pod length (cm)		Number of seeds per pod		1000-seed weight (g)		Seed yield (g)	
	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21
Sowing dates (D)														
15th October (D₁)	197A	170 A	6	6 A	374 A	238 A	6.80	7.43 A	17	23 A	4.14 A	4.32 A	1170 A	2606 A
1st November (D₂)	179 B	163 B	5	5 B	319 B	171 B	6.60	6.22 B	17	18 B	3.58 B	3.76 B	723 B	2232 B
HSD	10.79	4.19	-	0.16	10.79	17.14	-	0.34	-	1.26	1.06	1.24	43.56	32.66
Treatments (T)														
Control (T₁)	182	165	5	5	254 D	175 C	6.43 B	6.54 B	15 C	13 C	3.04 C	3.22 C	849 D	2037 D
50 kg K ha⁻¹ (T₂)	193	165	5	5	334 C	198 BC	6.63 B	6.68 AB	17 B	21 B	3.90 B	4.08 B	890 C	2278 C
0.25% K+50 kg K ha⁻¹ (T₃)	190	168	5	5	379 B	209 AB	6.58 B	6.99 A	17 B	23 A	4.29 A	4.38 A	951 B	2601 B
0.50% K+50 kg K ha⁻¹ (T₄)	190	168	5	6	419 A	235 A	7.17 A	7.07 A	19 A	25 A	4.20 A	4.47 A	1094 A	2760 A
HSD	-	-	-	-	26.86	31.37	0.52	0.44	1.52	1.78	0.18	0.18	41.23	87.11
D×T														
D₁T₁	167	170	5	6	234 d	219 ab	6.56	7.17	14	15	3.35	3.53	1156 bc	2280 de
D₁T₂	188	169	6	5	339 c	237 a	6.63	7.32	18	24	4.12	4.30	1097 c	2413 cd
D₁T₃	181	171	6	6	436 b	243 a	6.60	7.60	18	27	4.59	4.77	1186 ab	2752 b
D₁T₄	180	171	6	6	487 a	253 a	7.43	7.61	20	26	4.49	4.67	1240a	2980 a
D₂T₁	196	160	5	5	273 d	132 c	6.30	5.91	16	12	2.72	2.90	543 f	1794 f

D ₂ T ₂	197	160	5	5	328 c	160 c	6.63	6.04	16	18	3.69	3.87	684 e	2143 e
D ₂ T ₃	198	165	4	5	322 c	174b c	6.57	6.53	16	22	3.99	4.17	715 e	2450 c
D ₂ T ₄	199	165	5	5	351 c	216 ab	6.90	6.37	18	20	3.92	4.10	949 d	2540 c
HSD	-	-	-	-	46.3 0	53.82	-	-	-	-	-	-	31.2 5	150.1 5


208 Means sharing same case letter do not differ significantly at $P < 0.05$

209

210 Economic Analysis

211 When the crop was sowed on October 15th and treated with 0.50% K-foliar sprays at the pre- and
 212 post-flowering stages, compared to the control, there was a 23% increase in net revenue and benefit
 213 cost ratio according to the economic analysis (Fig. 6). Similarly, when crop was sowed on
 214 November 1st and treated with 0.50% K-foliar sprays at pre- and post-flowering stages, compared
 215 to control, 40% higher net income was obtained, likewise, when crop was sowed on November 1st
 216 and treated with 0.50% K-foliar sprays at pre- and post-flowering stages, compared to control, 32%
 217 higher benefit cost ratio was obtained (Fig. 6).

218



219 **Fig. 6:** Economic analysis of two sowing dates and foliar spray treatments of canola.

220

221 DISCUSSION

222 The findings of study demonstrated that K-foliar sprays resulted in improved physiological response and yield of canola plants which are likely to face high temperature at flowering under

223 late sowing conditions. Under higher temperature exposure, the plant metabolism as well as many
224 biochemical reactions is disrupted (Hasanuzzaman et al., 2013). Potassium enhances tissue water
225 potentiality, which aids in extreme temperature tolerance, and it helps to activate numerous
226 physiological and metabolic processes like photosynthesis, respiration, and nutrition homeostasis
227 (Hasanuzzaman et al., 2018). In present study the activity of enzymatic antioxidants (SOD, POD,
228 and CAT) was significantly improved by spraying K solution of canola plants sown on 1st
229 November (Fig.3). These enzymes are protein in nature and their function is the transformation of
230 ROS into the form which is non-toxic/harmful for plants, hence play crucial role in defense
231 mechanism of plant which enables plant to mitigate the harmful effect of oxidative stress caused
232 by many abiotic stresses (Saez and Estan-Capell, 2017). These ROS cause lipid peroxidation of the
233 cell membrane (Fang et al., 2022).

234 A higher level of enzymatic antioxidant activity in plants supplied with K-foliar sprays might be
235 due to the role of K in enhancing the production/activity of enzymatic antioxidants in stressed
236 plants (Fang et al., 2022). One way K enhances their production/activity is by the regulation of
237 enzymes activity present in antioxidant defense pathways (Anschütz et al., 2014). Potassium also
238 aids in maintaining cellular ion balance and osmotic potential, which are crucial for enzyme activity
239 and overall plant health under stress. By regulating osmotic balance, K helps prevent cellular
240 dehydration and maintains turgor pressure, which is essential for optimal enzyme function
241 (Siddiqui et al., 2021). In addition, fewer contents of MDA and less occurrence of RCI (Fig. 4) in
242 plants sprayed with K-foliar sprays were also recorded in this study. Malondialdehyde contents in
243 plant leaves are used to estimate the lipid peroxidation (Houmani et al., 2022). As discussed above
244 the K application increase the efficacy and contents of enzymatic antioxidants defense systems in
245 plants (Siddiqui et al., 2021) thus reduce the cellular damage in plants. Hence less content of MDA
246 and less prevalence of RCI might be due to this protective role of K against ROS.

247 Additionally higher level of TSS in plants (Fig. 3) sprayed with K was also recorded.
248 Accumulation of TSS is mainly observed in mild stress which retard the growth of plant but any
249 how the process of photosynthesis is not much inhibited (Keunen et al., 2013). Mainly the
250 production of TSS is taken place during osmotic stress and act as osmo-protectants to stabilize the
251 activity of cell membrane as well as to main the turgor of cell (Dien et al., 2019). Many researchers
252 reported that K enhances the accumulation and assembly of TSS in plants facing abiotic stress
253 through several mechanisms such as osmotic adjustment (Tränkner et al., 2018:), the stimulation

254 of numerous enzymes, like RuBisCO (Weng et al., 2007) and governing the functioning of
255 photosynthesis machinery (Tavakol et al., 2022). Translocation of photo-assimilates from source
256 to sink is also governed by K (Cakmak, 2005).

257 Net photosynthesis and stomatal conductance (Fig. 4) was also enhanced by the foliar application
258 K on canola. Stomatal opening and closing is governed by K ions (Shabala, 2003). Potassium ions
259 are actively pumped into guard cells causing them to swell and leading to stomatal opening (Lu et
260 al., 2019). This process is crucial for regulating loss of water by transpiration and facilitating the
261 uptake of CO₂ for photosynthesis (Lu et al., 2019). Moreover K also influence the photosynthesis
262 as it performs crucial part in the activation of enzymes vital for photosynthesis, such as Rubisco,
263 which catalyzes the fixation of CO₂ during the Calvin cycle (Rawat et al., 2022). Canola plants
264 sown on November 1st and subjected to 0.50% K-foliar sprays exhibited an augmentation in
265 chlorophyll levels compared to untreated plants (Fig. 5) sown on the same date. Potassium plays a
266 crucial part in the production of precursor molecules for chlorophyll pigments and enhances the
267 capacity to transform radiation energy into chemical energy within the chloroplasts (Zhao et al.,
268 2001).

269 The supplementation of external K caused a noticeable increase in several important parameters
270 of canola crop sown on November 1st. These parameters included the pod count per plant, seeds
271 per pod, and the weight of 1000 seeds (Table 1). This enhancement in pod and seed quantities can
272 likely be attributed to the ability of K to improve growth by its participation in numerous biological
273 functions, such as enzyme activation, assimilate and nitrate transport, water relations, stomatal
274 regulation and photosynthesis (Oosterhuis et al., 2014). The improvement in 1000-seed weight
275 (Table 1) due to the application of K, might be the function of K in photosynthesis and
276 transportation of sugars from source to sink (Oosterhuis et al., 2014). The increase in overall yield
277 (Table 1) observed in canola sown on November 1st, attributed to K application, can be attributed
278 to a combination of factors. These factors include an improvement in membrane stability, enhanced
279 efficiency of photosynthesis, increased accumulation of carbohydrates, and effective translocation
280 of these substances to developing seeds (Sarwar et al., 2022). These combined effects contribute
281 to an overall increase in seed yield and benefit cost ratio.

282 CONCLUSIONS

284 This study has demonstrated that the foliar application of K triggers antioxidant activity within
285 plants, as evidenced by increased production of SOD, POD, CAT, and TSS. This timely activation

286 of the antioxidant defense system, brought about by K-foliar spraying, led to a significant decrease
 287 in MDA levels, reduced cell injury, and the maintenance of optimal photosynthesis rates, stomatal
 288 conductance, and chlorophyll a/b contents. Furthermore, when a combination of 50 kg K ha⁻¹
 289 applied at sowing and foliar spraying with 0.50% K at both pre- and post-flowering stages was
 290 employed, higher crop yield and improved benefit-cost ratios were obtained. Based on these
 291 findings, it is recommended to apply 50 kg K ha⁻¹ at sowing and a foliar spray of 0.50% K at the
 292 pre and post-flowering stages to canola crop sown late in the season to achieve optimal and
 293 economical yield levels.

294

295 ACKNOWLEDGEMENTS

296 This research was not funded by any public, commercial, or non-profit organizations.

297

298 REFERENCES

299 Annisa, Chen, S., Turner, N. C., Cowling, W. A. 2013. Genetic Variation for Heat Tolerance
 300 During the Reproductive Phase in *Brassica rapa*. *J Agron Crop Sci*, **199**: 424–435.

301 Anshütz, U., Becker, D., Shabala, S. 2014. Going Beyond Nutrition: Regulation of Potassium
 302 Homeostasis as a Common Denominator of Plant Adaptive Responses to the Environment. *J Plant*
 303 *Physiol*, **171(9)**: 670–687.

304 Aslam, B., Khurshid, M., Arshad, M. I., Muzammil, S., Rasool, M., Yasmeen, N., Shah, T.,
 305 Chaudhry, T. H., Rasool, M. H., Shahid, A., Xueshan, X., Baloch, Z. 2021. Antibiotic Resistance:
 306 One Health One World Outlook. *Front Cellular Infection Microbiol*, **11**: 771510.

307 Byerlee, D. 1988. From Agronomic Data to Farmer's Recommendation, An Economics Training
 308 Manual, CIMMYT, Mexico, PP. 31–33.

309 Chen, J., Pan, A., He, S., Pin, S., Xiaoling, Y., Shengwei, Z., Zhi, L. 2020. Different microRNA
 310 families involved in regulating high temperature stress response during cotton (*Gossypium*
 311 *hirsutum* L) anther development. *Int J Mol Sci*, **21**: 1280.

312 Chen, S., Stefanova, K., Siddique, K. H. M., Cowling, W. A. 2021. Transient Daily Heat Stress
 313 during the Early Reproductive Phase Disrupts Pod and Seed Development in *Brassica napus* L.
 314 *Food Energy Security*, **10**: 262.

315 Dien, D. C., Mochizuki, T., Yamakawa, T. 2019. Effect of various drought stresses and
 316 subsequent recovery on proline, total soluble sugar and starch metabolisms in Rice (*Oryza sativa*
 317 L.) varieties. *Plant Prod Sci*, **22(4)**: 530–545.

- 318 Dubois, M., Gilles, K. A., Hamilton, J. K., Rebers, P. A., Smith, F. 1956. Colorimetric Method
 319 for Determination of Sugars and Related Substances. *Analy Chem*, **28(3)**: 350–356
- 320 Fang, S., Yang, H., Wei, G., Shen, T., Wan, Z., Wang, M., Wang, Z., Wu, Z. 2022. Potassium
 321 application enhances drought tolerance in sesame by mitigating oxidative damage and regulating
 322 osmotic adjustment. *Front Plant Sci*, **13(2022)**.
- 323 Hasanuzzaman, M., Nahar, K., Alam, M. M., Roychowdhury, R., Fujita, M. 2013. Physiological,
 324 biochemical, and molecular mechanisms of heat stress tolerance in plants. *Int J Mol Sci*, **14**: 9643–
 325 9684.
- 326 Hasanuzzaman, M., Bhuyan, M. H. M. B., Nahar, K., Hossain, M.S., Mahmud, J. A., Hossen, M.
 327 S., Masud, A. A. C., Moumita, Fujita, M. 2018. Potassium: A Vital Regulator of Plant Responses
 328 and Tolerance to Abiotic Stresses. *Agronomy*, **8(3)**: 31.
- 329 Hasanuzzaman, M., Borhannuddin Bhuyan, M. H. M., Zulfiqar, F., Raza, A., Mohsin, S. M.,
 330 Mahmud, J. A., Fujita, M., Fotopoulos, V. 2020. Reactive Oxygen Species and Antioxidant
 331 Defense in Plants under Abiotic Stress: Revisiting the Crucial Role of a Universal Defense
 332 Regulator. *Antioxidants*, **9(8)**: 681.
- 333 Houmani, H., Debez, A., Freitas-Silva, Ld, Abdelly, C., Palma, J. M., Corpas, F. J. 2022
 334 Potassium (K⁺) starvation-induced oxidative stress triggers a general boost of antioxidant and
 335 nadph-generating systems in the halophyte *Cakile maritima*. *Antioxidants*, **11(3)**: 401.
- 336 Intergovernmental Panel on Climate Change. 2018. The Special Report on Global Warming of
 337 1.5°C.
- 338 Jan, A. U., Hadi, F., Akbar, F., Yu, Q. 2019. Role of Potassium, Zinc, and Gibberellic Acid in
 339 Increasing Drought Stress Tolerance in Sunflower (*Helianthus annuus* L.). *Pak J Bot*, **51**: 809–
 340 815.
- 341 Kaushal, M., Wani, S. P. 2016. Plant-Growth-Promoting Rhizobacteria: Drought Stress
 342 Alleviators to Ameliorate Crop Production in Drylands. *Annals Microbiol*, **66**: 35–42.
- 343 Kawakami, A., Sato, Y. Yoshida, M. 2008. Genetic engineering of rice capable of synthesizing
 344 fructans and enhancing chilling tolerance. *J Exp Bot*, **59**: 793–802.
- 345 Keunen, E., Peshev, D., VanGronsveld, J., Van den Ende, W., Cuypers, A. 2013. Plant sugars are
 346 crucial players in the oxidative challenge during abiotic stress: extending the traditional concept.
 347 *Environ Exp Bot*, **36(7)**: 1242–1255.

- 348 Kirkegaard, J. A., Lilley, J. M., Bril, R. D., Sprague, S. J., Fettell, N. A., Pengilley, G. C. 2017.
 349 Re-Evaluating Sowing Time of Spring Canola (*Brassica napus* L.) in South-Eastern Australia—
 350 How Early is Too Early? *Crop Pasture Sci*, **67**: 381–396.
- 351 Li, J., Li, Z., Li, X., Tang, X., Liu, H., Li, J., Song, Y. 2023. Effects of spraying KH₂PO₄ on flag
 352 leaf physiological characteristics and grain yield and quality under heat stress during the filling
 353 period in winter wheat. *Plants*, **12**: 1801.
- 354 Lu, Z., Xie, K., Pan, Y., Ren, T., Lu, J., Wang, M., Shen, Q., Guo, S. 2019. Potassium mediates
 355 coordination of leaf photosynthesis and hydraulic conductance by modifications of leaf anatomy.
 356 *Plant Cell Environ*, **42(7)**: 2231–2244.
- 357 Lutts, S., Kinet, J. M., Bouharmont, J. 1999. Salinity Tolerance in Plants: What are the keys? In:
 358 “*Responses of Plants to Abiotic Stresses: Cellular and Molecular Biology*”, (Eds.): Kinet, J. M.,
 359 Lutts, S. CRC Press, PP. 21-46.
- 360 Murchie, E. H., Lawson, T. 2013. Chlorophyll Fluorescence Analysis: a Guide to Good Practice
 361 and Understanding some New Applications. *J Exp Bot*, **64(13)**: 3983–3998.
- 362 Mostafa, S., Tengyue, H., Khalid, J., Yu, Q., Muhammad, N. 2022. Role of Eco-Friendly Products
 363 in the Revival of Developing Countries’ Economies and Achieving a Sustainable Green Economy.
 364 *Front Environ Sci*, **10**: 2553–2562.
- 365 Nesar, A., Atal, H., Meenakshi, A., Rakesh, A., Ranjeet, B., Rahmat, G., Hritik, S. 2022. Terminal
 366 Heat Stress and its Mitigation Options Through Agronomic Interventions in Wheat Crop: a Review.
 367 *Inter J Environ Climate Change*, **12(11)**: 131–139.
- 368 Ogawa, K., Hirata, T., Ohashi, H., Ono, T. 1985. A New Method for the Determination of
 369 Peroxidase Activity by the Oxidation of Guaiacol. *Analy Biochem*, **149(2)**: 198–206.
- 370 Oosterhuis, D. M., Loka, D. A., Kawakami, E. M., Pettigrew, W. T. 2014. The physiology of
 371 potassium in crop production. In: “*Advances in Agronomy*”, (Ed.): Sparks, D. Academic Press,
 372 126 PP. 203-233.
- 373 Rawat, J., Pandey, N., Saxena, J. 2022. Role of potassium in plant photosynthesis, transport,
 374 growth and yield. In: “*Role of Potassium in Abiotic Stress*”, (Eds.): Iqbal, N., Umar, S. Springer,
 375 Singapore.
- 376 Sáez, G. T., Están-Capell, N. 2014. Antioxidant enzymes. In: “*Encyclopedia of Cancer*”, (Ed.):
 377 Schwab, M. Springer, Berlin, Heidelberg.

- 378 Sarwar, M, Saleem, M. F., Maqsood, H., Ullah, N., Khan, A., Waqas, M., Sattar, N., Tasneem,
 379 M., Xu, X., Zhangli, H. Shuang, Y. 2022. Strengthening Leaf Physiological Functioning and Grain
 380 Yield Formation in Heat-Stressed Wheat Through Potassium Application. *Front Plant Sci*, **13**:1–
 381 16.
- 382 Shabala, S. 2003. Regulation of potassium transport in leaves: from molecular to tissue level. *Ann*
 383 *Bot*, **92**(5): 627–34.
- 384 Shabala, S. 2017. Signalling by Potassium: Another Second Messenger to Add to the list? *J Exp*
 385 *Bot*, **68**: 4003–4007.
- 386 Siddiqui, M. H., Mukherjee, S., Kumar, R., Alansi, S., Shah, A. A., Kalaji, H. M., Javed, T., Raza,
 387 A. 2021. Potassium and Melatonin-Mediated Regulation of Fructose-1,6-Bisphosphatase (FBPase)
 388 and Sedoheptulose-1,7-Bisphosphatase (SBPase) Activity Improve Photosynthetic Efficiency,
 389 Carbon Assimilation and Modulate Glyoxalase System Accompanying Tolerance to Cadmium
 390 Stress in Tomato Seedlings. *Plant Physiol Biochem*, **171**: 49–65.
- 391 Sinha, A. K. 1972. Colorimetric Assay of Catalase. *Analy Biochem*, **47**(2): 389–394
- 392 Tavakol, E., Jákli, B., Cakmak, I., Dittert, K., Senbayram, M. 2022. Optimization of potassium
 393 supply under osmotic stress mitigates oxidative damage in barley. *Plants*, **11**(1): 55.
- 394 Tränkner, M., Tavakol, E., Jákli, B. 2018. Functioning of potassium and magnesium in
 395 photosynthesis, photosynthate translocation and photoprotection. *Physiol Plant*, **163**: 414–431.
- 396 Valluru, R. Van den Ende, W. 2008. Plant fructans in stress environments: emerging concepts
 397 and future prospects. *J Exp Bot*, **59**: 2905–2916.
- 398 Velikova, V., Yordanov, I., Edreva, A. 2000. Oxidative Stress and Some Antioxidant Systems in
 399 Acid Rain-Treated Bean Plants: Protective Role of Exogenous Polyamines. *Plant Sci*, **151**(1): 59–
 400 66
- 401 Weng, X. Y., Zheng, C. J., Xu, H. X., Sun, J. Y. 2007. Characteristics of photosynthesis and
 402 functions of the water-water cycle in rice (*Oryza sativa*) leaves in response to potassium deficiency.
 403 *Physiol Plant*, **131**: 614–621.
- 404 Winterbourun, C. C., Hawkins, R. E., Brian, M., Carrell, R. W., Lunec, J. 1993. The Estimation
 405 of Red Cell Superoxide Dismutase Activity. *J Clinical Pathol*, **46**(6): 563–566.
- 406 Xu, X., Sharma, P., Shu, S. 2021. Global Greenhouse Gas Emissions from Animal-Based Foods
 407 are Twice Those of Plant-Based Foods. *Nat Food*, **2**: 724–732.

- 408 Yousaf, M., Amir A., Jahangir, M., Naseeb, T. 2002. Effect of Different Sowing Dates on the
409 Growth and Yield of Canola (Sarson) Varieties. *Asian J Plant Sci*, **1**: 634–635.
- 410 Zaman, B. U., Ali, A., Arshadullah, M. 2019. Growth Response of Sunflower to Potassium
411 Sulphate Application In Saline-Sodic Soil. *Asian J Res Agric For*, **1**: 1–7.
- 412 Zamani, S., Naderi, M. R., Soleymani, A. 2020. Sunflower (*Helianthus annuus* L.) Biochemical
413 Properties and Seed Components Affected by Potassium Fertilization under Drought Conditions.
414 *Ecotox Environ Saf*, **190**: 11–17.
- 415 Zhao, D., Oosterhuis, D. M., Bednarz, C. W. 2001. Influence of Potassium Deficiency on
416 Photosynthesis, Chlorophyll Content, and Chloroplast Ultrastructure of cotton plants.
417 *Photosynthetica*, **39**: 103–109.
- 418