

Improving Growth and Yield of Wheat under Drought Stress via Application of SiO₂ Nanoparticles

F. Behboudi¹, Z. Tahmasebi Sarvestani¹, M. Zaman Kassae^{2*}, S. A. M. Modares Sanavi¹, and A. Sorooshzadeh¹

ABSTRACT

Silicon (Si) and its derivatives have beneficial effects on a wide variety of plant species, especially under both biotic and abiotic stresses. Yet, their effects on wheat (*Triticum aestivum* L.) plants under drought stress are not well known. Therefore, in order to evaluate the effects of SiO₂ NanoParticles (NPs) under drought stress, wheat seeds were separately sown in pots. Then, the SiO₂ NPs were added to them through soil and foliar application at three stages of plant growth. Results indicated that drought stress significantly decreased majority of the studied traits compared to the normal irrigation. Soil application of NPs, under drought stress, significantly increased leaf greenness (SPAD) and Relative Water Content (RWC) by 12.54 and 84.04%, respectively, compared to the control (NPs= 0 ppm). Moreover, under drought stress, wheat yield also increased by 25.35 and 17.81%, respectively, by foliar and soil application of NPs. Under the same irrigation regimes, soil application of NPs significantly increased plant height and biomass compared to the foliar application of NPs. Finally, our results highlight that usage of the SiO₂ NPs, especially at rates of 30 and 60 ppm, can mitigate adverse effects of drought stress in wheat plants.

Keywords: Biomass, Foliar application, Grain protein, Leaf area, Relative water content, SPAD.

INTRODUCTION

Nanotechnology employs NPs having at least one dimension between 1 and 100 nm (Auffan *et al.*, 2009). Nanomaterials hold great promise regarding their application in agriculture in terms of plant protection and nutrition due to their size-dependent qualities, high surface-to-volume ratio and unique optical properties (Jatav and Nirmal, 2013). SiO₂ NPs is one of the most popular nanomaterial that has been used in this field (Le *et al.*, 2014).

The Si is the second most abundant element in the earth's crust (Jones and Handereck, 1976). Although this element is not considered an essential nutrient for most

terrestrial plants, it plays an important role in enhancing the quality, quantity, and protection of some plants such as rice and wheat (Epstein, 2009). Also, it can be beneficial in mitigating biotic and abiotic stresses such as insect pest attack, diseases, salinity, drought, wounding, and high temperature (Liang *et al.*, 2007; Van Bockhaven *et al.*, 2013). For instance, silica improves photosynthesis parameters of some plants under drought stress (Ma, 2009; Zhang *et al.*, 2013). It has also been related to affect the antioxidant enzyme activity. Kamangar and Haddad (2016) stated that Si partially offset the negative effects of drought stress by increasing the tolerance of grapevine (*Vitis vinifera* L.) by rising the soluble protein content and antioxidant enzyme activities.

¹ Department of Agronomy, Faculty of Agriculture, Tarbiat Modares University, Tehran, Islamic Republic of Iran.

² Department of Chemistry, Collage of Sciences, Tarbiat Modares University, Tehran, Islamic Republic of Iran.

*Corresponding author; e-mail: kassaeem@modares.ac.ir



On the other hand, drought stress, as a major abiotic stress, strongly limits growth, development, and yield of plants especially in arid and semiarid regions (Mahajan and Tuteja 2005; Eneji *et al.*, 2008). This stress can damage plant cell membranes, and cell wall architecture, besides inhibiting photosynthesis and cell division (Taiz and Zeiger, 2006). Monakhova and Chernyadev (2002) stated that drought stress significantly decreased photochemical activities and inhibited activities of enzymes of Calvin cycle. There is little information about SiO₂ NPs effect on growth and yield of wheat under drought stress. Therefore, the purpose of this research was studying the effects of SiO₂ NPs on the wheat under drought stress.

MATERIALS AND METHODS

Growth Condition

A pot test was carried out in a factorial experiment in randomized complete block design with three replications. The experimental factors included SiO₂ NPs concentrations (0, 30, 60 and 90 ppm), application methods (foliar and root application), and irrigation regimes (normal irrigation and withholding irrigation for 15 days after pollination). The experiment was conducted during the growing season of 2014-2015 at the College of Agriculture, Tarbiat Modares University (35°43' N; 51°8' E; 1215 m sea level), Tehran, Iran.

Soil Characteristics

Results of the studied soil analysis are

presented in Table 1.

Plant Materials

The seeds of wheat (*Triticum aestivum* cv. pishtaz) were purchased from the Seed and Plant Improvement Institute, Karaj, Iran.

NPS

Specific surface area of SiO₂ NPs was >80 m² g⁻¹ and purity was >99%. The size of the NPs was determined through Field Emission-Scanning Electron Microscope (Figure 1, FE-SEM). Average primary particle size was about 40±9.5 nm.

SiO₂ NPS Suspension Preparation

The SiO₂ NPs were suspended directly in distilled water and dispersed by ultrasonic vibration (100W, 40 KHz) for 30 minutes. Small magnetic bars were placed in the suspensions for stirring before use to avoid aggregation of the particles (Adhikari *et al.*, 2013). Different doses of the SiO₂ NPs suspensions (0, 30, 60, and 90 ppm) were prepared for the pot experiment.

Treatments

Ten seeds of wheat were surface sterilized and sown in the plastic pot (27 cm in height and 26 cm in diameter) containing 10 kg of soil. Fertilizers were applied to the pots according to the soil analysis. Urea fertilizer was added at rate of 1.03 g N pot⁻¹ in two equal portions; the first during the seedling

Table 1. Physical and chemical properties of soil used in pot experiment.

Sandy loam ^a			pH	EC	OM ^b	TN ^c	P	K	Fe
Clay	Silt	Sand							
%			Ds m ⁻¹		%		ppm		
10.55	17.25	72.2	7.7	0.4	0.11	0.11	69.46	616.08	7.76

^a, ^b, ^c: Denotes the soil texture, Organic Matter and Total Nitrogen, respectively.




Figure 1. Field Emission-Scanning Electron Microscope (FE-SEM) image of SiO₂ NPs.

stage and the second at stem elongation stage.

Seedlings were thinned out to allow four plants per pot for data recording. Four concentrations of the SiO₂ NPs (0, 30, 60, and 90 ppm) were applied three times at stage of tillering, stem elongation, and heading in the root and through foliar application. For normal irrigation, soil was frequently irrigated and, in stressed plants, water was withheld for 15 days after pollination. During rainy days, a mobile rain shelter was used in the drought stress treatments to prevent infiltration of the rain.

Measurements

At the end of one-week stress period, three flag leaves of the four plants in each pot were labeled and some traits were determined *i.e.* SPAD, Leaf Area (LA), content of proline, SuperOxide Dismutase (SOD) activity, and Relative Water Content (RWC). Also, plant height, biomass, yield, and yield components were recorded at the harvest time.

RWC

The RWC was calculated using the method devised by Mata and Lamattina (2001) using the following equation:

$$RWC (\%) = \frac{(FW - DW)}{(TW - DW)} \times 100 \quad (1)$$

Where, *FW* is Fresh Weight, *DW* is Dry Weight and *TW* is Turgid Weight of leaf samples.

Leaf Area and SPAD

The Leaf Area (LA) was estimated using portable area meter model Li-3000A LI-COR. Also, the SPAD was measured by chlorophyll meter (SPAD-502, Minolta, Japan).

Proline and SOD Activity

On the 7th day after drought stress, three flag leaves of the four plants in each pot were harvested and frozen in liquid nitrogen immediately for the analysis of SOD activity and proline. Both traits were measured using a spectrophotometer (Specord 200, Analytical Jena, Germany). Proline was determined following Bates *et al.* (1973). The SOD activity was assayed following the method of Giannopolitis and Ries (1977) by monitoring the inhibition of photochemical reduction of Nitro Blue Tetrazolium (NBT) at 560 nm.

Biomass and Yields

The biomass, yield, and yield components were determined and analyzed when the



grains were mature. The wheat plants were harvested and oven dried at 80°C for 48 hours and then weighed (Gubbins *et al.*, 2011).

Seed Quality

Contents of phosphorous, potassium, and protein were determined in the dry seeds after harvesting, using Near Infrared Reflectance (NIR).

Statistical Analysis

Analysis of variance was evaluated by SAS (Version 9.1; SAS Institute Inc., Cary, NC, USA). The significance of differences among treatment means were compared by the LSD test ($P < 0.05$).

RESULTS

Analysis of Variance

Analysis of variance showed that NPs concentration, application methods, and irrigation regimes significantly affected all measured traits. The three-way interaction

among NPs concentration, application methods, and irrigation regimes was significant for LA, SPAD, RWC, 1000-grain weight, number of grains per spike, yield, protein, potassium, phosphorus, plant height, and biomass (Tables 2 and 3). There was a significant two-way interaction between NPs concentration and irrigation regimes on SOD activity. Furthermore, there was a significant two-way interaction between NPs concentration and application methods as well as NPs concentration and irrigation regimes interaction on content of proline.

LA, SPAD, and RWC

Usage of NPs, especially 90 ppm, increased SPAD and RWC compared to the control (NPs= 0 ppm). Application of 30 ppm NPs in plants under non-stressed conditions and no application of NPs in plants under drought stress led to the highest LA (27.00 cm²) and the lowest LA (9.19 cm²), respectively (Table 4). There was no significant difference between application methods of NPs on LA in the same irrigation regimes. A visible decline in SPAD (44.75) was obtained in the treatment without NPs in plants under drought stress. In contrast, the highest SPAD (55.65) was achieved with

Table 2. Analysis of variance for the effects of SiO₂ NPs application and irrigation regimes on some agronomic traits of wheat.

SOV	DF	Mean square					
		LA	SPAD	RWC	SOD	Proline	1000-Grain weight
Rep	2	1.58	1.34	35.34	178.38	65.32	4.26
NPs (A)	3	13.69 *	42.94 **	2141.11 **	58.50 **	921.70 **	7.59 ^{ns}
Application methods (B)	1	4.08 ^{ns}	81.53 **	62.49 ^{ns}	4.106 ^{ns}	228.50 **	37.47 **
Irrigation regimes (C)	1	2821.33 **	278.59 **	23818.65 **	252.54 **	22533.76 **	5333.87 **
A×B	3	5.47 ^{ns}	50.09 **	1262.17 **	2.57 ^{ns}	71.52 **	40.55 **
A×C	3	5.16 ^{ns}	39.00 **	906.17 **	54.87 **	589.09 **	14.07 *
B×C	1	5.33 ^{ns}	560.88 **	376.82 **	16.40 ^{ns}	3.56 ^{ns}	235.01 **
A×B×C	3	11.83 *	100.43 **	963.73 **	4.30 ^{ns}	7.69 ^{ns}	51.78 **
Error	30	3.91	3.66	4.28	7.03	15.21	4.38
CV (%)	-	11.17	3.82	4.86	11.04	6.94	8.36

*, **, and ns: Significant at 0.05, 0.01 probability level and not significant, respectively.

Table 3. Analysis of variance for the effects of SiO₂ NPs application and irrigation regimes on some agronomic traits of wheat.

SOV	DF	Mean square						
		No grain per spike	Yield	Protein	Potassium	Phosphorus	Plant height	Biomass
Rep	2	8.44	44.97	3.32	0.01810	0.00915	9.30	3.15
NPs (A)	3	202.13 *	14.34 **	0.81 **	0.25515 **	0.00726 **	9.62 ^{ns}	149.71 **
Application methods (B)	1	97.07 ^{ns}	56.55 **	1.55 **	0.38880 **	0.00046 **	362.23 **	332.01 **
Irrigation regimes (C)	1	0.71 ^{ns}	3827.04 **	12.98 **	0.36750 **	0.05266 **	46.41 **	465.50 **
A×B	3	411.84 **	12.86 **	20.98 **	0.17365 **	0.00931 **	95.16 **	244.72 **
A×C	3	818.00 **	13.30 **	2.93 **	0.02615 **	0.00351 **	44.35 **	450.14 **
B×C	1	782.95 **	62.83 **	13.94 **	0.06750 **	0.00226 **	1.54 ^{ns}	8.56 ^{ns}
A×B×C	3	222.50 *	8.39 **	4.53 **	0.03135 **	0.00511 **	49.90 **	38.44 *
Error	30	59.68	0.61	0.13	0.00006	0.00001	4.10	11.41
CV (%)	-	17.45	5.16	4.62	0.83	0.99	2.99	5.66

*, **, and ns: Significant at 0.05, 0.01 probability level and not significant, respectively.

application of 60 ppm NPs in plants under non-stress conditions. Moreover, application of 90 ppm NPs to plants under non-stressed conditions and in the treatment without NPs in plants under drought stress caused the highest RWC (88.27%) and the lowest RWC (21.94%), respectively.

On the other hand, drought stress significantly decreased mean of LA, SPAD and RWC compared to the normal irrigation, while the use of NPs in plants under drought stress significantly increased mean of LA, SPAD and RWC of leaves compared to the control. Also, in plants under drought stress, root application of NPs had better effect on SPAD and RWC of leaves compared to the foliar application of NPs.

Proline and SOD Activity

Content of proline was affected by NPs concentration and application methods (Figure 2-a), as well as NPs concentration and irrigation regimes (Figure 2-b). The highest content of proline was obtained in the treatment without soil application of SiO₂ NPs (Figure 2-a). In contrast, the lowest content of proline was observed at foliar application of 30 ppm NPs. Also, content of proline was significantly

increased in plants under drought stress compared to the plants under non-stressed conditions (Figure 2-b).

On the other hand, the highest SOD activity was obtained by using 30 and 60 ppm NPs in plants under drought stress. In plants under non-stressed conditions, usage of NPs had no significant effect on SOD activity compared to the control (Figure 3). Drought stress significantly increased SOD activity compared to the normal irrigation.

Grain Characteristics

The results demonstrated that usage of NPs significantly decreased potassium and phosphorus compared to the control. The highest grain protein (10.21%) was found when 30 ppm NPs was used as soil application in plants (Table 5). In contrast, the lowest grain protein (5.90%) was observed in the treatment without NPs soil application. Foliar application of NPs with 0 ppm showed the lowest potassium (0.76%) and phosphorus (0.29%) values in the grains. Also, the highest potassium (1.17%) and phosphorus (0.40%) were obtained in plants under drought stress without application of NPs. Drought stress significantly increased grain protein,

**Table 4.** Effects of SiO₂ NPs application and irrigation regimes on some agronomic traits of wheat.^a

NPs (ppm)	Application methods	Irrigation regimes	LA (cm ²)	SPAD	RWC (%)	1000-Grain weight (g)	No. grains per spike	Yield (g bush ⁻¹)
0	A		16.66 ^c	48.89 ^b	31.49 ^c	24.15 ^b	38.46 ^b	13.90 ^d
30			18.75 ^a	48.94 ^b	57.12 ^b	25.00 ^{ab}	44.92 ^a	16.49 ^a
60			18.50 ^{ab}	49.50 ^b	58.18 ^{ab}	24.92 ^{ab}	48.14 ^a	14.89 ^c
90			16.91 ^{bc}	52.85 ^a	59.17 ^a	26.09 ^a	45.48 ^a	15.55 ^b
	Foliar		18.00 ^a	48.74 ^b	50.35 ^b	24.16 ^b	45.67 ^a	14.12 ^b
			17.41 ^a	51.35 ^a	52.63 ^a	25.92 ^a	42.83 ^a	16.29 ^a
	Root		25.37 ^a	52.46 ^a	73.76 ^a	35.58 ^a	44.37 ^a	24.13 ^a
			10.04 ^b	47.64 ^b	29.21 ^b	14.50 ^b	44.13 ^a	8.28 ^b
0	Foliar	Normal	18.50 ^{ab}	46.20 ^e	44.55 ^e	21.48 ^d	39.74 ^{cd}	11.70 ^e
	Root		18.33 ^{ab}	46.42 ^e	43.94 ^e	23.50 ^{cd}	34.25 ^d	13.98 ^d
30	Foliar		18.50 ^{ab}	48.18 ^{de}	49.67 ^d	25.36 ^{bc}	46.86 ^{bc}	14.10 ^d
	Root		19.16 ^a	49.69 ^{cd}	50.88 ^{cd}	24.81 ^{bc}	40.10 ^{cd}	17.11 ^a
60	Foliar	Drought	15.50 ^c	54.17 ^a	52.92 ^c	27.01 ^{ab}	56.17 ^a	15.67 ^c
	Root		17.83 ^{abc}	52.81 ^{ab}	63.36 ^b	24.65 ^{cd}	42.67 ^{bcd}	16.09 ^{bc}
90	Foliar		17.33 ^{abc}	51.54 ^{bc}	65.43 ^{ab}	25.16 ^{bc}	44.10 ^{bc}	16.70 ^{ab}
	Root		16.50 ^c	51.36 ^{bc}	66.68 ^a	28.36 ^a	50.10 ^{ab}	16.29 ^{abc}
0		Normal	25.66 ^{ab}	50.06 ^b	41.04 ^c	34.01 ^b	39.29 ^{cd}	22.34 ^d
		Drought	9.16 ^d	44.75 ^d	21.94 ^e	13.73 ^c	30.69 ^d	7.04 ^g
30		Normal	27.00 ^a	50.35 ^b	83.97 ^b	34.50 ^b	39.46 ^{cd}	26.05 ^a
		Drought	10.00 ^{cd}	47.44 ^c	30.07 ^d	13.81 ^c	39.48 ^{cd}	7.45 ^g
60		Normal	24.16 ^b	55.65 ^a	81.77 ^b	36.28 ^{ab}	51.50 ^{ab}	23.29 ^c
		Drought	9.20 ^d	48.31 ^{bc}	32.38 ^d	14.63 ^c	46.24 ^{bc}	8.48 ^f
90		Normal	24.66 ^{ab}	54.26 ^a	88.27 ^a	37.55 ^a	56.98 ^a	24.85 ^b
		Drought	11.83 ^c	49.57 ^{bc}	32.46 ^d	15.83 ^c	50.36 ^{ab}	10.14 ^e
	Foliar	Normal	25.33 ^a	54.57 ^a	69.82 ^b	34.25 ^b	49.83 ^a	24.19 ^a
		Drought	9.41 ^b	42.92 ^d	27.55 ^d	11.40 ^d	41.51 ^{bc}	6.05 ^c
	Root	Normal	25.41 ^a	52.36 ^b	77.71 ^a	36.91 ^a	46.74 ^{ab}	24.08 ^a
		Drought	10.66 ^b	50.34 ^c	30.87 ^c	17.60 ^c	38.91 ^c	10.51 ^b

^a For a given main effect or two-way interaction, means within each column followed by the same letter are not significantly different according to *LSD* test ($P < 0.05$).




Figure 2. Effect of interaction between treatments on proline content. Means with different letters in each column are significantly different according to *LSD* test ($P < 0.05$).




Figure 3. Effect of interaction between NPs concentration and irrigation regimes on SOD activity. Means having different letters in each column are significantly different according to LSD test ($P < 0.05$).

potassium, and phosphorus compared to normal irrigation. In the same irrigation regime, protein and potassium were significantly more in foliar application of NPs than soil application of NPs.

Yield and Yield Components

The highest 1,000-grain weight (37.55 g) was obtained with application of 90 ppm NPs to plants under non-stressed conditions (Table 4). In contrast, the lowest 1,000-grain weight (13.73 g) was observed in the treatment without NPs in plants under drought stress. Foliar application of 60 ppm NPs and soil application of 0 ppm NPs led to the highest (56.17) and the lowest (34.25) number of grains per spike, respectively. Usage of 0 ppm NPs in plants under drought stress led to the lowest grain yield (7.04 g bush⁻¹). In contrast, the maximum grain yield (26.05 g bush⁻¹) was achieved by application of 30 ppm NPs in plants under non-stressed conditions. In the same irrigation regimes, mean of 1,000-grain weight and grain yield with soil application of NPs were significantly more than foliar application of NPs. In both application methods, drought stress significantly decreased yield and yield components compared to the normal irrigation. Moreover, usage of NPs

significantly increased yield and yield components compared to the control.

Plant Height and Biomass

The highest plant height was observed with soil application of 90 ppm NPs in plants under non-stressed conditions (Table 5). Root application of NPs had more effect on plants height and biomass. The lowest plant height (60.18 cm) and biomass (49.41 g bush⁻¹) were observed in the treatment without foliar application of NPs. In contrast, the highest biomass (70.22 g bush⁻¹) was achieved with soil application of 60 ppm NPs. Also, drought stress significantly decreased mean of biomass and plant height compared to the normal irrigation. In both irrigation regimes, usage of NPs, especially 90 ppm, significantly increased plant height and biomass compared to the control.

DISCUSSION

Drought stress decreased SPAD, LA, and RWC of leaves in plants under drought stress.

There was a close relationship between the SPAD and total chlorophyll concentration for wheat under drought stress (Ommen *et al.*, 1999). Chlorophyll concentration

**Table 5.** Effects of SiO₂ NPs application and irrigation regimes on some agronomic traits of wheat.^a

NPs (ppm)	Application methods	Irrigation regimes	Protein (%)	Potassium (%)	Phosphorus (%)	Plant height (cm)	Biomass (g bush ⁻¹)
0			7.76 ^b	1.14 ^a	0.37 ^a	66.48 ^b	55.73 ^c
30			8.29 ^a	0.87 ^b	0.35 ^{ab}	67.87 ^{ab}	59.03 ^b
60			7.75 ^b	0.83 ^b	0.32 ^b	67.55 ^{ab}	64.31 ^a
90			8.06 ^{ab}	0.84 ^b	0.32 ^b	68.64 ^a	59.60 ^b
	Foliar		8.14 ^a	1.01 ^a	0.34 ^a	64.89 ^b	57.04 ^b
	Root		7.78 ^b	0.83 ^b	0.34 ^a	70.38 ^a	62.30 ^a
		Normal	7.44 ^b	0.83 ^b	0.31 ^b	68.62 ^a	62.78 ^a
		Drought	8.48 ^a	1.01 ^a	0.37 ^a	66.65 ^b	56.55 ^b
0	Foliar		7.65 ^d	1.40 ^a	0.41 ^a	60.18 ^e	49.41 ^f
	Root		5.90 ^e	0.83 ^c	0.32 ^{bc}	63.84 ^d	56.24 ^e
30	Foliar		7.35 ^d	0.87 ^{bc}	0.32 ^{bc}	68.39 ^c	58.40 ^{cde}
	Root		10.21 ^a	0.86 ^{bc}	0.35 ^b	69.10 ^{bc}	57.39 ^{de}
60	Foliar		8.13 ^c	0.90 ^b	0.35 ^b	67.36 ^c	62.95 ^b
	Root		7.37 ^d	0.86 ^{bc}	0.34 ^b	71.27 ^{ab}	70.22 ^a
90	Foliar		8.94 ^b	0.76 ^d	0.29 ^c	68.18 ^c	60.67 ^{bcd}
	Root		8.17 ^c	0.87 ^{bc}	0.33 ^{bc}	72.78 ^a	62.06 ^{bc}
0		Normal	6.53 ^e	0.90 ^c	0.35 ^b	68.28 ^b	50.33 ^e
		Drought	7.58 ^d	1.17 ^a	0.40 ^a	63.77 ^d	53.40 ^{de}
30		Normal	7.63 ^{cd}	0.76 ^d	0.29 ^c	68.30 ^b	72.93 ^a
		Drought	8.99 ^a	0.98 ^b	0.35 ^b	65.88 ^{cd}	55.68 ^d
60		Normal	7.87 ^{cd}	0.71 ^d	0.29 ^c	68.98 ^{ab}	65.80 ^b
		Drought	8.54 ^b	1.11 ^a	0.35 ^b	67.08 ^{bc}	55.99 ^d
90		Normal	8.04 ^c	0.76 ^d	0.29 ^c	71.34 ^a	62.06 ^{bc}
		Drought	8.55 ^b	0.98 ^b	0.39 ^{ab}	67.46 ^{bc}	61.14 ^c
	Foliar	Normal	7.80 ^b	0.88 ^b	0.32 ^b	69.58 ^a	59.73 ^b
		Drought	7.77 ^b	1.13 ^a	0.37 ^a	63.73 ^c	54.34 ^c
	Root	Normal	7.08 ^c	0.78 ^c	0.30 ^b	71.19 ^a	65.83 ^a
		Drought	9.20 ^a	0.88 ^b	0.38 ^a	66.05 ^b	58.76 ^b

^a For a given main effect or two-way interaction, means within each column followed by the same letter are not significantly different according to *LSD* test ($P < 0.05$).

decreases under drought stress by chlorophyllase, peroxidase enzymes and phenolic components production (Abaaszadeh *et al.*, 2007). Also, decrease of RWC of leaves in plants under drought stress suggests less relative water absorption or water maintenance in wheat plants, when faced with drought. Moreover, reducing water use efficiency and RWC in plants under drought stress decreased turgor pressure and plant size. Thus, it may be a

reason for decline in LA of wheat plants under drought stress. Similar results were observed by Farooq *et al.* (2009), Zhao *et al.* (2010), and Mamnouie *et al.* (2010).

On the other hand, LA, SPAD, and RWC increased as the result of SiO₂ NPs application in both irrigation regimes, especially in plants under drought stress. The SiO₂ NPs may alleviate the water stress effect on photosynthetic pigments by enhancing endogenous levels of cytokinins,

which stimulate chlorophyll synthesis and improve chloroplast ultrastructure (Liang, 1998). Also, Si is deposited beneath cuticle of leaves, forming a Si-cuticle double layer, consequently, transpiration through cuticle may be decreased by Si deposition (Ma *et al.*, 2001). Therefore, it is suggested that a silica-cuticle double layer formed on leaf epidermal tissue is responsible for higher RWC of leaves. In agreement with our results, Gong *et al.* (2003) found that soil application of Na₂SiO₃ increased LA, dry mass, RWC, and leaf thickness of wheat plants under water stress. Also, Silica NPs improved water use efficiency, RWC, and chlorophyll content in maize crop (Yuvakkumar *et al.*, 2011; Suriyaprabha *et al.*, 2012).

Amino acid proline has been described as an osmo-protectant and is accumulated along with several abiotic stresses, such as drought stress (Moradshahi *et al.*, 2004), as seen in the present study (Figure 3). Proline accumulation may be due to the increase of proline synthesis or reduction of proline degradation in response to drought stress. It is responsible for the hydration of biopolymers serving as readily utilizable energy and nitrogen source compounds during periods of inhibited growth (Kala and Godara, 2011). Our results are supported by Afshari-Behbahanzadeh *et al.* (2016) and Sayed *et al.* (2012). Moreover, the proline content increased in wheat plant leaves under drought stress when SiO₂ NPs was applied as compared to the control. The Si enhances resistance to various abiotic stresses such as salt, nutrient imbalance, drought, high temperature, freezing by osmoregulation (Ma and Yamaji, 2006). The obtained results are supported by suggestions of Gunes *et al.* (2008) and Crusciol *et al.* (2009), who found silicon increased proline content in drought-stressed plants tissue.

Furthermore, exposure of plants to different biotic or abiotic stresses lead to deregulation or disruption of electric transport chain and, consequently, give rise to the generation of Reactive Oxygen

Species (ROS), which are considered as strong oxidizing and potentially harmful agents for the cells (Kumar *et al.*, 2011). Thus, plants protect cell systems from the cytotoxic effects of drought-accumulated active oxygen species using anti-oxidative enzymes such as SOD, glutathione peroxidase and catalase (Verhagen *et al.*, 2004). The SOD detoxifies superoxide anion free radicals (O²⁻) by forming H₂O₂, and then, the H₂O₂ can be eliminated by catalase and peroxidase (Hasheminasab *et al.*, 2012). These results are in agreement with prior reports that revealed high activities of antioxidant enzymes improved drought tolerance of olive cultivars (Ben Ahmed *et al.*, 2009) and canola (Abedi and Pakniyat, 2010). Also, a higher SOD activity via SiO₂ NPs, in the plants under drought stress, seems to indicate the effectiveness of this compound as an antioxidant system inductor of plants that protect plants from oxidative damage in drought-stressed plants. Silicon partially offsets the negative impact of drought on plants by increasing the activities of SOD, glutathione reductase (Gong *et al.*, 2005), and catalase (Zarafshar *et al.*, 2015). On the other hand, protein accumulation changes the response to drought stress. Drought stress reduces starch deposition in wheat grain, resulting in an increase in grain protein content (Gooding *et al.*, 2003). Some researchers stated that drought stress increased total protein content in rice (Fofana *et al.*, 2010). The role of SiO₂ NPs, especially that of soil application, in increasing ionic content in some of our concentrations, in both irrigation regimes, may be due to their effects on stabilizing cellular membranes through increasing antioxidant substances. This saves cell membranes from oxidative stress and improves plant cell permeability. Also, Si plays an important role in balancing the uptake, transport, and distribution of minerals in drought-stressed plants through water uptake and development of root growth (Hattori *et al.*, 2003; Ahmed *et al.*, 2008). In agreement with our results, Chen *et al.* (2011) stated that silicon improved



water use efficiency, photosynthesis and mineral nutrient absorption in rice plants, under drought stress. In another study, Si application increased water uptake in the plant under drought stress, thereby stimulated nutrient uptake, especially phosphorus nutrient (Sonobe *et al.*, 2011).

It is well known that drought stress affects plant growth and production by a multitude of molecular, biochemical, and physiological changes (Boutraa, 2010). For instance, closure of stomata and decrease in CO₂ concentration inhibited dry matter production. This was due to limitation of photosynthesis (Reddy *et al.*, 2004) which decreased the yield and its components in plants under drought stress. Moreover, drought stress caused excessive accumulation of intermediate compounds such as reactive oxygen species (Yazdanpanah *et al.*, 2011) which caused oxidative damage to DNA, lipids and proteins and, consequently, decreased plant growth and yield. Reduction in biomass, yield, and grain weight of wheat under drought stress was reported by some researchers (Anjum *et al.*, 2011; Abdoli and Saeidi, 2012).

Moreover, the results showed that using some concentration of SiO₂ NPs increased biomass, plant height, yield, and yield components in both irrigation regimes. Generally, positive effect of Si application in plants is not too obvious under optimum condition, but it is most evident when plant is under suboptimal condition (Henriet *et al.*, 2006). The role of SiO₂ NPs in alleviating the harmful effect of water stress on the growth, yield, and its components may be due to a change in transpiration, improvement in photosynthesis rate and plant water status, changes in ultra-structure of leaf organelles, activation of plant defense systems, maintenance of adequate supply of essential nutrients, and restriction in toxic ions uptake (Sacala, 2009; Parveen and Ashraf, 2010). These findings are in line with Sharifi Rad *et al.* (2014) and Shallan *et al.* (2016).

CONCLUSIONS

Results of this study showed that drought stress affected the growth and yield of wheat plants. Use of SiO₂ NPs, especially soil application of 30 and 60 ppm, decreased the adverse effects of drought stress. In normal irrigation, there was no significant difference between application methods of NPs. It can be concluded that SiO₂ NPs may produce various metabolites that cause reduction in transpiration, improve photosynthesis rate, affect stomatal conductance, and increase chlorophyll content and photochemical efficiency of leaf. Therefore, the results suggest that application of nano-scale nutrients, SiO₂ NPs, can be helpful to wheat plants either through soil or foliar application in normal irrigation, and soil applications in drought stress. However, further study is required to elucidate how SiO₂ NPs initiates these effects.

REFERENCES

1. Abaaszadeh, P., Sharifi, A., Lebaschi, H. and Moghadasi, F. 2007. Effect of Drought Stress on Proline, Soluble Sugars, Chlorophyll and RWC Level in *Melissa oggicinalis*. *Iran. J. Med. Plants Res.*, **23(4)**: 504-513.
2. Abdoli, M. and Saeidi, M. 2012. Using Different Indices for Selection of Resistant Wheat Cultivars to Postanthesis Water Deficit in the West of Iran. *Ann. Bio. Res.*, **3**: 1322-1333.
3. Abedi, T. and Pakniyat, H. 2010. Antioxidant Enzyme Changes in Response to Drought Stress in Ten Cultivars of Oilseed Rape (*Brassica napus* L.). *Czech J. Genet. Plant Breed.*, **46(1)**: 27-34.
4. Adhikari, T., Kundu, S. and Rao, S. A. 2013. Impact of SiO₂ and Mo Nano Particles on Seed Germination of Rice (*Oryza Sativa* L.). *Int. J. Agri. Food Sci. Technol.*, **4(8)**: 809-816.
5. Ahmed, A. H., Harb, E. M., Higazy, M. A. and Morgan, S. 2008. Effect of Silicon and Boron Foliar Applications on Wheat

- Plants Grown under Saline Soil Conditions. *Int. J. Agri. Res.*, **3**: 1–26.
6. Anjum, S. A., Xie, X. Y., Wang, L., Saleem, M. F., Man, C. and Lei, W. 2011. Morphological, Physiological and Biochemical Responses of Plants to Drought Stress. *Afr. J. Agri. Res.*, **6(9)**: 2026–2032.
 7. Afshari-Behbahanizadeh, S., Akbari, G. A., Shahbazi, M., Alahdadi, I., Farahani, L., Tabatabaee, S. A. and Ganji, M. 2016. Qualitative and Physical Properties of Barley Grains under Terminal Drought Stress Conditions. *J. Agr. Sci. Tech.*, **18**: 1303–1317.
 8. Auffan, M., Rose, J., Bottero, J. Y., Lowry, G. V., Jolivet, J. P. and Wiesner, M. R. 2009. Towards a Definition of Inorganic Nanoparticles from an Environmental, Health and Safety Perspective. *Nature Nanotechnol.*, **4**: 634–64.
 9. Bates, L. S., Waldran, R. P. and Teare, I. D. 1973. Rapid Determination of Free Proline for Water Stress Studies. *Plant Soil*, **39**: 205–208.
 10. Ben Ahmed, C., Ben Rouina, B., SensoyM, S., Boukhris, M. and Ben Abdallah, F. 2009. Changes in Gas Exchange, Proline Accumulation and Antioxidative Enzyme Activities in Three Olive Cultivars under Contrasting Water Availability Regimes. *Environ. Exp. Bot.*, **67**: 345–352.
 11. Boutraa, T. 2010. Improvement of Water Use Efficiency in Irrigated Agriculture: A Review. *J. Agron.*, **9**: 1–8.
 12. Chen, W., Yao, X. Q., Cai, K. Z. and Chen, J. 2011. Silicon Alleviates Drought Stress of Rice Plants by Improving Plant Water Status, Photosynthesis and Mineral Nutrient Absorption. *Biol. Trace Elem. Res.*, **142**: 67–76.
 13. Crusciol, C. A. C., Pulz, A. L., Lemos, L. B., Soratto, R. P. and Lima, G. P. P. 2009. Effects of Silicon and Drought Stress on Tuber Yield and Leaf Biochemical Characteristics in Potato. *Crop Sci.*, **49**: 949–954.
 14. Eneji, A. E., Inanaga, S., Muranaka, S., Li, J., Hattori, T., An, P. and Tsuji, W. 2008. Growth and Nutrient Use in Four Grasses under Drought Stress as Mediated by Silicon Fertilizers. *J. Plant Nutr.*, **31**: 355–365.
 15. Epstein, E. 2009. Silicon: Its Manifold Roles in Plants. *Ann. Appl. Biol.*, **155**: 155–160.
 16. Farooq, M., Wahid, A., Kobayashi, N., Fujita, D. and Basra, S. M. A. 2009. Plant Drought Stress: Effects, Mechanisms and Management. *Agron. Sustain. Dev.*, **29**: 185–212.
 17. Fofana, M., Cherif, M., Kone, B., Futakuchi, K. and Audebert, A. 2010. Effect of Water Deficit at Grain Repining Stage on Rice Grain Quality. *J. Agric. Biotech. Sustain. Dev.*, **2**: 100–107.
 18. Giannopolitis, C. N. and Ries, S. K. 1977. Superoxide Dismutases. I. Occurrence in Higher Plants. *Plant Physiol.*, **59**: 309–14.
 19. Gong, H. J., Chen, K. M., Chen, G. C., Wang, S. M. and Zhang, C. L. 2003. Effect of Silicon on Growth of Wheat under Drought. *J. Plant Nutr.*, **26(5)**: 1055–1063.
 20. Gong, H. J., Zhu, X. Y., Chen, K. M., Chen, G. C., Wang, S. M. and Zhang, C. L. 2005. Silicon Alleviates Oxidative Damage of Wheat Plants in Pots under Drought. *Plant Sci.*, **169**: 313–321.
 21. Gooding, M. J., Ellis, R. H., Shewry, P. R. and Schofield, J. D. 2003. Effects of Restricted Water Availability and Increased Temperature on the Grain Filling, Drying and Quality of Winter Wheat. *J. Cereal Sci.*, **37**: 295–309.
 22. Gubbins, E. J., Batty, L. C. and Lead, J. R. 2011. Phytotoxicity of Silver Nanoparticles to *Lemna minor* L. *Environ. Pollut.*, **159**: 1551–1559.
 23. Gunes, A., Pilbeam, D. J., Inal, A. and Coban, S. 2008. Influence of Silicon on Sunflower Cultivars under Drought Stress. I. Drowth, Antioxidant Mechanisms, and Lipid Peroxidation. *Commun. Soil Sci. Plant Anal.*, **39**: 1885–1903.
 24. Hasheminasab, H., Assad, M. T., Aliakbari, A. and Sahhafi, R. 2012. Influence of Drought Stress on Oxidative Damage and Antioxidant Defense Systems in Tolerant and Susceptible Wheat Genotypes. *J. Agric. Sci.*, **4(8)**: 20–30.
 25. Hattori, T., Inanaga, S., Tanimoto, E., Lux, A., Luxova, M. and Sugimoto, Y. 2003. Silicon Induced Changes in Viscoelastic Properties of Sorghum Root Cell Walls. *Plant Cell Physiol.*, **44**: 743–749.



26. Henriët, C., Draye, X., Oppitz, I., Swennen, R. and Delvaux, B. 2006. Effects, Distribution and Uptake of Silicon in Banana (*Musa* spp.) under Controlled Conditions. *Plant Soil*, **287**: 359–374.
27. Jatav, G. K. and Nirmal, D. E. 2013. Application of Nanotechnology in Soil-Plant System. *An. As. J. Soil Sci.*, **8(1)**: 176–184.
28. Jones, L. H. and Handereck, K. A. 1976. Silica in Soils and Plants. *Agron. J.*, **19**: 107–109.
29. Kala, S. and Godara, A. K. 2011. Effect of Moisture Stress on Leaf Total Proteins, Proline and Free Amino Acid Content in Commercial Cultivars of *Ziziphus mauritiana*. *J. Sci. Res.*, **55**: 65–69.
30. Kamangar, A. and Haddad, R. 2016. Effect of Water Stress and Sodium Silicate on Antioxidative Response in Different Grapevine (*Vitis vinifera* L.) Cultivars. *J. Agr. Sci. Tech.*, **18**: 1859–1870.
31. Kumar, R. R., Krishna, K. and Naik, G. R. 2011. Effect of Polyethylene-Glycol-Induced Water Stress on Physiological and Biochemical Responses in Pigeonpea (*Cajanus cajan* L. Millsp.). *Recent Res. Sci. Technol.*, **3(1)**: 148–152.
32. Le, V. N., Rui, Y., Gui, X., Li, X., Liu, S. and Han, Y. 2014. Uptake, Transport, Distribution and Bio-Effects of SiO₂ Nanoparticles in Bt-Transgenic Cotton. *J. Nano Biotechnol.*, **5**: 12:50.
33. Liang, Y., Sun, W., Zhu, Y. G. and Christie, P. 2007. Mechanisms of Silicon-Mediated Alleviation of Abiotic Stresses in Higher Plants: A Review. *Environ. Poll.*, **147(2)**: 422–428.
34. Liang, Y. C. 1998. Effects of Si on Leaf Ultrastructure, Chlorophyll Content and Photosynthetic Activity in Barley under Salt Stress. *Pedosphere*, **8**: 289–296.
35. Ma, J. F. 2009. Silicon Uptake and Translocation in Plants. In *The Proceedings of the International Plant Nutrition Colloquium XVI*, UC Davis, PP. 1–6.
36. Ma, J. F., Goto, S., Tamai, K. and Ichii, M. 2001. Role of Root Hairs and Lateral Roots in Silicon Uptake by Rice. *Plant Physiol.*, **127**: 1773–1780.
37. Ma, J. F. and Yamaji, N. 2006. Silicon Uptake and Accumulation in Higher Plants. *Trends Plant Sci.*, **11(8)**: 392–397.
38. Mahajan, S. and Tuteja, N. 2005. Cold, Salinity and Drought Stresses: An Overview. *Arch. Biochem. Biophys.*, **444**: 139–158.
39. Mamnouie, E., Fotouhi Ghazvini, R., Esfahani, M. and Nakhoda, B. 2010. The Effects of Water Deficit on Crop Yield and the Physiological Characteristics of Barley (*Hordeum vulgare* L.) Varieties. *J. Agr. Sci. Tech.*, **8**: 211–219.
40. Mata, C. G. and Lamattina, L. 2001. Nitric Oxide Induces Stomatal Closure and Enhances the Adaptive Plant Responses against Drought Stress. *Plant Physiol.*, **126**: 1196–1204.
41. Monakhova, O. F. and Chernyadev, I. I. 2002. Protective Role of Kartolin-4 in Wheat Plants Exposed to Soil Drought. *Appl. Biochem. Micro.*, **38**: 373–380.
42. Moradshahi, A., Eskandari, S. B. and Kholdebarin, B. 2004. Some Physiological Responses of Canola (*Brassica napus* L.) to Water Deficit Stress under Laboratory Conditions. *Iran. J. Sci. Technol. Trans A.*, **28(A1)**: 43–50.
43. Ommen, O. E., Donnelly, A., Vanhoutvin, S., Oijen, M. V. and Manderscheid, R. 1999. Chlorophyll Content of Spring Wheat Flag Leaves Grown under Elevated CO₂ Concentrations and Other Environmental Stresses within the ESPACE-Wheat Project. *Eur. J. Agron.*, **10**: 197–203.
44. Parveen, N. and Ashraf, M. 2010. Role of Silicon in Mitigating the Adverse Effects of Salt Stress on Growth and Photosynthetic Attributes of Two Maize (*Zea Mays* L.) Cultivars Grown Hydroponically. *Pak. J. Bot.*, **42(3)**: 1675–1684.
45. Reddy, A. R., Chaitanya, K. V. and Vivekanandanb, M. 2004. Drought-Induced Responses of Photosynthesis and Antioxidant Metabolism in Higher Plants. *J. Plant Physiol.*, **161**: 1189–1202.
46. Sacala, E. 2009. Role of Silicon in Plant Resistance to Water Stress. *J. Elementol.*, **14(3)**: 619–630.
47. Sayed, M. A., Schumann, H., Pillen, K., Naz, A. A. and Leon, J. 2012. AB-QTL Analysis Reveals New Alleles Associated to Proline Accumulation and Leaf Wilting

- Under Drought Stress Conditions in Barley (*Hordeum vulgare* L.). *BMC Genet.*, **13**(1): 61.
48. Shallan, M. A., Hassan, H. M. M., Namich, A. A. M. and Ibrahim, A. A. 2016. Effects of TiO₂ and SiO₂ Nanoparticles on Cotton Plant under Drought Stress. *Res. J. Pharm. Bio. Chem. Sci.*, **7**(4): 1540-1551.
 49. Sharifi Rad, J., Karimi, J., Mohsenzadeh, S., Sharifi Rad, M. and Moradgholi, J. 2014. Evaluating SiO₂ Nanoparticles Effects on Developmental Characteristic and Photosynthetic Pigment Contents of *Zeamays* L. *Bull. Environ. Pharm. Life Sci.*, **3**(6): 194-201.
 50. Sonobe, K., Hattori, T., An, P., Tsuji, W., Eneji, A. E., Kobayashi, S., Kawamura, Y., Tanaka, K. and Inanaga, S. 2011. Effect of Silicon Application on Sorghum Root Responses to Water Stress. *J. Plant Nutr.*, **34**: 71-82.
 51. Suriyaprabha, R., Karunakaran, G., Yuvakkumar, R., Prabu, P., Rajendran, V. and Kannan, N. 2012. Growth and Physiological Responses of Maize (*Zea mays* L.) to Porous Silica Nanoparticles in Soil. *J. Nanopart. Res.*, **14**: 1-14.
 52. Taiz, L. and Zeiger, E. 2006. *Plant Physiology*. 4th Edition, Sinauer Associates Inc. Publishers, Massachusetts, 690 PP.
 53. Van Bockhaven, J., De Vleeschauwer, D. and Hofte, M. 2013. Towards Establishing Broad-Spectrum Disease Resistance in Plants: Silicon Leads the Way. *J. Exp. Bot.*, **64**: 1281-1293.
 54. Verhagen, J., Put, M., Zaal, F. and Van Keulen, H. 2004. Climate Change and Drought Risks for Agriculture. *Environ. Poll.*, **39**: 49-59.
 55. Yazdanpanah, S., Baghizadeh, A. and Abbassi, F. 2011. The Interaction between Drought Stress and Salicylic and Ascorbic Acids on Some Biochemical Characteristics of *Satureja hortensis*. *Afr. J. Agric. Res.*, **6**(4): 798-807.
 56. Yuvakkumar, R., Elango, V., Rajendran, V., Kannan, N. S. and Prabu, P. 2011. Influence of Nanosilica Powder on the Growth of Maize Crop (*Zea mays* L.). *Int. J. Green Nanotechnol.*, **3**: 180-190.
 57. Zarafshar, M., Akbarinia, M., Askari, H., Hosseini, S. M., Rahaie, M. and Struve, D. 2015. Toxicity Assessment of SiO₂ Nanoparticles to Pear Seedlings. *Int. J. Nanosci. Nanotechnol.*, **11**(1): 13-22.
 58. Zhang, C., Moutinho-Pereira, J. M., Correia, C., Coutinho, J., Gonçalves, A., Guedes, A. and Gomes-Laranjo, J. 2013. Foliar Application of Silika Increases Chestnut (*Castanea* spp.) Growth and Ohotosynthesis, Simultaneously Increasing Susceptibility to Water Deficit. *Plant Soil*, **365**: 211-225.
 59. Zhao, J., Sun, H., Dai, H., Zhang, G. and Wu, F. 2010. Difference in Response to Drought Stress among Tibet Wild Barley Genotypes. *Euphytica*, **172**(3): 395-403.

بهبود رشد و عملکرد گندم تحت تنش خشکی از طریق کاربرد نانوذرات دی اکسید-سیلیسیم

ف. بهبودی، ز. طهماسبی سروستانی، م. زمان کسایی، س. ع. م. مدرس ثانوی، و ع. سروش زاده

چکیده

سیلیسیم و مشتقاتش اثرات سودمندی بر انواع گونه‌های گیاهی بخصوص در گیاهان تحت تنش خشکی دارد. اثرات آنها روی گندم تحت تنش خشکی به خوبی شناخته نشده است. بنابراین، به منظور



ارزیابی اثر نانوذرات دی‌اکسیدسیلیسیم بر گندم تحت تنش خشکی، بذور گندم به طور جداگانه در گلدان ها کشت شد. سپس نانوذرات از طریق محلول‌پاشی و مصرف خاکی (در سه مرحله) به آنها اضافه شد. نتایج نشان داد که تنش خشکی، اکثریت صفات اندازه گیری شده را به طور معنی‌داری نسبت به آبیاری نرمال کاهش داد. مصرف خاکی نانوذرات در گیاهان تحت تنش خشکی به طور معنی‌داری سبزی‌نگی برگ و محتوای نسبی آب برگ را به ترتیب ۱۲/۵۴٪ و ۸۴/۰۴٪ در مقایسه با کنترل (نانوذرات=۰) افزایش داد. در گیاهان تحت تنش خشکی، محلول‌پاشی و مصرف خاکی نانوذرات عملکرد را به ترتیب ۲۵/۳۵٪ و ۱۷/۸۱٪ در مقایسه با کنترل به طور معنی‌داری افزایش داد. در رژیم های آبیاری مشابه، کاربرد خاکی نانوذرات به طور معنی‌داری ارتفاع و بیوماس گیاه را در مقایسه با محلول‌پاشی نانوذرات افزایش داد. در نهایت، نتایج آشکار کرد که مصرف نانوذرات دی‌اکسیدسیلیسیم به‌ویژه غلظت ۶۰ و ۹۰ پی‌پی‌ام می‌تواند اثرات منفی تنش خشکی را در گندم کاهش دهد.