

Copper Effects on Growth, Lipid Peroxidation, and Total Phenolic Content of Rosemary Leaves under Salinity Stress

M. Hejazi Mehrizi^{1*}, H. Shariatmadari², A. H. Khoshgoftarmanesh², and F. Dehghani³

ABSTRACT

Plant growth is often limited by low levels of soil micronutrients such as copper (Cu), especially in calcareous salt-affected soils of arid and semiarid regions. The aim of this work was to investigate individual and combined effects of salinity and Cu on growth, leaf relative water content (LRWC), cell membrane permeability, lipid peroxidation, and total phenolic content (TPC) of rosemary (*Rosmarinus officinalis* L.) in a hydroponic experiment. A factorial arranged experiment with three salinity levels (i.e., 0, 50, and 100 mM NaCl), and three levels of copper (i.e., 0, 0.5, and 1.0 μM Cu^{2+} , as CuSO_4) was performed. The salt stress led to a significant decrease in leaf relative water content especially at 0 μM Cu^{2+} . Salt treatments (50 and 100 mM) were found to increase electrolyte leakage and malonyldialdehyde content of rosemary; however, this increase was greater at 0 μM than 1.0 μM Cu^{2+} . Regardless of Cu treatments, salinity (50 and 100 mM) induced significant increases in TPC by 13 and 29%, respectively. The application of 1.0 μM Cu at the 50 mM NaCl treatments increased total phenolic content by 19% compared to 0 μM Cu. Copper nutrition resulted in greater accumulation of phenolic compounds in plant roots and thereby decreasing lipid peroxidation under salt stress conditions.

Keywords: Cu, Medicinal plants, Rosemary, Salinity, Oxidative damage, Total phenolic content.

INTRODUCTION

Rosemary (*Rosmarinus officinalis* L., Lamiaceae) is an economically important herb known not only as a source of essential oils but also for its natural antioxidants such as polyphenolic compounds (Genena *et al.*, 2008). In the past few years, rosemary has been successfully cultivated in warm and dry climates of arid and semiarid regions (El-Rjoob *et al.*, 2008). However, as in the majority of cultivated plants growth and yield of rosemary can be affected by salinity (Baatour *et al.*, 2009) and nutritional disorders (Grattan and Greive, 1999).

Salinity reduces plant growth due to osmotic stress and ionic cytotoxicity (Saleh and Maftoon, 2008; Silva *et al.*, 2008; Dashti *et al.*, 2010) as well as oxidative damages to macromolecules and cell structure (Neto *et al.*, 2006; Eraslan *et al.*, 2007).

Copper deficiency is a significant nutritional disorder of plants grown on alkaline-saline soils of arid and semi arid climates (Whitehead, 2000). Cu is known as an essential micronutrient for the function of copper-zinc superoxide dismutases (SOD) and catalase (CAT) which are the most important ROS scavenger enzymes. Copper plays an important role in the synthesis of

¹ Department of Soil Science, College of Agriculture, Shahid Bahonar University of Kerman, Kerman, Islamic Republic of Iran.

* Corresponding author, e-mail: mhejazi@uk.ac.ir

² Department of Soil Science, College of Agriculture, Isfahan University of Technology, 84154, Isfahan, Islamic Republic of Iran.

³ Department of Soil Science, Yazd Agricultural and Natural Resources Research Center, Yazd, Islamic Republic of Iran.



phenolic compounds and its deficiency can decrease phenolics in the plants (Dicko *et al.*, 2006).

Changes in total phenolic content of medicinal plants have been reported to be influenced by salinity (Navaro *et al.*, 2006). Ksouri *et al.* (2007) reported an increase in TPC in *Cakile maritime* at 100 and 400 mM NaCl. Indeed, phenolic compounds participate in the defense against ROS which are inevitably produced when aerobic or photosynthetic metabolism is impaired by environmental stresses (Ksouri *et al.*, 2007).

In spite of its great importance, Cu nutrition of plants is much less investigated than other micronutrients such as zinc (Zn) and iron (Fe). On the other hand, there is a growing interest for the cultivation of rosemary in arid and semi-arid regions of Iran, where soils contain high levels of salts and low available Cu. The aim of this study was to investigate the effects of Cu on the biomass production, cell membrane damage, and total phenolic content of rosemary grown under saline condition.

MATERIALS AND METHODS

Plant Growth

The experiment was conducted in the greenhouses of Yazd Agricultural and Natural Resources Research Center. Cuttings of rosemary (*Rosmarinus officinalis* L.) were collected from Yazd medicinal plants field. To avoid genetic variability, cuttings were obtained from one individual plant and grown in 10 liter pots filled with washed sand and kept for 3 months in a greenhouse under controlled conditions (16/8 h light/darkness, 24/18°C day/night temperature). Plants were irrigated twice a week with distilled water and Hoagland nutrient solution (Ballesta *et al.*, 2004), alternately. After 3 months, the plants were transferred to 5 liter pots containing the nutrient solution. Pots were covered with black plastic to inhibit light exposure of the solution. The nutrient solution used in this

experiment contained (in µM): KNO₃ (3 mM), Ca(NO₃)₂ (2 mM), NH₄H₂PO₄ (0.5 mM), MgSO₄ (0.5 mM), KCl (50 µM), H₃BO₃ (25 µM), MnSO₄ (2.0 µM), ZnSO₄ (2.0 µM), CuSO₄ (0.5 µM), H₂MoO₄ (0.5 µM), Fe-ethylendiamino-di(o-hydroxyphenylacetic) acid (Fe-EDDHA) (20 µM). The pH of the nutrient solution was adjusted to 6.0 by adding HNO₃ 0.1 mM. Treatments consisted of three levels of salinity (0, 50, and 100 mM) and three levels of Cu (0, 0.5, and 1.0 µM) which were applied by dissolving NaCl and CuSO₄ in the nutrient solution, respectively.

Leaf Relative Water Content (LRWC)

Leaf relative water content was measured using the method of Yamasaki and Dillenburg (1999). Individual leaves were sampled from the mid section of each plant and then weighed to obtain fresh mass (FM). In order to determine the turgid mass (TM), whole leaves were floated in distilled water inside a closed Petri dish. During the imbibition period, leaf samples were weighed periodically to reach a constant weight, after gently wiping water off the leaf surfaces with tissue paper. At the end of the imbibition period, leaf samples were placed in a pre-heated oven at 80°C, for 48 hours and then weighed (DM). All mass measurements were made using an analytical balance with a precision of 0.0001 g. The LRWC was calculated using the below equation:

$$\text{LRWC} = (\text{FM} - \text{DM}) / (\text{TM} - \text{DM}) \times 100$$

Cell Membrane Permeability (Electrolyte Leakage)

Electrolyte leakage was used to assess cell membrane permeability according to the method described by Lutts *et al.* (1996). Five leaf samples of two plants per replicate were taken from each treatment. Leaf samples were cut into 1 cm segments and placed in individual vials containing 10 mL

of distilled water after three washes with distilled water to remove surface contamination. These samples were incubated at room temperature (25°C) on a shaker (100 rpm) for 24 hours. Electrical conductivity of samples (EC_1) was measured after incubation. The same samples were then placed in an autoclave at 120°C for 20 minutes and the second measurement (EC_2) was done after cooling the solution to room temperature. The electrolyte leakage was calculated using EC_1/EC_2 ratio and was expressed as percent.

Lipid Peroxidation

Malonyldialdehyde (MDA) - a product of lipid peroxidation- content has been considered as an indicator of oxidative damage (Neto *et al.*, 2006). One hundred milligrams of fresh leaf samples were homogenized in 2.5 mL of 0.1% (w/v) trichloroacetic acid (TCA). The homogenate was centrifuged at 3000×g for 20min. One ml of the extract was mixed with 2.25 mL of a 0.5% (w/v) thiobarbituric (TBA) acid solution containing 20% (w/v) trichloroacetic acid. The mixture was heated at 95°C for 30 minutes and then quickly cooled in an ice-bath. The mixture was centrifuged at 3,000×g for 10 minutes and the absorbance of the supernatant was measured at 532 and 600 nm. After subtracting the non-specific absorbance (600 nm), the MDA concentration was determined by using molar extinction coefficient of $155 \text{ mM}^{-1} \text{ cm}^{-1}$ and the results were expressed as nmol MDA g^{-1} fresh weight (FW).

Total Phenolic Content

250 mg of fresh leaves were powdered with liquid nitrogen in a mortar and extracted with 10 mL of 80% methanol at 37°C for 3 hours in a shaking water bath (Tawaha *et al.*, 2007). After cooling, the extract was centrifuged at 3,500 g and stored at 4°C. Total phenolic content was estimated by the Folin-Cicalteu colorimetric method

as described by Singleton and Rossi (1965). Gallic acid was used as a standard phenolic compound. One mL of the filtered extracts was mixed with 2.5 mL of 10% Folin-Cicalteu reagent. After 5 min, 2 mL of saturated sodium carbonate (75 mg L^{-1}) was added and the solution was incubated for 90 minutes at 30°C. The absorbance of the resulting blue-colored solution was measured at 765 nm. Quantitative measurements were performed based on a standard calibration curve of six points: 20, 100, 200, 300, 400 and 500 mg L^{-1} of gallic acid in 80% methanol. Total phenolic content of the plant was expressed as mg gallic acid equivalents (GAE) per gram of fresh weight.

Dry Weight Determination and Copper Analysis

At harvest, shoot and root were separated and washed twice with tap water and twice with distilled water. They were then dried in a thermostated oven at 70°C for 72 hours for dry weight determination. Copper analyses were carried out on a dry weight basis. Ground samples were dry-ashed at 550°C for 4 hours, mixed with hot 2M HCl, filtered, and then brought to a final volume of 50 ml with distilled water. Copper concentration was determined in the digest solution by atomic absorption spectrophotometry (Chapmann and Pratt, 1982).

Statistical Analysis

The experiment was set up in a factorial design with three levels of salinity and three levels of Cu in four replicates. Analysis of variance (ANOVA) was performed on all data using the SAS. Means were compared by the Lowest Significant Deviations (LSD) test at $P < 0.05$. Pearson's correlation between electrolyte leakage and MDA was also performed.



RESULTS AND DISCUSSION

The shoot and root dry weight (DW) of rosemary was significantly ($P < 0.05$) decreased by salinity stress (Figures 1 and 2). The application of 50 mM NaCl in the nutrient solution ($EC = 7 \text{ dS m}^{-1}$) caused only 7% reduction in the shoot and root dry matter production confirming that rosemary is a salt tolerant plant. Westerwelt (2003) also reported that rosemary could tolerate high levels of salinity ($4\text{--}8 \text{ dS m}^{-1}$) in the soil. The adverse effects of salinity on shoot and root growth of rosemary were more evident at the lower dose of Cu ($0 \mu\text{M}$) in comparison with 0.5 and $1.0 \mu\text{M}$ Cu levels (Figures 1 and 2). These results suggested that Cu was protective against NaCl toxicity in plants. Copper deficiency decreases the rate of photosynthesis and increases the rate of respiration which result in limited water availability for cell expansion and growth

(Olszewska *et al.*, 2008) especially under saline condition.

LRWC which directly reflects the water status of plants under stress condition (Saied *et al.*, 2005; Kaya *et al.*, 2007) was decreased in the leaves of rosemary grown under the 50 and 100 mM NaCl treatments (Figure 3). Salinity caused water deficit in rosemary leaves through an increase in soluble salts concentrations. Water deficiency decreases plant growth by limiting water availability for cell expansion and nutrient uptake (Yang *et al.*, 2009). For example, leaf Cu concentration of rosemary plants grown in the 0.5 and $1.0 \mu\text{M}$ Cu treatments was decreased in response to salt stress (Figure 4). Similar results were obtained by Izzo *et al.* (1991) and Villora (2000) who reported that salinity decreased leaf Cu concentration of maize and zucchini plants, respectively.

Increasing salinity was associated with a




Figure 1. Effect of increasing Cu and NaCl in levels on shoot dry biomass (Mean \pm S.E) of rosemary (g pot^{-1}) (Different letters above bars indicate significant difference at $P < 0.05$).




Figure 2. Effect of increasing Cu and NaCl in levels on root dry biomass (Mean \pm S.E) of rosemary (g pot^{-1}) (Different letters above bars indicate significant difference at $P < 0.05$).




Figure 3. Effect of increasing Cu and NaCl in levels on leaf relative water content (Mean \pm S.E) of rosemary (%) (Different letters above bars indicate significant difference at $P < 0.05$).




Figure 4. Effect of increasing Cu and NaCl levels on leaf Cu concentration (Mean \pm S.E) of rosemary (mg kg^{-1}) (Different letters above bars indicate significant difference at $P < 0.05$).




Figure 5. Effect of increasing Cu and NaCl levels on membrane permeability (Mean±S.E) of rosemary (%) (Different letters above bars indicate significant difference at $P < 0.05$).




Figure 6. Relationship between electrolyte leakage and malonyldialdehyde (MDA) in the leaves of Rosemary ($n = 36$).

Table 1. Effects of salinity and copper concentration on the malonyldialdehyde content of rosemary leaves.

NaCl (mM)	Cu (μ M)			Mean
	0	0.5	1	
	MDA (nmol g ⁻¹ FW)			
0	16.5±0.275 c	13.5±0.558 e	8.0±0.655 f	12.7±3.75 C
50	17.4±1.337 c	13.5±1.091 e	14.3±0.768 de	15.0±2.04 B
100	21.8±0.458 a	19.1±0.094 b	15.0±0.150 d	18.6±2.99 A
Mean	18.6±2.55 A	15.4±2.89 B	12.4±3.33 C	
Mean square				
NaCl	80.46 **			
Cu	85.28 **			
NaCl×Cu	9.44 **			

** Significant at $P < 0.01$.
Means followed by the same letter (small letters for means and capital letters for means of rows and columns) are not significantly different at 5% level of probability using LSD test.

significant ($P < 0.05$) increase in the electrolyte leakage (Figure 5) and lipid peroxidation (Table 1) of rosemary leaves. Salinity can cause oxidative damage to cell membrane and lipids (lipid peroxidation), leading to an increase in electrolyte leakage (Amor *et al.*, 2005; Kaya *et al.*, 2007; Arvin and Donnelly, 2008). Supporting this idea, the electrolyte leakage was positively correlated with the MDA content, indicating that the membrane injury induced by salt stress is a result of oxidative damage (Figure 6). The increase in electrolyte leakage and lipid peroxidation by salinity became more severe at 0 μ M Cu²⁺ than 0.5 and 1.0 μ M Cu²⁺. Since copper is directly involved in both gene expression and protein synthesis, it seems that copper deficiency inhibits the

activities of CAT, resulting in extensive oxidative damage to membrane lipids (Marschner, 1995). Leaf Cu concentration was increased in response to Cu application (Figure 4) and thus lower electrolyte leakage and MDA content at 1.0 μ M Cu²⁺ treatments could be attributed to a higher activity of CAT (Data not published). These findings are in agreement with those of Henryka (1999) who found that Cu inhibits lipid peroxidation in root nodule of yellow lupine plants.
In contrast to the reduced growth of rosemary, total phenolic content in the leaves was greatly enhanced by salinity (Table 2). The reduction in growth induced by salt stress may have resulted in a new pattern of resource partitioning providing

**Table 2.** Effects of salinity and copper concentration on the total phenolic content of rosemary leaves.

NaCl (mM)	Cu (μ M)			Mean
	0	0.5	1	
	TPC (mg gallic acid g ⁻¹ FW)			
0	14.45 \pm 0.050 f	14.69 \pm 0.120 d	17.58 \pm 0.106a	15.57 \pm 1.51 C
50	16.65 \pm 0.021 e	16.52 \pm 0.0150 d	19.75 \pm 0.064 a	17.64 \pm 1.59 B
100	20.18 \pm 0.035 c	20.12 \pm 0.097 b	20.21 \pm 0.031 a	20.17 \pm 0.07 A
Mean	17.09 \pm 2.50 B	17.11 \pm 2.40 B	19.18 \pm 1.22 A	
Mean square				
NaCl	47.7 **			
Cu	13.0 **			
NaCl \times Cu	3.1 **			

** Significant at $P < 0.01$.

Means followed by the same letter (small letters for means and capital letters for means of rows and columns) are not significantly different at 5% level of probability using LSD test.

additional carbon skeletons for phenolic biosynthesis. Regardless of copper concentration, the highest contents of TPC were found in the leaves of plants grown at the 100 mM NaCl treatment. Similarly, in pepper and *Cakile maritime* plants, Navarro *et al.* (2006) and Ksouri *et al.* (2007) found that salinity induced significant increases in TPC. Copper nutrition effect and its interaction with salinity on TPC were significant (Table 2). At the 50 mM NaCl treatment, plants exposed to 0.5 and 1.0 μ M of Cu, accumulated TPC by 12% and 37% greater than those grown in the Cu-free treatment. At the 100 mM NaCl treatment, these values changed to 12% and 15%, respectively. The increase in TPC by Cu became less distinct when TPC was also increased by salinity. This indicated that salinity was a more effective factor than Cu nutrition in the increase of TPC. Dicko *et al.* (2006) also reported that Cu increased TPC in sorghum by increasing biosynthesis of phenolic compounds.

CONCLUSIONS

The results showed that salinity induced oxidative stress, resulting in lipid peroxidation and increase in cell membrane permeability to toxic ions (e.g. Na⁺ and Cl⁻) which in turn reduced the plant growth.

Copper nutrition reduced lipid peroxidation and membrane permeability while it increased total phenol content of salt stressed plants. It seems that copper nutrition may effectively ameliorate salt-induced oxidative damage in rosemary.

ACKNOWLEDGEMENT

The authors wish to acknowledge the technical support of Yazd Agricultural and Natural Resources Research Center.

REFERENCES

1. Amor, N. B., Hamed, K. B., Debez, A., Grignon, C. and Abdelly, C. 2005. Physiological and Antioxidant Responses of the Perennial Halophyte *Crithmum maritimum* to Salinity. *Plant Sci.*, **168**: 889-899.
2. Arvin, M. J. and Donnelly, D. J. 2008. Screening Potato Cultivars and Wild Species to Abiotic Stresses Using an Electrolyte Leakage Bioassay. *J. Agric. Sci. Technol.*, **10**: 33-42.
3. Baatour, O., Kaddour, R., Aidi Wannes, W., Lachaal, M. and Marzouk, B. 2009. Salt Effects on the Growth, Mineral Nutrition, Essential Oil Yield and Composition of Marjoram (*Origanum majorana*). *Acta Physiologica Plantarum*, **32**: 45-51.

4. Ballesta, M. C., Martinez, V. and Carvajal, M. 2004. Osmotic Adjustment, Water Relations and Gas Exchange in Pepper Plants Grown Under NaCl or KCl. *Environ. Exp. Bot.* **52**: 161-174.
5. Chapman, H. D. and Pratt, P. F. 1982: *Methods of Analysis for Soils, Plants and Water*. Chapman Publishers, Riverside, CA. P.382.
6. Dashti, H., Naghavi, M. R. and Tajabadipour, A. 2010. Genetic Analysis of Salinity Tolerance in a Bread Wheat Cross. *J. Agric. Sci. Technol.*, **12**: 347-356.
7. Dicko, H. M., Gruppen, H., Traore A. S., Voragen, A. G. J. and Berkel, W. J. H. V. 2006. Phenolic Compounds and Related Enzymes as Determinants of Sorghum for Food Use. *Biotechnol. Mol. Biol. Rev.*, **1**: 21-38.
8. El-Rjoob, A. W. O., Massadeh, A. M. and Omari, M. N. 2008. Evaluation of Pb, Cu, Zn, Cd, Ni and Fe Levels in *Rosmarinus officinalis labiatae* (Rosemary) Medicinal Plant and Soils in Selected Zones in Jordan. *Environ. Monit. Assess.* **140**: 61-68.
9. Eraslan, F., Inal, A., Savasturk, O. and Gunes, A. 2007. Changes in Antioxidative System and Membrane Damage of Lettuce in Response to Salinity and Boron Toxicity. *Sci. Hortic.*, **114**: 5-10.
10. Genena, A. K., Hense, H., Junior, A. S. and de Souza, S. M. 2008. Rosemary (*Rosmarinus officinalis* L.): A Study of the Composition, Antioxidant and Antimicrobial Activities of Extracts Obtained with Supercritical Carbon Dioxide. *Science Technol. Alimentos.*, **28**:1-7.
11. Grattan, S. R. and Greive, C. M. 1999. Salinity-mineral Nutrient Relations in Horticultural Crops. *Sci. Hortic.*, **78**: 127-157.
12. Henryka, S. 1999. Antioxidant Activity of Copper in Root Nodules of Yellow Lupin Plants. *Acta. Physiol. Plantarum*, **21**: 427-431.
13. Izzo, R., Navari-Izoo, F. and Quartacci, M. F. 1991. Growth and Mineral Absorption in Maize Seedlings as Affected by Increasing NaCl Concentrations. *J. Plant Nutr.*, **14**:687-699.
14. Kaya, C., Tuna, A. L., Ashraf, M. and Altunlu, H. 2007. Improved Salt Tolerance of Melon (*cucumis melo* L.) by the Addition of Proline and Potassium Nitrate. *Environ. Exp. Bot.*, **60**: 397-403.
15. Ksouri, R., Megidiche, W., Debez, A., Falleh, H., Grignon, C. and Abdelly, C. 2007. Salinity Effects on Polyphenol Content and Antioxidant Activities in Leaves of the Halophyte *Cakile maritima*. *Plant. Physiol. Biochem.*, **45**: 244-249.
16. Lutts, S., Kinet, M. and Bouharmont, J. 1996. NaCl-induced Senescence in Leaves of Rice (*Oryza sativa* L.) Cultivar Differing in Salinity Resistance. *Ann. Bot.*, **78**: 389-398.
17. Marschner, H. 1995. *Mineral Nutrition of Higher Plants*. Academic Press, London. P.995.
18. Navarro J. M., Flores, P., Garrido, C. and Martinez, V. 2006. Changes in the Contents of Antioxidant Compounds in Pepper Fruits at Ripening Stages, as Affected by Salinity, *Food Chem.*, **96**: 66-73.
19. Neto, A. D. A., Prisco, J. T., Filho, J. E., Abreu, C. E. B. and Filho, E. G. 2006. Effect of Salt Stress on Antioxidative Enzymes and Lipid Peroxidation in Leaves and Roots of Salt-tolerant and Salt Sensitive Maize Genotypes. *Environ Exp. Bot.*, **56**: 87-94.
20. Olszewska, M., Grzegorzczak, S., Alberski, J., Baluch-Malecka, A. and Kozikowski, A. 2008. Effect of Copper Deficiency on Gas Exchange Parameters, Leaf Greenness (Spad) and Yield of Perennial (*Lolium Perenne* L.) and Orchard Grass (*Dactylis glomerata* L.), *J. Elementol.*, **13**: 597-604.
21. Saied, A. S., Keutgen, A. J. and Noga, G. 2005. The influence of NaCl Salinity on Growth, Yield and Fruit Quality of Strawberry cvs. 'Elsanta' and 'Korona'. *Sci. Hortic.*, **103**: 289-303.
22. Saleh, J. and Maftoon, M. 2008. Interactive Effects of NaCl Levels and Zinc Sources and Levels on the Growth and Mineral Composition of Rice. *J. Agric. Sci. Technol.*, **10**: 325-336.
23. Silva, E., Nogueira, R. J. M. C., Ara'ujo, F. P., Melo, N. F. and Neto, A. D. A. 2008. Physiological Responses to Salt Stress in Young Umbu plants. *Environ. Exp. Bot.*, **63**: 147-157.
24. Singleton, V. L. and Rossi, J. A. Jr. 1965. Colorimetry of Total Phenolics with Phosphomolybdic-phosphotungstic Acid Reagents. *Amer. J. Enol. Vertic.* **16**: 144-158.
25. Tawaha, K, Alali, F. Q., Gharaibeh, M., Mohammad, M. and El-Elmat, T. 2007. Antioxidant Activity and Total Phenolic



- Content of Selected Jordanian Plant Species. *Food Chem.* **104**: 1372-1378.
26. Villora, G., Moreno, D. A., Pulgar, G. and Romero, L. 2000. Yield Improvement in Zucchini under Salt Stress: Determining Micronutrient Balance. *Sci. Hortic.*, **86**:175-183.
27. Westerwelt, P. M. 2003. Greenhouse production of *Rosmarinus officinalis* L. MSc Thesis in Horticultural Science, Blacksburg, Virginia University.
28. Whitehead, D. C. 2000. *Nutrient Elements in Grassland, Soil-plant-animal Relationships*. CABI Publication, UK University Press, Cambridge. PP. 220-303.
29. Yamasaki, S. and Dillenburg, L. C. 1999. Measurements of Leaf Relative Water Content in *Araucaria angustifolia*. *R. Bras. Fisiol. Veg.*, **11**: 69-75.
30. Yang, F., Xiao, X., Zhang, S., Korpelainen, H. and Li, C. 2009. Salt Stress Responses in *Populus cathayana* Rehder. *Plant Sci.*, **176**: 669-677.

تأثیر مس بر رشد، پراکسیداسیون چربی و کل محتوای فنلی برگ رزماری در شرایط تنش شور

م. حجازی مهریزی، ح. شریعتمداری، ا. ح. خوشگفتار منش و ف. دهقانی

چکیده

کمبود عناصر ریزمغذی مانند مس موجب کاهش رشد گیاهان در خاک‌های شور و آهکی مناطق خشک و نیمه‌خشک می‌شود. هدف این پژوهش بررسی اثرات اصلی شوری و تغذیه مس بر رشد، رطوبت نسبی برگ، نفوذپذیری غشای سلولی، پراکسیداسیون چربی و ترکیبات فنلی برگ رزماری رشد یافته در کشت محلول بود. این پژوهش به صورت آزمایش فاکتوریل با سه سطح شوری (صفر، ۵۰ و ۱۰۰ میلی‌مولار کلرید سدیم) و سه سطح مس (صفر، ۰/۵ و ۱ میکرومولار) در قالب طرح کاملاً تصادفی انجام گرفت. شوری ۵۰ و ۱۰۰ میلی‌مولار موجب کاهش رطوبت نسبی برگ رزماری به خصوص در سطح صفر میکرومولار مس شد. افزایش شوری موجب افزایش معنی‌دار نفوذپذیری غشای سلولی و پراکسیداسیون چربی گردید که این افزایش در سطح صفر میکرومولار مس بیشتر از سطح ۱ میکرومولار بود. صرف‌نظر از غلظت مس، افزایش شوری به ۵۰ و ۱۰۰ میلی‌مولار به ترتیب موجب افزایش ۱۳ و ۲۹ درصدی غلظت ترکیبات فنلی برگ رزماری شد. به نظر می‌رسد که مس تأثیرات اکسیداتیو منفی را از طریق افزایش غلظت ترکیبات فنلی، حفظ نفوذپذیری غشای سلولی و کاهش پراکسیداسیون چربی بهبود بخشیده است.