

Screening Potato Cultivars and Wild Species to Abiotic Stresses Using an Electrolyte Leakage Bioassay

M. J. Arvin^{1*} and D. J. Donnelly²

ABSTRACT

Ten *Solanum tuberosum* cultivars and accessions from 11 wild *Solanum* species were evaluated for their tolerance to salt (200 mM NaCl or 100 mM Na₂SO₄), drought (35% PEG), cold (-4°C) and heat (37°C). Evaluation was based on electrolyte leakage from detached leaves of *in vitro* plantlets. Testing involved a rinsing treatment with three changes of distilled water to remove electrolytes from leaf and petiole surfaces, an exposure period of 24 hours for salt and PEG stress and four hours for temperature stress and a rehydration period of 24 hours in distilled water followed by measurement of electrical conductivity into distilled water to determine the effects of stress. Significant differences were observed between *Solanum* genotypes for all types of stresses. Among the *S. tuberosum* cultivars, stress tolerance was great to NaCl and PEG in Alpha and Bintje; to Na₂SO₄ in Agria and to heat and cold in Norland. Among the wild species, tolerance to all stresses was great in *S. demissum*, to all stresses except heat in *S. acaule* and to heat and cold in *S. commersonii*. The correlations among types of stress tolerance were significant for all stresses except for heat and drought. The level of tolerance in some wild species was significantly greater compared with the cultivars tested. There appears to be a wide genetic base available to improve the stress tolerance of cultivated potato.

Keywords: Abiotic stresses, Electrolyte leakage, Potato.

INTRODUCTION

Environmental stresses, including extreme temperatures, salinity and drought, affect plant growth and productivity in many crop species including potato. The loss to agricultural and horticultural industries as a result of exposure of plants to adverse environmental conditions is estimated in the billions of dollars annually (Senaratna, 2003).

Cold stress sensitivity is a major factor limiting potato production; freezing damage to foliage results in reduced tuber yield and limits cultivation. In the US, one sixth of the total cultivated area is subjected to cold-induced limitation (Boyer, 1982). It is not possible to adjust planting times to avoid

damage since frosts occur sporadically or throughout the growing season (Richardson and Estrada, 1971). The development of frost tolerant potato cultivars could greatly expand potato production; areas that are currently marginal due to low temperatures could be brought into production and the season extended in areas where production is limited by cold.

High temperatures are also a major limitation on potato production in many developing countries, which prevails during most of the year (Dodds, 1990). Reductions in leaf area, tuber number and tuber weight have been reported as symptoms of elevated temperatures during the growing season of potato plants (Menzel, 1985). However, potato

1. Department of Horticulture, College of Agriculture, Shahid Bahonar University, Kerman, Islamic Republic of Iran.

2. Department of Plant Science, Macdonald Campus of McGill University, 21,111 Lakeshore Road, Ste. Anne de Bellevue, Quebec, H9G 3 V9, Canada.

* Corresponding author, e-mail: mjarvin@mail.uk.ac.ir



cultivars or clones which are able to maintain relatively high yields at high temperatures have been identified in field trials (Levy, 1984; Malik *et al.*, 1992).

Salinity is also a serious problem for commercial agriculture, particularly in arid and semi-arid regions. Six of 14 billion ha of arable land available in the world are located in these areas, and out of this, about one billion ha are affected by excess salt (Christiansen, 1982). With increasing world population, the use of irrigation has increased by 300% during the last 3-4 decades (Iyengar and Reddy, 1994). The most common salts responsible for toxicity and associated with saline soils are NaCl and Na₂SO₄ (Cramer and Spuir, 1986).

Moisture stress also severely affect potatoes' growth and productivity as documented in reviews by Singh (1969). Water stress during tuber initiation has been reported to reduce tuber set (Lynch and Tai, 1989; Struik and Van Voorst, 1986). The development of drought tolerant cultivars is one of the major breeding objectives in hot tropical environments where moisture is insufficient during the growing season.

Examining the field performance of genotypes under a particular stress is the usual method for evaluation, but, the results are often inconclusive. In the case of salt, a field trial is normally associated with the spatial distribution of salt, non-uniform moisture availability and temperature fluctuations during the growing season. In the case of low or high temperatures, depending on the severity of a particular season for cold or heat evaluation, either the complete survival or death of all genotypes might result. For drought tolerance, the line source sprinkler irrigation technique, which creates a continuous and linear moisture gradient, has been used extensively to screen drought stress evaluations (Levy, 1986). This method not only depends on weather changes from season to season but also involves considerable space, time, labor, equipment and planting material resources.

Therefore, a number of evaluation methods have replaced costly and labor intensive

field trials. To predict freezing tolerance, controlled freezing is usually followed by observations on plant recovery; often a lengthy, tedious and destructive process requiring a large number of plants. Plant tissue water has also been used to screen for freezing tolerance in wheat and rye (Brule-Babel and Fowler, 1989). The water content percentage of fully acclimated plants (crowns and leaves) has been correlated with field evaluation, but the technique was not able to detect small differences between cultivars without excessive replication. Alternatively, the dielectric properties of plant tissues can be used as an indicator of cold survival. This involves measurement of the proportion of bound water on structural proteins in the plant's terminal meristem. This method is neither rapid nor simple and does not measure a particular physiological parameter associated with freezing stress. *In vitro* methods for the evaluation of potato genotypes to salt and drought were proposed as alternatives to the costly, labor-intensive field based evaluations. These have used single node cuttings (Zhang and Donnelly, 1997; Arvin, 1992), five node cuttings (Morpurgo, 1991), root tip segment or suspension culture (Naik and Widholm, 1993) or callus growth (Arvin, 1992) and measured one or more growth parameters at one or more salinity or osmoticum level *in vitro*. More recently, Khrais *et al.* (1998) determined the relative tolerance of cultivars based on a multivariate analysis of the relative means of six growth parameters of *in vitro* plantlets over a range of salinity levels. While promising, these *in vitro* assays are also labor intensive. Genetic improvement of crop plants requires the identification of appropriate stress tolerance mechanisms and particularly the development of suitable methodologies for their measurement in large breeding populations.

Electrolyte leakage tests have been widely used to assess the level of plant tolerance to various stresses. These tests determine the degree of cell membrane injury caused by stress based on electrolyte leakage from the cells. The technique is relatively simple, re-

peatable and rapid and requires inexpensive equipment, can be used on plant material from a variety of cultural systems and it is suitable for the analysis of large numbers of samples. For example, it has been used to quantify damage to cell membranes in various abiotic stress conditions such as high temperatures (Ismail and Hall, 1999; Maheshwary *et al.*, 1999; Saelim and Zwi-azekk, 2000; Rahman *et al.*, 2004; Blum, 1981), low temperatures (alfalfa; Sulc *et al.*, 1991), air pollution (Garty *et al.*, 2000), sodium chloride salt (Chen *et al.*, 1999), soil acidity (Spencer and Ksander, 1999), heavy metals (De and Mukherjee, 1996) and even in response to biotic stresses such as wheat leaf rust (Adam *et al.*, 2000) or rice sheath blight (Sriram *et al.*, 2000).

The objective of this study was to examine the relative tolerance of 10 *S. tuberosum* cultivars and accession from 11 wild *Solanum* species to abiotic stresses based on electrolyte leakage and also to examine the possible relationship between the various stress tolerances.

MATERIALS AND METHODS

Plant Materials

These experiments were conducted in the tissue culture lab of the Plant Science Department of McGill University in Canada. True potato seeds from 11 wild species (accession number in brackets) were received from the United States Department of Agriculture Research Service, Inter-Regional Potato Introduction Station, at Sturgeon Bay, WI., USA and held at fridge temperature (4°C) until required. *Solanum* species included: *S. acaule* (4n, PI 266386), *S. berthaultii* (2n, PI 473331), *S. chacoense* (2n, PI 209411), *S. commersonii* (2n, PI 472833), *S. demissum* (6n, PI 186562), *S. hjertingii* (4n, PI 251065), *S. lycopersicoides* (2n, PI 265378), *S. microdomtum* (2n, PI 473171), *S. phureja* (2n, PI 243461), *S. stenotomum* (2n, PI 195214), and *S. stoloniferum* (4n, PI 230477). Seeds were disinfested for 20 min-

utes in 10% household bleach (5.25% sodium hyperchlorite), rinsed several times with sterile distilled water, and aseptically transferred to a micropropagation medium for *in vitro* germination.

Ten potato cultivars were received as *in vitro* plantlets: Caribe, Carlton, Eramosa and Jemseg were received from Potato Gene Resources, New Brunswick, Agriculture and Agri-Food, Canada; Agria from Seed Potato Specialist, AAFRD, Crop Diversification Centre North, Edmonton and Kennebec; and Alpha, Norland, Bintje and Russet Burbank from the Plant Propagation Centre, Fredericton, NB, Canada. All were acquired in November, 2003. The *Solanum* spp. seedlings and the *S. tuberosum* plantlets were micropropagated using single node cuttings.

Culture Conditions and Medium

Cultures were incubated at 25±2°C with 16/8 h D/N at 40 µmol m⁻² s⁻¹ photon flux density (cool white fluorescent light). For micropropagation, MS (Murashige and Skoog, 1962) basal salt solution was used, supplemented with 3% sucrose and 0.7% agar. The medium was adjusted to pH 5.7 prior to autoclaving at 121°C for 20 minutes.

Sampling

All stress tolerance tests were performed using leaves with attached petioles of *in vitro* plantlets grown under non-stress conditions. A single sample consisted of 10 randomly selected leaves from two plantlets (four weeks old and approximately 10 cm long), avoiding the top and bottom leaves. The leaves were gently pinched by hand and introduced into test tubes.

Salt and Drought Tolerance Test

The method of Sullivan and Ross (1979) was used with minor modification according to the type of stress tested. Samples were



washed with three changes of distilled water to remove solutes from leaf surfaces and damaged areas. Ten leaves were placed in a Pyrex test tube (15 × 25mm) into 15 ml solution of a given concentration of PEG for drought and NaCl or Na₂SO₄ for salinity tests. The control tubes contained distilled water. Samples were incubated at 10°C for 24 hours, after which media were drained and samples washed with five changes of distilled water. Samples were incubated with 15 ml of distilled water and kept at 10°C for 24 hours in the dark. Tubes were then brought to 25°C and the electrical conductivity (EC) of the incubation medium was read using a conductivity meter (conduct meter; Radiometer, Copenhagen) after a vigorous mixing of the tube's contents by hand. Following the initial reading, samples were autoclaved for 15 minutes to kill leaf tissues, brought to 25°C and a final reading was obtained. Calculation of the percentage injury of was performed as follows:

$$\% \text{ Injury} = 1 - \left[\frac{1 - (T_1/T_2)}{1 - (C_1/C_2)} \right] \times 100$$

Where T and C refer to the EC values of stress treated and control tubes and 1 and 2 refer to the initial and final EC, respectively.

Preliminary experiments with six genotypes (three cultivars and three wild species) and four replications were conducted to select an appropriate treatment under which maximum genotypic differences could be observed for each stress (data are not presented). For PEG, concentrations of 0, 30, 35 and 40% of PEG 8000, for NaCl 0, 100, 150, 200 and 250 mM and for Na₂SO₄ 0, 50, 100, and 150 mM were tested. Maximum genotypic differences were observed with 35 % PEG, 200 mM NaCl and 100 mM Na₂SO₄. Therefore, those concentrations were used for the reported experiments.

Heat Tolerance Test

The method of Blum and Ebercon (1981) was used with minor modifications. The treatment temperature and duration test was

conducted as follows; The tubes containing leaves were placed in a glycol bath (NESLAB Instruments, INC. USA) at 33, 35, 37 and 39°C for a period of 2, 3, 4 and 5 hours after which tubes were removed and kept at 10°C for 24 hours before initial conductivity was measured at 25°C. For final reading, the tubes were autoclaved for 15 minutes at 121°C and brought to 25°C. The combination of temperature and duration treatments exposing maximum genotypic differences was 37°C for 4 hours, which was selected for the reported experiments.

Cold Tolerance Test

The method used for this test was that of Sekozawa *et al.* (2003) with modifications. Tubes containing leaves were submerged in a glycol bath at 5°C and then 0°C each for 30 minutes, cooled down at the rate of 1°C h⁻¹ and removed at -2, -3, -4 or -5°C from the bath, thawed on ice overnight in a fridge. Ice was added to each tube for initiating ice nucleation after 30 minutes at -1°C. Control tubes were kept in a fridge at 5°C without subjecting them to freeze-thaw treatment. Initial and final EC readings were performed as mentioned before. Leaves were suspended in 15 ml of distilled water and kept at 10°C for 12 hours before initial EC reading was made at 25°C. The final reading was made after autoclaving. Maximum genotypic differences were observed at -4°C, which was selected for the reported experiments.

Statistical Analysis

All experiments were conducted twice with four replicates each, using a completely randomized design. Data were combined and subjected to ANOVA using SAS (SAS, 1988) (Table 1) and the means were separated by Duncan's Multiple Range test at 5% level.

Table 1. ANOVA for percentage of ion leakage (percentage of injury) of potato cultivars and wild species in response to types of stress (VMS and EMS represent mean squares of stress treatment and error, respectively).

Source	VMS	EMS	F Value	CV
NaCl	676.09	52.94	12.77**	13.3%
Na ₂ SO ₄	541.56	28.04	19.31**	7.7%
PEG	803.82	51.69	15.55**	12.1%
Heat	386.35	51.36	7.52**	14.0%
Cold	612.10	80.89	7.57**	16.6%

** Significant at 1%.

RESULTS AND DISCUSSION

NaCl

Significant differences existed between genotypes (Table 2). Little difference was found between overall performance of cultivars and wild species (56.0 vs 53.6). Agria, Alpha, Bintje, Caribe and Norland showed above average tolerance compared with the other cultivars while the rest of the cultivars, especially Carlton and Russet Burbank, were less tolerant than the average. These results are in accordance with a number of published reports; Alpha was relatively tolerant to NaCl, both *in vitro* and in a greenhouse pot trial (Arvin, 1992). Bintje was in the most tolerant cultivar group based on *in vitro* growth measurements (Khrais *et al.*, 1998). Russet Burbank was not tolerant to NaCl in a greenhouse pot trial (Bilski *et al.*, 1988a).

S. acaule and *S. demissum* had significantly greater tolerance to NaCl, compared with the other wild *Solanum* spp. and cultivars tested. Both were highly salt tolerant *in vitro* and in a greenhouse pot experiment (Arvin, 1992). *S. lycopersicoides*, *S. phureja* and *S. stoloniferum* were comparatively salt sensitive while the rest of accessions were moderately salt tolerant.

Na₂SO₄

Agria was more tolerant to Na₂SO₄, compared with the other cultivars. More varia-

tion was found among the wild species than among the cultivars as judged by the calculated range (26 vs 55.3) (Table 2). *S. acaule* was the most tolerant of the species tested followed by *S. commersonii*, *S. demissum*, *S. stenotomum* and *S. stoloniferum*.

When genotype responses to NaCl and Na₂SO₄ were compared, it is clear that the effect of Na₂SO₄ is more severe (68.3 vs 54.7) (Table 2). The severity of this effect is likely to be due to the sulfate ion as the two salts are iso-osmotic with equal amounts of Na. The correlation between genotype tolerance to NaCl and Na₂SO₄ was formally significant ($r = 0.44^{**}$) (Table 4). This indicated that some genotypes responded differently to the two salts (Table 3). Cv. Carlton showed greater tolerance to Na₂SO₄, Bintje, Alpha, Eramosa and Norland showed relative susceptibility to Na₂SO₄ and the rest of the cultivars responded similarly to the two salts. Furthermore, *S. commersonii*, *S. lycopersicoides*, *S. stoloniferum*, *S. chacoense* and *S. phureja*, were relatively more tolerant to the sulfate salt, while *S. berthaultii* and *S. hjertingii* were relatively more tolerant to NaCl; *S. demissum*, *S. acaule* and *S. stenotomum*, on the other hand, responded similarly to both salts. Similar variation in response to different salts was reported for four potato cultivars by Bilski *et al.* (1988a, b). They found that cv. Russet Burbank and Norchip were more severely affected by Na₂SO₄ compared with NaCl whereas Red Pontiac and Norgold Russet responded similarly to NaCl and Na₂SO₄.



Drought (PEG)

Significant differences were observed between genotypes (Table 2). Alpha, Bintje, *S. acaule*, *S. demissum* and *S. stenotomum* were significantly more drought tolerant compared with the other genotypes. High levels of drought tolerance were also reported for *S. acaule* and *S. demissum* *in vitro* and in greenhouse pot trials (Arvin, 1992).

There were highly significant correlations between PEG and NaCl tolerance ($r=0.74^{**}$) (Table 4). It is likely therefore that genotype response to NaCl is via an osmotic effect. *S. commersonii* was the only exception; it was relatively salt tolerant but very drought sensitive (Table 3).

genotype response to heat. Compared with the other cultivars, Norland and Bintje showed more tolerance to heat. Among the wild species, *S. chacoense*, *S. commersonii*, *S. demissum*, and *S. hjertingii* showed more tolerance. No correlation was found between heat and drought stress tolerance. This indicated that different mechanism(s) are involved for heat and drought stresses. This supports the work of Bilski *et al.* (1988b) who reported that no relationship existed between the drought and salt tolerance of various accessions of wild potato species. A similar finding was also reported for wheat cultivars but not for sorghum cultivars (Blum and Ebercon, 1981).

Cold

Heat

Significant differences were found in

There were few differences among cultivar responses to cold (Tables 2 and 3). Cultivar Norland was the most tolerant and

Table 2. Means of percentage of ion leakage (percentage of injury) of potato cultivars and wild species as affected by types of stress.

Genotype	NaCl	Na ₂ SO ₄	PEG	Heat	Cold
Agria	44.5gh	58.1ih	55.5cd	50.0cdef	62.2abcd
Alpha	39.8h	78.1bcd	38.8e	53.4cde	55.4bcd
Bintje	39.4h	67.3fgh	36.4e	44.5cdefg	54.0bcd
Caribe	53.2efgh	73.9cdef	66.2abcd	59.0bc	75.5a
Carlton	72.1abc	70.5defg	72.7ab	60.5bc	60.5abcd
Eramosa	59.9bcde	84.7ab	56.3cd	55.4cd	65.4abc
Jesmeg	65.7bcde	78.6bcd	69.0abc	50.6cdef	55.4bcd
Kennebec	64.0bcde	80.0bcd	67.7abcd	56.2cd	75.5a
Norland	52.2efgh	77.5bcde	58.1cd	40.0efgh	50.4cd
Russet. B	69.2abc	81.7bc	57.5cd	72.9a	63.6abc
Range	(39.4-72.1)	(58.1-84.7)	(36.4-72.7)	(40.0-72.9)	(50.4-75.5)
Mean(cultivar)	56.0	75.0	57.8	54.2	61.9
<i>Acaule</i>	24.6i	37.5j	27.4e	57.0bcd	20.1e
<i>berthaultii</i>	64.2bcde	92.8a	75.9ab	59.6bc	55.0bcd
<i>chacoense</i>	58.3cdefg	61.5ghi	62.2bcd	36.0gh	63.4abc
<i>commersonii</i>	47.6fgh	74.0cdef	55.0d	37.5fgh	45.2d
<i>demissum</i>	25.4i	52.0i	29.5e	33.4h	28.4e
<i>hjertingii</i>	46.1fgh	74.0cdef	55.0d	37.5fgh	45.2d
<i>lycopersiocoide</i>	79.9a	68.0efg	74.4ab	69.8ab	65.4abc
<i>microdontum</i>	55.0defg	71.8cdef	74.8ab	43.3defgh	63.7abc
<i>phureja</i>	72.6ab	74.4a	72.6ab	67.1fgh	59.2bc
<i>stenotomum</i>	47.2fgh	54.7i	38.8e	58.7bc	50.2cd
<i>stoloniferum</i>	68.5abcd	53.0i	76.1ab	48.4cdefg	55.1bcd
Range	(24.6-79.9)	(37.5-92.8)	(29.5-78.8)	(32.1-69.8)	(20.1-65.4)
Mean (W.species)	53.6	62.2	60.7	51.3	47.9

Different letters in each column indicate significant differences by Duncan's multiple range test at 5%.

Kennebec and Caribe the least tolerant compared with the other cultivars. However, the wild species showed highly significant differences with respect to cold tolerance. *S. acaule*, *S. commersonii*, and *S. demissum* showed very high levels of cold tolerance relative to the other *Solanum* species and cultivars tested. These rankings, based on EC, support the field results of Vega and Bamberg (1995), who found high levels of cold tolerance in these three species. Cold tolerance showed significant correlations with other stresses, though the coefficient was not very high (Table 4).

CONCLUSION

This study showed that 1) The electrolyte leakage bioassay is able to detect differences among genotypes to many environmental stresses as the results are in agreement with many published results. 2) Variation existed between cultivars and, to a larger extent, between wild species for many stresses. 3) Generally, correlations existed among types of stress, especially between NaCl and PEG. This indicates that it might be possible to breed cultivars tolerant of more than one stress using multi stress tolerant wild species such as *S. acaule* and *S. demissum*. 4) Many

Table 3. Ranking of genotypes as affected by types of stress.

Genotype	NaCl	Na ₂ SO ₄	PEG	Heat	Cold
Agria	5	6	7	9	13
Alpha	4	16	3	12	11
Bintje	3	11	2	6	7
Caribe	10	13	12	16	20
Carlton	10	13	14	17	13
Eramosa	13	20	7	13	17
Jesmeg	16	17	13	11	12
Kennebec	14	18	19	14	21
Norland	9	15	10	7	5
Russet. B	17	19	9	21	17
<i>acaule</i>	2	1	1	15	1
<i>berthaultii</i>	14	21	18	17	7
<i>chacoense</i>	12	7	11	3	16
<i>commersonii</i>	8	2	20	1	2
<i>demissum</i>	1	3	5	2	3
<i>hjertingii</i>	6	13	6	4	4
<i>lycopersicoide</i>	21	9	15	20	17
<i>microdontum</i>	11	12	17	5	13
<i>phureja</i>	20	8	15	17	7
<i>stenotonum</i>	7	5	3	9	5
<i>stoloniferum</i>	17	4	21	8	7

Table 4. Pearson's coefficient correlations for genotypes to types of stress.

	NaCl	Na ₂ SO ₄	PEG	Heat	Cold
NaCl	-	0.44**	0.74**	0.45**	0.55**
Na ₂ SO ₄	-	-	0.33**	0.25**	0.53**
PEG	-	-	-	0.08 ^{Ns}	0.35**
Heat	-	-	-	-	0.39**

** Significant at 1%.

Ns: not significant



cultivars could be relatively easily evaluated by this method for immediate use in areas with environmental limitations. Similarly, more wild species and accessions could be screened for the identification of suitable parental lines with improved stress tolerance.

ACKNOWLEDGEMENTS

Financial support from Shahid Bahonar University of Kerman to the first author during his sabbatical leave in Canada is gratefully acknowledged. We also thank Dr. Shahrokh Khanizadeh from Agriculture and Agri-Food, Canada for supplying the glycol bath instrument.

REFERENCES

1. Adam, A. L., Gala, A. A., Manninger, K. and Bama, B. 2000. Inhibition of the Development of Leaf Rust (*Puccinia recondita*) by Treatment of Wheat with Allopurinol Production of a Hypersensitive Like Reaction in a Compatible Host. *Plant Pathol.*, **49**: 317-323.
2. Arvin, M. J. 1992. Tissue Culture in Potato and *in vitro* Selection for Salinity Tolerance. Ph. D. Thesis. Reading University, UK.
3. Bilski, J. J., Nelson, D. C. and Colon, R. L. 1988a. The Response of Four Potato Cultivars to Chloride Salinity, Sulphate, and Calcium in Pot Experiment. *Am. Potato J.*, **65**: 85-90.
4. Bilski, J. J., Nelson, D. C. and Colon, R. L. 1988b. Response of Six Wild Potato Species to Chloride and Sulfate Salinity. *Am. Potato J.*, **65**: 605-612.
5. Blum, A. and Ebercon, A. 1981. Cell Membrane Stability as a Measure of Drought and Heat Tolerance in Wheat. *Crop Sci.*, **21**: 43-47.
6. Boyer, J. S. 1982. Plant Productivity and Environment. *Science*, **18**: 443-448.
7. Brule-Babel, A. L. and Flower, D. B. 1989. Use of Controlled Environments for Winter Cereal Cold Hardiness Evaluation: Controlled Freeze Tests and Tissue Water Content as Prediction Tests. *Can. J. Plant Sci.*, **69**: 355-366.
8. Chen, Q., Zhang, W. H. and Liu, Y. L. 1999. Effect of NaCl, Glutathione and Ascorbic Acid on Function of Tonoplast Vesicles Isolated From Barley Leaf. *J. Plant Physiol.*, **155**: 685-690.
9. Christiansen, M. N. 1982. World Environmental Limitations to Food and Fiber Culture. In: "*Breeding Plants for Less Favourable Environment*". (Eds): Christiansen, M. N. and Lewis, C. F., John Wiley and Sons. New York, pp.1-12.
10. Cramer, G. R. and Spuir, A. R. 1986. Responses of Lettuce to Salinity. I. Effects of NaCl and Na₂SO₄ on Growth. *J. Plant Nutr.*, **9**(2): 115-130.
11. De, B. and Mukherjee, A. K. 1996. Mercuric Chloride Induced Membrane Damage in Tomato Cultured Cells. *Bio. Plant.*, **38**: 469-473.
12. Dodds, J. H. 1990. Molecular Biology of Potato: Current and Future Prospects for Developing Countries. In: "*The Molecular and Cellular Biology of the Potato Biotech Agriculture*". (Eds): Vaydr, M. E. and Park, W., **13**: 223-232.
13. Garty, J., Weissman, L., Tamir, Q., Beer, S., Cohen, Y. and Karnieli, A. 2000. Comparison of Five Physiological Parameters to Assess the Validity of the Lichen *Ramalina lacera* Exposed to Air Pollution. *Physiol. Plant*, **109**: 410-418.
14. Ismail, A. M. and Hall, A. E. 1999. Reproductive Stage, Heat Tolerance, Leaf Membrane Thermostability and Plant Morphology in Cowpea. *Crop Sci.*, **39**: 1762-1768.
15. Iyengar, E. R. R. and Reddy, M. P. 1994. Crop Response to Salt Stress: Sea Water Application and Prospects. In: "*Handbook of Plant and Crop Stress*". (Ed): Pesarakli, M., Marcel Dekker, New York, pp. 183-203.
16. Khrais, T., Leclerc, Y. and Donnelly, D. 1998. Relative Salinity Tolerance of Potato Cultivars Assessed by *in vitro* Screening. *Am. J. Potato Res.*, **75**: 207-210.
17. Levy, D. 1986. Tuber Yield and Tuber Quality of Several Potato Cultivars as Affected by Seasonal High Temperatures and by Water Deficit in a Semi-arid Environment. *Potato Res.*, **29**: 97-107.
18. Levy, D. 1984. Cultivated *Solanum tuberosum* as a Source for the Selection of Cultivars Adapted to Hot Climate. *Trop. Agri.*, **61**: 167-170.
19. Lynch, D. R. and Tai, G. C. C. 1989. Yield and Yield Component Response of Eight Po-

- tato Genotypes to Water Stress. *Crop Sci.*, **29**: 1207-1211.
20. Mahshawary, M. D., Joshi, K., Saha, R., Nagarajan, S. and Gambhir, P. N. 1999. Transverse Relaxation Time of Leaf Water Protons and Membrane Injury in Wheat (*Triticum aestivum*) in Response to High Temperature. *Ann. Bot.*, **84**: 741-745.
 21. Malik, N. J., Dwelle, R. B., Thonrton, M. K. and Pavék, J. J. 1992. Dry Matter Accumulation in Potato Clones Under Seasonal High Temperature Conditions in Pakistan. *Am. Potato J.*, **69**: 667-676.
 22. Menzel, C. M. 1985. Tuberization in Potato at High Temperatures: Interaction between Temperature and Irradiance. *Ann. Bot.*, **55**: 35-39.
 23. Morpurgo, R. 1991. Correlation between Potato Clones Grown *in vivo* and *in vitro* Under Sodium Chloride Stress Conditions. *Plant Breeding*, **107**: 80-82.
 24. Murashige, T. and Skoog, F. 1962. A Revised Medium for Rapid Growth and Bioassays with Tobacco Tissue Culture. *Physiol Plant.*, **15**: 473-497.
 25. Naik, P. S. and Widholm, J. M. 1993. Comparison of Tissue Culture and Whole Plant Responses to Salinity in Potato. *Plant Cell Tiss. Organ Cult.*, **33**: 273-280.
 26. Rahman, H. U., Maalik, S. A. and Saleem, M. 2003. Heat Tolerance of Upland Cotton during the Fruiting Stage Using Cellular Thermostability. *Field Crop Res.*, **85 (2-3)**: 149-158.
 27. Richardson, D. C. and Estreda, N. 1971. Evaluation of Frost Resistant Tuber Bearing Solanum Hybrids. *Am. Potato J.*, **48**: 339 - 343.
 28. Saelim, S. and Zwazek, J. J. 2000. Preservation of Thermal Stability of Cell Membranes and Gas Exchange in High Temperature Acclimated *Xylia xylocarpa*. *J. Plant Physiol.*, **156**: 380-385.
 29. Spencer, D. F. and Ksander, G. G. 1999. Influence of Dilute Acetic Acid Treatments on Survival of Monoecious *Hydrilla* Tuber in the Oregon House Canal, California. *J. Aqu. Plant Manag.*, **37**: 67-71.
 30. Srirama, S., Raguchander, T., Babu, S., Nandarkumar, R., Shanmugamm, V. and Vidhyasekaran, P. 2000. Inactivation of Phytotoxin Produced by the Rice Sheath Blight Pathogen *Rhizoctonia solani*. *Can. J. Microbiol.*, **46**: 520-524.
 31. Senaratana, T., Merritt, D., Kingsley, D., Eric, B., Darren, T. and Sivasithamparam, K. 2003. Benzoic Acid May Act as the Functional Group in Salicylic Acid and Derivatives in the Induction of Multiple Stress Tolerance in Plants. *Plant Growth Regul.*, **39**: 77-81.
 32. Singh, G. 1969. A Review of Soil Moisture Relationship in Potatoes. *Am. Potato J.*, **46**: 398-403.
 33. Sluc, R. M., Albrecht, K. and Duke, S. H. 1991. Leakage of Intracellular Substances as an Indicator of Freezing Injury in Alfalfa. *Crop Sci.*, **31**: 430-435.
 34. Struik, P. C. and Van Voorst, G. 1986. Effects of Drought on the Initiation, Yield and Size Distribution of *Solanum tuberosum* L. cv. Bintje. *Potato Res.*, **29**: 487-500.
 35. Sullivan, C. Y. and Ross, W. M. 1979. Selecting for Drought and Heat Resistance in Grain Sorghum. In: "Stress Physiology in Crop Plants". (Eds): Mussel, H., and Staples, R.C., John Wiley and Sons, New York, pp. 263-281.
 36. Vega, S.E. and Bamberg, J.B. 1995. Screening the U.S. Potato Collection for Frost Hardiness. *Am. Potato J.*, **72**: 13-21.
 37. SAS Institue. 1988. SAS Procedures Guide Release, 6.03 ed. SAS Institute, Cary, NC.
 38. Sekozawa, Y., Sugaya, S., Gemma, H. and Iwahori, S. 2003. Cold Tolerance in 'Kousui' Japanese Pear and Possibility for Avoiding Frost Injury by Treatment with n-Propyl Dihydrojasmonate. *HortScience.*, **38**: 288-292.
 39. Zhang, Y. and Donnelly, D. 1997. *In vitro* Bioassay for Salinity Tolerance Screening of Potato. *Potato Res.*, **40**: 285-295.



استفاده از روش نشت یونی برای غربال ارقام و گونه های وحشی سیب زمینی به تنشهای محیطی

م. ج. آروین و د. دونلی

چکیده

در یک سری آزمایش ها، دو رقم سیب زمینی و نژادهای مختلف از یازده گونه وحشی سیب زمینی برای مقاومت به تنشهای شوری (۲۰۰ میلی مول کلرورسديم یا ۱۰۰ میلی مول سولفات سدیم)، خشکی (۳۵ درصد پلی اتیلن گلیکول)، سرما (۴- درجه سانتی گراد) و گرما (۳۷ درجه سانتی گراد) ارزیابی گردید. ارزیابی مقاومت بر اساس میزان نشت یون از برگهای جدا شده از گیاهان درون شیشه ای بود. روش انجام آزمایش شامل سه بار شستشوی برگها با آب مقطر برای از بین بردن یونها از سطوح برگها و دمبرگها، قرار دادن برگها در معرض تنشهای شوری و خشکی برای مدت ۲۴ ساعت و مدت ۴ ساعت برای تنش های گرما و سرما، سپس نگهداری بافت ها در آب مقطر به مدت ۲۴ ساعت و نهایتاً اندازه گیری هدایت الکتریکی محلول به منظور تعیین مقدار تنش بود. نتایج آزمایش ها نشان داد اختلافات معنی داری بین ژنوتیپهای آزمایش شده برای انواع تنشها وجود دارد. از بین ارقام آزمایش شده، مقاومت به کلرور سدیم و خشکی در رقمهای آلفا و بیتزه، مقاومت به سولفات سدیم در رقم آگریا و مقاومت به گرما و سرما در رقم نورلند نسبت به اکثر ارقام بیشتر بود. در گونه های وحشی مقاومت به کلیه تنشها در *S. demissum*، به کلیه تنشها به غیر از گرما در *S. acaule* و مقاومت به گرما و سرما در *S. commersonii* بیشتر از اکثر گونه های وحشی بود. ضرایب همبستگی برای کلیه تنشها به جز برای گرما و خشکی معنی دار بود. میزان مقاومت به تنش ها در بعضی از گونه های وحشی بسیار بیشتر از ارقام بود و به نظر می رسد که منابع بسیار مهم مقاومت به تنش ها در ارقام وحشی وجود دارد که برای اصلاح ارقام می تواند مورد استفاده قرار گیرد.