

Dolichogenidea fernandeztrianai sp. nov. (Hymenoptera: Braconidae, Microgastrinae) from Iran

P. Abdoli¹, A. A. Talebi^{1*}, and S. Farahani²

ABSTRACT

The genus *Dolichogenidea* was studied from the northern Iran during 2010–2011. The specimens were collected using a set of Malaise traps. A new species *Dolichogenidea fernandeztrianai* Abdoli and Talebi sp. nov. is hereby described and illustrated. The new species can be distinguished by the following characters: fore wing vein R1 as long as or slightly longer than pterostigma and 4.90× as long as distance of vein R1 to vein 3RSb; T1 more or less parallel- to subparallel sided, smooth, with a distinct protuberance on central area and its length 1.70× posterior width; T2 smooth, transverse and rectangular; T3 longer than T2; ovipositor sheath with uniform width from base to apex and clearly longer than metatibia. An updated checklist for the Iranian species of this genus is provided in addition to their distribution data and references.

Keywords: New species, Species-groups, Taxonomy.

INTRODUCTION

Parasitic Hymenoptera is one of the extremely species-rich group of insects in terrestrial ecosystems (Shaw and Hochberg, 2001). The family Braconidae is the second largest family after Ichneumonidae in the order Hymenoptera (Aguiar *et al.*, 2013) include more than 21,220 species under 1,100 genera (Yu *et al.*, 2016). The braconid subfamily Microgastrinae is one of the most important groups of parasitoids in terms of both species richness and economic importance (Rodriguez *et al.*, 2013). This subfamily comprises 81 genera and over 2,700 species described worldwide (Yu *et al.*, 2016; Fernandez-Triana and Boudreault, 2018). The last estimate of species richness for Microgastrinae is likely 8–20 times that of currently described species. The extraordinary species richness of the group,

coupled with the occurrence of many cryptic species produces a significant taxonomic impediment that hinders the advancement of their study (Smith *et al.*, 2008, 2013). Rodriguez *et al.* (2013) suggested that number of described species could be about 5–15% of true global diversity of this subfamily.

Viereck (1911) initially described *Dolichogenidea* as a subgenus of *Apanteles* merely because of its elongate genae. Subsequently, it was treated as several species-groups of *Apanteles sensu lato* (Nixon, 1965), which was later raised to genus level by Mason (1981) in which concept of *Dolichogenidea* included three of Nixon's (1965) species-groups: the *ultor*-group, the *laevigata*-group, and the *longipalpis*-group.

Some genera (i.e. *Apanteles*, *Exoryza*, *Iconella* and *Illidops*) are the closest genera

¹ Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Islamic Republic of Iran.

² Research Institute of Forests and Rangelands, Agricultural Research Education and Extension Organization (AREEO), Tehran, Islamic Republic of Iran.

* Corresponding author; e-mail: talebia@modares.ac.ir



to *Dolichogenidea*, and *Apanteles* is the most controversial and difficult to separate from *Dolichogenidea*. Mason (1981) proposed several characters to distinguish *Dolichogenidea* from *Apanteles sensu stricto*, including punctures of anteromesoscutum typically distinctly separated and not breaking into aciculations posterolaterally; margin of vannal lobe convex and evenly setose; T1 parallel-sided or slightly wider apically; apicolateral areas of propodeum (if distinct) wider than high. The analysis of thousands of specimens by Fernandez-Triana et al. (2014) suggests that the only reliable character in *Dolichogenidea* is the presence of a convex to almost straight vannal lobe, which is uniformly fringed with setae, while in *Apanteles sensu stricto* the vannal lobe is strongly concave to almost straight and lacks setae medially. The lack of setae may be partial (there may be some small and sparse setae on the lobe) or complete (absence of setae). These two genera are also generally

resolved as separate monophyletic clades using molecular data and thus are distinguishable in DNA barcoding studies (Smith et al., 2013; Fagan-Jeffries et al., 2018).

The genus *Dolichogenidea* Viereck, 1911 is a large group consisting of over 235 species in the world and 84 species in the west Palearctic (Fernandez-Triana and Ward, 2015; Yu et al., 2016; Fagan-Jeffries et al., 2018; Fernandez-Triana et al., 2018; Liu et al., 2018). The Palearctic species of the genus *Dolichogenidea* have been revised and keyed by Nixon (1972), Papp (1978), Tobias (1986), Chen and Song (2004), and Liu et al. (2018).

The fauna of some subfamilies of Iranian Braconidae is already studied (Rakhshani et al., 2007; Farahani et al., 2012, 2014a,b, 2015, 2016; Ameri et al., 2015; Ghahari and Beyarslan, 2017; Rajabi Mazhar et al., 2018), but the Microgastrinae has been poorly studied (Farahani et al., 2014c, 2016; Gadallah et al., 2015; Ghafouri Moghaddam




Figure 1. Map of Iran: Mazandaran Province, where *Dolichogenidea fernandeztrianai* sp. nov. Specimens have been collected.

et al., 2018; Abdoli *et al.*, 2019). This group has a relatively wide range of hosts, including many lepidopteran families such as Aegeriidae, Choreutidae, Cochyliidae, Coleophoridae, Crambidae, Elachistidae, Gelechiidae, Momphidae, Oecophoridae, Plutellidae, Pterophoridae, Pyralidae, Tortricidae, and Yponomeutidae (Liu *et al.*, 2018).

The objective of this study was to improve our knowledge about the genus *Dolichogenidea* from Iran. We here describe a new species as a part of the ongoing research on the systematics of Microgastrinae from Iran.

MATERIALS AND METHODS

The specimens for the present study were collected using Malaise traps during March to November 2010–2011 in northern Iran (e.g., Alborz, Guilan, Mazandaran, Qazvin and Tehran Provinces) (Figure 1). Malaise traps were placed in different habitats such as forests, rangelands, and orchards.

The specimens were identified by the keys of Nixon (1972), Papp (1978), Tobias (1986), Chen and Song (2004) and Liu *et al.* (2018). Morphological terminology follows Wharton *et al.* (1997) for wing venation, and Karlsson and Ronquist (2012) for the other various body parts used in description of the new species. The abbreviations T1, T2, and T3 refer to the metasomal tergites 1, 2, and 3, respectively, and F 2, 3, 14, 15 refer to flagellomeres 2, 3, 14 and 15, respectively; OOL means Ocello-Ocular Line (minimum distance between posterior ocellus and eye), OD means Ocellus Diameter (transverse diameter of anterior ocellus) and POL means Posterior Ocellar Line (minimum distance between posterior ocelli). USNM means National Museum of Natural History, Washington, DC, USA.

Measuring ovipositor sheath length is usually difficult due to its curvature. Thus, the comparison of ovipositor sheaths is based on approximate (not exact) values of

its length with considering curvature (Abdoli *et al.*, 2019). We measured the ovipositor sheath length from the beginning of the setal area to the distal end of the sheath.

The specimens were photographed with a Keyence VHX-1000 Digital microscope, using a lens with a range of 13–130×. Multiple images through the focal plane later were combined to produce a single in-focus image. The software associated with the Keyence system produced the focused images taken with that camera. The measurements were done using an Olympus™ SZX9 stereomicroscope equipped with a graticule. The specimens of new species are deposited in the Insect Collection of the Department of Entomology, Tarbiat Modares University, Tehran, Iran (TMUC).

RESULTS

Present study describes one new *Dolichogenidea*. An updated checklist for the Iranian species of this genus in addition to their distribution data and references are provided (Table 1).

Dolichogenidea Viereck, 1911

Dolichogenidea Viereck, 1911: 173. Type species: *Apanteles (Dolichogenidea) banksi* Viereck, 1911, by original designation, (USNM).

Viereck (1911) initially described *Dolichogenidea* as a subgenus of *Apanteles*. Subsequently, it was treated as some species-groups of *Apanteles sensu lato* (Nixon, 1965), but was later raised to genus level by Mason (1981). Fagan-Jeffries *et al.* (2018), Fernandez-Triana *et al.* (2018), and Liu *et al.* (2018) considered *Dolichogenidea* as a valid genus. However, Yu *et al.* (2016) treated *Dolichogenidea* as a synonym of *Apanteles* Foerster (1863).

**Table 1.** Distribution and references for all species of the genus *Dolichogenidea* from Iran.

Number	Species	Distribution: Provinces (sampled regions)	References
1	<i>D. aethiopica</i> (Wilkinson, 1931)	Guilan (Fooman and Rasht)	(Ghahari et al., 2010)
2	<i>D. agilla</i> (Nixon, 1972)	East Azarbayjan (Arasbaran)	(Ghahari et al., 2011d)
3	<i>D. appellator</i> (Telenga, 1949)	Isfahan	(Kazemzadeh et al., 2014)
4	<i>D. candidata</i> (Haliday, 1834)	Ilam (Ilam) Lorestan (Noor Abad) Semnan (Damghan)	(Ghahari et al., 2011c) (Ghahari et al., 2012) (Samin and Fischer, 2015)
5	<i>D. cytherea</i> (Nixon, 1972)	Golestan (Kalaleh)	(Ghahari and Fischer, 2011)
6	<i>D. decora</i> (Haliday, 1834)	Qazvin (Danesfahan)	(Ghahari et al., 2011a)
7	<i>D. fernandeztrianai</i> Abdoli & Talebi sp. nov.	Mazandaran (Tangehvaz)	This study
8	<i>D. emarginata</i> (Nees, 1834)	Ardabil (Ardabil)	(Ghahari et al., 2009)
9	<i>D. halidayi</i> (Marshall, 1872)	Isfahan (Golpayegan)	(Ghahari et al., 2011b)
10	<i>D. iranica</i> (Telenga, 1955)	Kerman (Shurab - Zarudnyi), Sistan and Baluchistan (Degak, Megas - Zarudnyi).	(Telenga, 1955)
11	<i>D. lacteicolor</i> (Viereck, 1911)	Guilan (Bandar-e Anzali)	(Herard et al., 1979)
12	<i>D. longipalpis</i> (Reinhard, 1880)	Lorestan	(Pirhadi et al., 2008)
13	<i>D. seriphia</i> (Nixon, 1972)	Golestan (Galikesh)	(Ghahari and Fischer, 2011)
14	<i>D. sicaria</i> (Marshall, 1885)	Ilam (Darrehshahr)	(Ghahari et al., 2011a)

Diagnosis

Members of this genus are diagnosed by having moderate to large hypopygium, usually with a series of median longitudinal striae but at least sharply folded medially; ovipositor sheath usually long, setose throughout, sometimes with short ovipositor straight to gently down-curved; T1 length usually much longer than width, usually parallel-sided or barrel-shaped, but occasionally slightly wider or narrower apically and almost always bearing a median apical longitudinal depression; T2 always wider than its length and usually a little or much shorter than T3; propodeum coarsely sculptured to smooth and never with a median longitudinal carina, often with a more or less well-defined areola and

costulae, this carination often reduced to a U-shaped area or median depression extending forward from the abdominal fovea or carination completely suppressed, leaving a uniformly sculptured or smooth surface; margin of vannal lobe of hind wing uniformly convex and setose, rarely slightly flattened (Mason, 1981).

Dolichogenidea fernandeztrianai Abdoli and Talebi **sp. nov.** (Figures 2 and 3)

Type of Material

Holotype: Female, Mazandaran Province, Noor, Chamestan, Tangehvaz (36° 21' 55.68" N, 52° 06' 10.32" E, 702 m a.s.l.), 05.ix.2011 (TMUC). **Paratypes:** 2 females, same data as Holotype (TMUC).




Figure 2. *Dolichogenidea fernandeztrianai* Abdoli & Talebi **sp. nov.** (Holotype). Female: (A) Head; frontal view, (B) Head; lateral view, (C) Head; dorsal view, (D) Mesosoma, dorsal view; (E) Habitus; lateral view, (F) Metasoma; dorsal view, (G) Habitus; dorsal view.




Figure 3. *Dolichogenidea fernandeztrianai* Abdoli & Talebi sp. nov. (Holotype). Wings, female: (A) Fore wing, (B) Hind wing.

Diagnosis

The newly described species can be distinguished from all previously described species of the genus *Dolichogenidea* from the Palaearctic region by the following combination of characters: Antennal flagellomeres, all coxae and metafemur dark brown to black; tegula reddish yellow or yellow; pterostigma whitish yellow with brownish margin; propodeum with areola weakly defined by an impression; fore wing vein R1 as long as or slightly longer than pterostigma; vein R1 4.90× as long as distance of vein R1 to vein 3RSb (Figure 3-A); T1 more or less parallel- to subparallel-sided i.e., slightly narrower medially; T1 smooth, with a distinct protuberance on central area; T1 length 1.70× posterior width; T2 smooth, transverse and rectangular; T3 longer than T2 (Figure 2-F);

hypopygium pointed apically and slightly protruding beyond apical tergites; ovipositor sheath with uniform width from base to apex and clearly longer than metatibia (Figure 2-E).

Description (Female-Holotype)

Body Length without Ovipositor: 2.50 mm, antennae length 2.60 mm and forewing length 2.60 mm

Head: Antenna about as long as body; F 2, 3, 14, 15 length/width: 2.40, 2.40, 1.30 and 1.30, respectively; mouthparts not elongated (Figures 2-A and -B); head rounded or weakly constricted behind eyes (Figure 2-C); upper face (frons) smooth to rather finely punctate; gena, vertex and lower face (face) distinctly punctate; basal width of mandible 3.20–4.00× malar space length (Figure 2-A); OOL, OD, POL length: 0.13,

0.06 and 0.13 mm, respectively (Figure 2-C).

Mesosoma: Anteromesoscutum densely punctate and setose, notauli indistinct; scutoscuteellar suture narrow, distinctly crenulate; mesoscutellar disc finely and sparsely punctate to smooth and shiny; metanotum with uniform coarse wrinkles; anterior half of propodeum with few punctures and shiny, rest mostly smooth to slightly uneven; propodeum without distinct areola, weakly defined by an impression; pronotum weakly punctate; propleuron punctate; prepectal carina absent; mesopleuron medially smooth with some punctures on antero- and ventro-lateral parts; metapleuron weakly wrinkle to smooth, anteriodorsal corner rugulose, and with deep medial pit on anterior half (Figure 2-D).

Wings: Forewing length 2.60 mm; Forewing length/width: 2.90; vein R1 as long as or slightly longer than pterostigma; Pterostigma length/width: 2.50; vein R1 4.90× as long as distance of vein R1 to vein 3RSb; vein r as long as or slightly longer than vein 2RS; vein 1-CU_b slightly shorter than vein 1-CU_a (Figure 3-A).

Legs: Metafemur length/width: 3.60; length of metafemur, metatibial, metabasitarsus and sum of other metatarsal segments i.e., sum of segments 2, 3, 4 and 5: 0.75, 0.85, 0.40 and 0.60 mm, respectively; length of metatibial inner and outer spur 0.20 and 0.17 mm, respectively; metatibia inner spur length/metabasitarsus length: 0.50.

Metasoma: T1 more or less parallel- to subparallel-sided i.e. slightly narrower medially; T1 smooth, with a distinct central protuberance; T1 length 1.70× posterior width; T2 smooth, transverse and rectangular (T2 posterior width 2.75× its medial length); T3 longer than T2 (T3/T2 length: 1.25) (Figure 2-F); hypopygium pointed apically, slightly extending beyond the apex of metasoma; ovipositor sheath with uniform width and slightly curved downwards; ovipositor sheath clearly longer than metatibia (ovipositor sheath 1.20× as

long as metatibia), ovipositor sheath length 1.15 mm (Figure 2-E).

Colouration: Body black; except yellow or reddish yellow mouthparts, tegula, fore legs, apex of mesofemur, mesotibia, mesotarsus, basal one-third of metatibia and ovipositor; metatarsus brown; palpi whitish yellow; pterostigma whitish yellow, with brownish margin; wing venation with combination of whitish yellow and brown (Figures 2 and 3).

Male: Unknown.

Note: Although there is no available key that covers all Palaearctic species of *Dolichogenidea*, we provide comparisons of *Dolichogenidea fernandeztrianai* sp. nov. with all previous keys of Palaearctic *Dolichogenidea*.

Dolichogenidea fernandeztrianai sp. nov. runs close to *D. britannica* (Wilkinson, 1941), in the keys by Nixon (1972) but it differs from the latter as follows: in *D. fernandeztrianai* sp. nov. T1 more or less parallel- to subparallel-sided not narrowed apically and ovipositor sheath 1.20× as long as metatibia (whereas in *D. britannicus* T1 markedly narrowed towards apex and ovipositor sheath twice as long as metatibia). This species runs into *D. jilinensis* Chen and Song (2004) [this species is synonymized with *D. purda* (Papp, 1977) by Liu *et al.* (2018)] as per the key by Chen and Song (2004), nevertheless, it differs from *D. fernandeztrianai* sp. nov. as follows: in *D. fernandeztrianai* sp. nov. T1 length 1.70× posterior width; pterostigma whitish yellow, with brownish margin; anterior half of propodeum with some punctures, shiny, other areas mostly smooth to slightly uneven, propodeum without distinct areola, weakly defined by an impression; notauli not defined (whereas in *D. purda*, T1 1.50× its posterior width, pterostigma evenly brown; propodeum mostly smooth, with only a small trace of posterior areola carinae near nucha, notauli distinct by crowded punctation). This species runs in the key by Liu *et al.* (2018) to *D. poliobrevis* Liu & Chen, 2018, however, it differs from *D. fernandeztrianai* sp. nov. as follows: in *D. fernandeztrianai*



sp. nov. T1 more or less parallel- to subparallel-sided i.e., slightly narrower medially, T1 length $1.70\times$ posterior width; pterostigma whitish yellow, with brownish margin; ovipositor sheath $1.20\times$ as long as metatibia (whereas in *D. poliobrevis* T1 parallel-sided and as long as posterior width, pterostigma brown, ovipositor sheath nearly as long as metatibia).

Dolichogenidea fernandeztrianai **sp. nov.** is similar to *Dolichogenidea eleagnellae* Tobias, 1976, but it differs from *D. eleagnellae* as follows: in *D. fernandeztrianai* **sp. nov.** scutoscuteellar suture narrowed with 20 crenulate; ovipositor sheath clearly longer than metatibia, $1.20\times$ as long as metatibia; pterostigma whitish yellow, with brownish margin (whereas in *eleagnellae* scutoscuteellar suture wide and deep with 10 crenulate; ovipositor sheath slightly longer than metatibia; pterostigma entirely dark i.e. opaque brown to blackish brown, without pale basal spot).

Paratypes: F 2, 3, 14, 15 length/width: 2.40–2.70, 2.40–2.70, 1.16–1.30 and 1.20–1.25, respectively; OOL, OD, POL length: 0.12, 0.05–0.06 and 0.13–0.15 mm, respectively; Forewing length: 2.70–2.75 mm; Forewing length/width: 2.90–3.00; Pterostigma length/width: 2.60–2.70; length of metafemur, metatibial, metabasitarsus and sum of other metatarsal segments: 0.71–0.75, 0.83–0.92, 0.41–0.42 and 0.60 mm, respectively.

Etymology: The new species is named in the honor of Dr. Jose L. Fernandez-Triana, specialist on the systematics of Braconidae at the Canadian National Collection of Insects, for his significant contributions to the Microgastrinae knowledge of the world.

DISCUSSION

The new species was collected from deciduous forest of Mazandaran Province. The genus *Dolichogenidea* is newly recorded from Mazandaran Province. The sampled regions in this study belong to the

Irano-Anatolian hotspot that forms a natural barrier between the Mediterranean basin and the dry plateaus of western Asia (Paknia and Rajaei, 2015; Kiani et al., 2017). The mountains and basins that make up the Irano-Anatolian region contain many centers of local endemism. Most of north and west parts of Iran belong to this Irano-Anatolian region (Kiani et al., 2017; Mohammadi-Khoramabadi and Talebi, 2018), which is expected to have the highest biodiversity in Iran. Previously, the species of this genus were recorded from 11 provinces (Table 1), which shows that *Dolichogenidea* species are well adapted to a wide range of habitats in north, central, and south Iran. This study has added one new species of *Dolichogenidea*, thus increasing the total number to 85 in the west Palearctic region. The number of Iranian *Dolichogenidea* species has risen from 13 to 14. However, the majority of the species records from Iran are not confirmed, since they are sourced from the solely faunistic works, with no depository. Therefore, only an updated checklist is written based on previous literatures (Table 1). The number of Iranian *Dolichogenidea* species compares to neighboring countries as follows: 41 species in Russia (Telenga, 1955), 27 species in Turkey (Inanc and Beyarslan, 2001), 20 species in Azerbaijan (Tobias, 1976; 1986), eight species in Tajikistan (Telenga, 1955; Kotenko, 2007), four species in Afghanistan (Tobias, 1998) and one species in Pakistan (Watanabe, 1967). Further investigations are needed to obtain better knowledge on the genus and unveil distribution range of Iranian *Dolichogenidea* because many areas of the country are still unexplored.

ACKNOWLEDGEMENTS

Thanks are due to the Department of Entomology, Tarbiat Modares University for providing financial support for this study and to the Canadian National Collection of Insects (CNC) of Agriculture and Agri-Food

Canada for providing facilities. We cordially thank four anonymous reviewers for their critical review and constructive comments, which significantly helped the improvement of the manuscript. We are thankful to Drs. Khayrandish, Nadimi and Mohammadi for helping with the collection of the specimens. The senior author especially thanks Dr. Jose L. Fernandez-Triana (CNC) for his guidance in improving her knowledge on Microgastrinae.

REFERENCES

1. Abdoli, P., Talebi, A. A., Farahani, S. and Fernandez-Triana, J. 2019. Three New Species of the Genus *Choeras* Mason, 1981 (Hymenoptera: Braconidae, Microgastrinae) from Iran. *Zootaxa*, **4545(1)**: 77–92.
2. Aguiar, A., Deans, A. R., Engel, M.S., Forshage, M., Huber, J. T., Jennings, J. T., Johnson, N., Lelej, A. S., Longino, J. T., Lohrmann, V., Miko, I., Ohl, M., Rasmussen, C., Taeger, A. and Yu, D. S. K. 2013 Order Hymenoptera. *Zootaxa*, **3703(1)**: 51–62.
3. Ameri, A., Talebi, A. A., Rakhshani, E., Beyarslan, A. and Kamali, K. 2015. Additional Evidence and New Records of the Genus *Bracon* Fabricius, 1804 (Hymenoptera: Braconidae) in Southern Iran. *Turk. J. Zool.*, **39(6)**: 1110–1120.
4. Chen, J. and Song, D. 2004. *Systematic Studies on Microgastrinae of China (Hymenoptera: Braconidae)*. Fujian Scientific Publisher, Fuchow, China, PP. 1–354.
5. Fagan-Jeffries, E. P., Cooper, S. J. B. and Austin, A. D. 2018. Three New Species of *Dolichogenidea* Viereck (Hymenoptera, Braconidae, Microgastrinae) from Australia with Exceptionally Long Ovipositors. *J. Hymenopt. Res.*, **190**: 177–190.
6. Farahani, S., Talebi, A. A. and Rakhshani, E. 2014a. Wasps of the Subfamily Doryctinae (Hymenoptera: Braconidae) in Iran. *Zool. Middle East*, **60(1)**: 65–81.
7. Farahani, S., Talebi, A. A. and Rakhshani, E. 2016. Iranian Braconidae (Insecta: Hymenoptera: Ichneumonoidea): Diversity, Distribution and Host Association. *J. Ins. Biodiv. Syst.*, **2(1)**: 1–92.
8. Farahani, S., Talebi, A.A., Rakhshani, E. and van Achterberg, C. 2012. First Record of *Homolobus infumator* (Lyle, 1914) (Insecta: Hymenoptera: Braconidae: Homolobinae) from Iran. *Check List*, **8(6)**: 1350–1352.
9. Farahani, S., Talebi, A. A., van Achterberg, C. and Rakhshani, E. 2014b. A New Species of the Genus *Mirax* Haliday, 1833 (Hymenoptera: Braconidae: Miracinae) from Iran. *Ann. Zool.*, **64(4)**: 677–682.
10. Farahani, S., Talebi, A. A., van Achterberg, C. and Rakhshani, E. 2014c. First Record of *Microplitis rufiventris* Kokujev, 1914 (Braconidae: Microgastrinae) from Iran. *Check List*, **10(2)**: 441–444.
11. Farahani, S., Talebi, A. A., van Achterberg, C. and Rakhshani, E. 2015. A Review of the Subfamily Rogadinae (Hymenoptera: Braconidae) from Iran. *Zootaxa*. **3973(2)**: 227–250.
12. Fernandez-Triana, J. and Boudreault, C. 2018. Seventeen New Genera of Microgastrine Parasitoid Wasps (Hymenoptera, Braconidae) from Tropical Areas of the World. *J. Hymenopt. Res.*, **64**: 25–140.
13. Fernandez-Triana, J., Sakagami, K. and Shimizu, S. 2018. *Dolichogenidea maetoi* sp. nov. (Hymenoptera: Braconidae) from Japan, the First Parasitoid Wasp Recorded from *Hyblaea fortissima* (Lepidoptera). *Acta Ent. Mus. Nat. Pra.*, **58(1)**: 167–75.
14. Fernandez-Triana, J., Whitfield, J. B., Rodriguez, J. J., Smith, M. A., Janzen, D. H., Hallwachs, W. D., Hajibabaei, M., Burns, J. M., Solis, M. A., Brown, J., Cardinal, S., Goulet, H. and Hebert, P. D. 2014. Review of *Apanteles sensu stricto* (Hymenoptera, Braconidae, Microgastrinae) from Area de Conservación Guanacaste, Northwestern Costa Rica, with Keys to All Described Species from Mesoamerica. *ZooKeys*, **565**: 1–565.
15. Gadallah, N. S., Ghahari, H. and Peris-Felipo, F. J. 2015. Catalogue of the Iranian Microgastrinae (Hymenoptera: Braconidae). *Zootaxa*, **4043(1)**: 1–69.
16. Ghafouri Moghaddam, M., Rakhshani, E., van Achterberg, C. and Mokhtari, A. 2018. A study of the Iranian species of



- Choeras* Mason (Hymenoptera: Braconidae: Microgastrinae), with the Description of a New Species. *Zootaxa*, **4446(4)**: 455–476.
17. Ghahari, H. and Beyarslan, A. 2017. A Faunistic Study on Braconidae (Hymenoptera: Ichneumonoidea) from Iran. *Nat. Somogy.*, **30**: 39–46.
 18. Ghahari, H. and Fischer, M. 2011. A Contribution to the Braconidae (Hymenoptera) from Golestan National Park, Northern Iran. *Z. Arbeitsgem. Oesterr. Entomol.*, **63(1)**: 78–80.
 19. Ghahari, H., Fischer, M. and Papp, J. 2011a. A study on the Braconidae (Hymenoptera: Ichneumonoidea) from Qazvin Province, Iran. *Entomofauna*, **32(9)**: 197–204.
 20. Ghahari, H., Fischer, M. and Papp, J. 2011b. A Study on the Braconid Wasps (Hymenoptera: Braconidae) from Isfahan Province, Iran. *Entomofauna*, **32(16)**: 261–272.
 21. Ghahari, H., Fischer, M. and Papp, J. 2011c. A Study on the Braconidae (Hymenoptera: Ichneumonoidea) from Ilam Province, Iran. *Calodema*, **160**: 1–5.
 22. Ghahari, H., Fischer, M., Cetin Erdogan, O., Beyarslan, A., Hedqvist, K. J. and Ostovan, H. 2009. Faunistic Note on the Braconidae (Hymenoptera: Ichneumonoidea) in Iranian Alfalfa Fields and Surrounding Grasslands. *Entomofauna*, **30(24)**: 437–444.
 23. Ghahari, H., Fischer, M., Papp, J. and Tobias, V. 2012. A Contribution to the Knowledge of Braconids (Hymenoptera: Braconidae) from Lorestan Province, Iran. *Entomofauna*, **33(7)**: 65–72.
 24. Ghahari, H., Fischer, M., Sakenin, H. and Imani, S. 2011d. A contribution to the Agathidinae, Alysinae, Aphidiinae, Braconinae, Microgastrinae and Opiinae (Hymenoptera: Braconidae) from Cotton Fields and Surrounding Grasslands of Iran. *Linz. Biol. Beitr.*, **43(2)**: 1269–1276.
 25. Ghahari, H., Erdogan, O.C., Sedivy, J. and Ostovan, H., 2010. Survey of the Ichneumonoidea and Chalcidoidea (Hymenoptera) Parasitoids of Saturniidae (Lepidoptera) in Iran. *Efflatounia*, **10**: 1–6.
 26. Herard, F., Mercadier, G. and Abai, M. 1979. Situation de *Lymantria dispar* (Lep.: Lymantriidae) et son Complexe Parasitaire en Iran, en 1976. *Entomophaga*, **24(4)**: 371–384.
 27. Inanc, F. and Beyarslan, A. 2001. A Study on Microgastrinae (Hymenoptera: Braconidae) Species in Gokceada and Bozcaada. *Turk. J. Zool.*, **25(3)**: 287–296.
 28. Karlsson, D. and Ronquist, F. 2012. Skeletal Morphology of *Opius dissitus* and *Biosteres carbonarius* (Hymenoptera: Braconidae), with a Discussion of Terminology. *PLoS ONE*, **7(4)**: 1–38.
 29. Kazemzadeh, Z., Shaw, M. R. and Karimzadeh, J. 2014. A New Record for Iran of *Dolichogenidea appellator* (Hym.: Braconidae: Microgastrinae), a Larval Endoparasitoid of Diamondback Moth, *Plutella xylostella* (Lep.: Plutellidae)(Short communication). *J. Entomol. Soc. Iran*, **33**: 81–82.
 30. Kiani, M., Mohammadi, S., Babaei, A., Sefidkon, F., Naghavi, M. R., Ranjbar, M., Razavi, S. A., Saeidi, K., Jafari, H., Asgari, D. and Potter, D. 2017. Iran supports a great share of biodiversity and floristic endemism for *Fritillaria* spp. (Liliaceae): A review. *Plant Diversity*, **39(5)**: 245–262.
 31. Kotenko, A. G. 2007. Review *Palaeartic* Species of the Genus *Iconella* (Hymenoptera, Braconidae, Microgastrinae): Species with Propodeum Lacking the Median Longitudinal Carina. *Vestn. Zool.*, **41(4)**: 315–325.
 32. Liu, Z., He, J. and Chen, X. 2018. The *Laevigata*-Group of the Genus *Dolichogenidea* Mason, 1981 from China, with Descriptions of 26 New Species. *Zootaxa*, **4436(1)**: 001–074.
 33. Mason, W. R. M. 1981. The Polyphyletic Nature of *Apanteles* Foerster (Hymenoptera: Braconidae): A Phylogeny and Reclassification of Microgastrinae. *Mem. Entomol. Soc. Can.*, **113(115)**: 1–147.
 34. Mohammadi-Khoramabadi, A. and Talebi, A.A. 2018. Study of Three Genera of the *Orthocentrus* Genus-Group (Hymenoptera: Ichneumonidae, Orthocentrinae) in Northern Iran. *J. Entomolo. Soc. Iran*, **37(4)**: 441–460.
 35. Nixon, G. E. 1965. A Reclassification of the Tribe Microgastrini (Hymenoptera: Braconidae). *Bull. Br. Mus. Nat. Hist. Entomol.*, **2(Suppl.)**: 1–284.

36. Nixon, G. E. 1972. A Revision of the North-Western European Species of the *Laevigatus*-Group of *Apanteles* Förster (Hymenoptera, Braconidae). *Bull. Entomol. Res.*, **61**: 701–743.
37. Paknia, O. and Rajaei, Sh., H. 2015. Geographical Patterns of Species Richness and Beta Diversity of Larentiinae Moths (Lepidoptera: Geometridae) in Two Temperate Biodiversity Hotspots. *J. Insect Conserv.*, **19(4)**: 729–739.
38. Papp, J. 1978. A Survey of the European Species of *Apanteles* Först. (Hymenoptera, Braconidae: Microgasterinae), II. The *Laevigatus*-Group, 1. *Annls Hist. -Nat. Mus. Nat. Hung.*, **70**: 265–301.
39. Pirhadi, A., Rajabi, G., Ebrahimi, E., Ostovan, H., Shekarian Moghaddam, B., Mohiseni, A.A., Mozaffarian, F. and Ghavami, S. 2008. Natural Enemies of Cereal Leaf Miner, *Syringopais temparetella* Led. (Lep.: Elachistidae) in Lorestan Province. *Proceedings of 18th Iranian Plant Protection Congress*, P. 71, Bu-Ali Sina University, Hamadan, Iran.
40. Rajabi Mazhar, A., Goldasteh, S., Farahani, S. and van Achterberg, C. 2018. New Species and New Record of Chelonini (Hymenoptera: Braconidae) from Western Iran. *Turk. J. Zool*, **42(4)**: 416–421.
41. Rakhshani, E., Talebi, A. A., Sary, P. S., Tomanovic, Z. T. and Manzari, S. M. 2007. Aphid-Parasitoid (Hymenoptera, Braconidae, Aphidiinae) Associations on Willows and Poplars in Iran. *Acta Zool. Acad. Sci. Hung.*, **53(3)**: 281–292.
42. Rodriguez, J. J., Fernandez-Triana, J. L., Smith, M. A., Janzen, D. H., Hallwachs, W., Erwin, T. L. and Whitfield, J. B. 2013. Extrapolations from Field Studies and Known Faunas Converge on Dramatically Increased Estimates of Global Microgastrine Parasitoid Wasp Species Richness (Hymenoptera: Braconidae). *Insect. Conserv. Diver.*, **6(4)**: 530–536.
43. Samin, N. and Fischer, M. 2015. A contribution to the Study on the Fauna of Braconidae (Hymenoptera, Ichneumonoidea) from the Province of Semnan, Iran. *Arquivos Entomoloxicos*, **13**: 429–433.
44. Shaw, M. and Hochberg, M. 2001. The Neglect of Parasitic Hymenoptera in Insect Conservation Strategies: The British Fauna as a Prime Example. *J. Insect Conserv.*, **5**: 253–263.
45. Smith, M. A., Andez-triana, J. L. F., Eveleigh, E. and Omez, J. G. 2013. DNA Barcoding and the Taxonomy of Microgastrinae Wasps (Hymenoptera, Braconidae): Impacts after 8 Years and Nearly 20000 Sequences. *Mol. Ecol. Resour.*, **13(2)**: 168–176.
46. Smith, M. A., Rodriguez, J. J., Whitfield, J. B., Deans, A. R., Janzen, D. H., Hallwachs, W. and Hebert, P. D., .2008. Extreme Diversity of Tropical Parasitoid Wasps Exposed by Iterative Integration of Natural History, DNA Barcoding, Morphology and Collections. *Proc. Natl. Acad. Sci. U S A*, **105**: 12359–12364.
47. Telenga, N. A., 1955. Hymenoptera. Vol. 5 No 4. Family Braconidae: Subfamily Microgasterinae, Subfamily Agathidinae. *Fauna SSSR (NS)*, **61**: 1–312.
48. Tobias, V. I. 1976. [*Braconids of the Caucasus (Hymenoptera, Braconidae)*.] (in Russian) Opred. Faune SSSR, Nauka Press, Leningrad, **110**: 286.
49. Tobias, V. I. 1986. Subfamily Microgasterinae. Part IV. In: (Ed.): “*Keys to the Insects of the European Part of the USSR*”, Medvedev, G. S. Volume III, Amerind Publishing Co. Pvt. Ltd., 883 PP.
50. Tobias, V. I., Capek, M. and Lauterer, P. 1998. Braconidae (Hymenoptera) from Afghanistan in the Collections of the Moravian Museum. *Acta Mus. Morav. Sci. Biol.*, **82**: 173–190.
51. Viereck, H. L. 1911. Descriptions of Six New Genera and Thirty-One New Species of *Ichneumon* flies. *Proceedings of the United States National Museum*, **40(1812)**: 173–196.
52. Watanabe, C. 1967. Notes on Braconidae Caught in a Sweep-Net at Paddy Fields: Part I. *Mushi*, **40**: 189–198.
53. Wharton, R. A., Marsh, P. M. and Sharkey, M. J. 1997. Manual of the New World Genera of the Family Braconidae (Hymenoptera). *Intl. Soc. Hymenopterists*, **1**: 439.
54. Yu, D. S. K., Van Achterberg, C. and Horstmann, K. 2016. *Taxapad 2016, Ichneumonoidea 2015*. Database on Flash-Drive. Nepean, Ontario, Canada.



توصیف گونه جدید *Dolichogenidea fernandeztrianai* sp. nov. (Hymenoptera: Braconidae: Microgastrinae) از ایران

پ. عبدلی، ع. ا. طالبی، و س. فراهانی

چکیده

جنس *Dolichogenidea* در طی سالهای ۱۳۸۹-۱۳۹۰ در شمال ایران مورد مطالعه قرار گرفت. نمونه ها با استفاده از یک سری از تله های مالایز جمع آوری شد. گونه جدید *Dolichogenidea fernandeztrianai* Abdoli and Talebi sp. nov. توصیف و شکل های مربوط به ویژگی های تاکسونومیک آن ارائه گردید. گونه جدید با استفاده از خصوصیات زیر تشخیص داده می شود: در بال جلو طول رگبال R1 برابر یا اندکی بلندتر از پترواستیگما و ۴.۹ برابر فاصله رگبال R1 تا رگبال RSb^۳؛ حاشیه های جانبی ترزیت اول شکم کم و بیش موازی، صاف و طول آن ۱.۷ برابر عرض ناحیه عقبی آن؛ ترزیت دوم شکم صاف، عریض و مستطیلی شکل؛ ترزیت سوم شکم بلندتر از ترزیت دوم؛ غلاف تخم‌ریز از قاعده تا انتها تقریباً یک شکل و طول آن بیشتر از طول ساق پای عقب است. چک لیست به روز شده برای گونه های ایرانی این جنس همراه با اطلاعات پراکنش جغرافیایی و مراجع مرتبط ارائه شده است.