

Utilizing Deficit Irrigation to Enhance Growth Performance and Water-use Efficiency of Eggplant in Arid Environments

O. Mohawesh¹

ABSTRACT

The objective of this research was to investigate the effects of deficit irrigation on physiological and agronomic terms of eggplant to maximize the Water Use Efficiency (WUE) without affecting the final yield and fruit quality parameters under arid environment. Therefore, two field experiments were conducted at two different sites: Ghor Al-Safi, Jordan Valley and Sail Al-Karak, Karak Valley, Karak Province, Jordan, using a common eggplant cultivar (Classic) using five irrigation levels: 20, 40, 60, 80, and 100% based on field capacity. The most stressful Deficit Irrigation (DI) treatments (40 and 20%) resulted in significant effects on leaf area, leaf relative water content, leaf water potential and leaf mineral content. Biochemical parameters also showed an increase in proline and a decrease in chlorophyll content under water deficit conditions. Fruit weight and total yield decreased with DI. The control (100% irrigation treatment) plants revealed higher nutrient contents than the water-stressed plants. The fruit TSS and titratable acidity were increased at both sites as the irrigation regime decreased from 100 to 20%. Fruit nutrient content decreased with increasing water deficit. However, the differences were not significant between the control (100% irrigation treatment) and the 80% irrigation treatment. The 80% treatment showed high water use efficiency with relatively small effects on plant growth performance compared with the control. As a result, DI level at 80% can be utilized to increase WUE without a significant effect on crop growth performance.

Keywords: Arid and semi-arid environments, Crop quality, Deficit irrigation, Water use efficiency, Yield.

INTRODUCTION

Water shortages in Mediterranean countries, mainly in arid and semi-arid zones are one of the main limiting factors in irrigated agriculture. Jordan is considered to be one of the ten poorest countries worldwide in water resources (Shatanawi *et al.*, 2007). The agricultural irrigation amount allocated for irrigation decreased through the period 1985–2008 (78% in 1985 to 60% in 2008) (Shatanawi *et al.*, 2007). Although irrigated areas increase to satisfy high population growth, a great amount of water resources will be diverted from agriculture to balance the growing water demand from

municipal and industrial sectors (Correia, 1999). Because of this, it is necessary to implement efficient irrigation management strategies.

The shortage of water resources may prevent additional growth of irrigated agriculture (Mohawesh and Al-Absi, 2009). In several countries, renewable water resources have already been exceeded. This has resulted in declining groundwater levels and water quality deterioration (Shatanawi *et al.*, 2007). The overuse of water resources possibly will reach a crisis degree (Gleich 2000; Mohawesh and Karajeh, 2014). Consequently, for sustainable irrigation water resources, it has become an imperative concern to enhance crop Water Use

¹ Department of Plant Production, Faculty of Agriculture, Mutah University P.O. Box 7, Karak 61710, Jordan. e-mail: osama.mohawesh@gmail.com



Efficiency (WUE) through suitable irrigation design and management. Deficit Irrigation (DI) has been commonly used as a suitable strategy in arid and semi-arid regions (Mohawesh *et al.*, 2010; Mohawesh and Karajeh, 2014) where water is the most limiting natural resource. DI has been applied productively on field crops. Kirnak *et al.* (2001) evaluated the effects of irrigation regimes (100, 80, 60, and 40% of Pot Capacity (PC)) on eggplant. They showed that plants grown under a lower irrigation regime had less fruit yield and quality, considerable decreases in chlorophyll content, lower leaf Relative Water Content (RWC), and less vegetative growth than those in the control treatment. Abd El-Aal *et al.* (2008) investigated the effects of irrigation interval (10 and 21 days intervals) on eggplant growth performance. They found that plant growth, total yield, and fruit physical and chemical properties were better when eggplant was irrigated at 10-day intervals. Since DI affects plant growth performance, yield, and fruit quality, it is an urgent issue to examine the degree of water deficit that can be applied without major losses in crop production under different conditions. Therefore, the aims of this work were to study the main effects of water stress on physiological and agronomic terms, and to define the threshold values of water stress to maximize the water use efficiency without affecting the final yield and fruit quality parameters.

MATERIALS AND METHODS

Description of the Experimental Sites

Two experiments were conducted at two sites: Ghor Al-Safi (31° 3' 38" N; 35° 29' 13" E, -384 m asl) from September 15, 2012 until February 15, 2013 and Sail Al-Karak (31° 12' 1" N; 35° 41' 37" E, 681 m asl) from February 15, 2013 until July 15, 2013, Karak Province of Jordan at private farms. The Ghor Al-Safi site has a subtropical climate with an annual average temperature

and rainfall of about 25°C and 83 mm (1986–2010), respectively. The Sail Al-Karak site has long-term averages of temperature and rainfall of about 18°C and 250 mm (1986–2010), respectively.

Description of the Experimental Design and Treatments

Two field experiments were conducted using a common eggplant cultivar (Classic). The sites were prepared for planting by plowing, disking, and leveling. A drip irrigation system was used. Polyethylene drip laterals (20 mm inside diameter) were installed before planting, with emitters (rated at 8 l h⁻¹ discharge) spaced every 0.3 m on the laterals. A buffer zone with spacing of 1.0 m was provided between each treatment. Experimental treatments were arranged using a Randomized Complete Block Design (RCBD). The eggplant crop was fertigated 5 times during the growing season according to the traditional agricultural practices in the area. A total of 450 kg urea ha⁻¹ (46% N), 250 kg ha⁻¹ ammonium sulfate, and 720 kg ha⁻¹ NPK (20:20:20 with trace elements) fertilizers was applied by fertigation. In the application of the fertilizer, a pressure differential was formed by throttling the water flow in the control head and diverting a fraction of the water through a tank containing the fertilizer solution. The fertigation period was dependent on the lowest DI level duration (Mahadeen *et al.*, 2011). 5-week-old (0.10 to 0.15 m height) eggplant transplants were planted on 15 September and 15 February at Ghor Al-Safi and Sail Al-Karak, respectively. Each experiment consisted of five treatments with three replications per treatment. The laterals were mulched with 80 cm wide black plastic. Row spacing was 160 cm, while the plant spacing was 30 cm. A single row of plants was grown in the middle of the plastic mulch with a single lateral for each row, with 100 plants per row and 500 plants per plot, thus the total


number of plants were 1,500 for the whole experiment.

Experimental treatments consisted of five irrigation regimes (20, 40, 60, 80, and 100% based on Field Capacity (FC)) based on Crop Consumptive Use (ET_c). All treatments were irrigated initially with the same amount of water for two weeks to achieve uniform initial water content and to reach FC . Irrigation treatments were commenced two weeks after transplanting. The measured meteorological data from the installed


weather station inside the field were used to calculate the reference Evapotranspiration (ET_o) [Figure 1 (a, b)]. The ET_o was calculated using the locally calibrated Hargreaves model (Mohawesh and Talazi, 2012) [Equation (1)].

$$ET_o = 0.408 \times 0.6957 \times 0.0023 (T_{mean} + 16.6) (T_{max} - T_{min})^{0.58} R_a \quad (1)$$

Where, ET_o is the reference evapotranspiration in mm day^{-1} , T_{max} , T_{min} ,



(a)



(b)

Figure 1. Meteorological data during the experiment at Ghor Al-Safi (a) and Sail Al-Karak (b) sites during the experiment period.



and T_{mean} are the maximum, minimum, and mean air temperatures ($^{\circ}\text{C}$), respectively, and R_a is the extraterrestrial radiation ($\text{MJ m}^{-2} \text{ day}^{-1}$). The ET_c was calculated by multiplying ET_o by the Crop Coefficient (K_c): $ET_c = K_c \times ET_o$, as recommended by Allen *et al.* (1998). The local estimated K_c values for plastic mulched drip irrigated open-field eggplant used in this study were 0.60, 0.85, 1.10, and 0.95 during the four growth stages (30 days per stage) (Jordan valley authority, 2006) [Equation (2)]. These K_c values were used at both sites of experiments.

Soil Properties Measurements

Soil samples from three depths (0–15, 15–30, 30–60 cm) from the eggplant fields were collected in triplicate. The soil samples were dried and crushed, then sieved using 2 mm sieve. Soil texture was determined by the hydrometer test (Klute, 1986). Table 1 shows the main soil chemical properties analyzed according to the standard procedures of the United States Salinity Laboratory Staff (USSL, 1954). The soil moisture content was measured using Moisture Sensor-ML2x (Delta-T Devices Ltd) and profile time domain reflectometry PR2 (Delta-T Devices Ltd). The soil water potential was measured weekly before and after each irrigation event using one tensiometer at the depth of 20–30 cm (Soil Moisture

Equipment Corporation, Santa Barbara, Calif.) for each treatment. The soil moisture measurements were done using three probes for each treatment (one probe for each replicate) once per week. The PR2 was used to measure soil moisture at a 60 cm soil depth. Soil properties of the soil samples used in this study are presented in Table 1. The soil textures were sandy loam and clay loam in Ghor Al-Safi and Sail Al-Karak, respectively. The alkalinity (pH), soil Electrical Conductivity (EC), soil Nitrogen Content (N), and soil Phosphorous Content (P) at Ghor Al-Safi showed higher values than at the Sail Al-Karak site. The soil samples had EC values ranging from 2.46 to 6.34 and 1.51 to 2.22 dS m^{-1} at the Ghor Al-Safi and Sail Al-Karak sites, respectively. The EC , N, P, K, and OM tended to decrease with soil depth at both sites (Table 1).

Plant Properties Measurements

Two months after plant transplanting, leaf samples were collected at midday to analyze the chloroplast pigments, proline, Leaf Nitrogen Content (NL), Leaf Phosphorus Content (PL), and Leaf Potassium Content (KL) in triplicates. Leaf samples were stored in plastic bags with wet tissue paper at 4°C . The measurements were carried out on the leaves within two days of cutting to estimate total chlorophyll content (Inskeep and Bloom, 1985). Leaf proline content was

Table 1. Physical and chemical properties of soil at Ghor Al-Safi and Sail Al-Karak sites.

Soil sample/Depth (cm)	pH ^a (-)	EC ^b (dS m^{-1})	N ^c (%)	P ^d (ppm)	K ^e (ppm)	OM ^f (%)	Sand (%)	Silt (%)	Clay (%)	FC ^g (%)	PWP ^h (%)
Ghor Al-Safi											
0-15	8.29	6.31	1.45	445.62	15.08	1.12	62	16	17	34	22
15-30	8.37	3.50	1.38	433.42	14.54	1.01	64	18	18	34	23
30-60	8.53	2.36	1.43	435.32	8.25	0.39	65	15	20	35	23
Sail Al-Karak											
0-15	7.10	2.26	0.24	364.23	18.14	1.24	42	20	38	38	23
15-30	7.11	1.45	0.20	362.35	18.42	0.98	45	25	30	35	22
30-60	7.10	1.43	0.20	297.36	11.46	0.83	40	30	30	34	23

^a Alkalinity (-); ^b Electrical conductivity (dS m^{-1}); ^c Nitrogen content (%); ^d Phosphorus content (ppm); ^e Potassium content (ppm); ^f Organic matter (%); ^g Field capacity(%); ^h Permanent Wilting Point (%).

measured using Bates *et al.* (1973) method based on leaf dry weight (mg g^{-1}). Subsequently, each leaf sample was dried at 75°C for three days then ground. The ground leaf samples were used to determine the total NL, PL, and KL in the plant leaves. The PL was analyzed by a vanadate-molybdate method (Kirnak *et al.*, 2001) using a UV/visible spectrophotometer (Bausch and Lomb, Belgium). The KL was analyzed using a Flame photometer (Corning 400, UK) (Kirnak *et al.*, 2001). The NL was determined using the Kjeldahl method (Chapman and Pratt, 1982). Additional leaves were examined for Relative Water Content (RWC), Leaf Water Potential (ψ_w), and Leaf Area (LA) measurements for each treatment in triplicate at the same time (two months from plant transplanting). The RWC was calculated according to Barrs and Weatherley (1968) method. The leaf was weighed directly [Fresh Weight (FW)] after collection. Then, it was placed in a petri dish containing wet filter paper and stored at 4°C . After 24 hours, the leaf Turgid Weight (TW) was measured. Then, the leaves were oven-dried for 24 hours at 75°C and weighed [Dry Weight (DW)]. Midday Leaf Water Potential (ψ_w) was measured using a pressure chamber (PMS Instruments Co., Corvallis). The leaves were mature, similar in age, fully expanded, and exposed to solar radiation. The LA was measured using the photoelectric method (Cox, 1972).

Yield Quantity and Quality Analysis

The yield was harvested six times during the growing season. The growth season was four months at both sites. The yield and fruit weight were recorded for each treatment. Ten fruits were used to calculate the average fruit weight for each treatment. The total yield for each treatment was calculated by summing the total harvested yield during the experiment. Three fruits were used to measure Total Soluble Solids (TSS) in triplicates. Fruits were pressed to get juice, which was used to find out TSS with a

digital refractometer (Atago Co., Ltd, Tokyo, Japan). A 5 g sample of the same juice was diluted with 50 ml of deionized water. The diluted samples were titrated with 0.1N NaOH to determine titratable acidity. Five random fruit samples were selected from each harvest time and replicate. The five fruit samples were combined for each replicate. In total, there were fifteen fruit samples for each treatment. Each combined sample was dried at 75°C for one week. Finally, fruit samples were ground. The powdered samples were used to determine Fruit Nitrogen Content (NF), Fruit Phosphorus Content (PF), and Fruit Potassium Content (KF). The analysis procedures were the same as the leaf analysis methods. Plant Water Use Efficiency (WUE) was calculated as the total harvested yield divided by the amount of irrigation water applied during the growth period (Mohawesh and Karajeh, 2014).

Statistical analysis

Data were analyzed using a General Linear Model (GLM) procedure (SPSS software version 11.5; SPSS Inc., Chicago, USA). A Least Significance Difference (LSD) test was used for mean separation at a level of significance of 0.05. Mean values of leaf chlorophyll content, proline, NL, PL, KL, RWC, ψ_w , LA, yield weight, fruit weight, TSS, titratable acidity, NF, PF, KF, and WUE were compared between five irrigation levels (100, 80, 60, 40, and 20% based on FC).

RESULTS AND DISCUSSION

Description of Soil Properties

The soil water content and soil water potential variations during the growth period are shown, for example, in Figure 2 for Sail Al-Karak site. The mean soil water contents were about 23.0, 26.0, 29.0, 31.0, and 33.0% at 10–15 cm soil depth measured using




Figure 2. Soil moisture content and soil water potential during the experiment at Sail Al-Karak sites.

Moisture Sensor-ML2x (Delta-T Devices Ltd) for 20, 40, 60, 80, and 100% irrigation levels at Sail Al-Karak site, respectively. The soil water potential decreased with decreasing irrigation water (Figure 2). The soil water potential values were less than -100, -60 to less than -80, -35 to -50, -20 to -35 and -10 to -20 kPa at 20–30 cm depth for 20, 40, 60, 80, and 100% irrigation levels at Sail Al-Karak site, respectively (Figure 2).

Plant Properties Measurements and Analysis

Table 2 shows the effect of irrigation levels on *LA*, *RWC*, ψ_w , proline, chlorophyll, *NL*, *PL*, and *KL* content. The differences were not significant between the 100% irrigation level and the lower level of water stress (80 and 60%) except for *LA* at Ghor

Table 2. Effect of deficit irrigation regimes on eggplant leaf area, relative water content, plant water potential, proline, chlorophyll, nitrogen, phosphorus and potassium content at Ghor Al-Safi and Sail Al-Karak sites.

Irrigation regime (%)	LA ^a (cm ²)	RWC ^b (%)	ψ_w ^c (MPa)	Proline (ppm)	Chlorophyll (ppm)	NL ^d (%)	PL ^e (ppm)	KL ^f (ppm)
Ghor Al-Safi								
100	60.22a	78.33a	0.250b	10.23b	2386.85a	4.83a	341a	7730a
80	44.49b	76.67a	0.250b	9.38b	2486.21a	4.59a	342a	7553a
60	33.40c	72.77ab	0.333ab	10.89ab	2312.67a	4.67a	349a	6553ab
40	31.19c	72.43b	0.433a	16.16a	2196.57b	4.63a	322b	6506b
20	26.19c	72.64b	1.133a	13.96a	1945.47b	3.70b	325b	6096b
Sail Al-Karak								
100	193.36a	86.30a	0.83c	6.90b	1416.57a	4.41a	1062a	4143a
80	184.15a	83.01a	1.07bc	9.39b	1329.74ab	4.56a	365b	4080a
60	130.57ab	81.91ab	1.13b	10.89ab	1198.00bc	4.42ab	365b	4313a
40	109.48b	81.67b	1.47a	16.16ab	1064.51cd	4.21ab	185c	4253a
20	106.77b	80.68b	1.57a	20.63a	923.70d	3.91b	263c	4040b

^a Leaf Area (cm²); ^b Relative Water Content (%); ^c Plant Water Potential (bar); ^d Leaf Nitrogen Content (%); ^e Leaf Phosphorus Content (ppm), ^f Leaf Potassium Content (ppm). Means within columns followed by the same letters are not significantly different at 0.05 probability level using LSD test.

Al-Safi. However, high levels of water stress (40 and 20%) showed a significant effect on LA, RWC, ψ_w , proline, chlorophyll, NL, PL, and KL content comparing to control treatment (100%) (Table 2). The water deficit reduced the growth of eggplant components. The LA was reduced from 60.22 and 193.36 cm² for 100% irrigation level to 26.19 and 106.77 cm² for 20% irrigation level at Ghor Al-Safi and Sail Al-Karak, respectively. The results of LA obtained at the two sites were different because plant growth depends on the season, climatological conditions, and many other factors including soil fertility and salinity. This is in agreement with Mohawesh and Karajeh (2014) who showed a decrease of LA under water deficit. The highest RWC values were for a lower level of water stress (100 and 80%). The decrease in plant RWC under drought stress may possibly depend on plant vigor reduction (Liu *et al.*, 2002). The RWC decreased from 78.33 and 86.30 for 100% irrigation level to 72.64 and 80.68% for the 20% irrigation level at Ghor Al-Safi and Sail Al-Karak, respectively. These results are in agreement with Mohawesh and Karajeh (2014) who found

that leaf RWC decreased with deficit irrigation treatments under greenhouse conditions. Plants at different irrigation levels showed obvious variations in ψ_w . The ψ_w decreased with decreasing irrigation quantity. The 100% and 80% irrigation regimes showed higher ψ_w than those of the stressed irrigation levels. The ψ_w was -0.25, -0.25, -0.333, -0.433 and -1.133 MPa at Ghor Al-Safi site and -0.83, -1.07, -1.13, -1.47 and -1.57 MPa at Sail Al-Karak site for 100, 80, 60, 40, and 20% irrigation levels, respectively. Our results are in agreement with Javadi *et al.* (2008) who also found a similar decrease of ψ_w as a result of water deficit. Similar observations were also reported for eggplant during and after repetitive water stress (Sarker *et al.*, 2005).

Biochemical parameters showed an increase in proline and a decrease in chlorophyll content under water deficit. Proline concentration in leaves increased significantly with increasing water deficit. The proline concentration increased by 36.46 and 198.98% for the 20% irrigation level at Ghor Al-Safi and Sail Al-Karak, respectively (Table 2). Water deficit induced an increase in proline synthesis with

**Table 3.** Effect of deficit irrigation regimes on eggplant yield, fruit weight, total soluble solids, nitrogen, phosphorus and potassium content at Ghor Al-Safi and Sail Al-Karak sites.

Irrigation regime (%)	Yield (Kg ha ⁻¹)	Fruit Wt. (Kg)	TSS ^a (%)	Acidity (%)	NF ^b (%)	PF ^c (ppm)	KF ^d (ppm)	WUE ^e (Kg m ⁻³)
Ghor Al-Safi								
100	27840a	0.63a	5.33b	0.1117c	3.38a	389a	7017a	5.13c
80	26020a	0.50ab	6.33b	0.1340a	3.20a	357a	6935a	6.00c
60	16687b	0.36b	7.00a	0.1787a	3.10a	377a	7017a	7.58b
40	11138c	0.22c	7.00a	0.1117c	3.01ab	358a	5704ab	7.13b
20	10057c	0.20c	6.67a	0.1227b	2.24b	317b	4718b	8.49a
Sail Al-Karak								
100	28290a	0.32a	3.00c	0.12b	3.56a	1930ab	5760a	2.78c
80	21000a	0.29ab	3.65bc	0.14b	3.25a	1755a	5300b	3.56c
60	15600b	0.22b	4.00b	0.16b	3.17ab	1611ab	4445c	3.85c
40	12540b	0.14c	4.33b	0.23a	3.12b	1493c	3870c	3.07b
20	9630c	0.10c	5.00a	0.24a	2.58c	1438c	3865c	4.73a

^a Total Soluble Solids (%); ^b Fruit Nitrogen Content (%); ^c Fruit Phosphorus Content (ppm), ^d Fruit Potassium Content (ppm). Means within columns followed by the same letters are not significantly different at 0.05 probability level using *LSD* test.

decreasing irrigation level. An increase in leaf proline content with an increase in water deficit levels was also reported in several studies (Yoshiba *et al.*, 1997; Heuer and Nadler, 1998). Chlorophyll content decreased as proline content increased. Water deficit significantly reduced leaf nutrient (NL, PL, and KL) content (Table 2). The control plants showed higher nutrient contents than the water-stressed plants. The highest significant content was found at 100% and the lowest was at the 20% irrigation level. For example, the NL concentration was 4.83 and 4.4% for the 100% irrigation level while it was 3.70 and 3.91% for the 20% irrigation level at the Ghor Al-Safi and Sail Al-Karak sites, respectively. Bharambe and Joshi (1993) and Honda (1971) found that the plant uptake of N, P, and K was adversely affected under water deficit.

Yield Quantity and Quality Analysis

Table 3 shows the effect of DI regimes on yield, fruit weight, TSS, titratable acidity, nutrient content (NF, PF, and KF), and *WUE*. A significant reduction occurred

between 100, and 80%, and 60, 40, and 20% treatments at both sites. Eggplant yield weight for the 100% irrigation level was 27,840 and 28,290 kg ha⁻¹ and for the 20% irrigation level was 10,057 and 9,630 kg ha⁻¹ at the Ghor AL-Safi and Sail Al-Karak sites, respectively. These results are in agreement with Ebrahim *et al.* (2012), Aziz *et al.* (2013) and Demirel *et al.* (2014) who reported a similar decrease in eggplant yield under water deficit. Fruit weight also decreased significantly with water deficit.

The eggplant yield decreased with decreasing irrigation level (Figure 3). While the highest yield was obtained from the control treatment, the 80% treatment was in the same class in terms of yield values. The lowest yield was obtained from 20% irrigation level to which the lowest irrigation was applied. The relationship between the yield loss and irrigation water savings is shown in Figure 3. It shows a linear relationship between irrigation water savings and yield loss for the eggplant. Higher water savings result in high yield losses compared with control. Figure 3 also shows a similar response of eggplant yield to water stress at both sites.




Figure 3. Yield loss with irrigation water amount saving at Ghor Al-Safi and Sail Al-Karak sites.

For fruit weight trait, the highest irrigation level treatment (100%) gave heavier fruits (630 and 320 g) in comparison with fruit weight values (200 and 100 g) of higher water stress (20%) treatment at Ghor Al-Safi and Sail Al-Karak sites, respectively (Table 3). These results were in harmony with Abd El-Aal *et al.* (2008) who found that fruit weight was significantly affected by DI regimes.

On the contrary, fruit quality traits (TSS and titratable acidity) increased as irrigation water decreased from the 100 to 20% level at both sites. There was no significant difference between the control (100%) and low water deficits (80%), however, significant differences were found between the control and the high water deficits (60, 40, and 20%). The TSS and titratable acidity increased significantly as the amount of irrigation water decreased from the 100% irrigation level to the 20% irrigation level at Ghor Al-Safi and Sail Al-Karak (Table 3). This is in agreement with Kirnak *et al.* (2002) who investigated the effects of DI on fruit yield and the quality of eggplant. They reported that TSS and titratable acidity increased with decreasing irrigation amounts from the 100 to the 20% irrigation level.

The fruit nutrient content of NF, PF, and KF decreased with water deficit. The highest significant differences were found between

the 100% irrigation regime and the two lowest irrigation regimes (20 and 40%). The KF was 7,017 and 5,760 ppm for 100%, while it was 4,718 and 3,865 ppm for 20% irrigation treatments at the Ghor Al-Safi and Sail Al-Karak sites, respectively. These results are consistent with the findings of Simonne *et al.* (1998) and Kirnak *et al.* (2002) who reported for several vegetable crops that water deficit has a main function in decreasing fruit nutrient content. The least *WUE* was 5.13 and 2.78 kg m⁻³ at 100% irrigation level while the highest value was for irrigation at 20% with 18.49 and 4.73 Kg m⁻³ at Ghor Al-Safi and Sail Al-Karak, respectively. Amiri *et al.* (2012) showed the same trend as the highest *WUE* was obtained with the lowest irrigation treatment which was under no irrigation treatment.

CONCLUSIONS

DI can be used to improve crop productivity per water unit to a certain extent. DI regimes affected plant physiological and agronomic parameters of eggplant. A significant reduction in plant growth performance occurred at 40 and 20% treatments at both sites as compared to the control. The *WUE* also decreased with increasing irrigation



amount. The effect of low DI regimes (80%) was not significant compared to control, while high DI levels (60, 40, and 20%) significantly affected plant growth performance as compared to the control. A linear relationship between irrigation water savings and the yield loss for the eggplant was found. Higher water savings result in high yield losses compared with control. A similar response of eggplant yield to water stress was found at both sites. Based on our results, a DI level at 80% can be utilized to increase *WUE* without a significant effect on crop growth performance.

ACKNOWLEDGEMENTS

This work is a part of the research project financially supported by the Scientific Research Support Fund of Ministry of Higher Education and Scientific Research/Jordan.

REFERENCES

1. Abd El-Aal, F. S., Abdel Mouty, M. M. and Ali, A. H. 2008. Combined Effect of Irrigation Intervals and Foliar Application of Some Antitranspirants on Eggplant Growth, Fruits Yield and Its Physical and Chemical Properties. *J. Agric. Biol. Sci.*, 4(5): 416-423.
2. Allen, R. G., Pereira, L. S., Raes, D. and Smith, M. 1998. Crop Evapotranspiration: Guidelines for Computing Crop Requirements: Irrig. Drain. Paper No. 56. FAO, Rome, Italy.
3. Amiri, E., Abdzad Gohari, A. and Esmailian, Y. 2012. Effect of Irrigation and Nitrogen on Yield, Yield Components and Water Use Efficiency of Eggplant. *J. Biotechnol.*, **11(13)**: 3070-3079.
4. Aziz, S. A., Ahmad, S. T., Rasheed, B. A., Sharef, H. B. and Talb, D. H. 2013. The Impacts of Deficit Irrigation Levels and Intervals on Tomatoes and Eggplants Yield in Unconditions Plastic House. *J. Zankoy Sulaimani Part A*, **15(2)**: 125-135.
5. Barrs, H. D. and Weatherley, P. E. 1962. A Re-examination of the Relative Turgidity Technique for Estimating Water Deficits in Leaves. *Aust. J. Biol. Sci.*, **15**: 413-28.
6. Bates, L. S., Waldren, R. P. and Teare, I. D. 1973. Rapid Determination of Free Proline for Water Stress Studies. *Plant Soil*, **39**: 205-207.
7. Bharambe, P. R. and Joshi, P. S. 1993. Effect of Soil Water Potential on Growth, Yield and Some Biochemical Changes in Sorghum. *J. Indian Soc. Soil Sci.*, **41(2)**: 342-343.
8. Chapman, H. D. and Pratt, P. F. 1982. *Method and of Analysis of Soil, Plant and Water*. 2nd Edition, California University Agricultural Division, California, 170 PP.
9. Correia, F. N. 1999. Water Resources in Mediterranean Region. *Water Int.*, **24(1)**: 22-30.
10. Cox, E. F. 1972. A Photoelectric Digital Scanner for Measuring Leaf Area. *New Phytol.*, **71 (5)**: 819-823.
11. Demirel, K., Genc, L., Bahar, E., Inalpulat, M., Smith, S. and Kizil, U. 2014. Yield Estimate Using Spectral Indices in Eggplant and Bell Pepper Grown Under Deficit Irrigation. *Fresen. Environ. Bull.*, **23 (5)**: 1232-1237.
12. Ebrahim, A., Gohari, A. A. and Esmailian, Y. 2012. Effect of Irrigation and Nitrogen on Yield, Yield Components and Water Use Efficiency of Eggplant. *Afr. J. Biotechnol.*, **11(13)**: 3070-3079.
13. Gleick, P. H. 2000. *The World's Water 2000-2001*. Island Press, Washington, DC.
14. Heuer, B. and Nadler, A. 1998. Physiological Response of Potato Plants to Soil Salinity and Water Deficit. *Plant Sci.*, **137**: 43-51.
15. Honda, N. 1971. Influence of Soil Moisture on Growth and Leaf Properties of Tobacco. III. On the Absorption of Mineral Nutrients. *Bull. Okayama Prefectural Agric. Exp. Station*, **30**: 37-41.
16. Inskeep, W. P. and Bloom, P. R. 1985. Extinction Coefficients of Chlorophyll a and b in N, N-dimethylformamide and 80% Acetone. *Plant Physiol.*, **77**: 483-485.
17. Javadi, T., Arzani, K. and Ebrahimzadeh, H. 2008. Study of Proline, Soluble Sugar, and Chlorophyll a and b Changes in nine Asian and One European Pear Cultivar under Drought Stress. *Acta Hortic.*, **769**: 241-246.
18. Jordan Valley Authority (JVA). 2006. Reclaimed Water Irrigation in the Jordan Valley, Practical Recommendations for

- Farmers and Extension Workers. GTZ Project, Jordan.
19. Kirnak, H., Kaya, G., Tas, I. and Higgs, D. 2001. The Influence of Water Deficit on Vegetative Growth, Physiology, Fruit Yield and Quality in Eggplant. *Bulg. J. Plant Physiol.*, **27(3-4)**: 34-46.
 20. Kirnak, H., Tas, I., Kaya, C. and Higgs, D. 2002. Effects of Deficit Irrigation on Growth, Yield, and Fruit Quality of Eggplant under Semi-arid Conditions. *Aust. J. Agric. Res.*, **53**: 1367-1373.
 21. Klute, A. 1986. Methods of Soil Analysis. Part I. Physical and Mineralogical Properties. 2nd Edition, Agronomy, American Society of Agronomy and Soil Science of America, Madison, WI, **9**.
 22. Liu, Y., Fiskum, G. and Schubert, D. 2002. Generation of Reactive Oxygen Species by Mitochondrial Electron Transport Chain. *J. Neurochem.*, **80**: 780-787.
 23. Mahadeen, A., Mohawesh, O., Al-Absi, K. and Al-Shareef, W. 2011. Effect of Irrigation Regimes and Frequency on Water Use Efficiency and Tomato Fruit (*Lycopersicon esculentum* Mill.) Grown under an Arid Environment. *Arch. Agron. Soil Sci.*, **57**: 105-114.
 24. Mohawesh, O. and Al-Absi, K. 2009. Physiological Response of Two Apple Genotypes to Different Water Regimes under Semiarid Conditions. *Adv. Hortic. Sci.*, **23(3)**: 158-165.
 25. Mohawesh, O., Al-Absi, K. and Tadros, M. 2010. Effect of Antitranspirant Application on Physiological and Biochemical Parameters of Three Orange Cultivars Grown under Progressive Water Deficit. *Adv. Hortic. Sci.*, **24(3)**: 183-194.
 26. Mohawesh, O. and Talazi, S. 2012. Comparison of Hargreaves and FAO56 Equations for Estimating Monthly Evapotranspiration for Semiarid Environment. *Arch. Agron. Soil Sci.*, **58(3)**: 321-334.
 27. Mohawesh, O. and Karajeh, M. 2014. Effects of Deficit Irrigation on Tomato and Eggplant and Their Infection with the Root-knot Nematode under Controlled Environmental Conditions. *Arch. Agron. Soil Sci.*, **60**: 1019-1102.
 28. Sarker, B. C., Hara, M. and Uemura, M. 2005. Proline Synthesis, Physiological Responses and Biomass Yield of Eggplants during and after Repetitive Soil Moisture Stress. *Scientia Hortic.*, **103**: 387-402.
 29. Shatanawi, M., Duwayri, M., Haddad, N. and Naber, S. 2007. Analysis of Water, Agriculture and Food Security Policies in Jordan. *Jordan J. Agric. Sci.*, **2**: 270-279.
 30. Simonne, E. H., Joseph, D. E. and Harris, C. E. 1998. Effects of Irrigation and Nitrogen Rates on Foliar Mineral Composition of Bell Pepper. *J. Plant Nutr.*, **21**: 2545-2555.
 31. SPSS Software Products. 1997. *SigmaStat Version 2.0 for Windows*. SPSS Inc., Chicago, IL, USA.
 32. US Salinity Laboratory Staff. 1954. *Diagnosis and Improvement of Saline and Alkali Soils*. US Department of Agricultural Hand Book 60, US Department of Agricultural Soils, Washington, DC.
 33. Yoshiba, Y., Kiyosue, T., Nakashima, K., Yamaguchi-Shinozaki, K. and Shinozaki, K. 1997. Regulation of Levels of Proline as an Osmolyte in Plants under Water Stress. *Plant Cell Physiol.*, **38**: 1095-1102.



استفاده از کم آبیاری به منظور افزایش عملکرد رشد و کارایی مصرف آب گیاه بادمجان در محیط های خشک

۱. موحوش

چکیده

هدف از این پژوهش، بررسی اثرات کم آبیاری بر ویژگی های فیزیولوژیکی و زراعی بادمجان به منظور به حداکثر رساندن راندمان مصرف آب (WUE) بدون تاثیر منفی بر پارامترهای عملکرد و کیفیت میوه نهایی در محیط مناطق خشک بود. بنابراین، دو آزمایش میدانی در دو محل مختلف: Ghor Al-Safi در دره اردن، اردن و Sail Al-Karak در دره کراک، استان کراک، اردن با استفاده از یک رقم بادمجان مشترک (کلاسیک) در پنج سطح آبیاری: ۲۰، ۴۰، ۶۰، ۸۰ و ۱۰۰٪ ظرفیت زراعی انجام شد. شدیدترین تیمارهای کم آبیاری (DI) (۴۰٪ و ۲۰٪) منجر به اثرات قابل توجهی در سطح برگ، محتوای نسبی آب برگ، پتانسیل آب برگ و محتوای مواد معدنی برگ شدند. همچنین پارامترهای بیوشیمیایی افزایش پرولین و کاهش در میزان کلروفیل در شرایط کمبود آب را نشان دادند. وزن میوه و عملکرد کل با DI کاهش یافت. گیاهان تیمار شاهد (آبیاری ۱۰۰٪) محتویات مواد مغذی بالاتری نسبت به گیاهان قرار گرفته در معرض تنش آب را نشان داد. TSS و اسیدیته قابل تیتراسیون میوه در هر دو سایت با تغییر رژیم آبیاری از ۱۰۰٪ به ۲۰٪ افزایش یافت. محتوای مواد مغذی میوه با افزایش کمبود آب کاهش یافت. با این حال، تفاوت ها بین شاهد (تیمار آبیاری ۱۰۰٪) و تیمار آبیاری ۸۰٪ معنی دار نبود. تیمار ۸۰٪ راندمان مصرف آب بالا با اثرات نسبتا اندکی بر عملکرد رشد گیاه در مقایسه با شاهد نشان داد. در نتیجه، سطح DI ۸۰٪ می تواند برای افزایش کارایی مصرف آب بدون اثر قابل توجهی بر عملکرد رشد محصول مورد استفاده قرار گیرد.