

RESEARCH NOTES

Sensitivity of the Nematophagous Fungus *Arthrobotrys oligospora* to Fungicides, Insecticides and Crop Supplements Used in the Commercial Cultivation of *Agaricus bisporus*

E. Mohammadi Goltapeh^{1*}, M. Shams-bakhsh¹ and B. S. Pakdaman²

ABSTRACT

The effect of various pesticides (diflubenzuro, malathion, mancozeb and carbendazim), disinfectants (calcium hypochlorite and formaldehyde) and oil cakes (sunflower and soybean oil cakes) commonly used as supplements in mushroom cultivation on the growth of the nematophagous fungus, *Arthrobotrys oligospora*, was studied under *in vitro* conditions. Carbenazim caused 99% inhibition of radial mycelial growth in Petri dishes at all concentrations tested (10-40 µg a. i. ml⁻¹) in comparison to non treated dishes. Mancozeb caused 43% and 23% inhibition at 250 and 500 µg a. i. ml⁻¹ respectively and 99% inhibition at concentration of 1000 µg a. i. ml⁻¹ and above. Diflubenzuro and malathion at 10-40 µg a. i. ml⁻¹ caused 30-41% and 24-54% inhibition, respectively. Formalin (0.5-2.0% v/v) inhibited growth of *A. oligospora* completely. However, calcium hypochlorite, sunflower and soybean oil cake at concentrations of up to 2.0% w/v caused less than 3.5% inhibition.

Keywords: *Agaricus bisporus*, *Arthrobotrys oligospora*, Biocontrol, Disinfectant and Oil cakes.

INTRODUCTION

White button mushroom [*Agaricus bisporus* (Lange) Singer], an edible fleshy fungus, is a commercially cultivated crop with a total world production of more than two million metric tonnes (Chang and Miles, 2004; Mohammadi Goltapeh and Pourjam, 2005). The annual production in Iran is approximately 30,000 metric tonnes (Mohammadi Goltapeh and Pourjam, 2005). Mushroom cultivation is adversely affected by many pests and diseases caused

by various organisms, namely bacteria, fungi, viruses, insects and nematodes. Among all these natural enemies, nematodes are distinguished as the most destructive and harmful ones and, once introduced, they become the main limiting factors in the successful cultivation of the mushroom crop (Cayrol, 1967). Several cases have been reported where nematodes have limited mushroom productivity or resulted in total crop failures (Sharma *et al.*, 1984; Garcha *et al.*, 1986; Mohammadi Goltapeh and Rezaei Danesh, 2006; Sharma *et al.*,

¹ Department of Plant Pathology, College of Agriculture, Tarbiat Modares University, Tehran, Islamic Republic of Iran.

² Department of Microorganisms, Bio-resources and Biosafety, Agricultural Biotechnology Research Institute of Iran, Karaj, Islamic Republic of Iran.

* Corresponding author, email: emgoltapeh@modares.ac.ir



2002; Staunton, 1996; Fletcher *et al.*, 1986; Heydari *et al.*, 2006).

There is increasing awareness about the harmful effects of pesticides to human health, non target organisms and the environment. Furthermore the efficacy of the pesticides can be compromised by the development of resistance in pests. Pesticides can also lead to residue problems in food, especially in mushrooms where they have been shown to be translocated to the mushrooms from treated composts (Bahl and Agnihorti, 1989). There is therefore a need to reduce pesticide inputs and develop an alternative strategies, such as cultural or biological control, such as with nematophagous fungi. *A. oligospora* is the most frequently associated nematophagous fungus encountered in the casing soils used in the commercial production of white button mushroom (Grewal, 1993). Species of *Arthrobotrys* have been commercially developed for the control of *Ditylenchus myceliophagus* in mushroom cultivation in France (Cayrol and Frankowski, 1979). *Arthrobotrys* forms adhesive hyphal networks to catch and kill the nematodes and can parasitize various nematodes parasitic on the mushroom crop, namely *D. myceliophagus*, *Aphelanchoides composticola*, *A. bicaudatus*, *Aphelenchus avenae*, as well as saprophagous species like *Rhabditis* and entomoparasitic nematodes of the genus *Steinernema* (Grewal, 1993). Therefore, the fungus may have potential for incorporation into integrated control programs developed for mushroom cultivation. *A. oligospora* produces a range of nematocidal compounds, including linoleic acid (Anke *et al.*, 1995) and oligosporons (4', 5'-dihydro-oligosporon, hydroxyoligosporon, and 10', 11'-epoxyoligosporon) (Anderson *et al.*, 1995). However, for it to be successfully integrated into mushroom cultivation, there is a need to assess the compatibility of the nematophagous fungus *A. oligospora* with the various pesticides, disinfectants and crop supplements commonly used in a commercial mushroom production.

MATERIALS AND METHODS

The sensitivity of *A. oligospora* was tested *in vitro* to the insecticides diflubenzuron (Dimlin 25% WP) and malathion (Malathion 50% EC), the fungicides carbendazim (Bavistin 50% WP) and mancozeb (Indofil M-45) and the disinfectants formaldehyde (Formalin 37%) and bleaching powder (calcium hypochlorite). In addition, *A. oligospora* was tested against sunflower cake and soybean cake which are commonly used for compost supplement in mushroom crop production. Carbendazim was tested at 10, 20, 30 and 40 $\mu\text{g a. i. ml}^{-1}$, Indofil M-45 at 500, 1,000 and 2,000 $\mu\text{g a. i. ml}^{-1}$, diflubenzuron at 10, 20, 30 and 40 $\mu\text{g a. i. ml}^{-1}$ and Malathion at 10, 20, 30 and 40 $\mu\text{g a. i. ml}^{-1}$. Stock solutions of the chemicals were prepared in sterilized distilled water and volumes added under aseptic conditions into 250 ml flasks, containing half-strength CMA (corn meal agar), so as to achieve the required final concentrations. Formalin was tested at 1, 2, 3 and 4% v/v and calcium hypochlorite at 0.5, 1, 1.5 and 2% w/v, and the required amount of the chemical was directly added to the medium (CMA) in the flasks. For the sunflower and soybean oil cakes, either 0.5, 1, 1.5 and 2.0g of the crushed cake was added to 100 ml media (corn meal broth), and this was then churned in a waring blender for 2-3 minutes. Agar was added to the churned media and autoclaved at 15 psi for 45 minutes. The amended media were poured into 9 cm sterilized Petri dishes, under aseptic conditions and allowed to cool. Plates were inoculated in the center with a 5 mm mycelial disc, taken from the edge of an actively growing three day-old *A. oligospora* culture. Petri dishes containing non-amended medium served as the control (check). For each concentration and treatment there were three replicate Petri dishes. The inoculated Petri dishes were incubated at $25\pm 1^\circ\text{C}$. The colony diameter was recorded daily until colonies had grown to the edge of the Petri dish. The diameter of growth zone was measured in treated and untreated cultures, and inhibi-

tion percentages were recorded daily (every 24 hours) by comparing the growth rates according to the following equation (Vincent, 1947):

Percentage of fungal growth inhibition = $(C-T)/C \times 100$

Where: C = Growth of the fungus in control and T = Growth of the fungus in treatment

Data were subjected to transformation ($\arcsin+0.1$) and analysed by analysis of variance using MSTATC, SAS and means separated by Duncan's multiple range test.

RESULTS AND DISCUSSION

Effect of Pesticides on the Growth of *A. oligospora*

At all concentrations tested, the pesticides diflubenzuron, malathion, mancozeb and carbendazim caused significant ($P = 0.05$) growth inhibition of *A. oligospora* in comparison to the control (Table 1). Carbendazim at all the concentrations tested ($10-40 \mu\text{g a. i. ml}^{-1}$) caused almost complete inhibition of fungal growth, with significantly ($P = 0.05$) greater inhibition than all other pesticides except for mancozeb at 1,000 and 2,000 $\mu\text{g a. i. ml}^{-1}$. Diflubenzuron and malathion caused the maximum inhibition of 41.2% and 54.1%, respectively at the highest concentration tested ($40 \mu\text{g a. i. ml}^{-1}$) and mancozeb caused 49.4% inhibition at the lowest concentration tested ($250 \mu\text{g a. i. ml}^{-1}$). These results suggest that low concentrations of carbendazim ($10 \mu\text{g a. i. ml}^{-1}$) are almost completely inhibitory to *A. oligospora*, as are higher concentrations of mancozeb ($>1000 \mu\text{g a. i. ml}^{-1}$). However, concentrations of diflubenzuron and malathion of $40 \mu\text{g a. i. ml}^{-1}$ caused only moderate inhibition of radial growth ($<54\%$).

The nematocidal effect of some insecticides has been known, and used practically in agriculture. For example, Diazinon is listed as a means of foliar nematode control in greenhouse florist crops (LaMondia, 2004). Furthermore, it has been shown that some fungicides, like Maneb, and Mancozeb

are toxic to certain nematode species, such as *Caenorhabditis elegans*, whereas their potentially toxic metabolite, ethylenethiourea (ETU) had of minimal anti-nematode toxicity only at high concentrations (Easton *et al.*, 2001). Additionally, there is some information in the literature indicating the anti-fungal activity of some insecticides, such as Thiodan, Diazinon and Basudin that inhibit mycelial growth of the white button mushroom; however, others such as Dimilin, and Trigard do not show any mycotoxic effect on mycelial growth of the mushroom (Dmoch *et al.*, 1989). Here, we have indicated the antifungal activity of two insecticides, namely Malathion and Diflubenzuron, of which the former showed a greater inhibitory effect against growth of the biocontrol agent, *A. oligospora*. Dimilin (Benzuran) is believed to inhibit chitin synthase activity through deactivation of its activator proteases in insects, and the same mechanism of action may also be true for protease (B) in fungi. Also, inhibition of DNA synthesis has been attributed to this insecticide that may also be true for fungi (Matsumura, 1985).

Grewal and Sohi (1988) evaluated the effect of commonly used pesticides on the mycelial growth of *Arthrobotrys conoides* and *A. oligospora*, and found that most of the pesticides inhibited the mycelial growth of both of the fungi.

Pullen *et al.* (1990) reported that benomyl at the rates of 5 and $10 \mu\text{g a. i. ml}^{-1}$ completely inhibited hyphal growth of the nematophagous fungus *Hirsutella rhossiliensis*; similarly, Chauhan *et al.* (2002) reported 80 percent growth inhibition with *A. musiformis* by carbendazim at $2.0 \mu\text{g ml}^{-1}$ on water agar.

Based on the findings given in Table 1, it seems that the application of diflubenzuron instead of malathion will lead to improved survival of the *A. oligospora* populations, although the population of other fungi, including mushrooms, and parasitic fungi may also increase in the absence of strong chemical growth inhibitors. Also, considering the antifungal impact imposed by the tested insecticides, the application of slightly higher doses of these insecticides can lead to eco-

**Table 1.** Effect of various pesticides on the growth of *Arthrobotrys oligospora* on corn meal agar at 10 days after inoculation.

Pesticides	Concentration ($\mu\text{g ml}^{-1}$)	Radial growth (mm)	Growth inhibition (%)
Diflubenzuron	10	59	30.6(0.60) h
	20	57	32.9 (0.62)g
	30	55	35.3(0.63)f
	40	50	41.2(0.67)e
Malathion	10	65	23.5 (0.55)i
	20	55	35.3(0.63)f
	30	43	49.4 (0.71)d
	40	39	54.1 (0.73)c
Mancozeb	250	43	49.4 (0.71)d
	500	23	72.9 (0.82)b
	1000	1	98.8 (0.91)a
	2000	1	98.8 (0.91)a
Carbendazim	10	1	98.8 (0.91)a
	20	1	98.8 (0.91)a
	30	1	98.8 (0.91)a
	40	1	98.8 (0.91)a
Control	-	85	0.0 (0.31)j

Within column, mean followed by the same letter are not significant at $P=0.05$ (Duncan's multiple range test).

Values in parenthesis are arc sin transformed values after adding 0.1 in all values.

Growth inhibition percentage= $(C-T)/C \times 100$.

nomical control of fungal parasites, in addition to the control of other target pests of importance in third world countries where farmers do not have access to most agro-chemicals. However, many more studies are needed in relation to the effects of higher doses on human health and the environment. It can be concluded that the first application of a low doses of malathion followed by the inoculation with *A. oligospora* propagules after some days can lead to the suppression of the parasitic nematodes.

Effects of Disinfectants and Oil Cakes on the Growth of *Arthrobotrys oligospora*

Calcium hypochlorite, sunflower oil cake and soybean oil cake caused little inhibition in the growth of *A. oligospora* at all the three concentrations (Table 2) without showing any significant difference among them.

However, formalin completely inhibited the growth of the fungus at all the four rates tested. The interactions between disinfectant/oil cakes and their concentrations were not significant.

Kerry and Crump (1998) reported that soil drenching with formalin reduced significantly the population of the nematophagous fungi *Verticilium chlamydosporium* and *Nematophthora gynophila*.

The disinfectant, calcium hypochlorite was found compatible with *A. oligospora* and oil cakes did not inhibit the growth of *A. oligospora* compared to non-amended controls. The reduction of fungal growth through soil amendment with oil cakes (groundnut, mustard, sesamum, and binola/cotton seed) has been reported with some fungi such as *Fusarium oxysporum* f. sp. *lycopersici*, where a positive correlation is found between the magnitude of fungal growth inhibition and

Table 2. Effect of disinfectants and oil cakes on the growth of *Arthrobotrys oligospora* at 3 days after inoculation.

	Concentration (%)	Radial growth (mm)	Growth inhibition (%)
Calcium hypochloride	0.5	85	0.0 (0.31)c
	1.0	85	0.0 (0.31)c
	1.5	83	2.35(0.34)b
	2.0	83	2.35(0.34)b
Formalin 37%	0.5	1	98.82(0.91)a
	1.0	1	98.82(0.91)a
	1.5	1	98.82(0.91)a
	2.0	1	98.82(0.91)a
Sunflower oil cake	0.5	85	0.00(0.31)c
	1.0	85	0.00(0.31)c
	1.5	83	2.35(0.34)b
	2.0	82	3.52(0.36)b
Soybean oil cake	0.5	85	0.00(0.31)c
	1.0	85	0.00(0.31)c
	1.5	83	2.35(0.34)b
	2.0	82	3.52(0.36)b
Control		85	0.00(0.31)c

Within column, mean followed by the same letter are not significant at $p = 0.05$ (Duncan's Multiple Range Test).

Values in parenthesis are arc sin transformed values after adding 0.1 in all values.

Growth inhibition percentage = $(C-T) / C \times 100$.

the increase in decomposition over a 45 day period (Raj and Kapoor, 1996). Among the four oil cakes, groundnut and mustard at 2.0% concentration of soil (w/w) were the most effective in reducing pathogen population ($> 70\%$) and disease incidence. However, groundnut seed was found superior to mustard seed, as it not only controlled the disease better, but also improved plant growth (Raj and Kapoor, 1996). Similar effects have been reported with other filamentous pathogenic fungi, such as *Thielaviopsis basicola*, *Rhizoctonia solani*, and *Sclerotinia sclerotiorum* (Papavizas *et al.*, 1970; Singh, 1968).

REFERENCES

1. Anderson, M.G., Jarman, T. B. and Rickards, R.W. 1995. Structures and Absolute Configurations of Antibiotics of Oligosporon Group from Nematode-trapping Fungus *Arthrobotrys oligospora*. *J. Antibiot.*, **48**: 391-398.
2. Anke, N., Stadler, M., Mayer, A. and Sterner, O. 1995. Secondary Metabolites with Nematicidal and Antimicrobial Activity from Nematophagous Fungi and Ascomycetes. *Can. J. Bot.*, **73**: S932-S939.
3. Bahl, N. and Agnihotri, N. P. 1989. Translocation of Nematicide Residues in Mushroom Following their Application at Composting. *Mush. Sci.*, **XII**: 883-888.
4. Cayrol, J. C. 1967. Eludes Preliminaires Des Relations Entre Quelques Champignons Pathogènes et Deux Espèces de Nématodes Mycophages *Ditylenchus myceliophagus* et *Aphelenchoides composticola*. *Ann. Epi. Phyc.*, **18**: 317-329.
5. Cayrol, J. C. and Frankowski, J. P. 1979. Lutte Contre les Nématodes Mycophages, Parasites des Champignons Omnicides à l'Aide d'un Hyménoptère Prédateur du Genre *Arthrobotrys*. *Mush. Sci.*, **10**: 407-413.
6. Chauhan, J. B., Subramanian, R. B. and Sanyal, P. K. 2002. Influence of Heavy



- Metals and a Fungicide on Growth Profiles of Nematophagous Fungus, *Arthrobotrys musiformis*, a Potential Biocontrol Agent against Animal Parasitic Nematodes. *India Environ. Toxicol.*, **12**: 22-25.
7. Dmoch, J., Jacewicz, E. and Bykowska, A. 1989. Effect of Some I. G. R. Insecticides on Mushroom Mycelial Growth. In: "*Mushroom Sci., XII (Part II)*", Proceedings of the Twelfth International Congress on the Science and Cultivation of Edible Fungi. Braunschweig, Germany (FRG), 1987, PP. 877-882.
 8. Easton, A., Guven, K. and de Pomerai, D. I. 2001. Toxicity of the Dithiocarbamate Fungicide Mancozeb to the Non-target Soil Nematode, *Caenorhabditis elegans*. *J. Biochem. Mol. Toxicol.*, **15**: 15-25.
 9. Fletcher, J. T., White, P. F. and Gaze, R. H. I. 1986. *Mushroom Pest and Disease Control*. Intercept Ponteland New Castle Upon Tyne, U. K. 147 PP.
 10. Garcha H S., Khanna, P. and Sandhu, G. S. 1986. Status of Pests in the Cultivated Mushroom in India. In: "*Cultivating Edible Fungi*". (Eds.) Wuest, P. J., Royse, D. J. and Belman, R. D. Elsevier Pub., New York (1987), PP. 649 – 665.
 11. Grewal, P. S. 1993. Nematodes of Cultivated Mushroom: Diversity Bionomics and Control. *Nematol. Absrt.*, **62**: 175-191.
 12. Grewal, P. S. and Sohi, H. S. 1988. A New and Cheaper Technique for Rapid Multiplication of *Arthrobotrys conoides* and Its Potential as a Bio-nematicide in Mushroom Culture. *Curr. Sci.*, **57**: 44-46.
 13. Heydari, R., Pourjam, E. and Mohammadi Goltapeh, E. 2006. Antagonistic Effect of Some Species of *Pleurotus* on the Root-knot Nematode, *Meloidogyne javanica* *in vitro*. *Plant Pathol. J.*, **5**: 173 – 177.
 14. Kerry, B. R. and Crump, D. H. 1998. The Dynamics of the Decline of the Cereal Cyst Nematode, *Heterodera avenae*, in Four Soils under Intensive Cereal Production. *Funda. Appl. Nematol.*, **21**: 617-625.
 15. LaMondia, J. 2004. *Plant Parasitic Nematodes*. GPN 14
 16. Matsumura, F. 1985. Toxicology of Insecticides, Second Edition. Plenum Press, New York, USA. 598p.
 17. Mohammadi Goltapeh, E., and Pourjam, E. 2005. *Principles of Mushroom Cultivation* 5th edition, Tarbiat Modarres University Press, Tehran, Iran. 626 PP.
 18. Mohammadi Goltapeh, E. and Rezaei Danesh, Y. 2006. Pathogenic Interaction between *Trichoderma* Species and *Agaricus bisporus*. *Inter. J. Agric. Technol.*, **2(1)**: 29-37.
 19. Papavizas, G. C., Lewis, J. A. and Adams, P. B., 1970. Survival of Root Infecting Fungi in Soil. XIV. Effect of Amendments and Fungicides on Bean Root Rot Caused by *Thielaviopsis basicola*. *Pl. Dis. Reprtr.*, **54**: 114–118.
 20. Pullen, M. P., Zehr, E. I. and Carter, G. E Jr. 1990. Influence of Certain Fungicides on Parasitism of the Nematode *Cricone-mella xenoplax* by the fungus *Hirsutella rhossiliensis*. *Phytopathol.*, **80**: 1142-1146.
 21. Raj, H. and Kapoor, I. J. 1996. Effect of Oil Cake Amendment of Soil on Tomato Wilt Caused by *Fusarium oxysporum* f. sp. *lycopersici*. *Indian Phytopath.*, **49**: 355-361.
 22. Sharma, A., Chauhan, R. and Singh, V. 2002. Some Common Nematodes of Mushroom and Their Management. *Ann. Agri. Bio. Res.*, **7**: 51-52.
 23. Sharma, N. K., Thapa, C. D. and Damanjeet, 1984. Estimation of Loss in Mushroom Yield Due to Myceliophagous Nematode, *Aphelenchoides sacchari*. *Indian J. Nematol.*, **14**: 121-124.
 24. Stauntan, L. 1996. Disease and Pest Control in Mushroom. *Food. Horticulture*, **1**: 13-17.
 25. Vincent, J. H. 1947. Distortion of Fungal Hyphae in the Presence of Certain Inhibitors. *Nature*, **159**: 850.

حساسیت قارچ نماتد خوار *Arthrobotrys oligospora* به قارچ کشتهای، حشره کشتهای و مکملهای غذایی استفاده شده در کشت تجاری قارچ خوراکی دکمه‌ای *Agaricus bisporus*

۱. محمدی گل تپه، م. شمس بخش و ب. س. پاکدامن

چکیده

تأثیر آفت کشتهای مختلف دیفلوبنزرو، مالاتیون، مانکوزب و کاربندازیم، ترکیبات ضد عفونی کننده (هیپوکلریت کلسیم و فرمالدئید) و کنجاله‌ها (آفتابگردان و سویا) که به طور معمول در کشت و پرورش قارچ خوراکی به عنوان مکمل های غذایی به کار می روند، بر رشد قارچ نماتد خوار (*Arthrobotrys oligospora*) در شرایط درون شیشه‌ای (*in vitro*) بررسی شد. کاربندازیم در شرایط درون شیشه‌ای و در تمامی غلظت‌های کاربردی (۱۰ تا ۴۰ میکروگرم در میلی لیتر ($\mu\text{g a.i. ml}^{-1}$)) در مقایسه با نمونه‌های تیمار نشده سبب ۹۹٪ بازدارندگی از رشد شعاعی میسلومی قارچ شد. مانکوزب در غلظت‌های ۲۵۰ و ۵۰۰ میکروگرم ($\mu\text{g a. i. ml}^{-1}$) به ترتیب باعث ۴۳٪ و ۲۳٪ بازدارندگی و در غلظت ۱۰۰۰ میکروگرم و بیشتر نیز سبب ۹۹٪ بازدارندگی شد. دیفلوبنزرو و مالاتیون نیز در غلظت ۱۰-۴۰ میکروگرم ($\mu\text{g a. i. ml}^{-1}$) به ترتیب باعث ۳۱-۴۱ و ۲۴-۵۴٪ بازدارندگی شدند. فرمالین با غلظت ۰/۵-۲/۰٪ (حجمی) نیز به طور کامل سبب ممانت از رشد قارچ *A. oligospora* شد. در حالی که، هیپوکلریت کلسیم، کنجاله آفتابگردان و سویا در غلظت‌های بیش از ۲/۰٪ (w/v) کمتر از ۳/۵٪ بازدارندگی را سبب می شدند.