

Comparison of 17 Rapeseed Cultivars under Terminal Water Deficit Conditions Using Drought Tolerance Indices

H. Eyni Nargeseh¹, M. Aghaalikhani^{1*}, A. H. Shirani Rad², A. Mokhtassi-Bidgoli¹, and S. A. M. Modarres Sanavy¹

ABSTRACT

Drought stress is one of the environmental factors influencing crops growth, development, and production. Two field experiments were performed in Karaj, Iran, to evaluate the drought tolerance indices of 17 winter rapeseed genotypes in 2015-2016 and 2016-2017 growing seasons. The factorial arrangement of treatments was set up as RCBD with three replications. To identify drought tolerant genotypes, several indices were used based on grain yield under normal and deficit irrigation conditions. Yield results showed that cultivars Artist (504.325 g m⁻²) and L72 (391.525 g m⁻²) were the superior treatments under normal and deficit irrigation conditions, respectively. According to correlation results, 3-D graphs were drawn based on Geometric Mean Productivity (GMP) and grain yield under normal irrigation and deficit irrigation to categorize the winter rapeseed genotypes in both years. In the first year, Zorica and Lauren were in group A, while in the second year; Mercure, SW102, L72, and HL3721 were in group A. Therefore, they had superior performance and stable grain yield under both irrigation conditions. Biplot diagram showed Lauren (first year) and Mercure (second year) were superior regardless of stress conditions. Altogether, under normal irrigation, Artist genotype, and under stress condition, Mercure, L72 and HL3721 genotypes could be used for cultivation.

Keywords: Biplot, Principal component analysis, Stress susceptibility index.

INTRODUCTION

Drought or water deficit is one of the environmental stresses that severely influence the crops growth, development, and production (Werteker *et al.*, 2010; Ongom *et al.*, 2016). The response of crops to water deficit stress is a function of genotypes, intensity and duration of stress, weather conditions and stages of plant growth and development. It should be noted that stress occurrence time is more important than drought stress intensity.

Due to different genetic makeup, genotypes usually vary in their responses to environment, which is called genotype-environment interaction (Mansour *et al.* 2017). The interaction

between genotype and environment further complicates breeding work because of difficulties in predicting how genotypes will perform under different environmental conditions (Ceccarelli, 1989; Shakhatareh *et al.*, 2001). Drought susceptibility of a genotype is often measured as a function of the reduction in yield under water deficit stress (Blum, 2012). In this context, Rashidi *et al.* (2017) evaluated the response of 36 Brassica genotypes belonging to seven famous species of Brassica. Results of this study showed that moisture, environments, and genotypes have significant influence on grain yield and yield components of *Brassica* species. Mansour *et al.* (2017) reported that grain yield and yield components of barely were affected by genotypes and water deficit stress. Grain yield

¹Department of Agronomy, Faculty of Agriculture, Tarbiat Modares University, PO Box 14115-336, Tehran, Islamic Republic of Iran.

²Seed and Plant Improvement Institute (SPII), Agricultural Research, Education and Extension Organization (AREEO), Karaj, Islamic Republic of Iran.

* Corresponding author; e-mail: maghaalikhani@modares.ac.ir



significantly increased by drought-tolerant genotypes followed by moderate tolerant genotypes in comparison with drought-sensitive genotypes. In another research, Samarah *et al.* (2009) reported a 73-87% grain yield reduction as a result of severe drought in various genotypes of barley in a Mediterranean environment. They discussed that drought tolerant cultivars could play a significant role in mitigating the negative impacts of water stress on plants.

There are several indices to evaluate the susceptibility or tolerance of a crop genotype to stress conditions compared to normal condition (Fernandez, 1992). These drought tolerance indices provide a measure of drought based on loss of yield under stress condition in comparison to the non-stress condition that has been used to select drought tolerant genotypes (Bahrami *et al.*, 2014). Several indices have been utilized to evaluate the drought tolerance of genotypes based on grain yield under stress and non-stress conditions. Mohammadi (2016) reported that discrimination among the genotypes based on mean values was better under severe stress than mild stress conditions. Rashidi *et al.* (2017) investigated the response of *Brassica* species to water deficit stress. Results of this study based on correlation coefficients showed that Geometric Mean Productivity (GMP), Stress Tolerance Index (STI), and Mean Productivity (MP) were the most appropriate criteria for selecting high-yield genotypes under stress and non-stress conditions.

Brassica napus L. (2n= 38), commonly called rapeseed or colza in many European countries, is an annual crop belonging to Brassicaceae (Cruciferae) family. It is one of the most important oilseed crops worldwide (Zhang *et al.*, 2013; Nowosad *et al.*, 2016; Eyni-Nargeseh *et al.*, 2019) with over 36 million hectares cultivation area in 2014 (FAO 2017). The total area under cultivation of rapeseed in Iran has decreased by 50% compared to the previous years, mainly due to water shortage (Ministry of Agriculture, 2017). Therefore, new researches to increase the area under cultivation of rapeseed are important and essential. Quantitative measurement of drought tolerance criteria has an important role in evaluating different cultivars for drought tolerance (Clarke *et al.*, 1992). Also, considering rapeseed as an important oilseed crop and its sensitivity to late season drought,

rapeseed genotypes should be evaluated in terms of adaptability and drought tolerance. These findings could be useful to find the appropriate solutions for crop production in semi-arid regions. Therefore, the objective of the current study was to evaluate drought tolerance of 17 new rapeseed genotypes based on drought tolerance indices as well as select and introduce the most drought tolerant genotypes in semi-arid regions.

MATERIALS AND METHODS

Location, Experimental Design and Treatments

Two field experiments were performed at the Research Field of Seed and Plant Improvement Institute (SPII), Karaj, Iran, to evaluate drought tolerance of 17 new winter rapeseed genotypes (cultivar, hybrid, and line) (Table 1) during 2015-2017 growing seasons. The studied genotypes included two Hungarian cultivars (G1- Zorica and G2- Zlanta), seven French hybrids (G3-Artist, G4-Mercure, G5-Kamilo, G6-Lauren, G7-Darko, G8-Alonso and G9-Hydromel), two German hybrids (G10-Rohan and G11-Garou), four Iranian lines (G12-SW102, G13-HL2012, G14-L72 and G15-HL3721), an Iranian cultivar (G16-Ahmadi) and a French cultivar (G17-Okapi). The experimental treatments consisted of two irrigation regimes [normal irrigation during the growing season and withholding irrigation from silique setting stage (69, BBCH-scale) until the end of the growing season] and the aforementioned 17 new winter genotypes of rapeseed. The BBCH-scale is a system for uniform coding of phenologically similar growth stages of all mono- and dicotyledonous plant species. According to this system, "69, BBCH-scale" is considered the end of flowering. More details are given by Lancashire *et al.* (1991). Factorial arrangement of the treatments was set up as a Randomized Complete Block Design (RCBD) with three replications. Each experimental plot consisted of 6 rows, 5 m length with an inter-row distance of 30 cm and inter-plant distance of 4 cm. A two m distance was kept to eliminate all influence of lateral water movement between plots. Irrigation intervals were adjusted based on 80 mm

Table 1. Genetic status, pollination status and maturity group for different genotypes of rapeseed.


| Genotype | Genetic Status | Pollination status | Maturity group |
|--------------|----------------|--------------------|----------------|
| G1- Zorica | Cultivar | Open pollinating | Late maturing |
| G2- Zlanta | Cultivar | Open pollinating | Late maturing |
| G3- Artist | Hybrid | - | Mid maturing |
| G4- Mercure | Hybrid | - | Mid maturing |
| G5-Kamilo | Hybrid | - | Mid maturing |
| G6- Lauren | Hybrid | - | Mid maturing |
| G7- Darko | Hybrid | - | Mid maturing |
| G8-Alonso | Hybrid | - | Mid maturing |
| G9- Hydromel | Hybrid | - | Mid maturing |
| G10- Rohan | Hybrid | - | Mid maturing |
| G11- Garou | Hybrid | - | Mid maturing |
| G12- SW102 | Line | Open pollinating | Mid maturing |
| G13- HL2012 | Line | Open pollinating | Mid maturing |
| G14- L72 | Line | Open pollinating | Mid maturing |
| G15- HL3721 | Line | Open pollinating | Mid maturing |
| G16- Ahmadi | Cultivar | Open pollinating | Mid maturing |
| G17- Okapi | Cultivar | Open pollinating | Late maturing |

evaporation from Class A evaporation pan (Safavi Fard *et al.*, 2018). Water volume entering the field was measured by a water meter. Experimental plots under normal irrigation received 5,760 and 5,120 m³ water ha⁻¹ in 2016 and 2017, respectively, while in deficit irrigation treatments, 1,280 m³ water ha⁻¹ were saved in both years.

The experimental site is located at 50° 75' E longitude, 35° 59' N latitude and 1,321 m in a

semi-arid area. Based on the long-term average (from 1985 to 2015), average annual precipitation is 253 mm, which occurs mainly during late autumn to early spring. More details including the mean monthly precipitation, minimum and maximum temperatures are given in Figure 1.

According to the results of soil analysis, soil texture was clay loam. Nitrogen fertilizer was applied in three splits (one-third pre-plant, one-

**Figure 1.** Rainfall (mm), maximum and minimum temperatures (°C) during growing season of rapeseed in 2015-2016 and 2016-2017.



third in stemming stage (30-31, BBCH-scale) and one-third in flowering stage (60, BBCH-scale)) but all P and K fertilizers were applied pre-planting. Weeds were controlled with an application of 2.5 L ha⁻¹ trifluralin (48% EC) pre-plant and hand weeding in both growing seasons. Finally, rapeseed seeds were planted on 2 October in both years.

Agronomic Traits

To measure the grain yield, the final harvest was conducted by harvesting the four middle rows at physiological maturity (at 14% humidity) when 50% of the grains in the main siliques and primary branches turned brown (Ozer, 2003). Silique number per plant and grain number per silique were counted from 50 randomly selected siliques. The 1,000-grain weight was determined by measuring the weight of eight random samples, each of which consisted of 100 grains, from each plot and multiplying it by 10 to express it to 1,000 grain.

Drought Tolerance Indices

To categorize different genotypes, drought tolerance indices were calculated based on grain yield of genotypes under normal irrigation (non-stress) and withholding irrigation (stress) conditions. In the current study, nine drought tolerance indices including Stress Susceptibility Index (SSI, Fischer and Maurer, 1978), stress Tolerance (TOL, Hossain *et al.*, 1990), Geometric Mean Productivity (GMP, Fernandez 1992), Stress Tolerance Index (STI, Fernandez 1992), Drought Susceptibility Index (DSI, Fischer and Maurer 1978), Harmonic Mean (HARM, Fernandez 1992), Mean Productivity (MP, Hossain *et al.* 1990), Yield Stability Index (YSI, Bouslama and Schapaugh,

1984), and Yield Index (YI, Gavuzzi *et al.*, 1997) were used.

Statistical Procedures

Combined analysis of variance, mean comparison (Least significant difference, *P<0.05) and correlation were done using SAS software (version 9.2). Finally, biplot and genotypes distribution graphs were drawn by OriginPro 9.1 software package. In the genotypes distribution graph, x, y, and z are grain yield under deficit irrigation condition (Ys), grain yield under normal irrigation condition (Yp) and stress tolerance indices, respectively. It is also worth noting biplot was obtained from principal component analysis using the grain yield of 17 rapeseed genotypes under normal irrigation (Yp) and deficit irrigation (Ys) conditions, and drought tolerance indices.

RESULTS AND DISCUSSION

Grain Yield and Yield Components

Deficit irrigation had a significant influence on grain yield and yield components of different genotypes (data not shown). According to the results of combined analysis of variance, there were no significant differences between the two years (2015-2016 and 2016-2017) in terms of grain yield and stress tolerance indices (Table 2), but the two-way interaction between genotype and year was statistically significant on grain yield and stress tolerance indices. Therefore, the response of rapeseed genotypes was different in both years, and results are presented for each year separately.

In the first year (Table 3), under normal irrigation condition, Zorica, Lauren and Artist

Table 2. Summary of combined *F* significance from analysis of variance for grain yield under normal irrigation (Yp) and deficit irrigation (Ys) conditions and drought tolerance indices of 17 rapeseed genotypes.^a

| SOV | DF | Yp | Ys | SSI | STI | TOL | DSI | GMP | HAM | YSI | YI | MP |
|-----|----|----|----|-----|-----|-----|-----|-----|-----|-----|----|----|
| Y | 1 | ns | ns | ns | ns | ns | ns | ns | ns | ns | ns | ns |
| G | 16 | ** | ** | * | ** | * | ** | ** | ** | * | ** | ** |
| Y×G | 16 | ** | ** | * | ** | * | ** | ** | ** | * | ** | ** |

^a Y: Year, G: Genotype, * and **: Significant at the 5 and 1% levels of probability, respectively; ns: Not significant.

genotypes produced higher grain yield at 532.42, 505.56 and 504.94 g m⁻², respectively, while Hydromel genotype produced the lowest grain yield (373.27 g m⁻²). In deficit irrigation condition, Lauren and Alonso with grain yield at 385.43 and 377.63 g m⁻², respectively, were in the top group, while HL2012 genotype had the lowest grain yield (269.63 g m⁻²) (Table 3). It is worth noting that the average grain yield of genotypes under normal irrigation (440.278 g m⁻²) was higher compared to deficit irrigation (332.764 g m⁻²). The response of rapeseed genotypes was different in the second year. Among studied genotypes under normal irrigation condition, HL3721 produced the highest grain yield (532.77 g m⁻²) followed by Artist, L72, Garu, Mercure, and Hydromel, while Zlanta genotype produced the lowest grain yield (366.58 g m⁻²) (Table 4). Superior genotypes under deficit irrigation condition were different, such that L72 genotype had the highest grain yield (430.8 g m⁻²) followed by Mercure, HL3721, HL2012, and Rohan genotypes, while Zorica genotype produced the minimum grain yield (213.57 g m⁻²) (Table 4).

Averaged over both years, Artist (504.325 g m⁻²) and (391.525 g m⁻²) was superior genotype under normal irrigation condition, but L72, HL3721, and Mercure (391.525, 389.245 and 375.445 g m⁻², respectively) were most resistant genotypes under deficit condition. High yield in drought-tolerant genotypes could be explained by higher yield components for those genotypes under water stress conditions (Mansour *et al.*, 2017). According to Diepenbrock (2000), rapeseed grain yield is a function of different traits, consisting of the number of silique per plant, the number of grains per silique and the individual grain weight. In general, the high grain yield of Artist genotype compared to other genotypes in both years can be attributed to the number of silique per plant (210.8), the number of grains per silique (25.6) and the 1,000-grain weight (4.6 g) (Tables 5 and 6). Results indicated that L72, HL3721, and Mercure genotypes produced more grain yield than other genotypes under deficit irrigation condition due to highest silique per plant (159.8, 165.8 and 139.7, respectively), number of grains per silique (16.8, 17.1 and 13.6,

respectively) and 1,000-grain weight (3.22, 3.27 and 2.69 g, respectively) (Tables 5 and 6). According to the results of correlation analysis (Table 7), there was a strong positive and significant correlation between grain yield and yield components (silique plant⁻¹, number of grains silique⁻¹ and 1,000-grain weight) under normal irrigation and deficit irrigation conditions in both years. The number of silique per plant during the course of development is ultimately determined by reduction in the number of branches, buds, flowers, and young siliques by source capacity, the supply of nutrients, water, and hormonal factors rather than by the potential numbers of flowers and siliques (Diepenbrock, 2000). An increase in the number of grains per silique results in higher source size and, finally, it leads to increased performance (Tayo and Morgan, 1979). The grain weight is the last yield component to be accomplished over development (Diepenbrock, 2000). Overall, the grain weight depends on the rate and duration of the grain filling, and it is the resultant of two sources of current photosynthesis and remobilization. Regarding the decreased grains weight under deficit irrigation treatments, rate and duration of the grain filling would be considered as the main effective factors for grains weight of rapeseed (Sinaki *et al.*, 2007).

Drought Tolerance Indices

As seen in Table 3, Lauren and Alonso were the most drought tolerant genotypes in the first year of the experiment. Lauren genotype had the highest values of STI (1.0), GMP (4,397.3), HAM (4,340.7), YI (1.15) and MP (4,455.0). It should be noted that this genotype had no significant difference in terms of other indices (SSI, TOL, and YSI) with superior genotypes (Table 3). Hydromel and Ahmadi genotypes had the lowest values of STI (0.55 and 0.56), GMP (3,268.4 and 3,317.6), DSI (3,739.5 and 3,843.1), HAM (3,232.5 and 3,273.7), YI (0.86) and MP (3,305.0 and 3,362.5), so, they were identified as susceptible genotypes (Table 3). In addition, these genotypes had low grain yield under normal irrigation condition. In the second year of the current study, Mercure, L72, and

Table 3. Mean of grain yield and drought tolerance indices of rapeseed genotypes under normal irrigation and deficit irrigation conditions in 2015-2016.^a

| Genotypes | Yp (g m ⁻²) | Ys (g m ⁻²) | SSI | STI | TOL | GMP | DSI | HAM | YSI | YI | MP |
|-----------|-------------------------|-------------------------|---------|---------|-----------|-----------|-----------|-----------|---------|---------|-----------|
| Zorica | 532.42 (±25.04)ja | 360.71 (±13.52)ab | 1.29ab | 0.98a | 1717.1abc | 4374.6ab | 5327.6a | 4285.9ab | 0.68bc | 1.08ab | 4465.6a |
| Zlanta | 461.51 (±11.37)bc | 357.45 (±26.15)abc | 0.91abc | 0.84ab | 1040.6a-e | 4053.6abc | 4620.7a-d | 4013.0abc | 0.77abc | 1.07abc | 4094.8ab |
| Artist | 504.94 (±22.43)ab | 302.24 (±16.30)bc | 1.62a | 0.78bcd | 2027.0a | 3899.7cde | 5052.0abc | 3769.0c-f | 0.60c | 0.90b-e | 4035.9abc |
| Mercure | 447.41 (±22.59)bcd | 336.61 (±36.54)ja-d | 0.99abc | 0.77b-e | 1107.9a-e | 3865.3cde | 4483.0b-f | 3811.7cde | 0.75abc | 1.01a-d | 3920.1b-e |
| Kamillo | 383.49 (±14.32)de | 351.90 (±29.75)ja-d | 0.31c | 0.69b-f | 315.2e | 3661.4c-g | 3793.1f | 3646.4c-g | 0.92a | 1.05a-d | 3676.7b-f |
| Lauren | 505.56 (±11.20)ab | 385.43 (±40.27)ja | 0.95abc | 1.00a | 1201.3a-e | 4397.3a | 5066.5ab | 4340.7a | 0.77abc | 1.15a | 4455.0a |
| Darko | 404.98 (±10.01)cde | 293.22 (±29.98)de | 1.11abc | 0.61ef | 1117.7a-e | 3434.2fg | 4054.9def | 3378.8efg | 0.72abc | 0.88cde | 3491.0ef |
| Alonso | 437.57 (±18.05)bc | 377.63 (±5.36)ja | 0.55bc | 0.85ab | 599.4de | 4063.3abc | 4384.1b-f | 4050.7abc | 0.86ab | 1.13a | 4076.0ab |
| Hydromel | 373.27 (±12.93)bc | 287.73 (±29.52)de | 0.93abc | 0.55f | 855.4b-e | 3268.4g | 3739.5f | 3232.5g | 0.77abc | 0.86de | 3305.0f |
| Rohan | 455.62 (±41.65)bc | 304.53 (±5.20)bc | 1.31ab | 0.71b-f | 1510.9a-d | 3718.5c-f | 4559.5b-e | 3638.5c-g | 0.67bc | 0.91b-e | 3800.7b-e |
| Garou | 380.73 (±3.72)de | 332.60 (±27.81)ja-e | 0.52bc | 0.65e-f | 481.3de | 3553.8d-g | 4269.5def | 3541.0d-g | 0.87ab | 0.99a-e | 3566.7def |
| SW102 | 459.88 (±39.30)bc | 340.15 (±27.96)ja-d | 1.01abc | 0.80bc | 1197.2a-e | 3940.6cd | 4605.8a-d | 3882.4bcd | 0.75abc | 1.02a-d | 4000.2bc |
| HL2012 | 455.37 (±45.93)bc | 269.59 (±19.65)je | 1.60a | 0.63def | 1857.8ab | 3487.5efg | 4556.6b-e | 3357.0fg | 0.60c | 0.81e | 3624.8c-f |
| L72 | 429.11 (±29.67)cde | 352.25 (±7.80)ja-d | 0.70bc | 0.77b-e | 768.6cde | 3882.3cde | 4298.7c-f | 3858.1bcd | 0.82ab | 1.05a-d | 3906.8b-e |
| HL3721 | 431.15 (±21.83)cde | 367.04 (±27.33)ab | 0.56bc | 0.81bc | 641.1de | 3963.3bcd | 4308.4c-f | 3936.1a-d | 0.86ab | 1.10ab | 3990.9bcd |
| Ahmadi | 383.49 (±30.63)de | 289.02 (±14.33)de | 0.95abc | 0.56f | 944.7b-e | 3317.6fg | 3843.1ef | 3273.7g | 0.76abc | 0.86de | 3362.5f |
| Okapi | 438.23 (±33.48)bc | 348.90 (±13.04)ja-d | 0.81abc | 0.79bcd | 893.3b-e | 3908.1cde | 4387.9b-f | 3880.8bcd | 0.80abc | 1.04a-d | 3935.6bcd |

^a Means within a columns followed by similar letters are not significantly different at 5% probability level. Note: The value after the (±) is standard error.

Table 4. Mean of grain yield and drought tolerance indices of rapeseed genotypes under normal irrigation and deficit irrigation conditions in 2016-2017.^a

| Genotypes | Yp (g m ⁻²) | Ys (g m ⁻²) | SSI | STI | TOL | GMP | DSI | HAM | YSI | YI | MP |
|-----------|-------------------------|-------------------------|---------|---------|-----------|-----------|-----------|-----------|---------|---------|-----------|
| Zorica | 423.04 (±19.90)c-g | 213.59 (±8.01)h | 1.93a | 0.45h | 2094.5a | 3000.6h | 4232.5c-f | 2829.6h | 0.50e | 0.64h | 3183.1h |
| Zlanta | 366.58 (±9.03)g | 306.57 (±22.43)efg | 0.63cde | 0.57e-h | 600.0de | 3345.7fgh | 3678.0ef | 3325.9fg | 0.83abc | 0.92efg | 3365.7fgh |
| Artist | 503.71 (±22.37)ab | 305.79 (±16.49)efg | 1.53ab | 0.78cd | 1979.3ab | 3917.7de | 5039.8abc | 3792.9de | 0.61de | 0.92efg | 4047.5cde |
| Mercure | 476.17 (±24.04)ja-d | 414.28 (±44.97)ab | 0.48e | 1.00ab | 619.0de | 4423.7abc | 4761.3bcd | 4395.5abc | 0.87a | 1.25ab | 4452.2abc |
| Kamillo | 417.68 (±15.60)c-f | 254.04 (±21.48)gh | 1.52ab | 0.53fgh | 1636.4a-d | 3246.9gh | 4179.5c-f | 3140.4gh | 0.61de | 0.76gh | 3358.6fgh |
| Lauren | 408.23 (±9.05)d-g | 338.78 (±35.40)def | 0.65b-e | 0.70e-f | 694.5cde | 3704.6def | 4077.8def | 3674.4def | 0.83a-d | 1.02def | 3735.1def |
| Darko | 424.67 (±10.50)c-g | 230.59 (±23.58)h | 1.78a | 0.49gh | 1940.8ab | 3118.6h | 4249.0c-f | 2970.1gh | 0.54e | 0.69h | 3276.3gh |
| Alonso | 398.76 (±16.44)efg | 280.96 (±3.99)fgh | 1.15a-e | 0.57e-h | 1178.0a-e | 3345.8fgh | 3991.0def | 3293.9fg | 0.70a-e | 0.85fgh | 3398.6fgh |
| Hydromel | 473.53 (±16.41)ja-d | 301.04 (±30.89)efg | 1.42abc | 0.72cde | 1724.9abc | 3765.9def | 4738.3bcd | 3661.9def | 0.63cde | 0.91efg | 3872.8de |
| Rohan | 432.47 (±39.53)c-g | 368.20 (±6.29)ja-d | 0.53de | 0.81cd | 642.7de | 3983.6cde | 4336.1c-f | 3964.0cde | 0.86ab | 1.11a-e | 4003.4de |
| Garou | 476.46 (±4.66)ja-d | 309.83 (±25.91)efg | 1.37a-d | 0.75cd | 1666.3a-d | 3837.0de | 4767.6bcd | 3745.3def | 0.64b-e | 0.93efg | 3931.5de |
| SW102 | 456.20 (±38.98)bc | 359.54 (±29.55)bc | 0.78b-e | 0.83bcd | 966.5b-e | 4035.1cde | 4625.5bcd | 3992.2cde | 0.80a-d | 1.08b-e | 4078.7cd |
| HL2012 | 449.28 (±45.31)bc | 380.66 (±27.74)ja-d | 0.51de | 0.87bc | 686.2cde | 4116.4bcd | 4492.7b-e | 4083.3bcd | 0.86ab | 1.15a-d | 4149.7bcd |
| L72 | 482.70 (±33.37)abc | 430.80 (±9.54)ja | 0.38e | 1.06a | 519.0e | 4553.6ab | 5904.9a | 4539.9ab | 0.90a | 1.30a | 4567.5ab |
| HL3721 | 532.78 (±26.97)ja | 411.45 (±30.64)abc | 0.85bc | 1.11a | 1213.3a-e | 4664.7a | 5337.5ab | 4609.5a | 0.78a-d | 1.24abc | 4721.1a |
| Ahmadi | 419.72 (±19.33)c-g | 359.70 (±22.24)bc | 0.54de | 0.76cd | 600.2de | 3877.3de | 4209.2c-f | 3857.7de | 0.86ab | 1.09b-e | 3897.1de |
| Okapi | 379.98 (±29.03)fg | 344.36 (±12.87)c-f | 0.34e | 0.67d-g | 356.0e | 3615.4efg | 3606.0f | 3609.0ef | 0.91a | 1.04c-f | 3621.7efg |

^a Means within a columns followed by similar letters are not significantly different at 5% probability level. Note: The value after the (±) is standard error.

Table 5. Mean of silique plant⁻¹, number of grain plant⁻¹ and 1000-grain weight (g) of rapeseed genotypes under normal irrigation condition in 2015-2016 and 2016-2017.^a

| Genotypes | Silique plant ⁻¹ | | Number of grain plant ⁻¹ | | 1000-Grain weight (g) | |
|-----------|-----------------------------|-----------|-------------------------------------|----------|-----------------------|---------|
| | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 |
| Zorica | 215.09a | 170.90gh | 26.87a | 21.35c-g | 4.82a | 3.83de |
| Zlanta | 180.86f-i | 143.65j | 21.94c-g | 17.43h | 3.86d-g | 3.07f |
| Artist | 211.12ab | 210.60b | 25.66ab | 25.60ab | 4.6ab | 4.59ab |
| Mercure | 195.47cde | 208.04bcd | 23.42a-e | 24.93abc | 4.16b-e | 4.43b |
| Kamilo | 173.45hij | 188.95ef | 20.39e-h | 22.21b-f | 3.81d-h | 4.15bcd |
| Lauren | 200.07bcd | 161.55hi | 24.34a-d | 19.65fgh | 4.4abc | 3.55ef |
| Darko | 164.73j | 172.74gh | 19.95e-h | 20.93d-h | 3.63fgh | 3.80de |
| Alonso | 189.02d-g | 172.26gh | 21.86c-g | 19.92e-h | 4.11cd-e | 3.75de |
| Hydromel | 164.61j | 208.82bc | 19.02gh | 24.13bcd | 3.47gh | 4.41b |
| Rohan | 205.56abc | 195.12def | 24.79abc | 23.53b-e | 4.61ab | 4.38bc |
| Garou | 175.46g-i | 219.58b | 19.74fgh | 24.71abc | 3.69e-h | 4.62ab |
| SW102 | 197.63b-d | 196.05cde | 24.28a-d | 24.08bcd | 4.28bcd | 4.24bcd |
| HL2012 | 184.57e-h | 182.10fg | 21.27d-h | 20.98d-h | 3.96c-f | 3.90cde |
| L72 | 170.30ij | 191.57fe | 19.11gh | 21.49d-g | 3.41gh | 3.84de |
| HL3721 | 194.06c-f | 239.81a | 22.65b-f | 27.99a | 4.08c-f | 5.04a |
| Ahmadi | 169.54ij | 166.63hi | 18.08h | 17.33h | 3.37h | 3.23f |
| Okapi | 180.83f-i | 156.80ij | 20.66e-h | 17.91gh | 3.74e-h | 3.24f |

^a Means within a columns followed by similar letters are not significantly different at 5% probability level.

Table 6. Mean of silique plant⁻¹, number of grain plant⁻¹ and 1000-grain weight (g) of rapeseed genotypes under deficit irrigation condition in 2015-2016 and 2016-2017.^a

| Genotypes | Silique plant ⁻¹ | | Number of grain plant ⁻¹ | | 1000-Grain weight (g) | |
|-----------|-----------------------------|-----------|-------------------------------------|----------|-----------------------|---------|
| | 2016 | 2017 | 2016 | 2017 | 2016 | 2017 |
| Zorica | 136.6bcd | 80.87j | 14.5c-f | 8.61h | 2.86b-e | |
| Zlanta | 141.6bc | 121.44fg | 15.2bcd | 13.06ef | 2.98a-d | 2.55def |
| Artist | 132.1cd | 133.71def | 13.9c-g | 14.05cde | 2.8b-e | 2.83cde |
| Mercure | 125.3def | 154.27bc | 12.2fgh | 15.01cde | 2.41ef | 2.97bcd |
| Kamilo | 146.3ab | 105.64hi | 15.9bc | 11.52fg | 3.2abc | 2.31f |
| Lauren | 158.6a | 139.49de | 18.6a | 16.37abc | 3.44a | 3.02bcd |
| Darko | 119.7ef | 94.15i | 11.5gh | 9.04gh | 2.2f | 1.73g |
| Alonso | 156.7a | 116.58gh | 17.3ab | 12.88ef | 3.28ab | 2.43ef |
| Hydromel | 128.3de | 134.29de | 13d-h | 13.65def | 2.5def | 2.61def |
| Rohan | 132.8cd | 160.58b | 13.2d-h | 15.99bcd | 2.7c-f | 3.27abc |
| Garou | 136.6bcd | 127.30efg | 14.1c-f | 13.19ef | 2.81b-e | 2.62def |
| SW102 | 127.3def | 134.57de | 12.6e-h | 13.30ef | 2.48def | 2.62def |
| HL2012 | 117.6ef | 166.09ab | 11.5gh | 16.24abc | 2.26f | 3.19abc |
| L72 | 143.8bc | 175.87a | 15.1b-e | 18.56a | 2.9b-e | 3.54a |
| HL3721 | 156.3a | 175.29a | 16.2abc | 18.14ab | 3.08abc | 3.46ab |
| Ahmadi | 115.9f | 161.04b | 10.7h | 15.37cde | 2.26f | 3.21abc |
| Okapi | 147ab | 145.12cd | 15.5bcd | 15.27cde | 2.97a-d | 2.93cde |

^a Means within a columns followed by similar letters are not significantly different at 5% probability level.

**Table 7.** Correlation coefficient between grain yield of rapeseed genotypes and yield attributes under normal irrigation and deficit irrigation conditions in 2015-2016 and 2016-2017.^a

| Yield component | 2015-2016 | | | 2016-2017 | | |
|--------------------|-----------|--------|--------|-----------|--------|--------|
| | Y1 | Y2 | Y3 | Y1 | Y2 | Y3 |
| Normal irrigation | | | | | | |
| Y2 | 0.86** | | | 0.92** | | |
| Y3 | 0.88** | 0.96** | | 0.88** | 0.95** | |
| Y4 | 0.84** | 0.96** | 0.98** | 0.86** | 0.96** | 0.98** |
| Deficit irrigation | | | | | | |
| Y2 | 0.87** | | | 0.95** | | |
| Y3 | 0.85** | 0.98** | | 0.91** | 0.96** | |
| Y4 | 0.84** | 0.96** | 0.98** | 0.92** | 0.97** | 0.98** |

^a Y1: Grain yield; Y2: Silique plant⁻¹; Y3: Number of grain plant⁻¹, Y4: 1000-grain weight.** Significant 1% levels of probability.

HL3721 were considered as the most suitable genotypes because of high grain yield under both normal and deficit irrigation conditions (Table 4). In contrast, Zorica, Kamilo, and Darko were considered as the most susceptible genotypes under both normal and deficit irrigation conditions (Table 4). L72 genotype had high values of STI (1.06), TOL (519.0), GMP (4,553.6), DSI (5,904.9), HAM (4,439.9), YSI (0.90), YI (1.30) and MP (4,567.5) (Table 4). Also, this genotype had a low value of SSI (0.38). It should be noted that L72 genotype was not superior in terms of some indices (SSI, STI, TOL, GMP, HAM, YSI, MP), but there was no significant difference between L72 and other superior genotypes (Mercurie and HL3721) (Table 4). Zorica had the lowest values of STI (0.45), GMP (3,000.6), HAM (2,829.6), YSI (0.64) and MP (3,183.1). On the other hand, this genotype had the highest values of SSI (1.93) and TOL (2,094.5) (Table 4); therefore, it was identified as the most susceptible genotype under deficit irrigation and normal irrigation conditions.

Stress Susceptibility Index (SSI) is used to measure the yield stability, i.e. the changes in both stress and non-stress conditions (Fischer and Maurer, 1978). The high value of SSI represents relatively more sensitivity to stress, while the low value is favoured. Tolerance (TOL) index is the difference in grain yield between non-stress (Yp) and stress (Ys) conditions, and the low value of TOL shows

higher tolerance to stress. Thus, selection based on this criterion resulted in the selection of low-yielding genotypes under the non-stress condition and high-yielding genotypes under stress condition (Fernandez, 1992). Mean Productivity (MP) is defined as the average of Yp and Ys, but it has an upward bias when there are larger differences between Yp and Ys. The Stress Tolerance Index (STI) and Geometric Mean Productivity (GMP) proposed by Fernandez (1992) to identify genotypes with higher yield potential and stress tolerance under stress and non-stress conditions. The GMP index is less sensitive to extreme values, and it is a better index relative to MP index to separate superior genotypes under stress and non-stress conditions (Rosielle and Hamblin, 1981). Breeders interested in relative performance have used the GMP to evaluate the drought tolerant genotypes (Ramirez and Kelly, 1998). The STI was defined as a useful criterion to determine the high yield and stress tolerance potential of genotypes. The genotypes with high values of STI and GMP are superior in terms of grain yield under both stress and non-stress conditions (Fernandez, 1992). The Yield Index (YI) (Gavuzzi *et al.*, 1997) and Yield Stability Index (YSI) (Bousslama and Schapaugh, 1984) were defined as tools to evaluate the stability of genotypes under stress and non-stress conditions. Higher values of YI and YSI indicate more performance stability under stress condition. AS expected, selection based

on high values of YI and YSI leads to the selection of high-yielding genotypes under both stress and non-stress conditions (Mohammadi, 2016). The genotypes have higher performance under both stress and non-stress conditions if Harmonic Mean (HAM) (Fernandez, 1992) and Drought Sensitivity Index (DSI) (Fischer and Maurer, 1978) have high values. It is worth noting that in DSI, the contribution of performance under non-stress condition is more than that of stress condition. In the same way, Dorostkar *et al.* (2015) concluded that superior genotypes could be selected based on high values of STI, MP and GMP and low value of SSI. El-Rawy and Hassa (2014) reported that there was a positive correlation among wheat grain yield under stress condition and STI, YSI, and HAM.

3-D Graph Based on Stress Tolerance Criterion

The correlation coefficients were calculated between grain yield under normal irrigation (Yp) and deficit irrigation (Ys) conditions with drought tolerance indices to determine the most desirable stress tolerance criterion (Tables 8 and 9). Grain yields (Yp and Ys) were found to have a highly significant positive correlation with GMP and STI in both years (Tables 8 and 9).

Considering that the aim of the current study was selection of the high yield genotypes under deficit irrigation, thus, three dimensional graphs were drawn based on GMP and grain yield under normal irrigation (Yp) and deficit irrigation (Ys) conditions to categorize the 17 rapeseed genotypes in both

Table 8. Correlation coefficient between grain yield of rapeseed genotypes and drought tolerance indices under normal irrigation (Yp) and deficit irrigation (Ys) conditions in 2015-2016.

| Genotype | Yp | Ys | SSI | STI | TOL | GMP | DSI | HAM | YSI | YI |
|----------|--------------------|---------------------|----------------------|--------------------|---------------------|--------------------|--------------------|--------------------|---------------------|--------------------|
| Ys | 0.32 ^{ns} | | | | | | | | | |
| SSI | 0.57 [*] | -0.57 [*] | | | | | | | | |
| STI | 0.80 ^{**} | 0.81 ^{**} | -0.006 ^{ns} | | | | | | | |
| TOL | 0.72 ^{**} | -0.41 ^{ns} | 0.97 ^{**} | 0.18 ^{ns} | | | | | | |
| GMP | 0.80 ^{**} | 0.82 ^{**} | -0.01 ^{ns} | 0.99 ^{**} | 0.17 ^{ns} | | | | | |
| DSI | 0.97 ^{**} | 0.33 ^{ns} | 0.53 [*] | 0.79 ^{**} | 0.68 ^{**} | 0.79 ^{**} | | | | |
| HAM | 0.73 ^{**} | 0.87 ^{**} | -0.12 ^{ns} | 0.99 ^{**} | 0.06 ^{ns} | 0.99 ^{**} | 0.73 ^{**} | | | |
| YSI | -0.56 [*] | 0.58 [*] | -0.99 ^{**} | 0.02 ^{ns} | -0.97 ^{**} | 0.02 ^{ns} | -0.52 [*] | 0.13 ^{ns} | | |
| YI | 0.33 ^{ns} | 0.99 ^{**} | -0.57 [*] | 0.82 ^{**} | -0.40 ^{ns} | 0.82 ^{**} | 0.34 ^{ns} | 0.88 ^{**} | 0.58 [*] | |
| MP | 0.86 ^{**} | 0.75 [*] | 0.09 ^{ns} | 0.99 ^{**} | 0.28 ^{ns} | 0.99 ^{**} | 0.85 ^{**} | 0.97 ^{**} | -0.08 ^{ns} | 0.75 ^{**} |

* and **: Significant at 5 and 1% levels of probability, respectively; ns: Not significant.

Table 9. Correlation coefficient between grain yield of rapeseed genotypes under normal irrigation (Yp) and deficit irrigation (Ys) conditions and drought tolerance indices in 2016-2017.

| Index | Yp | Ys | SSI | STI | TOL | GMP | DSI | HAM | YSI | YI |
|-------|---------------------|---------------------|---------------------|---------------------|---------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| Ys | 0.43 ^{ns} | | | | | | | | | |
| SSI | 0.11 ^{ns} | -0.84 ^{**} | | | | | | | | |
| STI | 0.76 ^{**} | 0.93 ^{**} | -0.59 [*] | | | | | | | |
| TOL | 0.28 ^{ns} | -0.73 ^{**} | 0.98 ^{**} | -0.44 ^{ns} | | | | | | |
| GMP | 0.71 ^{**} | 0.93 ^{**} | -0.60 ^{**} | 0.99 ^{**} | -0.45 ^{ns} | | | | | |
| DSI | 0.89 ^{**} | 0.51 [*] | -0.02 ^{ns} | 0.75 ^{**} | 0.12 ^{ns} | 0.74 ^{**} | | | | |
| HAM | 0.66 ^{**} | 0.96 ^{**} | -0.66 ^{**} | 0.99 ^{**} | -0.52 [*] | 0.99 ^{**} | 0.69 ^{**} | | | |
| YSI | -0.10 ^{ns} | 0.84 ^{**} | -0.99 ^{**} | 0.59 [*] | -0.97 ^{**} | 0.61 ^{**} | 0.03 ^{ns} | 0.67 ^{**} | | |
| YI | 0.42 ^{ns} | 0.99 ^{**} | -0.84 ^{**} | 0.92 ^{**} | -0.74 ^{**} | 0.93 ^{**} | 0.51 [*] | 0.96 ^{**} | 0.85 ^{**} | |
| MP | 0.78 ^{**} | 0.89 ^{**} | -0.52 [*] | 0.99 ^{**} | -0.36 ^{ns} | 0.99 ^{**} | 0.79 ^{**} | 0.98 ^{**} | 0.53 [*] | 0.89 ^{**} |

* and **: Significant at the 5% and 1% levels of probability, respectively; ns: Not significant.



years Figure 2 (a -b). Sio-Se Mardeh *et al.* (2006) stated that MP, GMP, and STI were more suitable indices for identifying high yielding cultivars under moderate stress. Results of Naderi and Emam (2014) showed that there were significant positive correlations between rapeseed yield with several drought indices such as STI, GMP, MP, and HAM under both deficit irrigation and normal irrigation conditions. They suggested these indices were suitable to identify the drought tolerance of rapeseed cultivars.

Three dimensional graphs divide the genotypes into four groups and each division represents one combination of the genotypes: high yields under both environments (Group A); high yield in a normal environment (Group B); high yield in a stressful environment (Group C); and low yield under both environmental conditions (Group D) (Fernandez, 1992; Bahrami *et al.*, 2014). In the first year, Zorica and Lauren genotypes were in Group A, all of which had superior performance and stable grain yield under both normal irrigation and deficit irrigation conditions. Darko, Hydromel and Ahmadi genotypes were in Group D and performed poorly in both conditions. As previously

explained, the response of rapeseed genotypes was different in both years of the experiment. In the second year, Mercure, SW102, L72, and HL3721 genotypes were in Group A and Zorica, Zlanta, Kamilo, Darko, and Alonso genotypes were in Group D. Likewise, in different studies, the same method was used to categorize the genotypes into four groups based on their performance under stress and non-stress conditions (Fernandez, 1992; Rashidi *et al.*, 2017; Kamrani *et al.*, 2018).

Principal Component Analysis Using Drought Tolerance Indices and Grain Yield

Considering that 3-D graph categorizes all genotypes based on three variables (Yp, Ys, and GMP), the biplot diagrams were drawn to investigate and compare the genotypes as well as the interrelationship among all drought tolerance indices Figure 3 (a-b). As previously explained, biplot diagram was obtained from principal component analysis using the grain yield under normal irrigation (Yp) and deficit irrigation (Ys) conditions, and drought tolerance indices in 17 rapeseed genotypes (Table 10). Given that Eigenvalues were greater than or equal to 1.0, the first and second components, in total, explained more than 98.7% of the variation of the drought tolerance indices in both years (Table 10).

The principal component analysis indicated




Figure 2. Three-dimensional diagram for identifying drought tolerance genotypes based on grain yield under normal irrigation (Yp) and deficit irrigation (Ys) conditions as well as the GMP in (a) 2015-2016 (b) 2016-2017. Numbers inside the chart are according to the codes of Table 1 (i.e. G1 to G17).




Figure 3. Biplot diagram based on the first and second components obtained from PCA using the grain yield under normal irrigation (Yp) and deficit irrigation (Ys) conditions, and drought tolerance indices in 17 rapeseed genotypes in (a) 2015-2016, (b) 2016-2017. Numbers inside the chart are according to the codes of Table 1 (G1 to G17).

that the first Component (PC1) explained 60.53% of the total yield variation in the first year and it was positively correlated with STI, GMP, HAM, and MP (Table 10). Hence, the first component (PC1) could be named as the yield potential component (Figure 3-a). The second Component (PC2) explained 38.86% of the total variation in the first year, and it showed a high and positive correlation with SSI and TOL as well as a negative correlation with Ys, YSI, and YI. Thus, PC2 could be named as the stress susceptibility component, which can identify the drought tolerant genotypes from drought-sensitive ones (Figure 3-a).

According to biplot diagram, genotypes that had high PC1 (high productivity) and low PC2 (low susceptibility) are suitable under normal irrigation and deficit irrigation conditions. Accordingly, the results of the current study showed that Lauren (G6) was superior genotype under both deficit irrigation and normal irrigation conditions. In contrast, genotypes with low PC1 (low productivity) and high PC2 (high susceptibility) are susceptible under normal irrigation and deficit irrigation conditions. Therefore, their cultivation is not recommended. These genotypes included Darko (G7), Rohan (G10) and HL2012 (G13). As a result, Kamilo (G5)

and Garou (G11) with both low PC1 and PC2 had low sensitivity to deficit irrigation and can be used in breeding programs for drought tolerance (Dorostkar *et al.*, 2015). Based on the biplot diagram, indices are positively correlated if the angle between their vectors is $< 90^\circ$, negatively correlated if the angle is $> 90^\circ$, and independent if the angle is 90° (Yan and Kang, 2003). According to Figure 3-a, Yp positively correlated with the TOL, DSI, MP, GMP, STI, SSI and HAM indices, as shown by the acute angle between their vectors, while Ys positively correlated with the MP, GMP, STI, HAM, YSI and YI indices. Ys had a high negative correlation with SSI and TOL indices as shown by the obtuse angle between their vectors.

In the second year, the first Component (PC1) explained 72.85% of the total variation and exhibited a strong and positive correlation with Ys, STI, GMP, HAM, YI, and MP; while the second Component (PC2) explained 25.86% of the total variation and exhibited a high positive correlation with Yp, SSI, TOL, and DSI (Table 10; Figure 3-b). Therefore, PC1 and PC2 were named drought tolerance and stress susceptibility components, respectively. Rohan (G10), HL2012 (G13) and Ahmadi (G16) had high PC1 and low PC2; therefore, these genotypes are suitable under

**Table 10.** Results of PCA for grain yield of rapeseed genotypes under normal irrigation (Yp) and withholding irrigation (Ys) conditions and drought tolerance indices in 2015-2016 and 2016-2017 growing seasons.

| Component | Eigenvalue | Variance (%) | Yp | Ys | SSI | STI | MP | |
|-------------|------------|--------------|-------|-------|-------|-------|--------|-------|
| First year | | | | | | | | |
| PC1 | 6.65 | 60.53 | 0.32 | 0.30 | 0.01 | 0.38 | 0.38 | |
| PC2 | 4.27 | 38.86 | 0.26 | -0.29 | 0.48 | -0.02 | 0.02 | |
| Second year | | | | | | | | |
| PC1 | 8.01 | 72.85 | 0.19 | 0.34 | -0.26 | 0.34 | 0.33 | |
| PC2 | 2.84 | 25.86 | 0.48 | -0.08 | 0.38 | 0.13 | 0.18 | |
| Component | Eigenvalue | Variance (%) | TOL | GMP | DSI | HAM | YSI | YI |
| First year | | | | | | | | |
| PC1 | 6.65 | 60.53 | 0.08 | 0.38 | 0.32 | 0.38 | -0.009 | 0.30 |
| PC2 | 4.27 | 38.86 | 0.47 | -0.02 | 0.24 | -0.07 | -0.48 | -0.29 |
| Second year | | | | | | | | |
| PC1 | 8.01 | 72.85 | -0.22 | 0.34 | 0.22 | 0.34 | 0.27 | 0.34 |
| PC2 | 2.84 | 25.86 | 0.45 | 0.12 | 0.42 | 0.07 | -0.37 | -0.08 |

normal irrigation and deficit irrigation conditions, but Mercure (G4) was the best genotypes due to values of PC1 and PC2. In contrast, Zorica (G1), Artist (G3), Darko (G7), Hydromel (G9) and Garou (G11) had low PC1 and high PC2, so, these genotypes are susceptible under normal irrigation and deficit irrigation conditions, and their cultivation is not recommended. The genotypes that had high PC1 were superior under normal irrigation condition such as L72 (G14) and HL3721 (G15). Finally, Zlanta (G2) and Alonso (G8) with both low PC1 and PC2 had low sensitivity to deficit irrigation and can be used in breeding programs for drought tolerance. Based on the angle between vectors, Yp positively correlated with the DSI, MP, STI, GMP, HAM, TOL and YI indices, as shown by the acute angle between their vectors, while Ys positively correlated with the YI, HAM, GMP, STI, MP, YSI and DSI indices. It should be noted that Ys had a high negative correlation with TOL and SSI, as shown by the obtuse angle between their vectors, while Yp had near zero correlation with SSI and YSI indices, as shown by their nearly perpendicular vectors. Several researchers (PCA) have used the same method to investigate and select the superior genotypes under stress and non-stress conditions (Bennani *et al.*, 2017; Khalili *et al.*, 2016). Kaya *et al.* (2002) and Kamrani *et al.* (2018) reported that stable genotypes had greater PC1 but lower PC2 values. Results of Rashidi *et al.*

(2017) using PCA showed that *B. napus* and *B. carinata* were known as superior species for both normal and mild drought-stress conditions. In this research, Species of *B. oleracea* and *B. rapa* showed very low susceptibility and productivity under mild and intense stress conditions. *B. fruticulosa* was also recognized as a specie with high susceptibility and low productivity in both mild and intense stress. Bahrami *et al.* (2014) reported that Kermanshah47, IL, Hamedan38, Syrian, and Kordestan5 were known as superior safflower genotypes with high PC1 but low PC2 values.

CONCLUSIONS

Generally, the results illustrated that cultivar Artist (on average 504.325 g m⁻²) was a superior genotype under normal irrigation condition, but L72 (on average 391.525 g m⁻²), HL3721 (on average 389.245 g m⁻²), and Mercure (on average 375.445 g m⁻²) were the most tolerant genotypes under deficit irrigation condition. Based on the 3-D graphs, in the first year, Zorica and Lauren genotypes were in group A, but in the second year, Mercure, SW102, L72, and HL3721 genotypes were in group A, all of which had superior performance and stable grain yield under both normal irrigation and deficit irrigation conditions. According to biplot diagram, Lauren (first year) and Mercure (second year)

were superior genotypes under both deficit irrigation and normal irrigation conditions. In conclusion, the results of the present study using different approaches indicated that, under normal irrigation condition, Artist genotype, and under deficit irrigation condition, Mercure, L72, and HL3721 genotypes had high grain yield, and their cultivation is recommended in areas with similar climates.

ACKNOWLEDGEMENTS

This study is a part of a PhD. dissertation. The authors gratefully acknowledge the support provided for this survey by the Tarbiat Modares University, Iran.

REFERENCES

- Bahrami, F., Arzani, A. and Karimi V. 2014. Evaluation of Yield-Based Drought Tolerance Indices for Screening Safflower Genotypes. *Agro. J.*, **106**: 1219-1224.
- Bennani, S., Nsarellah, N., Jlibene, M., Tadesse, W., Birouk, A. and Ouabbou H. 2017. Efficiency of Drought Tolerance Indices under Different Stress Severities for Bread Wheat Selection. *Aust. J. Crop. Sci.*, **11(4)**: 395-405.
- Blum, A. 2012. Drought Resistance. In: *Plant Breeding for Water-Limited Environments*, (Eds.): Blum, A. Springer, New York, Dordrecht Heidelberg London.
- Bousslama, M. and Schapaugh, W. T. 1984. Stress Tolerance in Soybean. Part 1. Evaluation of Three Screening Techniques for Heat and Drought Tolerance. *Crop Sci.*, **24**: 933-937.
- Ceccarelli, S. 1989. Wide Adaptation: How Wide? *Euphytica*, **40**: 197-205.
- Clarke, J. M., Ronald, M. D. and Townly-smith, T. F. 1992. Evaluation of Method for Quantification of Drought Tolerance in Wheat. *Crop Sci.*, **32**: 723-728.
- Diepenbrock, W. 2000. Yield Analysis of Winter Oilseed Rape: A Review. *Field Crops Res.*, **67**: 35-49.
- Dorostkar, S., Dadkhodaie, A. and Heidari, B. 2015. Evaluation of Grain Yield Indices in Hexaploid Wheat Genotypes in Response to Drought Stress. *Arch. Agron. Soil Sci.*, **61(3)**: 397-413.
- El-Rawy, M. A. and Hassa, M. I. 2014. Effectiveness of Drought Tolerance Indices to Identify Tolerant Genotypes in Bread Wheat (*Triticum Aestivum* L.). *J. Crop Sci. Biotech.*, **17(4)**: 255-266.
- Eyni-Nargeseh, H., Aghaalkhani, M., Shirani Rad, A. H., Mokhtassi-Bidgoli, A. and Modares Sanavy, S. A. M. 2020a. Late Season Deficit Irrigation for Water-Saving: Selection of Rapeseed (*Brassica napus*) Genotypes Based on Quantitative and Qualitative Features. *Arch. Agron. Soil. Sci.* **66(1)**, 126-137. doi.org/10.1080/03650340.2019.1602866
- FAO. 2017. *Food and Agriculture Organization of the United Nations*. FAOSTAT Data, www.faostat.fao.org.
- Fernandez, G. C. J. 1992. Effective Selection Criteria for Assessing Stress Tolerance, In: *Proceedings of the International Symposium on Adaptation of Vegetables and Other Food Crops in Temperature and Water Stress*, (Ed.): Kuo, C. G. Publication, Tainan, Taiwan.
- Fischer, R. A. and Maurer, R. 1978. Drought Resistance in Spring Wheat Cultivars. I. Grain Yield Responses. *Aust. J. Agric. Res.*, **29**: 897- 912.
- Gavuzzi, P., Rizza, F., Palumbo, M., Campanile, R. G., Ricciardi, G. L. and Borghi, B. 1997. Evaluation of Field and Laboratory Predictors of Drought and Heat Tolerance in Winter Cereals. *Can. J. Plant Sci.*, **77**: 523-531.
- Hossain, A. B. S, Sears, A. G., Cox, T. S. and Paulsen, G. M. 1990. Desiccation Tolerance and Its Relationship to Assimilate Partitioning In Winter Wheat. *Crop Sci.*, **30**: 622-627.
- Kamrani, M., Hoseini, Y. and Ebadollahi, A. 2018. Evaluation for Heat Stress Tolerance in Durum Wheat Genotypes Using Stress Tolerance Indices. *Arch. Agron. Soil Sci.*, **64(1)**: 38-45.
- Khalili, M., Pour-Aboughadareh, A., and Naghavi, M. R. 2016. Assessment of Drought Tolerance in Barley: Integrated Selection Criterion and Drought Tolerance Indices. *Environ. Exp. Biol.*, **14**: 33-41.
- Kaya, Y., Palta, C. and Taner, S. 2002. Additive Main Effects and Multiplicative Interactions Analysis of Yield Performances in Bread Wheat Genotypes



- across Environments. *Turk. J. Agric. For.*, 26: 275–279.
19. Lancashire, P. D., Blieholder, H., Langeluddecke, P., Stauss, R., Van den boom, T., Weber, E. and Witzgen-Berger, A. 1991. An Uniform Decimal Code for Growth Stages of Crops and Weeds. *Ann. Appl. Biol.*, **119**: 561-601.
 20. Mansour, E., Abdul-Hamid, M. I., Yasin, M. T., Qabil, N. and Attia, A. 2017. Identifying Drought-Tolerant Genotypes of Barley and Their Responses to Various Irrigation Levels in a Mediterranean Environment. *Agric Water Manag.*, **194**: 58-67.
 21. Ministry of Agriculture Iran. 2017. *Agricultural Statistics, 2013-2014*. Volume 1. Available at: <http://www.maj.ir/Portal/Home/.pdf>
 22. Mohammadi, R. 2016. Efficiency of Yield-Based Drought Tolerance Indices to Identify Tolerant Genotypes in Durum Wheat. *Euphytica*, **211**: 71-89.
 23. Naderi, R. and Emam, Y. 2014. Evaluation of Rapeseed (*Brassica Napus* L.) Cultivars Performance under Drought Stress. *Aust. J. Crop Sci.*, **8(9)**: 1319-1323.
 24. Nowosad, K., Liersch, A., Poplawska, W. and Bocianowski, J. 2016. Genotype by Environment Interaction for Seed Yield in Rapeseed (*Brassica Napus* L.) Using Additive Main Effects and Multiplicative Interaction Model. *Euphytica*, **208**: 187-194.
 25. Ongom, P. O., Volenec, J. J. and Ejeta, G. 2016. Selection for Drought Tolerance in Sorghum Using Desiccants to Simulate Post-Anthesis Drought Stress. *Field Crops Res.*, **198**: 213-321.
 26. Ozer, H. 2003. Sowing Date and Nitrogen Rate Effects on Growth, Yield and Yield Components of Two Summer Rapeseed Cultivars. *Eur. J. Agro.*, **19(3)**: 453–463.
 27. Ramirez, P. and Kelly, J. D. 1998. Traits Related to Drought Resistance in Common Bean. *Euphytica*, **99**: 127–136.
 28. Rashidi, F., Majidi, M. M. and Pirboveyry, M. 2017. Response of Different Species of Brassica to Water Deficit. *Int. J. Plant Prod.*, **11(1)**: 1-16.
 29. Rosielle, A. A. and Hamblin, J. 1981. Theoretical Aspects of Selection for Yield in Stress and Non-Stress Environments. *Crop Sci.*, **21**: 943-946.
 30. Safavi Fard, N., Heidari Sharif Abad, H., Shirani Rad, A. H., Majidi Heravan, E. and Daneshian, J. 2018. Effect of Drought Stress on Qualitative Characteristics of Canola Cultivars in Winter Cultivation. *Ind. Crops Prod.*, **114**: 87–92.
 31. Samarah, N., Alqudah, A., Amayreh, J. and McAndrews, G. 2009. The Effect of Late-Terminal Drought Stress on Yield Components of Four Barley Cultivars. *J. Agron. Crop Sci.*, 195: 427–441.
 32. Shakhathreh, Y., Kafawin, O., Ceccarelli, S. and Saoub, H. 2001. Selection of Barley Lines for Drought Tolerance in Low-Rainfall Areas. *J. Agron. Crop Sci.*, **186**: 119–127.
 33. Sinaki, J., Majidi Heravan, M. E., Shirani Rad, A. H., Noormohammadi, Gh. and Zarei, Gh. 2007. The Effects of Water Deficit during Growth Stages of Canola (*Brassica Napus* L.). *Amer-Eurasian. J. Agric. Environ. Sci.*, **2**: 417-422.
 34. Sio-Se Mardeh, A., Ahmadi, A., Poustini, K. and Mohamadi, V. 2006. Evaluation of Drought Resistance Indices under Various Environmental Conditions. *Field Crops Res.*, **98**: 222-229.
 35. Tayo, T. O. and Morgan, D. G. 1979. Factor Influencing Flower and Pod Development in Oilseed Rape. *J. Agric. Sci.*, **92**: 363-373.
 36. Werteker, M., Lorenz, A., Johannes, H., Berghofer, E. and Findlay, C. S. 2010. Environmental and Varietal Influences on the Fatty Acid Composition of Rapeseed, Soybean and Sunflowers. *J. Agron. Crop Sci.*, **196**: 20-27.
 37. Yan, W. and Kang, M. S. 2003. GGE Biplot Analysis: A Graphical Tool for Breeders, Geneticists and Agronomists. CRC Press LLC, Boca Roton, Florida.
 38. Zhang, H., Berger, J. D and Milroy, S. P. 2013. Genotype×Environment Interaction Studies Highlight the Role of Phenology in Specific Adaptation of Canola (*Brassica Napus*) to Contrasting Mediterranean Climates. *Field Crops Res.*, **144**: 77-88.

مقایسه هفده رقم کلزا تحت شرایط کم آبی انتهای فصل با استفاده از شاخص‌های تحمل به خشکی

ح. عینی نرگسه، م. آقاعلیخانی، ا. ح. شیرانی‌راد، ع. مختصی بیدگلی، و س. ع. م. مدرس ثانوی

چکیده

تنش خشکی یکی از عوامل زیست محیطی است که رشد، نمو و تولید محصولات زراعی را تحت تأثیر قرار می‌دهد. به منظور ارزیابی شاخص‌های تحمل به خشکی در ۱۷ ژنوتیپ زمستانه کلزا دو آزمایش مزرعه‌ای در سال‌های ۱۳۹۴-۱۳۹۵ و ۱۳۹۵-۱۳۹۶، در کرج، ایران انجام شد. آزمایش به صورت فاکتوریل و در قالب طرح بلوک‌های کامل تصادفی با سه تکرار اجرا شد. برای شناسایی ژنوتیپ‌های متحمل به خشکی، چندین شاخص مبتنی بر عملکرد دانه تحت شرایط قطع آبیاری و آبیاری معمولی استفاده شد. نتایج عملکرد نشان داد که ژنوتیپ‌های Artist (۵۰۴۳/۲۵ کیلوگرم در هکتار) و L72 (۳۹۱۵/۲۵ کیلوگرم در هکتار) به ترتیب تیمارهای برتر تحت شرایط آبیاری معمولی و قطع آبیاری بودند. با توجه به نتایج همبستگی برای طبقه‌بندی ژنوتیپ‌های زمستانه کلزا، نمودارهای سه بعدی بر اساس متوسط بهره‌وری هندسی (GMP) و عملکرد دانه تحت شرایط آبیاری معمولی و قطع آبیاری در هر دو سال رسم شدند. در سال اول، ژنوتیپ‌های Zorica و Lauren در گروه A قرار گرفتند، در حالی که در سال دوم، این جایگاه به ژنوتیپ‌های Mercure، SW102، L72 و HL3721 اختصاص یافت. بر این اساس می‌توان اذعان داشت که این ژنوتیپ‌ها تحت هر دو تیمار آبیاری برتری و عملکرد دانه پایدار داشتند. نمودار بای‌پلات نشان داد Lauren (سال اول) و Mercure (سال دوم) در شرایط بدون تنش برتر بودند. در مجموع، ژنوتیپ Artist تحت شرایط آبیاری معمولی و ژنوتیپ‌های L72، Mercure و HL3721 تحت شرایط تنش برای کشت قابل توصیه می‌باشند.