## New Data of the Genus *Opius* Wesmael (Hymenoptera: Braconidae, Opiinae) from Northern Iran

S. Dolati<sup>1</sup>, A. A. Talebi<sup>1\*</sup>, S. Farahani<sup>2</sup>, and M. Khayrandish<sup>3</sup>

#### ABSTRACT

Species of the genus *Opius* Wesmael, 1835 (Hym.: Braconidae, Opiinae) was faunistically studied in northern provinces of Iran. Malaise traps were used to obtain adult specimens from various habitats during 2010–2011. A total of 17 species, belonging to 12 subgenera, were identified, of which six species including *O.* (*Hypocynodus*) larissa Fischer, 1968; *O.* (*Allotypus*) saevus Haliday, 1837; *O.* (*Merotrachys*) schmidti Fischer, 1960; *O.* (*Misophthora*) pulicariae Fischer, 1969; *O.* (*Thoracosema*) repentinus Papp, 1980; and *O.* (*Nosopaeopius*) ochrogaster Wesmael, 1835 were recorded for the first time from Iran. A key for identification of the collected species is provided.

Keywords: Iranian fauna, Leaf miner, Parasitic wasps, Parasitoid.

#### INTRODUCTION

Species of the subfamily Opiinae Blanchard, 1845 (Hym.: Braconidae) comprise one of the largest groups of parasitic wasps, with about 2,063 valid species in 39 genera, worldwide (Yu et al., 2016). Opius Wesmael, 1835 is the largest genus of Opiinae with 34 subgenera and more than 1,000 described species (Yu et al., 2016). The subfamily Opiinae is one of the important groups of parasitic wasps for biological control of Diptera. Most species of the genus Opius are endoparasitoids of leaf mining Agromyzidae, and to a lesser extent Anthomyiidae and Tephritidae (Shaw and Huddleston, 1991). Some species have great importance for biological control of agromyzid leaf miners (Belokobylskij et al., 2004; Yu et al., 2016).

The Iranian fauna of Braconidae includes 780 species in 141 genera and 26

subfamilies (Farahani et al., 2016). Most of these species have been recorded within the last decade, resulting in a four-fold increase species number from the 202 in (Fallahzadeh and Saghaei, 2010) to 780 within only six years (Rakhshani et al., 2007; Ameri et al., 2013, 2015, Farahani et al., 2014a, b, c; 2015; 2016). By 2010, only seven species of Opiinae, belonging to four genera, were known from Iran (Fallahzadeh and Saghaei, 2010). Recent taxonomic studies have led to discovery of many new records and species in this country (Farrar et al., 2009; Talebi et al., 2009; Lashkari et al., 2011; Khajeh et al., 2014; Ameri et al., 2014; Peris-Felipo et al., 2014; 2015; 2018). An updated checklist of Iranian Opiinae, including 101 species in 11 genera, was recently published by Gadallah et al. (2016), of which 66 species (65%) belonged to the genus Opius. This species number is much less than 182 species recorded from the

<sup>&</sup>lt;sup>1</sup>Department of Agricultural Entomology, Faculty of Agriculture, Tarbiat Modares University, 14115-336, Tehran, Islamic Republic of Iran.

<sup>&</sup>lt;sup>2</sup>Research Institute of Forests and Rangelands, Agricultural Research Education and Extension Organization (AREEO), Tehran, Islamic Republic of Iran.

<sup>&</sup>lt;sup>3</sup>Department of Plant Protection, Faculty of Agriculture, Shahid Bahonar University, Kerman, Islamic Republic of Iran.

<sup>\*</sup>Corresponding author: email: atalebi@modares.ac.ir.

neighboring country (Turkey) (Beyarslan and Fischer, 2013), indicating that further studies are required to better understand the real fauna of Opiinae in Iran. This paper presents a faunistic study on the genus *Opius* in the north of Iran, including six new species records.

#### MATERIALS AND METHODS

Adult wasps were collected using Malaise traps from various habitats (orchards, rangelands, field crops, and forests) with different altitudes in northern parts of Iran including Mazandaran, Guilan, Qazvin, Tehran and Alborz Provinces (Figure 1) during March-November 2010-2011.

The specimens were extracted from the Malaise traps weekly (Figure 2). They were then preserved in 70% ethanol. After drying on a filter paper, the specimens were card-mounted and labeled. Images from specimens were taken using an Olympus<sup>TM</sup> AX70 microscope and an Olympus<sup>TM</sup> SZX9 stereomicroscope equipped with a Canon (EOS 550D) digital camera. A series of 4–5

captured images were stacked into a single in-focus image using Combine ZP 1.0 terminology software. The for and morphological characters wing venations follows Sharkey and Wharton (1997) (Figure 3). The term precoxal sulcus is employed instead of sternaulus accordingly to Wharton (2006). The specimens were identified at species level using the keys provided by van Achterberg (1997), Fischer (1972, 1974, 1991, 1995, 1996, 1998, 1999), Fisher and Beyarslan (2005a, b), Papp (1978, 1979, 1982a), and Tobias and Jakimavicius (1986). The specimens were deposited in the insect collection of the Entomology, Department of Tarbiat Modares University, Tehran, Iran (TMUC).

#### RESULTS

In this study, a total of 17 species of the genus *Opius* Wesmael were identified, of which six species, namely: *O.* (*Allotypus*) saevus Haliday, 1837, *O.* (*Hypocynodus*) larissa Fischer, 1968, *O.* (*Merotrachys*)



Figure 1. Geographical location of four provinces (Alborz, Tehran, Guilan, Mazandaran, and Qazivin) in northern region of Iran.



**Figure 2.** Habitats of northern Iran, where the braconid specimens were collected: (A) Gaznasara (Mazandaran Province), (B and C) Qazichak (Guilan Province), and (D) Zereshk Road (Qazvin Province).



**Figure 3.** Front wing of *Opius saevus* illustrating longitudinal and cross veins (A), and wing cells (B), parastigma (psi), stigma (stg), basal cell (bas), sub-basal cell (sba), 1<sup>st</sup> sub-marginal cell (1sm), 2<sup>nd</sup> sub-marginal cell (2sm), 3<sup>rd</sup> sub-marginal cell (3sm), marginal cell (mar), 1<sup>st</sup> discal cell (1di), 1<sup>st</sup> sub-discal cell (1sd), 2<sup>nd</sup> discal cell (2di), 2<sup>nd</sup> sub-discal cell (2sd), anal cell (ana), jugal lobe (jug).

schmidti Fischer, 1960, O. (Misophthora) pulicariae Fischer, 1969, O. (Thoracosema) repentinus Papp, 1980 and O. (Nosopaeopius) ochrogaster Wesmael, 1835 are reported for the first time from Iran. A key for the identification of the collected species as well as the general distribution of each species, either in the world or within Iran, are provided.

#### Genus Opius Wesmael, 1835

#### Subgenus Agnopius Fischer, 1982 O. (Agnopius) basirufus Fischer, 1958 [Figures 12 (A-C) and 14-B]

Materials Examined: Iran, Alborz Province: Sarziarat ( $35^\circ 55' 10''$  N,  $51^\circ 06' 51''$  E, 1,980 masl), 06.VII.2010, 2  $\bigcirc$ ; leg. M. Khayrandish; Guilan Province: Orkom ( $36^\circ 45' 44''$  N  $50^\circ 18' 11''$  E, 1,201 masl), 22.VI.2010, 1  $\bigcirc$ ; leg. M. Khayrandish; Mazandaran Province: Chamestan, Gaznasara road ( $36^\circ 18' 54''$  N,  $52^\circ 07' 48''$ E, 1,291 masl), 05.XI.2011, 1  $\bigcirc$ ; leg. M. Khayrandish.

Distribution in Iran: Alborz, Guilan, and Mazandaran (current study), Golestan Province (Sakenin *et al.*, 2012), Lorestan Provinces (Farahani *et al.*, 2016).

General Distribution: Palaearctic Region (Yu *et al.*, 2016).

#### *O.* (*Agnopius*) *periclymenii* Fischer, 1964 [Figures 10 (D-F) and 15-E]

Examined: Materials Mazandaran Province: Chamestan, Gaznasara Road (36° 18' 54" N, 52° 07' 48" E, 1,291 masl), 28.VI.2011, 1  $\mathcal{Q}$ ; Chamestan, Gaznasara Road (36° 18' 54" N, 52° 07' 48" E, 1,291 07.VI.2011, 1  $\bigcirc$ ; Chamestan, masl), Gaznasara Road (36° 18' 54" N, 52° 07' 48" E, 1,291 masl), 16.VIII.2011, 1  $\mathcal{Q}$ ; Tangehvaz (36° 21′ 54″ N, 52° 06′ 12″ E, 687 masl), 13.VII.2011, 1  $\bigcirc$ ; Chamestan, Gaznasara Road (36° 18' 54" N, 52° 07' 48" E, 1,291 masl), 16.VIII.2011, 1  $\bigcirc$ ; leg. M. Khavrandish.

Distribution in Iran: Golestan (Sakenin *et al.*, 2012), Lorestan (Farahani *et al.*, 2016) and Mazandaran Provinces (current study).

General Distribution: Palaearctic Region (Yu *et al.*, 2016).

*O.* (*Agnopius*) *tirolensis* Fischer, 1958 [Figures 11 (C-D) and 15-H]

Materials Examined: Iran, Guilan Province: Orkom (36° 45' 44" N, 50° 18' 11" E, 1,201 masl), 17.V.2010, 1  $\bigcirc$ ; leg. M. Khayrandish.

Distribution in Iran: Guilan (Farahani *et al.*, 2016, current study) and Hormozgan Provinces (Queshm Island) (Ameri *et al.*, 2014).

General Distribution: Palaearctic Region (Yu *et al.*, 2016).

Subgenus Allophlebus Fischer, 1972

*O.* (*Allophlebus*) *singularis* Wesmael, **1835** (Figures 13-I and 14-H)

Materials examined: Guilan Province: Orkom (36° 45' 42" N, 50°18'12" E, 1,235 masl), 22.IV.2010, 1 $\Im$ ; Iran, Mazandaran Province: Tangehvaz (36° 21' 55" N, 52° 06' 10" E, 702 masl), 29.IV.2011, 1 $\Im$ ; leg. A. Nadimi.

Distribution in Iran: Guilan (Farahani *et al.*, 2016; current study) and Mazandaran Provinces (current study).

General Distribution: Palaearctic Region (Yu *et al.*, 2016).

Subgenus Allotypus Forster, 1862

*O.* (*Allotypus*) saevus Haliday, 1837 [Figure 4 (A-I) and 14-G]

Materials Examined: Iran, Mazandaran Province: Nour (36° 34' 52" N, 52° 02' 45" E, 14 masl 1.), 12.IV.2011, 1  $\bigcirc$ ; leg. A. Nadimi.

Distribution in Iran: Mazandaran Province (current study), new record in Iran.

General Distribution: Palaearctic Region (Yu *et al.*, 2016).

Diagnosis: Body 2.8 mm; antennae 1.5 times as long as body, with 28 segments in females, flagellar segments short; oral cavity not developed (Figure 4-A); mandibles basally uniformly broadened (Figure 4-A); temples behind eyes roundly narrowed (Figure 4-C); thorax 1.2 times as long as high (Figure 4-D); postero-median area of the mesoscutum with rounded midpit (Figure 4-E); propodeum rugose or with granulate sculpture (Figure 4-F); precoxal sulcus absolutely smooth (Figure 4-D); mcu vein on forewing interstitial (Figure 4-I); 3-1A vein directed obliquely outward; marginal cell on forewing reaching apex (Figure 4-I); 3RSa vein longer than 2RS vein (Figure 4-I); stigma cuneate (Figure 4-I); hind femur 4.5 times as long as wide (Figure 4-H); metasoma monochromatic, dark (Figure 4-G); first metasomal tergite longer than its width at apex, lacking distinct spiraclular tubercle (Figure 4-G); ovipositor exerted to 1/3 length of abdomen.

#### Subgenus Cryptonastes Forster, 1862

*O.* (*Cryptonastes*) *gracilis* Fischer, 1957 [Figures 11 (E-F) and 15-B]

Materials Examined: Alborz Province: Karaj ( $35^{\circ} 46' 20'' N$ ,  $50^{\circ} 56' 44'' E$ , 1,278 masl), 13.X.2010, 1  $\bigcirc$ ; Guilan Province: Ziaz ( $36^{\circ} 52' 27'' N$ ,  $50^{\circ} 13' 24'' E$ , 490



**Figure 4.** *Opius saevus*: (A) Frontal aspect of head, (B) Lateral aspect of head and compound eye, (C) Dorsal aspect of head, (D) Lateral aspect of mesosoma, (E) Dorsal aspect of mesosoma, (F) Propodeum, (G) Dorsal aspect of abdomen, (H) Hind leg, and (I) Fore wing.

masl), 29.VIII.2010, 1  $\bigcirc$ ; Tehran Province: Shahriar (35° 40' 03" N, 50° 56' 52" E, 1,168 masl), 01.VI.2010, 1  $\bigcirc$ ; leg. M. Khayrandish.

Distribution in Iran: Kerman (Safahani *et al.*, 2018), Alborz, Guilan and Tehran Provinces (current study).

General Distribution: Palaearctic Region (Yu *et al.*, 2016).

#### Subgenus Gastrosema Fischer, 1972 O. (Gastrosema) docilis Haliday, 1837 [Figures 13 (E-F) and 14-C]

Materials Examined: Mazandaran Province: Tangehvaz ( $36^{\circ} 21' 54'' N$ ,  $52^{\circ} 06' 12'' E$ , 687 masl), 05.XI.2011,  $1 \ Q$ ; Nour ( $36^{\circ} 34' 54'' N$ ,  $52^{\circ} 02' 48'' E$ , -14 masl), 10.X.2011,  $1 \ Q$ ; Jourband ( $36^{\circ} 26' 18'' N$ ,  $52^{\circ} 07' 12'' E$ , 275 masl), 26.V.2011,  $1 \ Q$ ; Tangehvaz ( $36^{\circ} 21' 54'' N$ ,  $52^{\circ} 06' 12'' E$ ,



**Figure 5.** *Opius larissa*: (A) Frontal aspect of head, (B) Lateral aspect of head and compound eye, (C) Dorsal aspect of head, (D) Lateral aspect of mesosoma, (E) Dorsal aspect of mesosoma, (F) Propodeum, (G) Dorsal aspect of abdomen, (H) Hind leg, and (I) Fore wing.

687 masl), 16.VIII.2011, 1 ♀; (36° 26' 18" N, 52° 07' 12" E, 275 masl), 13.VII.2011, 1 ♀; (36° 26′ 18″ N, 52° 07′ 12″ E, 275 masl), 25.IV.2011, 1 ♀; Tangehvaz (36° 21' 54" N, 52° 06′ 12″ E, 687 masl), 13.VII.2011, 1 ♀; Tangehvaz (36° 21' 54" N, 52° 06' 12" E, 687 masl), 22.IV.2011, 1 ♀; (36° 26' 18" N, 52° 07′ 12″ E, 275 masl), 26.X.2011, 1 ♀; (36° 26' 18" N, 52° 07' 12" E, 275 masl), 26.IX.2011, 1 ♀; Tangehvaz (36° 21' 54" N, 52° 06′ 12″ E, 687 masl), 26.V.2011, 1 ♀; Nour (36° 34' 54" N, 52°02'48" E, -14 masl), 25.IX.2011, 1 ♀; Tangehvaz (36° 21' 54" N, 52° 06' 12" E, 687 masl), 28.VI.2011, 1 ♀; Tangehvaz (36° 21' 54" N, 52° 06′ 12″ E, 687 masl), 13.VII.2011, 2 ♀; Tangehvaz (36° 21′ 54″ N, 52° 06′ 12″ E,



– Dolati et al.



**Figure 6.** *Opius schmidti*: (A) Frontal aspect of head, (B) Lateral aspect of head and compound eye, (C)D aspect of head, (D) Lateral aspect of mesosoma, (E) Dorsal aspect of mesosoma, (F) Propodeum, (G) Dorsal aspect of first metasomal tergite, (H) Hind leg, (I) Fore wing, and (J) Dorsal aspect of abdomen.

**Figure 7.** *Opius pulicariae*: (A) Frontal aspect of head, (B) Lateral aspect of head and compound eye, (C) Dorsal aspect of head, (D) Lateral aspect of mesosoma, (E) Dorsal aspect of mesosoma, (F) Propodeum, (G) Dorsal aspect of abdomen, (H) Hind leg, and (I) Fore wing.

07' 12" E, 275 masl), 26.V.2011, 1  $\bigcirc$ ; leg. M. Khayrandish.

Distribution in Iran: Kermanshah (Farahani *et al.*, 2016) and Mazandaran Provinces (current study).

General Distribution: Palaearctic Region (Yu et al., 2016).

Subgenus Hypocynodus Forster, 1862 O. (Hypocynodus) larissa Fischer, 1968 [Figures 5 (A-I) and 14-D]

Materials Examined: Iran, Alborz Province: Sarziarat ( $35^{\circ} 55' 10''$  N,  $51^{\circ} 06' 51''$  E, 1,980 masl), 15.VI.2010, 2  $\bigcirc$ ; 22.VI.2010, 3  $\bigcirc$ ; 28.VII.2010, 1  $\bigcirc$ ; Guilan Province: Ziaz ( $36^{\circ} 52' 27''$  N,  $50^{\circ} 13' 24''$  E, 490 masl), 21.XI.2010, 1  $\bigcirc$ ; leg. M. Khayrandish.

Distribution in Iran: Alborz and Guilan Provinces (current study), new record for Iran.

General Distribution: Palaearctic Region (Papp, 1982b).

Diagnosis: Body 1.5-1.9 mm; clypeus narrow, falcate (Figure 5-A); antennae barely longer than body, 23-25 segmented; oral cavity developed (Figure 5-A); mandibles basally distinctly broadened (Figure 5-A), thorax 1.3-1.4 times as long as high (Figure 5-D); postero-median area of mesoscutum without midpit (Figure 5-E);



**Figure 8.** *Opius ochrogaster*: (A) Frontal aspect of head, (B) Lateral aspect of head and compound eye, (C) Dorsal aspect of head, (D) Lateral aspect of mesosoma, (E) Dorsal aspect of mesosoma, (F) propodeum, (G) Dorsal aspect of first metasomal tergite, (H) Hind leg, (I) Fore wing, and (J) Dorsal aspect of abdomen.



**Figure 9.** *Opius repentinus*: (A) Frontal aspect of head, (B) Lateral aspect of head and compound eye, (C) Dorsal aspect of head, (D) Lateral aspect of mesosoma, (E) Dorsal aspect of mesosoma, (F) Propodeum, (G) Dorsal aspect of first metasomal tergite, (H) Hind leg, (I) Fore wing, and (J) Dorsal aspect of abdomen.



**Figure 10.** (A-C) *Opius wachsmanni*: (A) Frontal aspect of head, (B) Fore wing, (C) Dorsal aspect of first metasomal tergite; (D-F) *Opius periclymenii*: (D) Dorsal aspect of mesosoma, (E) Dorsal aspect of first metasomal tergite, (F) Lateral aspect of mesosoma.



**Figure 11.** (A-B) *Opius basalis*: (A) Front wing, (B) Dorsal aspect of mesosoma; (C-D) *Opius tirolensis*: (C) Front wing, (D) hind leg; (E-F) *Opius gracilis*: (E) Frontal aspect of head, (F) dorsal aspect of first metasomal tergite.



**Figure 12.** (A-C) *Opius basirufus*: (A) Frontal aspect of head, (B) Dorsal aspect of abdomen, (C) Dorsal aspect of first metasomal tergite; (D-F) *Opius maculipes*: (D) Dorsal aspect of mesosoma, (E) Dorsal aspect of head, (F) Lateral aspect of mesosoma.

New Records of the Genus Opius from Iran here -



**Figure 13.** (A-D) *Opius ambiguous*: (A) Dorsal aspect of mesosoma, (B) Dorsal aspect of abdomen, (C) Fore wing, (D) Frontal aspect of head; (E-F) *Opius docilis*: (E) Propodeum, (F) Fore wing; (G-H) *Opius propodealis*: (G) Propodeum, (H) Frontal aspect of head; (I) *Opius singularis*: (I) Fore wing.



**Figure 14.** Dorsal habitus of female: (A) *Opius ambiguus*, (B) *Opius basirufus*, (C) *Opius docilis*, (D) *Opius larissa*, (E) *Opius propodealis*, (F) *Opius repentinus*, (G) *Opius saevus*, (H) *Opius singularis*, and (I) *Opius wachsmanni*.

precoxal sulcus smooth or not developed (Figure 5-D); marginal cell on forewing terminating apical wing; r vein originating from basal third of stigma, 3RSa 1.5 times as long as 2RS vein, 3RSb curved, 3 times as long as 3RSa (Figure 5-I); legs light yellow, hind femur 3-3.5 times as long as wide (Figure 5-H); propodeum smooth (Figure 5-F); metasoma red (Figure 5-G); metasomal posterior to 1<sup>st</sup> tergite smooth; first metasomal tergite as long as its width at apex (Figure 5-G); ovipositor slightly exerted.

Subgenus *Merotrachys* Fischer, 1972 O. (*Merotrachys*) schmidti Fischer, 1960

[Figures 6 (A-J) and 15-G]

Materials Examined: Iran, Guilan Province: Orkom (36° 45' 44" N, 50° 18' 11" E, 1,201 masl), 07.VI.2010, 1  $\bigcirc$ ; leg. A. Nadimi.

Distribution in Iran: Guilan Province (current study), new record for Iran.

General Distribution: Hungary and Turkey (Yu *et al.*, 2016).

Remark: *Opius schmidti* has very narrow distribution range in the Palaearctic Region, so far known from Hungary and Turkey, and Iran is added in the current study.

Diagnosis: Body length 1.5 mm; antennae 1.5 times as long as body, 24 segmented; oral cavity developed (Figure 6-A); temples shorter than eyes (Figure 6-C); postero-



**Figure 15.** Lateral habitus of female: (A) *Opius basalis*, (B) *Opius gracilis*, (C) *Opius maculipes*, (D) *Opius ochrogaster*, (E) *Opius periclymenii*, (F) *Opius pulicariae*, (G) *Opius schmidti*, and (H) *Opius tirolensis*.

median area of mesoscutum without midpit (Figure 6-E); precoxal sulcus smooth (Figure 6-D); m-cu vein on forewing interstitial, 3RSa vein longer than 2RS vein (Figure 5-I); 1cu-a slightly postfurcal (Figure 6-I); hind femur 5 times as long as wide (Figure 6-H); propodeum sculptured (Figure 6-F); spiracles on propodeum and first metasomal tergite distinctly raised on tubercles (Figure 6-G); metasoma posterior to 1<sup>st</sup> tergite sculptured (Figure 6-J); body black (Figure 6).

#### Subgenus Misophthora Forster, 1862

*O.* (*Misophthora*) basalis Fischer, 1958 [Figures 11 (A-B) and 15-A]

Materials Examined: Iran, Alborz Province: Karaj (35° 46' 08" N, 50° 56' 55" E, 1277 masl), 24.IX.2010, 1  $\bigcirc$ ; Mazandaran Province: (36° 26' 18" N, 52° 07' 12" E, 275 masl), 10.X.2011, 1  $\bigcirc$ ; leg. M. Khayrandish.

Distribution in Iran: Mazandaran (Khajeh *et al.*, 2014) and Alborz Provinces (current study).

General Distribution: Palaearctic Region (Yu *et al.*, 2016).

*O.* (*Misophthora*) *pulicariae* Fischer, **1969** [Figures 7 (A-I) and 15-F]

Materials Examined: Iran, Guilan Province: Ziaz (36° 52' 27" N, 50° 13' 24" E, 490 masl), 08.XI.2010, 3  $\bigcirc$ ; 26.IX.2010, 2  $\bigcirc$ ; 18.X.2010, 1  $\bigcirc$ ; 19.IV.2010, 2  $\bigcirc$ ; 19.IX.2010, 1  $\bigcirc$ ; 04.X.2010, 3  $\bigcirc$ ; 25.IV.2010, 1  $\bigcirc$ ; Orkom (36° 45' 44" N, 50° 18' 11" E, 1,201 masl), 04.X.2010, 4  $\bigcirc$ ; 31.V.2010, 1  $\bigcirc$ ; Qazichak (36° 45' 57" N, 50° 19' 35" E, 1,803 masl), 06.VI.2010, 3  $\bigcirc$ ; 28.V.2010, 2  $\bigcirc$ ; leg. M. Khayrandish.

Distribution in Iran: Guilan Province (current study), new record for Iran.

General Distribution: Palaearctic Region (Yu *et al.*, 2016).

Diagnosis: Body 1.9-2.2 mm; head more than 2.0 times as wide as long (Figure 7-C); oral cavity not developed (Figure 7-A); mandibles basally steplike broadened (Figure 7-A); antennae 1.4-1.5 times as long as body, 24-27 segmented; mesosoma lacking red coloration, 1.3 times as long as high (Figure 7-D); notaulus distinct only in middle (Figure 7-E); precoxal sulcus absolutely smooth or not developed (Figure 7-D); postero-median area of mesoscutum with elongated midpit (Figure 7-E); stigma cuneate (Figure 7-I); hind wing lacking stigma thickening; 3RSa vein longer than 2RS vein (Figure 7-I); 3RSb of radial vein terminating or almost terminating at wing apex (Figure 7-I); propodeum at least in anterior part smooth or only with ridges (Figure 7-F); first metasomal tergite rugose, 1.3 times as long as its width at apex (Figure 7-G); ovipositor barely exerted from abdominal apex (Figure 7-G).

O. (*Misophthora*) wachsmanni Szepligeti, 1898 [Figures 10 (A-C) and 14-I) Materials Examined: Alborz Province: Arangeh ( $35^{\circ} 55' 07''$  N,  $51^{\circ} 05' 09''$  E, 1,891 masl), 31.VIII.2010, 1  $\bigcirc$ ; Guilan Province: Orkom ( $36^{\circ} 45' 44''$  N,  $50^{\circ} 18'$ 11'' E, 1,201 masl), 07.VI.2010, 1  $\bigcirc$ ; 17.V.2010, 1  $\bigcirc$ ; Qazvin Province: Zereshk ( $36^{\circ} 25' 42''$  N,  $50^{\circ} 06' 36''$  E, 2,035 masl), 04.IV.2011, 1  $\bigcirc$ ; leg. M. Khayrandish.

Distribution in Iran: Lorestan (Farahani *et al.*, 2016), Alborz, Guilan, and Qazvin Provinces (current study).

General Distribution: Palaearctic Region (Yu et al., 2016).

Subgenus Nosopoea Forster, 1862

*O.* (*Nosopoea*) *ambiguus* Wesmael, 1835 [Figures 13 (A-D) and 14-A]

Materials Examined: Mazandaran Province: Tangehvaz (36° 21' 54" N, 52° 06' 12" E, 687 masl), 07.VI.2011, 1  $\bigcirc$ ; leg. M. Khayrandish.

Distribution in Iran: Isfahan (Farahani *et al.*, 2016) and Mazandaran Provinces (current study).

General distribution: Palaearctic region (Yu *et al.*, 2016).

*O.* (*Nosopoea*) *maculipes* Wesmael, 1835 [Figures 12 (D-F) and 15-C]

Materials Examined: Guilan Province: Orkom (36° 45' 44" N, 50° 18' 11" E, 1,201 masl), 07.VI.2010, 1  $\bigcirc$ ; Ziaz (36° 52' 27" N, 50° 13' 24" E, 490 masl), 17.V.2010, 1  $\bigcirc$ ; Mazandaran Province: (36° 26' 18" N, 52° 07' 12" E, 275 masl), 26.IX.2011, 1  $\bigcirc$ ; leg. A. Nadimi.

Downloaded from jast.modares.ac.ir on 2024-09-25

Distribution in Iran: East Azarbaijan, Guilan, and Tehran (Gadallah *et al.*, 2016), and Mazandaran Provinces (current study).

General distribution: Palaearctic Region (Yu et al., 2016).

#### Subgenus Nosopaeopius Fischer, 1972 O. (Nosopaeopius) ochrogaster Wesmael, 1835 [Figures 8 (A-J) and 15-D]

Materials Examined: Alborz Province: Arangeh (35° 55' 07" N, 51° 05' 09" E, 1,891 masl), 16.VIII.2010, 1  $\bigcirc$ ; 17.IX.2010, 1 ♀; Sarziarat (35° 55' 10" N, 51° 06' 51" E, 1,980 masl), 08.VI.2010, 1 ♀; Guilan Province: Orkom (36° 45' 44" N, 50° 18' 11" E, 1,201 masl), 31.V.2010, 2  $\Im$ ; 07.VI.2010, 1 ♀; Qazichak (36° 45' 52" N, 50° 20′ 01″ E, 1,787 masl), 17.V.2010, 1 ♀; Mazandaran Province: Tangehvaz (36° 21' 54" N, 52° 06' 12" E, 687 masl), 10.X.2011, 1 ♀; (36° 26' 18" N, 52° 07' 12" E, 275 masl), 03.X.2011, 1 2; Gaznasara (36° 16' 54" N, 52° 10' 54" E, 2,031 masl), 26.VII.2011, 1 ♀; (36° 26' 18" N, 52° 07' 12" E, 275 masl), 10.X.2011, 1 ♀; (36° 26' 18" N, 52° 07' 12" E, 275 masl), 05.XI.2011, 1  $\bigcirc$ ; leg. M. Khayrandish.

Distribution in Iran: Alborz, Guilan and Mazandaran Provinces (current study), new record for Iran.

General Distribution: Palaearctic Region (Yu *et al.*, 2016).

Diagnosis: Body 1.8-2.3 mm; oral cavity not developed (Figure 8-A); antennae 1.5 times as long as body, 24-29 segmented; mesonotum with midpit (Figure 8-E); scutellum smooth or pubescent or posteriorly finely rugose (Figure 8-E); midpit somewhat elongate (Figure 8-E); propodeum sculptured, rugose or granulate (Figure 8-F); precoxal sulcus absolutely smooth or not developed (Figure 8-D); 3RSa vein up to 1.5 times as long as 2RS vein, 3RSb straight or sometimes arcuate, terminating preapically (Figure 8-I); m-cu vein postfurcal (Figure 8-I); first metasomal tergite longitudinally rugose and black (Figure 8-G), other tergites yellow or dark brown (Figure 8-J), apical tergites often with black transverse stripes (Figure 8-J), in males sometimes entirely black.

Subgenus Opiognathus Fischer, 1972

*O.* (*Opiognathus*) *propodealis* Fischer, **1958** [Figures 13 (G-H) and 14-E]

Materials Examined: Guilan Province: Ziaz (36° 52' 27" N, 50° 13' 24" E, 490 masl), 10.V.2010, 1  $\bigcirc$ ; Mazandaran Province: Tangehvaz (36° 21' 54" N, 52° 06' 12" E, 687 masl), 05.IX.2011, 1 ♀; (36° 26' 18" N, 52° 07' 12" E, 275 masl), 05.XI.2011, 1 ♀; (36° 26' 18" N 52° 07' 12" E, 275 masl), 26.IX.2011, 1 ♀; Tangehvaz (36° 21' 54" N, 52° 06′ 12″ E, 687 masl), 05.IX.2011, 1 ♀; (36° 26' 18" N, 52° 07' 12" E, 275 masl), 26.IX.2011, 1 ♀; Tangehvaz (36° 21' 54" N, 52° 06′ 12″ E, 687 masl), 16.VIII.2011, 1 ♀; (36° 26' 18" N, 52° 07' 12" E, 275 masl), 26.IX.2011, 1 ♀; (36°26′18″ N, 52°07′12″ E, 275 masl), 05.IX.2011, 1 ♀; (36° 26' 18" N, 52° 07′ 12″ E, 275 masl), 16.VIII.2011, 1 ♀; leg. M. Khayrandish.

Distribution in Iran: Kermanshah (Farahani *et al.*, 2016), Guilan and Mazandaran Provinces (current study).

General Distribution: Palaearctic Region (Yu *et al.*, 2016).

Subgenus *Thoracosema* Fischer, 1972

*O.* (*Thoracosema*) *repentinus* Papp, 1980 [Figures 9 (A-J) and 14-F]

Materials Examined: Mazandaran Province: Tangehvaz (36° 21′ 54″ N 52° 06′ 12″ E, 687 masl), 16.VIII.2011, 1  $\bigcirc$ ; Tangehvaz (36° 21′ 54″ N, 52° 06′ 12″ E, 687 masl), 16.VIII.2011, 1  $\bigcirc$ ; Tangehvaz (36° 21′ 54″ N, 52° 06′ 12″ E, 687 masl), 16.VIII.2011, 1  $\bigcirc$ ; Chamestan-Gaznasara road (36° 18′ 54″ N, 52° 07′ 48″ E, 1,291 masl), 12.VII.2011, 1  $\bigcirc$ ; Chamestan-Gaznasara road (36° 18′ 54″ N, 52° 07′ 48″ E, 1,291 masl), 16.VIII.2011, 1  $\bigcirc$ ; Tangehvaz (36° 21′ 54″ N, 52° 06′ 12″ E, 687 masl), 16.VIII.2011, 1  $\bigcirc$ ; leg. M. Khayrandish.

Distribution in Iran: Mazandaran Province (current study), new record for Iran.

General Distribution: Palaearctic Region (Yu *et al.*, 2016).

Diagnosis: Body 1.2-1.5 mm, very dark brown (Figure 9); head dorsally smooth, 2 times as wide as long (Figure 9-C); clypeus 3 times as wide as high (Figure 9-A); antennae longer than body, 15-16 segmented; oral cavity developed (Figure 9-A); mandibles basally steplike broadened (Figure 9-A); thorax slightly longer than high; posteromedian area of mesoscutum without midpit (Figure 9-E); notaulus distinct only anteriorly (Figure 9-E); precoxal sulcus sculptured (Figure 9-D); stigma cuneate (Figure 9-I); r vein short, 3RSa 1.7-1.8 times as long as 2RS vein (Figure 9-I); legs yellow; hind femur 4-5 times as long as wide (Figure 9-H); propodeum lacking transverse ridge, smooth or only on posterior margin slightly sculptured, finely rugose in posterior margin (Figure 9-F); first metasomal tergite slightly longer than its width at apex, smooth or with slightly wrinkles in apical part, with slightly developed light colored patterns (Figure 9-G); ovipositor slightly exerted.

Key to the Species of the Genus *Opius* Collected from Northern Iran

1- Postero-median area of mesoscutum with rounded or elongated midpit (Figures 4-E, 7-E, 8-E,) .....

2- Postero-median area of mesoscutum without midpit (Figures 5-E, 6-E, 9-E) .... 12

3 (2)- Propodeum at least in anterior part smooth or only with ridges (Figure 7-F); [midpit is rounded (Figure 4E, 8E); mesosoma usually 1.3 times as long as high (Figures 4-D and 7D); stigma cuneate (Figures 4-I and 7-I)] ......4

4 (3)- Hind wings of males with pterostigma thickening between costal and radial veins; mandibles basally uniformly broadened (Figure 10-A); [antennae with 24-26 segments; marginal cell on forewing almost reaching its apex (Figure 10B); first metasomal tergite somewhat longer than its width at apex (Figure 10-C)]

..... Opius (Misophthora) wachsmanni

- Hind wings of male without pterostigma thickening between costal and radial veins .. 5

- Mesosoma without red coloration or only prothorax sometimes reddish (Figure 7-E)

6 (5)- First metasomal tergite coarsely sculptured, 1.3 times as long as its width at apex (Figure 7-G); [antennae 24-27 segmented]....

7 (6)- Radial vein on forewing not reaching its apex (Figure 11-A); [3RSa vein 1.7 times as long as 2RS vein, 3RSb 2.4 times as long as 3RSa (Figure 11-A); midpit punctiform (Figure 11-B)] *Opius (Misophthora) basalis* 

- 3RSa vein 2 times or more as long as 2RS vein; hind femur 5 times as long as wide .

9 (8) Mandibles basally steplike (Figure 11-E); first metasomal tergite as long as its width at apex, yellow or yellowish dark brown (Figure 11-F); [clypeus 2.5 times as wide as high (Figure 11-E); Ovipositor as long as 1<sup>st</sup> metasomal tergite or longer] .....

.....*Opius (Cryptonastes) gracilis* - Mandibles basally uniformly broadened (Figure 12-A); [metasoma dark brownish red (Figure 12-B); antennae almost 2 times as long as body, with 24-27 segments; first metasomal tergite 1.5 times as long as its width at apex (Figure 12-C); ovipositor as long as 1<sup>st</sup> abdominal tergite] .....

.....Opius (Agnopius) basirufus 10 (3)- m-cu vein interstitial; 3-1A vein directed obliquely outward (Figure 4-I); [first metasomal tergite without distinct specular tubercles (Figure 4-G); ovipositor 1/3 as long as of abdomen; antennae 28-31 segmented]

11 (10)- Mesonotum fairly densely pilose (Figure 12-D); [antennae slightly longer than body, with 25-35 segments; head behind eyes slightly narrowed (Figure 12-E); mesosoma 1.3 as long as high (Figure 12-F); ovipositor slightly extends byond the apex of metasoma or concealed] ... Opius (Nosopoea) maculipes - Mesonotum absolutely glabrous or with hairs mostly on anterior sloping part (Figure 13-A); [metasoma black (Figure 13-B); 3RSa vein usually 2 times as long as 2RS vein, 3RSb 1.7-1.8 times as long as 3RSa (Figure 13-C); antennae longer than body, with 26-31 segments; face smooth (Figure 13-D)] ..... Opius (Nosopoea) ambiguous......12 (1)- Precoxal sulcus sculptured (Figure 9-D) - Precoxal sulcus smooth or not developed (Figures 5-D and 6-D)..... 15 13 (12)- Propodeum smooth, sometimes only on posterior margin slightly sculptured or almost smooth (Figure 9-F).....14 - Propodeum sculptured (Figure 13E); stigma large, triangular (Figure 13-F) ..... ..... Opius (Gastrosema) docilis 14 (13)- Propodeum with transverse ridge (Figure 13-G); [face with reddish pattern (Figure 13H)] ..... ..... Opius (Opiognathus) propodealis

15 (12)- Metasoma posterior to 1<sup>st</sup> tergite sculptured (Figure 6-G); [propodeum sculptured (Figure 6-F); temples shorter than eye (Figure 6-C); fore wing with m-cu vein interstitial (Figure 6-I); antennae 1.5 times as long as body, with 24 segments; body black.] ...... *Opius (Merotrachys) schmidti* - Metasoma posterior to 1<sup>st</sup> tergite smooth (Figure 5-G) ...... 16 16 (15)- 3RSa vein in fore wing 2 times as long as 2RS vein (Figure 13-I); Ovipositor barely exserted; antennae 27-32 segmented

- 3RSa vein in fore wing 1.5 times as long

as 2RS vein in fore wing 1.5 times as long as 2RS vein, 3RSb curved, 2 times as long as 3RSa (Figure 5-I); [legs light yellow (Figure 5-H); r vein originating from basal third of stigma (Figure 5-I); metasoma red (Figure 5-G); mandibles basally distinctly broadened (Figure 5-A); ovipositor slightly exserted].....*Opius (Hypocynodus) larissa* 

#### DISCUSSION

According to the previous studies and current research, a total of 107 species of Opiinae are known from Iran, of which 72 species belong to the cosmopolitan genus Opius. Our investigations in northern parts of Iran resulted in collection of 17 species of the genus Opius, of which six species are new records for the fauna of Iran: O. (Allotypus) saevus Haliday, 1837, O. (Hypocynodus) larissa Fischer, 1968, O. (Merotrachys) schmidti Fischer, 1960, O. (Misophthora) pulicariae Fischer, 1969, O. (Thoracosema) repentinus Papp, 1980, and O. (Nosopaeopius) ochrogaster Wesmael, 1835. Apart from this, five species (i.e., O. (*Cryptonastes*) gracilis, О. (Agnopius) periclymenii, O. (Gastrosema) docilis, O. (Nosopoea) ambiguous, and O. (Opiognathus) propodealis) are reported from the studied area for the first time. The geographical distribution of all species is limited to the Palearctic region, especially the West Palearctic and the Middle East (Tobias, 1995). According to the literature (Shaw and Huddleston, 1991; Tobias, 1995), most species of the genus, including those reported in this study, are associated with phytophagous cyclorrhaphan Diptera, especially leaf miner flies in the family Agromyzidae. At least, some species within the identified genera of host flies, such as Amauromyza, Nemorimyza, Ophiomyza, Liriomyza, Agromyza, Cerodontha, and Phytomyza, are known to associate with agricultural and forest plants, giving Opius a relatively high status in term of pest biological control (Shaw and Huddleston, 1991). Further studies may evaluate the potential of these parasitic wasps for applied biological control of leaf miner flies on agricultural and greenhouse products. Additionally, further faunistic investigations are needed to improve understanding of biodiversity and our geographical distribution of Opiine parasitic wasps in Iran.

#### ACKNOWLEDGEMENTS

Thanks are due to the Department of Entomology, Tarbiat Modares University, for providing financial support for this study. The authors would like to appreciate Dr. Nadimi for his great helps with sampling.

#### REFERENCES

- Ameri, A., Talebi, A. A., Beyarslan, A., Kamali, K. and Rakhshani, E. 2013. Study of the Genus *Bracon* Fabricius, 1804 (Hymenoptera: Braconidae) of Southern Iran with Description of a New Species. *Zootaxa*, 3754: 353–380.
- Ameri, A., Talebi, A. A., Rakhshani, E., Beyarslan, A. and Kamali, K. 2014. Study of the Genus *Opius* Wesmael (Hymenoptera: Braconidae: Opiinae) in Southern Iran, with Eleven New Records. *Zootaxa*, 3884: 1–26.
- Ameri, A., Talebi, A. A., Rakhshani, E., Beyarslan, A. and Kamali, K. 2015. Additional Evidence and New Records of the Genus *Bracon* Fabricius, 1804 (Hymenoptera: Braconidae) in Southern Iran. *Turk. J. Zool.*, **39**: 1110–1120
- Belokobylskij, S. A., Wharton, R. A. and La Salle, J. 2004. Australian Species of the Genus *Opius* Wesmael (Hymenoptera: Braconidae) Attacking Leaf-mining Agromyzidae, with the Description of a New Species from South-East Asia. *Austral. J. Entomol.*, 43: 138–147.

- Beyarslan, A. 2015. Taxonomical Investigations on the Fauna of Opiinae (Hymenoptera: Braconidae) in Ardahan, Erzurum, Iğdır, and Kars Provinces of the Turkish Northeastern Anatolian Region. *Turk. J. Zool.*, **39**: 820–831.
- Beyarslan, A. and Fischer, M. 2011. Contributions to the Opiinae Fauna of Turkey (Hymenoptera: Braconidae). *Turk. J. Zool.*, 35: 293–305.
- 7. Beyarslan, A. and Fischer, M. 2013. Checklist of Turkish Opiinae (Hymenoptera, Braconidae). *Zootaxa*, **3721**: 401–454.
- 8. Fallahzadeh, M. and Saghaei, N. 2010. Checklist of Braconidae (Insecta: Hymenoptera) from Iran. *Munis Entomol. Zool.*, **5**: 170–186.
- Farahani, S., Talebi, A. A. and Rakhshani, E. 2014a. Wasps of the Subfamily Doryctinae (Hymenoptera: Braconidae) in Iran. Zool. Middle East, 60: 65–81.
- Farahani, S., Talebi, A. A. and van Achterberg, C. 2014b. A Review of Species of the Genus Ascogaster Wesmael (Hymenoptera: Braconidae, Cheloninae) from Iran. Far East. Entomol., 275: 1–12.
- Farahani, S., Talebi, A. A., van Achterberg, C. and Rakhshani, E. 2014c. A New Species of the Genus *Mirax* Haliday, 1833 (Hymenoptera: Braconidae: Miracinae) from Iran. *Ann. Zool.*, 64: 677–682.
- Farahani, S., Talebi, A. A., van Achterberg, C. and Rakhshani, E. 2015. A Review of the Subfamily Rogadinae (Hymenoptera: Braconidae) from Iran. *Zootaxa*, **3973**: 227– 250.
- Farahani, S., Talebi, A. A. and Rakhshani, E. 2016. Iranian Braconidae (Insecta: Hymenoptera: Ichneumonoidea): Diversity, Distribution and Host Association. J. Insect Biodiv. Syst., 2: 1–92.
- Farrar, N., Golestaneh, R., Askari, H. and Assareh, M. H. 2009. Studies on Parasitism of *Fopius carpomyie* (Silvestri) (Hymenoptera: Braconidae), an Egg-Pupal Parasitoid of Ber (Konar) Fruit Fly, *Carpomya vesuviana* Costa (Diptera: Tephritidae), in Bushehr Iran. *Acta Hortic.*, 840: 431–438.
- 15. Fischer, M. and Beyarslan, A. 2005a. New Maggot Wasps from the Turkish Fauna (Hymenoptera, Braconidae, Opiinae). *Pol. Pismo Entomol.*, **74**: 379–421.
- Fischer, M. and Beyarslan, A. 2005b. A Survey of Opiinae (Hymenoptera, Braconidae) of Turkey. *Fragm. Faun.*, 48: 27–62.

Downloaded from jast.modares.ac.ir on 2024-09-25

- 17. Fischer, M. 1972. Hymenoptera, Braconidae (Opiinae I). *Das Tierreich*, **91**: 1–620.
- Fischer, M. 1974. Opiinae aus dem Ötztal (Österreich) (Hymenoptera: Braconidae). *Beitr. Entomol.*, 24: 73–86.
- Fischer, M. 1991. Wiederbeschreibungen und Neubeschreibungen von Opiinae aus der Alten Welt (Hymenoptera: Braconidae). *Ann. Nat. His. Mus. Wien*, **92**: 139–203.
- Fischer, M. 1995. Korrekturen und Ergänzungen zur Taxonomie altweltlicher Opiinae und Neufassung eines Bestimmungsschlüssels für die Paläarktischen Arten des Subgenus Opiothorax Fischer, M(1972) des Genus Opius Wesmael, 1835 (Hymenoptera, Braconidae). Entomofauna, 16: 217–244.
- Fischer, M. 1996. Beschreibungen und Wiederbeschreibungen von Einigen Europäischen und Kanarischen Opiinae (Hymenoptera: Braconidae). Z. Arbeitsgem. Österr. Entomol., 48: 49–62.
- Fischer, M. 1998. Neue Taxonomische Untersuchungen über Madenwespen der Alten Welt mit Besonderer Berücksichtigung der Gattungen Eurytenes Förster, Aulonotus Ashmead, <u>Biosteres</u> Förster und der Untergattung Gastrosema Fischer (Hymenoptera, Braconidae: Opiinae). Linzer Biol. Beitr., 30: 21–51.
- Fischer, M. 1999. Zur Evolution und zum System der Opiusverwandten Gatt ungen der Unterfamilie Opiinae Miteiner Erweiterten Auft Eilung Dieses Gattungs-Komplexes (Hymenoptera, Braconidae, Opiinae. *Linzer Biol. Beitr.*, 31: 277–336.
- Gadallah, N. S., Ghahari, H., Peris-Felipo F. J. and Fischer, M. 2016. Updated Checklist of Iranian Opiinae (Hymenoptera: Braconidae). *Zootaxa*, 4066: 1–40.
- Khajeh, N., Rakhshani, E., Peris-Felipo, F. J. and Žikić, V. 2014. Contributions to the Opiinae (Hymenoptera: Braconidae) of Eastern Iran with Updated Checklist of Iranian Species. *Zootaxa*, **3784**: 131–147.
- Lashkari Bod, A., Rakhshani, E., Talebi, A. A., Lozan, A. and Zikic, V. 2011. Contribution to the Knowledge of Braconidae (Hym., Ichneumonoidea) of Iran. *Biharean Biol.*, 5: 147–150.
- Li, X. Y., van Achterberg, C. and Tan, J. C. 2013. Revision of the Subfamily Opiinae (Hymenoptera, Braconidae) from Hunan (China), Including Thirty-Six New Species and Two New Genera. *ZooKeys*, 268: 1–168.

- Papp, J. 2010. Contribution to the Braconid Fauna of the Former Yugoslavia VI (Hymenoptera: Braconidae). *Entomofauna*, 25: 389–404.
- Papp, J. 1978. Braconidae (Hymenoptera) from Korea. 3. Folia Entomol. Hung., 24: 133– 148.
- Papp, J. 1979. Braconidae (Hymenoptera) from Tunisia. I. Folia Entomol. Hung., 32: 175–187.
- Papp, J. 1982a. Braconidae (Hymenoptera) from Greece, 1. Ann. Mus. Goulandris, 5: 183– 193.
- Papp, J. 1982b. Taxonomical and Faunistical Noveltiesm of the Opiinae in the Arctogaea (Hymenoptera, Braconidae). *Ann. His-Nat. Mus. Natl. Hung.*, 74: 241–253.
- 33. Peris-Felipo1, F. J., Ameri, A., Talebi, A. A. and Belokobylskij, S. A. 2015. Review of the Genus Aristelix Nixon, 1943 (Hymenoptera, Braconidae, Alysiinae), with Description of a New Species from Iran and Clarification of the Status of Antrusa chrysogastra (Tobias, 1986). J. Hym. Res., 45: 97–111.
- Peris-Felipo, F. J., Rahmani, Z., Belokobylskij, S. A. and Rakhshani, E. 2014. Genus *Indiopius* Fischer, 1966 (Hymenoptera, Braconidae, Opiinae) in Iran with a Key to the World Species. *ZooKeys*, 368: 37–44.
- Peris-Felipo, F. J., Belokobylskij, S. A., Derafshan, H. A. and Rakhshani, E. 2018. Revision of the Genus *Hoplocrotaphus* Telenga, 1950 (Hymenoptera, Braconidae, Opiinae). J. Hym. Res. 62: 55–72.
- Rakhshani, E., Talebi, A.A., Starý, P., Tomanović, Ž. and Manzari, S. 2007. Aphid-Parasitoid (Hymenoptera, Braconidae, Aphidiinae) Associations on Willows and Poplars in Iran. *Acta Zool. Acad. Sci. Hung.* 53: 281-292.
- Safahani, S., Madjdzadeh, S. M. and Peris-Felipo, F. J. 2018. Contribution to the Fauna and Phenological Knowledge of High Mountains Opiinae (Hymenoptera, Braconidae) in Kerman Province (Iran). J. Insect Biodiv. Syst. 4: 73–83.
- Sakenin, H., Naderian, H., Samin, N., Rastegar, J., Tabari, M. and Papp, J. 2012. On a Collection of Braconidae (Hymenoptera) from Northern Iran. *Linzer Biol. Beitr.*, 44: 1319–1330.
- Sharkey, M. J. and Wharton, R. A. 1997. Morphology and Terminology. In: "Manual of the New World genera of the family Braconidae (Hymenoptera)", (Eds.): Wharton,

R. A., Marsh, P. M. and Sharkey, M. J. Special Publication of the International Society of Hymenopterists, No 1, Washington DC, PP. 19–37.

- 40. Shaw, M. R. and Huddleston, T. 1991. Classification and Biology of Braconid Wasps (Hymenoptera: Braconidae). Handbook for the Identification of British Insects, The Natural History Museum, London, 126 PP.
- Talebi, A.A., Rakhshani, E., Fathipour, Y., Starý, P., Tomanović, Ž. And Rajabi-Mazhar, N. 2009. Aphids and their parasitoids (Hym., Braconidae: Aphidiinae) associated with medicinal plants in Iran. *Am.-Eurasian J. Sustain. Agric.*, 3(2): 205-219.
- 42. Tobias, V. I. 1995. *Keys to the Fauna of the USSR III, Hymenoptera V.* Science Publishers, Lebanon, New Hampshire, USA, 883 PP.
- Tobias, V. I. and Jakimavicius, A. 1986. Alysiinae & Opiinae. In: "Opredelitel Nasekomych Evrospeiskoi Tsasti SSSR 3,

*Peredpontdatokrylye 4*", (Eds.): Medvedev, G. S., Opr. Faune SSSR, PP. 7–231.

- van Achterberg, C. 1988. Revision of the Subfamily Blacinae Foerster (Hymenoptera, Braconidae). *Zool. Verh.*, 249: 1324.
- van Achterberg, C. 1993. Illustrated Key to the Subfamilies of the Braconidae (Hymenoptera: Ichneumonoidea). *Zool. Verh.*, 283: 1–189.
- van Achterberg, C. 1997. Revision of the Haliday Collection of Braconidae (Hymenoptera). *Zool. Verh.*, **314**: 1–115.
- Wharton, R.A. 2006. The Species of Sternaulopius Fischer (Hymenoptera: Braconidae, Opiinae) and the Braconid Sternaulus. J. Hym. Res., 15: 317–347.
- 48. Yu, D. S., van Achterberg, C. and Horstmann, K. 2016. Taxapad 2016, Ichneumonoidea 2015. *Database on Flash-Drive*. http://www.taxapad.com, Nepean, Ontario, Canada.

# Opius Wesmael (Hemenoptera: اطلاعات جدید از گونه های جنس Braconidae, Opiinae)

ث. دولتی، ع. ا. طالبی، س. فراهانی، و م. خیراندیش

### چکیدہ

در این مطالعه، فون گونههای جنس (Opius Wesmael (Hymenoptera: Braconidae در طول سالهای Opius Wesmael (Hymenoptera: Braconidae) در استانهای شمالی ایران مورد بررسی قرار گرفت. حشرات بالغ زنبورها در طول سالهای ۱۳۹۹ تا ۱۳۹۱ ۱۹ استفاده از تله مالیز از زیستگاههای مختلف جمع آوری شدند. در مجموع، ۱۷ گونه متعلق به ۱۲ ۱۹ زیرجنس جمع آوری و شناسایی شدند که از این تعداد، شش گونه شامل (Hypocynodus) . ۱۹68 (Hypocynodus) saevus Haliday, 1837 (Arissa Fischer, 1968 O. (Allotypus) saevus Haliday, 1837 (Scher, 1968 O. (Misophthora) pulicariae (Merotrachys) schmidti Fischer, 1960 O. و . (Thoracosema) repentinus Papp, 1980 (Fischer, 1969 O. (Thoracosema) برای اولین بار از ایران گزارش می شوند. کلید شناسایی برای گونههای جمع آوری شده در این مطالعه تهیه شده است.

1887