

Influence of Selected Salts and Sugars on the Rheological Behavior of Quince Seed Mucilage

A. Farahmand¹, S. Naji-Tabasi^{1*} and, S. Shahbazizadeh¹

ABSTRACT

Properties of solvent/cosolute in a food system obviously can influence the rheological properties of hydrocolloids. In this work, the impact of various concentrations of some sugars (sucrose and glucose at 10, 20, and 30% w/w concentration) and salts (CaCl₂ and NaCl, 50, 100, and 150 mM) on the rheological properties of Quince Seed Mucilage (QSM) were investigated. QSM produced high viscosity dispersion with yield stress, which was affected by concentrations and types of salt and sugar. The salts altered the electrostatic interactions between the QSM chains and led to the viscosity reduction. Increasing the salt concentration, decreased the yield stress (τ_0) and consistency coefficient (k) values, but led to increase in the flow behavior index (n) value. Sucrose and glucose increased the apparent viscosity of QSM dispersions. High sugar concentrations resulted in higher viscosity. Assessment of the time-dependent flow behavior of QSM dispersions proved that the structural kinetic model had the best data description. All the samples showed thixotropic behavior. The Atomic Force Microscope (AFM) results exhibited a proper understanding of the chain interactions of QSM and changes in the flow behavior. The results of this research showed that CaCl₂ and sucrose had more pronounced effect on the rheological properties of QSM.

Keywords: Hydrocolloid, Quince mucilage, Rheology, Structural kinetic model.

INTRODUCTION

Plant seed mucilage can be applied in the food products due to its interesting characteristics (texturizing, emulsifying, and thickening) and dietary properties. The physicochemical and functional characteristics of mucilage are a function of chemical composition, molecular configuration, and extraction and processing circumstances (Williams and Phillips, 2000).

Quince (*Cydonia oblonga* Miller, Rosaceae family) is a fruit with high nutritional and therapeutic properties. This fruit has some functional characteristics such as anticarcinogenic, antimicrobial, antioxidant, and anti-allergic activities, which are relevant to the existence of phenolic compounds (Wojdylo *et al.*, 2013; Legua *et al.*, 2013;

Sharma *et al.*, 2011).

Quince seed also has a high content of mucilage (11.58%) and is known as biocompatible and biodegradable mucilage (Jouki *et al.*, 2014; Elboutachfaiti *et al.*, 2017). Moreover, its low production cost, ready availability, easy extraction and high viscosity make Quince Seed Mucilage (QSM) attractive. QSM could be highly valuable for the development of stimuli-responsive intelligent/smart formulations for the targeted and sustained release of drug delivery systems on commercial scale because of its biocompatible nature, high swelling index, pH, salt responsiveness, and on-off switching properties (Hussain *et al.*, 2019). Average molecular weight of QSM is recorded as 9.61×10^6 g mol⁻¹, which is greater than commercially available gums such as guar gum (1.45×10^6), gellan gum (1.64×10^6) and locust bean gum (1.6×10^6) (Rezaghali *et*

¹ Department of Food Nanotechnology, Research Institute of Food Science and Technology (RIFST), P. O. Box: 91895-157.356, Mashhad, Islamic Republic of Iran.

*Corresponding author; e-mail: s.najitabasi@rifst.ac.ir



al., 2018). It can be applied in the medical, food industry, cosmetic, and pharmaceutical industries as a stabilizer (Soukoulis *et al.*, 2009). Recently, the emulsifying properties of quince seed mucilage have been extensively studied and confirmed for a pH range of 6-8 (Ritzoulis *et al.*, 2014). Jouki *et al.*, (2014) also reported that QSM consisted of 2.71% protein and its emulsion stability was 94.89%.

Rezagholi *et al.* (2018) have reported that β -1,4-D-xylan with the fraction of glucuronic acid residues is the main water-soluble polysaccharide of QSM. A mixture of aldobionic acids, arabinose, and a cellulosic segment is found in QSM structure. According to Abbastabar *et al.* (2015), QSM exhibits shear-thinning behavior and has considerable gelling capacity. They found that the high hydrodynamic volume of QSM (due to high critical concentration of 0.077 g 100 mL⁻¹ and intrinsic viscosity of 1530 mL g⁻¹), result in improving its gel production in aqueous solution. Furthermore, it is proposed that QSM has stiff conformation based on its salt tolerance value (Rezagholi *et al.*, 2018).

The pH, concentration, molecular weight, temperature, and counter-ions are the important variables that affect the rheological behavior of hydrocolloids (Kar and Arsalan, 1999). The most abundant low-molecular ingredients that are generally used as food additives are NaCl and CaCl₂. For the investigation of functional and rheological attributes of hydrocolloid, we should determine the impact of the concentration and type of salt on the functional characteristics of hydrocolloid (Koochehi *et al.*, 2009). The arrangement of polysaccharide chains and their physicochemical properties are related to the electrostatic forces (Lin and Lai, 2009).

Sucrose and glucose are the most prevalent sugars that are applied in the food products. Glucose is extensively applied in the food industry, especially in syrups, due to the higher water activity, lower kinematic viscosity and molecular weight compared to the sucrose (Imeson, 2010). The sucrose modifies the texture and flavor of the product, extends the freshness, and plays an important

role in moisture retention (Torres *et al.*, 2013).

The influence of salt and sugar types on the flow behavior of hydrocolloids is reported by other researchers. According to Farahmandfar and Naji-Tabasi (2020), alteration of electrostatic interactions between chains of basil seed gum led to the viscosity reduction in the presence of CaCl₂, NaCl, and KCl. They investigated the time-dependent rheological properties of BSG. The time-independent properties of semi-dilute sage seed gum solution were affected by the different sugars and salts (Yousefi *et al.*, 2016). Steady-shear flow behavior of semi-dilute guar gum was investigated by Chenlo *et al.* (2011) and a synergic impact on the apparent viscosity due to the addition of sugar was reported. This effect was correlated with sugar concentration. Zhang *et al.* (2007) also reported that the addition of salts (CaCl₂ and KCl) decreased the initial viscosity of semi-dilute solutions of hydroxypropyl guar gum, which caused the lowering of thixotropic properties.

Hydration and molecular hydrodynamic volume of soluble polysaccharides depend on the type of interactions (attraction or repulsion) among chain fragments. The configuration of chains in the polysaccharide structure determines the molecular volume. Inclusion of additives such as sugars and salts influence the electrostatic repulsion among chains and chains interactions (Van Aken, 2006).

Additives such as sugars and salts can alter the viscosity of hydrocolloids, which changes depend on the type and concentration of the additives. Despite great potential of additive impact on the rheological characteristics of mucilage in complex systems such as food, no comprehensive research has been reported about QSM. The objective of this work was to study the influence of different concentrations of NaCl, CaCl₂, sucrose and glucose on the rheological characteristics (shear-rate dependency and time-dependency) of QSM and to investigate the molecular structure and interactions of QSM chains with salts and sugars by using AFM.

MATERIALS AND METHODS

The quince seeds were purchased from a local market in Mashhad, Iran. Different salts (sodium chloride and calcium chloride) and sugars (sucrose and glucose) were purchased from Merck Co., Germany.

Quince Seed Mucilage Extraction

Quince seeds were cleaned manually to remove all the foreign matters. The quince seed mucilage was extracted according to the optimized conditions reported by Farahmand *et al.* (2016). Briefly, the optimized conditions were: Water/Seed ratio of 32.5:1, temperature of 47°C and time of 45 minutes. An electrical mixer (1,000 rpm) was used to mix the seed-water slurry. After the extraction, seeds were separated and recovering by mixing with three volumes of 96% ethanol. The precipitates were collected and dried overnight at 50°C, then, it was milled and sieved through a mesh 18 sifter. Then, the mucilage powders were packed and kept in the dry and cold conditions.

Preparation of QSM Dispersion

The QSM dispersions (0.25% w/w) were prepared in de-ionized water with different concentrations of salts (NaCl and CaCl₂) (50, 100, and 150 mM) and sugars (glucose and sucrose) (10, 20, and 30% w/w). In order to complete hydration, the dispersions were mixed using a roller mixer for 24 hours at 25°C.

Rheological Experiments

The rheological tests were performed using a rotational viscometer (Model RVDV-II, Brookfield, Inc. USA). In order to equilibrate the temperature of the dispersions at 25°C, a jacket with a water circulating system (Model ULA-40Y, Brookfield, Inc. USA) was used. The steady

shear data of QSM were measured by spindle 18 at 25°C (±0.1°C). All the tests were carried out twice and the data was reported as an average of each test value.

Time-Independence

The steady shear rheological behavior was assessed in the shear rate range of 10-220 s⁻¹. Fitting the data of shear stress (τ)-shear rate (γ̇) was conducted with the Herschel-Bulkley model (Equation 1):

$$\tau = k \dot{\gamma}^n + \tau_0 \quad (1)$$

Where, τ is the shear stress (Pa), γ̇ is known as the shear rate (s⁻¹), k is the consistency coefficient (Pa.sⁿ), τ₀ is the yield stress, and n is the flow behavior index (dimensionless).

Time-Dependence

Investigation of the time-dependent characteristics was performed at a constant shear rate (50 s⁻¹) as a simulated oral shear rate and the shear stress (τ) was reported as a function of shearing time (t) until shear stress achieved a constant value. Fitting the obtained data was carried out with the structural kinetic model (Equation 2):

$$\left[\left(\frac{\eta - \eta_\infty}{\eta_0 - \eta_\infty} \right) \right]^{1-n} = (n-1)kt + 1 \quad (2)$$

Where, η₀ is the initial apparent viscosity at t = 0 (structured state), and η_∞ is the steady-state apparent viscosity at t_∞ (non-structured state). n and k are the order of the structure breakdown and the breakdown rate constant, respectively.

AFM Measurement

AFM images (ARA-AFM, Iran) were obtained in tapping mode at a constant temperature (25°C). The AFM sample was prepared by drying the QSM dispersions (0.25% w/w) in presence of CaCl₂ (150 mM)

**Table 1.** The rheological parameters of Herschel–Bulkley models obtained for quince seed mucilage dispersion (0.25% w/w) at different salts and sugars concentrations. ^a

Salt (mM)	concentration	Herschel–Bulkley model					
		k (Pa s ⁿ)	n	τ_0 (Pa)	R^2	RMSE	SSE
Control	0	0.19± 0.01 ^{a, E}	0.59± 0.10 ^{e, A}	0.14± 0.07 ^{a, D}	0.99	0.04	0.06
NaCl	50	0.11± 0.02 ^b	0.60± 0.20 ^e	0.12± 0.15 ^b	0.99	0.04	0.07
	100	0.08± 0.01 ^c	0.63± 0.10 ^d	0.11± 0.19 ^c	0.99	0.01	0.01
	150	0.07± 0.05 ^d	0.65± 0.30 ^c	0.11± 0.04 ^c	0.99	0.01	0.01
CaCl ₂	50	0.08± 0.02 ^c	0.65± 0.12 ^c	0.12± 0.09 ^b	0.99	0.02	0.02
	100	0.07± 0.03 ^d	0.67± 0.16 ^b	0.11± 0.16 ^c	0.99	0.02	0.02
	150	0.05± 0.05 ^f	0.69± 0.20 ^a	0.10± 0.08 ^d	0.99	0.02	0.03
<hr/>							
Sugar concentration (% w/w)							
Glucose	10	0.18± 0.01 ^E	0.59± 0.25 ^A	0.13± 0.03 ^D	0.98	0.05	0.12
	20	0.24± 0.01 ^D	0.59± 0.43 ^A	0.21± 0.24 ^C	0.99	0.02	0.022
	30	0.43± 0.00 ^C	0.55± 0.18 ^C	0.30± 0.06 ^B	0.99	0.28	0.31
Sucrose	10	0.24± 0.01 ^D	0.57± 0.39 ^B	0.14± 0.07 ^D	0.99	0.10	0.44
	20	0.47± 0.01 ^B	0.55± 0.42 ^C	0.22± 0.21 ^C	0.99	0.22	0.89
	30	0.62± 0.01 ^A	0.53± 0.47 ^D	0.38± 0.18 ^A	0.99	0.02	0.02

^a (a-f) and (A-E): Different letters show significant differences ($P < 0.05$) (small letters for salts and capital letters for sugars), ns: Not significant.

and sucrose (30% w/w) on a mica surface at room temperature.

RESULTS AND DISCUSSION

Statistical Analysis

Rheological parameters were analyzed with MATLAB (2016a). The data were statistically analyzed by the Analysis Of Variance (ANOVA) based on a completely randomized design. Then, the Duncan Multiple Range Test was used to determine the difference between means ($P < 0.05$). All the experiments were done at least in triplicate.

Time-Independent Rheological Properties of QSM

The best description of the steady-shear flow behavior of QSM dispersions in the presence of salts and sugars was acquired by the Herschel-Bulkley model. For this model, the ranges of R^2 and RMSE were 0.99 and 0.012-0.281, respectively. The rheological properties of QSM dispersions are presented in Table 1.

According to the results, increasing the shear rates ($10\text{--}220\text{ s}^{-1}$) led to the viscosity reduction, which signified the shear-thinning behavior of the QSM dispersions. As shown in Figure 1, a drastic reduction occurred with increasing the shear rates from 10 to 40 s^{-1} . This decrement trend has been observed at higher shear rates ($40\text{--}220\text{ S}^{-1}$), but the viscosity reduction was not pronounced. Intertwining the chains of the polysaccharide can produce a network, making a unique viscosity in solution. When the shear is applied, this network can be disrupted easily, which can explain the shear-thinning behavior of QSM (Morris *et al.*, 1981; Yanbei *et al.*, 2015).

Influence of Salts

Shear-dependent rheological characteristics of the QSM dispersions at various concentrations of the salts are presented in Table 1 and Figure 1. The aqueous mucilage dispersions also showed the pseudoplastic properties in the presence of salts (n values were lower than 1 for all the samples).

Adding salts (50, 100 and 150 mM) reduced the apparent viscosity noticeably,

and CaCl_2 compared to NaCl decreased the viscosity of the QSM dispersion to a greater extent. The amounts of viscosity reduction compared to the control sample at shear rate of 100 s^{-1} were 3.51 and 12.19% for NaCl and CaCl_2 at 50 mM, respectively (Figure 1).

As shown in Table 1, the flow behavior index (n) increased significantly in the presence of salts ($P > 0.05$). According to the Tukey test (ANOVA), the impact of type and concentration of salt and also their interactions on k , τ_0 and n values were significant ($P < 0.05$). The higher concentration of salts had more influence on the flow behavior index. Farahmandfar and Naji-Tabasi (2020) also reported a similar trend for n value in the presence of salt addition. The consistency coefficient value (k) decreased from 0.193 to 0.056 Pa s^n in the presence of cations compared to the control ($P < 0.05$). Increasing the salt concentration had a negative impact on the amount of τ_0 value, which decreased from 0.14 to 0.11 Pa at 150 mM NaCl and from 0.14 to 0.10 Pa at 150 mM CaCl_2 ($P < 0.05$) (Table 1). According to Yousefi *et al.* (2016), the effect of KCl and MgCl_2 on the sage seed gum (SSG) dispersions was similar to our results. They reported that




Figure 1. Impact of the various concentrations (50, 100 and 150 mM) of NaCl and CaCl_2 on the apparent viscosity of QSM dispersions.



Sisko ($R^2 = 0.995-0.999$) and Herschel-Bulkley ($R^2 = 0.994-0.999$) models showed suitable results to describe the steady-shear behavior of SSG. According to their study, salts decreased the yield stress and consistency coefficient of SSG.

By adding a monovalent salt (NaCl), the values of k were higher than those that were obtained by adding the divalent salt (CaCl_2) to the QSM dispersion (Table 1). These alterations have an important role in viscosity reduction. The reduction of k value in a salty solution is related to the sulfate group and carboxylic acids in the polysaccharide chains (Alpizar-Reyes *et al.*, 2018). The contraction of polysaccharide conformation induced by the salts led to viscosity reduction. A similar trend has been observed for chia mucilage (Velázquez-Gutiérrez *et al.*, 2015), and Balangu seed gum (Razavi *et al.*, 2016). The rheological characteristics of *E. contortisiliquum* gum by the addition of Na^+ , Ca^{2+} , and Al^{3+} were evaluated by Oliveira *et al.* (2001). They reported that positive ions affinity of gum structure was a function of charge to ionic radius ratio of counter ions. The lower amounts of charge to ionic radius ratio resulted in the lower chain compression for hydrocolloid molecules in the solution. Therefore, gum solutions (at infinite dilution) containing sodium or aluminum ions caused the highest or least viscosity, respectively. The larger ions (Ca^{2+}) shield the electrostatic interaction among the QSM chains more strongly compared to the smaller cations (Na^+), as they have larger hydrated radius. (Harrison *et al.*, 1999). Salehi *et al.* (2014) mentioned that the type of salt extremely affected the rheological characteristics of Balangu seed gum. These researchers found that the viscosity reduction by CaCl_2 was more obvious than NaCl.

Increasing the salts concentration caused the rapid reduction of viscosity. The reduction of viscosity was more predominate at high concentration of the salt (Figure 1 and Table 1). It seems that when the concentration of added salts was low,

expanding the gum structure was obviously limited, which led to the viscosity reduction. Enhancing the salt concentration helps conformation change of hydrocolloid to a more compact coil. This is because of neutralization of the negatively charged residues of gum that were exposed to the cations. These processes reduced the hydrodynamic volume and electrostatic repulsion of hydrocolloid chains to a greater extent, which led to the more viscosity reduction.

It seemed that the QSM was anionic mucilage like a polyelectrolyte. Therefore, the addition of cations decreased the repulsion and then mucilage conformation collapsed. Our results confirmed that the QSM is a negatively charged polyelectrolyte in the solution. Drastic intermolecular electrostatic repulsion and more expanded molecule are responsible for inherent negative charge of the QSM. According to Ritzoulis *et al.* (2014), hydrocolloids extracted from the quince seed at four different pHs had negative charges and zeta potential of the hydrocolloid extracted at $\text{pH} = 7$ was about -35 mV. The existence of inorganic salt and positive ions with the opposite charge (via capability to form the salt bridges among acid groups) cover the electrostatic repulsion among charges along the structure of polyelectrolyte. Folding up the chain is a result of screening of the electrostatic interactions and assumed a smaller and more compact structure. As a result, the viscosity of the QSM dispersions decreased (Wyatt *et al.*, 2011). According to Mazza and Biliaderis (1989), joining of the counter-ions with polymers structure could persuade NaCl to reduce the flaxseed mucilage viscosity; therefore, the electrostatic repulsion among charged groups of polymer chain was decreased.

According to Carrington *et al.* (1996), the molecular conformation of xanthan gum extremely changed due to the influence of salt addition on its rheological characteristics. The highest viscosity belongs to the control sample, whose structure was unfolded in the solution.




Figure 2. Impact of the various concentrations (10, 20, and 30% w/w) of glucose and sucrose on the apparent viscosity of QSM dispersions.

Alpizar-Reyes *et al.* (2018) declared that rheological characteristics of tamarind seed mucilage were influenced by the salts and their concentrations owing to the addition of cations, which decreases repulsion and allows molecular expansion, thereby promoting a significant reduction in viscosity. Lai *et al.* (2000) also reported that the impact of salt on the viscosity reduction of hsian-tsao leaf gum was more obvious when using divalent positive ions. According to Razavi *et al.* (2011), the ionized groups on gum structure will be neutralized at the higher contents of salt, hence its structure was more compact.

Influence of Sugars

Figure 2 demonstrates the flow behaviour curves of QSM (0.25% w/w) at the presence of two selected sugars (glucose and sucrose) and at different concentrations (10, 20, and 30 % w/w). It can be seen that all samples exhibited the shear-thinning behaviour. Yousefi *et al.* (2016) also reported similar results for the effect of sugars on the flow behaviour of sage seed gum. They found a rapid decrease in the viscosity at 10-50 s⁻¹

shear rates. As shown in Figure 2, the sugars had a synergistic influence on the viscosity of QSM dispersion. The apparent viscosity increased by enhancing the sugars concentration. Sucrose had considerable influence on the viscosity increment in comparison with the glucose. The viscosity of control sample was the lowest. The viscosity enhancing of 30% w/w glucose and sucrose compared to the control at 100 s⁻¹ were 165.48 and 192.50%, respectively. The *n* values of the all samples were lower than 1, which proved the shear-thinning behavior of the QSM in the presence of sugars (Table 1). The *n* values of QSM dispersion were decreased by increasing the concentration of sugars. The type and concentration of sugars also influenced *k* value, so, increment in the *k* value was observed when the sugar concentration was increased (from 0.185 to 0.433 Pa sⁿ for glucose and from 0.244 to 0.629 Pa sⁿ for sucrose). The yield stress of QSM dispersion increased as the sugar concentration increased from 10 to 30% w/w and the sucrose had more prominent effect on τ_0 (Table 1).

By adding sucrose/glucose into the polysaccharide network, the density of QSM



dispersion noticeably increased, which subsequently resulted in increase in viscosity. Moreover, the sugar in the solution could bind to the polysaccharide structure of the QSM via hydrogen bonding and then promoting the cross-linking strength of the polysaccharide network (Lei *et al.*, 2001, 2001; Richardson and Norton, 1998), which also causes the viscosity increment. Salehi and Kashaninejad (2014) mentioned that the competition between the sugar and basil seed gum for water-binding will enhance by the increase in sugar concentration. Chenlo *et al.* (2011) proposed that the polymer-polymer interactions are influenced by the sugars and hydrogen bonds created among the hydrocolloid and hydroxyl groups of the sugar, which could increase the number of junction zones and stabilize the polymer chain associations. Finally, these interactions result in structural changes.

A number of researches have reported the positive effect of sugars on the viscosity of other hydrocolloids such as oat gum (Cancela *et al.*, 2005), cress seed gum (Behrouzian *et al.*, 2013), Balangu seed gum (Salehi and Kashaninejad, 2015) and sage seed gum (Yousefi *et al.*, 2016). According to Chenlo *et al.*, (2011), the viscosity of guar gum was enhanced by adding sucrose and glucose, which was related to the interactions between the polymer and sugars. As reported by Wu *et al.* (2015), the addition of high concentration of sucrose (30 mg mL⁻¹) had the comprehensive influence on the flow behaviour of tara gum solution. They concluded that by adding sucrose, a compact conformation of tara gum appeared.

According to Capitani *et al.* (2015), the addition of sugar to aqueous dispersion of chia mucilage leads to decrease in the pseudoplastic behaviour. Other researchers also observed similar results by adding sugar into the hydrocolloid solution (Salehi *et al.*, 2014 in case of glucose; Yanes *et al.*, 2002 in case of sucrose; Galmarini *et al.*, 2010; Bayari *et al.*, 2004 and Sato *et al.*, 2004). In contrast, some authors explained that the flow behavior index (*n*) of hydrocolloid

decreased in the presence of sugar (Elfak *et al.*, 1980; Salehi *et al.*, 2014 for sucrose; Yousefi *et al.*, 2016). Moreover, the additive concentrations and compositions are the main factors influencing the rheological properties of particular hydrocolloid.

Time-Dependent Rheological Properties of QSM

Influence of Salts

After initial shearing, the viscosity of dispersions attained a constant value corresponding to a steady-state just about 350 s. When a polymer solution is exposed to shearing, the rearrangement or breakdown of links among polymer chains could happen, which is related to the thixotropy (Mao and Chen, 2006). This behavior proved that the QSM chains have the possibility to create a three-dimensional structure and shearing force could change the polysaccharide configuration. Other researchers concluded similar trends for Balangu seed gum, *Alyssum homolocarpum* seed gum and green Laver mucilage (Razavi and Karaizhyan, 2009; Koocheki and Razavi, 2009; Huei Chen and Yuu Chen., 2001).

Figure 3 (a, b) demonstrates the time-dependent rheological characteristics of the QSM in the presence of cations (Na⁺ and Ca²⁺). By increasing the time of shearing, the viscosity of all samples decreased in comparison with the control, which revealed their thixotropic behavior. Basil seed gum and its fractions also showed thixotropic behavior during the shearing (Naji-Tabasi and Razavi, 2017). This behavior is also reported for the other mucilage like cress seed gum and its fractions (Karazhian *et al.*, 2009; Razmkhah *et al.*, 2017).

The structural kinetics model was used to describe the time-dependency of the flow behavior of QSM and the obtained data are presented in Table 2. It was found that their transient viscosity data could be fitted satisfactory with a second order structural

Table 2. Viscosity of QSM dispersions (0.25 w/w) in the presence of salts (NaCl and CaCl₂) and sugars (glucose and sucrose) as a function of shearing time (shear rate 50 L s⁻¹).^a

Salt (mM)	concentration	η_0 (Pa s)	η_0/η_∞	$K \times 10^{-3}$ (s ⁻¹)	R ²	RMSE	SSE
Control	0	45.02± 0.21 ^{a, F}	1.21± 0.03 ^{c, C}	12.34± 0.00 ^{ns}	0.99	0.08	0.28
	NaCl 50	22.44± 0.43 ^b	1.21± 0.05 ^c	12.30± 0.00 ^{ns}	0.99	0.05	0.11
	100	21.87± 0.17 ^c	1.22± 0.06 ^c	12.17± 0.01 ^{ns}	0.98	0.06	0.16
	150	20.30± 0.13 ^d	1.26± 0.09 ^b	12.55± 0.01 ^{ns}	0.99	0.06	0.15
CaCl ₂	50	21.92± 0.19 ^c	1.19± 0.06 ^d	12.16± 0.00 ^{ns}	0.99	0.06	0.16
	100	20.71± 0.11 ^e	1.26± 0.04 ^b	12.19± 0.01 ^{ns}	0.98	0.07	0.19
	150	19.88± 0.14 ^f	1.77± 0.17 ^a	12.03± 0.01 ^{ns}	0.98	0.12	0.59
Sugar concentrations (% w/w)							
Glucose	10	43.37± 0.30 ^G	1.14± 0.17 ^D	12.44± 0.00 ^{ns}	0.98	0.03	0.26
	20	49.26± 0.32 ^E	0.92± 0.64 ^F	12.09± 0.00 ^{ns}	0.98	0.15	0.11
	30	94.68± 0.39 ^B	1.36± 0.06 ^A	12.23± 0.01 ^{ns}	0.99	0.06	0.69
Sucrose	10	63.57± 0.41 ^D	1.28± 0.19 ^B	12.11± 0.00 ^{ns}	0.99	0.28	0.14
	20	66.66± 0.27 ^C	1.10± 0.18 ^E	12.10± 0.00 ^{ns}	0.97	0.06	0.96
	30	105.20± 0.44 ^A	1.14± 0.14 ^D	12.68± 0.01 ^{ns}	0.98	0.07	0.54

^a (a-f) and (A-G): Different letters show significant differences (P< 0.05) (small letters for salts and capital letters for sugars), ns: Not significant.




Figure 3. Viscosity of QSM dispersions (0.25 % w/w) in the presence of NaCl (a) and CaCl₂ (b), glucose (c), and sucrose (d) as a function of shearing time (shear rate 50 L s⁻¹).



kinetic model ($n=2$). Fitting the data with this model showed the highest R^2 and the lowest RMSE and SSE values, which proved the acceptable capability of the structural kinetic model for prediction of the time-dependent behavior of the QSM dispersion. Shear-induced breaking of internal configuration was responsible for the changes of rheological characteristics. On the other hand, breakdown rate of QSM structure during the shearing is determined by the kinetics changes of structured to non-structured state (Abu-Jdayil, 2003). Koocheki and Razavi (2009) also reported that this second-order model was the best for fitting the time-dependent viscosity of *A. homolocarpum* seed gum at a constant shear rate.

According to Table 2, the highest initial apparent viscosity (η_0) belonged to the control sample. This parameter declined by enhancing the salt concentration. The lowest value of initial viscosity was shown at 150 mM CaCl_2 sample (19.884 Pa s), which showed 55.84% reduction compared with the control. The salts had no significant impact on the breakdown rate constant (k), which shows how fast the QSM dispersions reach the equilibrium viscosity under the action of shearing. Similar results were reported by Farahmandfar and Naji-Tabasi (2020). The ratio of initial to equilibrium viscosity (η_0/η_∞) can be presumed as a relative assessment of the extent of structural breakdown (Koocheki and Razavi, 2009). The result of the η_0/η_∞ is given in Table 2. The ratio η_0/η_∞ is a relative measure of the extent of thixotropy of the QSM dispersions. According to Table 2, the addition of salts led to increase in the η_0/η_∞ values ($P < 0.05$) and also led to enhancing the thixotropic behavior. Seemingly, the salts caused weak links among the chains of QSM structure and reduced the amount of structuring.

Influence of Sugars

Figure 3 (c, d) present the time-dependent characteristics of QSM in the presence of glucose and sucrose (10, 20, and 30 % w/w).

Increasing the shearing times resulted in the viscosity reduction, which indicated thixotropic behavior of QSM in all samples. The viscosity of QSM dispersions increased after adding sucrose and glucose (Figure 3). The rheological parameters of the structural kinetic model are summarized in Table 2. As shown, by adding 30 % w/w sucrose, the viscosity elevated (the highest value). In contrast to salt, the sugar addition had synergistic effect on the η_0 , likewise, the highest amount of initial viscosity belonged to the 30% w/w sucrose (57.21% increase compared to the control). There was no significant change in case of breakdown rate constant (k), because this index was a function of ion change. In case of the ratio of initial to equilibrium viscosity (η_0/η_∞), the systematic trend was not observed. Razavi and Kashaninejad, (2009) reported that k value (as a relative measure of thixotropic breakdown rate) for Balangu seed gum in shear rate of 50 s^{-1} was 0.0643. They also stated that this value was independent of the shear rate. According to their research, the extent of thixotropy (η_0/η_∞) at a similar shear rate was 2.44, which was higher than the data obtained for the QSM dispersions containing sugars.

Atomic Force Microscopy (AFM)

As shown in AFM Figures (Figure 4, a-b), addition of CaCl_2 (150 mM) had some effect on fibrils and lengthened them to a certain direction. A tightly pack continuous network structure with some inhomogeneity was observed in the presence of CaCl_2 . On the contrary, the direction of fibers in the QSM sample was not definite and the structure was heterogeneous. Generally, addition of cations (like calcium ions) leads to inter-helical association of the QSM chains. The degree of structural shrinkage was determined by stiffness of the polysaccharide chain (Wyatt *et al.*, 2011). These observations were in agreement with the rheological behavior result, which thixotropic behavior increased behavior




Figure 4. Typical AFM topographic images of the QSM dispersions: (a) 2D view; (b) 3D view; control (0.25% w/w) (A), in presence of 150 mM CaCl_2 (B), and 30% w/w Sucrose (C).

increased after adding salt. Similar results were concluded for Gellan gum in the presence of various cations (Funami *et al.*, 2008; 2009). They reported that the reduction of repulsion was distinguishable among cations, which explained the difference in the network formation, and the effect of CaCl_2 was more obvious, at least 10 times more than the other cations. Adding CaCl_2 promoted inter-helical association of Carageenan and formation of the network-like structures. Mean vertical height was longer than the control sample (Funami *et al.*, 2007).

QSM molecules, as the polyelectrolyte (opposite charges), were in the extended form in dispersion, so the fibrils were shown in the different directions in AFM images (Figure 4-a). The presence of a high concentration of sucrose (30% w/w) decreased considerably the water available for hydration, so, it had an impact on the flow behavior of mucilage. Moreover, by dissolving the sucrose into the polysaccharide network, density of solution increased. The sucrose also could bridge among the QSM chains via hydrogen bonds and strengthened the QSM network (Lei *et*

al., 2001; Richardson and Norton, 1998), which condensed the molecular structure. The fibrils were not imaged clearly due to the surface roughness (Figure 4, C). These results were in agreement with those reported by Yanbei *et al.* (2015) for tara gum.

CONCLUSIONS

In this research, the impact of different kinds of salt (NaCl and CaCl_2) and sugar (glucose and sucrose) on the time-dependent and time-independent rheological characteristics of QSM were studied. It was found that the Herschel-Bulkley model was appropriate for characterizing the rheological properties of QSM as a function of the type and concentrations of the selected salts and sugars. The consistency coefficient (k) value of QSM dispersion was higher in the presence of NaCl compared to CaCl_2 . In comparison with the control, the salt addition caused increase in the flow behavior index (n) and reduction in yield stress (τ_0) and k value. Typically, the impact of Ca^{2+} on the rheological characteristics



was stronger than Na^+ . By increasing the concentration of glucose and sucrose, the viscosity was increased. The structural kinetic model showed the acceptable ability to fit the time-dependent data. The data obtained by this model proved that the rate of thixotropic breakdown was higher in presence of CaCl_2 . The viscosity of salty solution decreased by increasing the time of shearing. There were no systematic trends in the ratio of the initial to equilibrium viscosity in the presence of sucrose and glucose. The AFM results demonstrated the mechanism of QSM chains interactions in the presence of salt and sugar that justify the viscosity changes. According to the results of this work, the rheological characteristics of QSM dispersions were strongly influenced by the type and concentration of salts and sugars. Hence, adjusting the salt and sugar content could change the rheological properties of food products. The rheological and sensory properties of mucilage applied in the food products are determined by its microstructure and the degree of interactions. This issue is prominent, especially in the development of food formulations with proper sensory, functional, and rheological properties.

REFERENCES

1. Abbastabar, B., Azizi, M. H., Adnani, A. and Abbasi, S. 2015. Determining and Modeling Rheological Characteristics of Quince Seed Gum. *Food Hydrocoll.*, **43**: 259-264.
2. Abu-Jdayil, B. 2003. Modelling the Time-Dependent Rheological Behavior of Semisolid Foodstuffs. *J. Food Eng.*, **57**: 97-102.
3. Alpizar-Reyes, E., Román-Guerrero, A., Gallardo-Rivera, R., Varela-Guerrero, V., Cruz-Olivares, J. and Pérez-Alonso, C. 2018. Rheological Properties of Tamarind (*Tamarindus indica* L.) Seed Mucilage Obtained by Spray-Drying as a Novel Source of Hydrocolloid. *Int. J. Biol. Macromol.*, **107**: 817-824.
4. Behrouzian, F., Razavi, S. M. A. and Karazhiyan, H. 2013. The Effect of pH, Salts and Sugars on the Rheological Properties of Cress Seed (*Lepidium sativum*) Gum. *Int. J. Food Sci. Tech.*, **48**: 2506-2513.
5. Bergman, R., Afifi, A. K. and Heidgerip, P. M. 1996. *Text of Histology*. 9th Edition, WB Saunders Company, Montréal, PP. 159-68.
6. Bayarri, S., Duran, L., and Costell, E. 2004. Influence of Sweeteners on the Viscoelasticity of Hydrocolloids Gelled Systems. *Food Hydrocoll.*, **18**: 611-619.
7. Cancela, M. A., Alvarez, E., and Maceiras, R. 2005. Effects of Temperature and Concentration on Carboxymethylcellulose with Sucrose Rheology. *J. Food Eng.*, **71**: 419-424.
8. Capitani, M. I., Corzo-Rios, L. J., Chel-Guerrero, L. A., Betancur-Ancona, D. A., Nolasco, S. M., and Tomás, M. C. 2015. Rheological Properties of Aqueous Dispersions of Chia (*Salvia hispanica* L.) Mucilage. *J. Food Eng.*, **149**: 70-77.
9. Carrington, S., Odell, J., Fisher, L., Mitchell, J. and Hartley, L. 1996. Polyelectrolyte Behavior of Dilute Xanthan Solutions: Salt Effects on Extensional Rheology. *Polymer*, **37(13)**: 2871-2875.
10. Chenlo, F., Moreira, R. and Silva, C. 2011. Steady-Shear Flow of Semi-Dilute Guar Gum Solutions with Sucrose, Glucose and Sodium Chloride at Different Temperatures. *J. Food Eng.*, **107** (2): 234-240.
11. Elboutachfai, R., Delattre, C., Quéro, A., Roulard, R., Duchêne, J., Mesnard, F., and Petit, E. 2017. Fractionation and Structural Characterization of Six Purified Rhamnogalacturonans Type from Flaxseed Mucilage. *Food Hydrocoll.*, **62**: 273-279.
12. Farahmand, A., Varidi, M. and Koocheki, A. 2016. Investigation of Functional Properties of Quince Seed Mucilage Extracted by Ultrasound. *Iran. Food Sci. Technol. Res. J.*, **12(1)**: 163-181.
13. Farahmandfar, R. and Naji-Tabasi, S. 2020. Influence of Different Salts on Rheological and Functional Properties of Basil (*Ocimum bacilicum* L.) Seed Gum. *Int. J. Biol. Macromol.*, **149**: 101-107.
14. Funami, T., Hiroe, M., Noda, S., Asai, I., Ikeda, S. and Nishinari, K. 2007. Influence of Molecular Structure Imaged with Atomic Force Microscopy on the Rheological Behavior of Carrageenan Aqueous System in the Presence or Absence of Cations. *Food Hydrocoll.*, **21**: 617-629.
15. Funami, T., Noda, S., Nakauma, M., Ishihara, S., Takahashi, R., Al-Assaf, S.,

- Ikeda, S., Nishinari, K. and Phillips, G. O. 2008. Molecular Structures of Gellan Gum Imaged with Atomic Force Microscopy in Relation to the Rheological Behavior in Aqueous Systems in the Presence or Absence of Various Cations. *J. Agric. Food Chem.*, **56**: 8609-8618.
16. Funami, T., Noda, S., Nakauma, M., Ishihara, S., Takahashi, R., Al-Assaf, S., Ikeda, S., Nishinari, K. and Phillips, G. O. 2009. Molecular Structures of Gellan Gum Imaged with Atomic Force Microscopy (AFM) in Relation to the Rheological Behavior in Aqueous Systems in the Presence of Sodium Chloride. *Food Hydrocoll.*, **23**: 548-554.
 17. Galmarini, M. V., Baeza, R., Sanchez, V., Zamora, M. C., and Chirife, J. 2011. Comparison of the Viscosity of Trehalose and Sucrose Solutions at Various Temperatures: Effect of Guar Gum Addition. *LWT-Food Sci, Technol.*, **44**: 186-190.
 18. Harrison, G., Franks, G. V., Tirtaatmadja, V. and Boger, D. V. 1999. Suspensions and Polymers-Common Links in Rheology. *Korea-Aust. Rheol. J.*, **11(3)**: 197-218.
 19. Huei Chen, R. and Yuu Chen, W. 2001. Rheological Properties of the Water Soluble Mucilage of a Green Laver, *Monostroma nitidum*. *J. Appl. Phycol.*, **13**: 481-488.
 20. Hussain, M. A., Muhammad, G., Haseeb, M. T. and Tahir, M. N. 2019. Quince Seed Mucilage: A Stimuli-Responsive/Smart Biopolymer. In: "*Functional Biopolymers. Polymers and Polymeric Composites: A Reference Series*", (Eds.): Jafar Mazumder, M., Sheardown, H. and Al-Ahmed, A. Springer, Cham.
 21. Imeson, A. 2010. Food Stabilisers, Thickeners and Gelling Agents. Wiley-Blackwell, Oxford, UK.
 22. Jouki, M., Mortazavi, S. A., Tabatabaei Yazdi, F. and Koocheki, A. 2014. Optimization of Extraction, Antioxidant Activity and Functional Properties of Quince Seed Mucilage by RSM. *Int. J. Biol. Macromol.*, **66**: 113-124.
 23. Kar, F. and Arslan, N. 1991. Characterization of Orange Peel Pectin and Effect of Sugars, Ascorbic Acid, Ammonium Persulfate, Salts on Viscosity of Orange Peel Pectin Solutions *Carbohydr. Polym.*, **40**: 285-291.
 24. Karazhiyan, H., Razavi, S. M., Phillips, G. O., Fang, Y., Al-Assaf, S., Nishinari, K., and Farhoosh, R. 2009. Rheological Properties of *Lepidium sativum* Seed Extract as a Function of Concentration, Temperature and Time. *Food Hydrocoll.*, **23**: 2062-2068.
 25. Koocheki, A. and Razavi, S. M. A. 2009. Effect of Concentration and Temperature on Flow Properties of Alyssum homolocarpum Seed Gum Solutions: Assessment of Time Dependency and Thixotropy. *Food Biophys.*, **4**: 353-364.
 26. Koocheki, A., Mortazavi, S. A., Shahidi, F., Razavi, S. M. A., and Taherian, A. R. 2009. Rheological Properties of Mucilage Extracted from Alyssum Homolocarpum Seed as a New Source of Thickening Agent. *J. Food Eng.*, **91**: 490-496.
 27. Lai, L. S., Tung, J. and Lin, P. S. 2000. Solution Properties of Hsian-Tsao (*Mesona procumbens* Hems L) Leaf Gum. *Food Hydrocoll.*, **14**: 287-94.
 28. Legua, P., Serrano, Melgarejo, P. M., Valero, D., Martínez, J. J., Martínez, R., and Hernández, F. 2013. Quality Parameters, Biocompounds and Antioxidant Activity in Fruits of Nine Quince (*Cydonia oblonga* Miller) Accessions. *Sci. Hortic.*, **154**: 61-65.
 29. Lei, M., Lu, X. L. and Xiao, K. 2001. Study on Influencing Factors of Viscosity of Low Concentration Food Gum Slutions (2). *J. Sichuan Univ. (Eng. Sci. Ed.)*, **33(1)**: 78-81. (in Chinese with English Abstract)
 30. Lin, H. Y. and Lai, L. S. 2009. Isolation and Viscometric Characterization of Hydrocolloids from Mulberry (*Morus alba* L.) Leaves. *Food Hydrocoll.*, **23(3)**: 840-848.
 31. Mao, C. F. and Chen, J. C. 2006. Interchain Association of Locust Bean Gum in Sucrose Solutions: An Interpretation Based on Thixotropic Behavior. *Food Hydrocoll.*, **20**: 730-739.
 32. Mazza, G. and Biliaderis, C. G. 1989. Functional Properties of Flax Seed Mucilage. *J. Food Sci.*, **54**: 1302-1305.
 33. Morris, E. R., Cutler, A. N., Ross-Murphy, S. B., Rees, D. A. and Price, J. 1981. Concentration and Shear Rate Dependence of Viscosity in Random Coil Polysaccharide Solutions. *Carbohydr. Polym.*, **1**: 5-21.
 34. Naji-Tabasi, S. and Razavi, S. M. A. 2017. New Studies on Basil (*Ocimum bacilicum* L.) Seed Gum. Part III. Steady and Dynamic



- Shear Rheology. *Food Hydrocoll.*, **67**: 243-250.
35. Oliveira, J. D., Silva, D. A., Paula, R. C. M., Feitosa, J. P. A. and Paula, H. C. B. 2001. Composition and Effect of Salt on Rheological and Gelation Properties of *Enterolobium contortisiliquum* Gum Exudate. *Int. J. Biol. Macromol.*, **29**(1): 35-44.
 36. Razavi, S. M. A and Karazhiyan, H. 2009. Flow Properties and Thixotropy of Selected Hydrocolloids: Experimental and Modeling Studies. *Food Hydrocoll.*, **23**: 908-912
 37. Razavi, S. M. A., Cui, S. W. and Ding, H. 2016. Structural and Physicochemical Characteristics of a Novel Water-Soluble Gum from *Lallemantia royleana* Seed. *Int. J. Biol. Macromol.*, **83**: 142-151.
 38. Razavi, S. M. A., Taheri, H. and Quinchia, L. A. 2011. Steady Shear Flow Properties of Wildsage (*Salvia macrosiphon*) Seed Gum as a Function of Concentration and Temperature. *Food Hydrocoll.*, **25**: 451-458.
 39. Razmkhah, S., Razavi, S. M. A., and Mohammadifar, M. A. 2017. Dilute Solution, Flow Behavior, Thixotropy and Viscoelastic Characterization of Cress Seed (*Lepidium sativum*) Gum Fractions. *Food Hydrocoll.*, **63**: 404-413.
 40. Rezagholi, F., Hashemi, S. M. B., Gholamhosseinpour, A., Sherahi, M. H., Hesarinejad, M. A., and Ale, M. T. 2019. Characterizations and Rheological Study of the Purified Polysaccharide Extracted from Quince Seeds. *J. Sci. Food Agric*, **99**: 143-151.
 41. Richardson, P. H. and Norton, I. T. 1998. Gelation Behavior of Concentrated Locust Bean Gum Solutions. *Macromolecules*, **31**: 1575-1583.
 42. Ritzoulis, C., Marini, E., Aslanidou, A., Georgiadis, N., Karayannakidis, P., Koukiotis, C., Filotheou, A., Lousinian, S. and Tzimpilis, E. 2014. Hydrocolloids from Quince Seed: Extraction, Characterization, and Study of Their Emulsifying/Stabilizing Capacity. *Food Hydrocoll.*, **42**: 178-186.
 43. Salehi, F. and Kashaninejad, M. 2015. Static Rheological Study of *Ocimum basilicum* Seed Gum. *Int. J. Food Eng.*, **11**(1): 97-103.
 44. Salehi, F., Kashaninejad, M. and Behshad, V. 2014. Effect of Sugars and Salts on Rheological Properties of Balangu Seed (*Lallemantia royleana*) Gum. *Int. J. Biol. Macromol.*, **67**: 16-21.
 45. Sato, Y., Kawabuchi, S., Irimoto, Y., and Miyawaki, O. 2004. Effect of Water Activity and Solvent-ordering on Intermolecular Interaction of High-methoxyl Pectins in Various Sugar Solutions. *Food Hydrocoll.*, **18**: 527-534.
 46. Sharma, R., Joshi, V. K. and Rana, J. C. 2011. Nutritional Composition and Processed Products of Quince (*Cydonia oblonga* Mill.). *IJNPR*, **2**: 354-357.
 47. Soukoulis, C., Lebesi, D. and Tzia, C. 2009. Enrichment of Ice Cream with Dietary Fiber: Effects on Rheological Properties, Ice Crystallization and Glass Transition Phenomena. *Food Chem.*, **115**(2): 665-671.
 48. Soukoulis, Ch., Gaiani, C. and Hoffmann, C. 2018. Plant Seed Mucilage as Emerging Biopolymer in Food Industry Applications. *Curr. Opin. Food Sci.*, **22**: 28-42.
 49. Torres, M. D., Raymundo, A. and Sousa, I. 2013. Effect of Sucrose, Stevia and Xylitol on Rheological Properties of Gels from Blends of Chestnut and Rice Flours. *Carbohydr. Polym.*, **98**: 249-256.
 50. Van Aken, G. A. 2006. Polysaccharides in food emulsions. In: "Food Polysaccharides and Their Applications", (Eds.): Stephen, A. M., Phillips, G. O. and Williams, P. A. 2nd Edition, Taylor and Francis, London, PP. 521-539.
 51. Velázquez-Gutiérrez, S. K., Figueira, A. C., Rodríguez-Huezo, M. E., Román-Guerrero, A., Carrillo-Navas, H. and Pérez-Alonso, C. 2015. Sorption Isotherms Thermodynamic Properties and Glass Transition Temperature of Mucilage Extracted from Chia Seeds (*Salvia hispanica* L.), *Carbohydr. Poly.*, **121**: 411-419.
 52. Williams, P. A. and Phillips, G. O. 2000. Introduction to Food Hydrocolloids. In: "Handbook of Hydrocolloids", (Eds.): Phillips, G. O. and Williams, P. A. CRC Press, New York, NY, PP. 1-19.
 53. Wojdyło, A., Oszmianski, J. and Bielicki, P. 2013. Polyphenolic Composition, Antioxidant Activity, and Polyphenol Oxidase (PPO) Activity of Quince (*Cydonia oblonga* Miller) Varieties. *J. Agric. Food Chem.*, **61**: 2762-2772.
 54. Wu, Y., Ding, W., Jia, L. and He, Q. 2015. The Rheological Properties of Tara Gum (*Caesalpinia spinosa*), *Food Chem.*, **168**: 366-371.

55. Wyatt, N. B., Gunther, C. M. and Liberatore, M. W. 2011. Increasing Viscosity in Entangled Polyelectrolyte Solutions by the Addition of Salt. *Polymers*, **52**: 2437-2444.
56. Yanbei, W., Wei, D., Lirong, J. and Qiang, H. 2015. The Rheological Properties of Tara Gum (*Caesalpinia spinosa*). *Food Chem.*, **168**: 366-371.
57. Yanes, M., Durán, L., and Costell, E. 2002. Effect of Hydrocolloid Type and Concentration on Flow Behavior and Sensory Properties of Milk Beverages Model Systems. *Food Hydrocoll*, **16**: 605-611.
58. Yousefi, A.R., Eyyvazloo, R. and Razavi, S. M. A. 2016. Steady Shear Flow Behavior of Sage Seed Gum Affected by Various Salts and Sugars: Time-Independent Properties, *Int. J. Biol. Macromol.*, **91**: 1018-1024.
59. Zhang, L. M., Kong, T. and Hui, P. S. 2007. Semi-Dilute Solutions of Hydroxypropyl Guar Gum: Viscosity Behaviour and Thixotropic Properties, *J. Sci.*, **87**: 684-688.

اثر برخی نمک و قندها بر رفتار رئولوژیکی موسیلاژ دانه به

ع. فرهمند، س. ناجی طبسی، و س. شهبازی زاده

چکیده

خواص حلال / حل شونده در سامانه های غذایی به طور قابل توجهی بر ویژگی های رئولوژیکی هیدروکلوئیدها موثر می باشد. در این پژوهش، اثر غلظت های مختلف دو قند (ساکارز و گلوکز در ۱۰، ۲۰ و ۳۰ درصد وزنی-وزنی) و دو نمک (کلرید کلسیم و کلرید سدیم در غلظت های ۵۰، ۱۰۰ و ۱۵۰ میلی مولار) بر ویژگی های رئولوژیکی موسیلاژ دانه به ارزیابی گردید. موسیلاژ به، محلولی با ویسکوزیته و تنش تسلیم (τ_0) بالا تولید کرد که تابع نوع و غلظت نمک و قند بود. نمک ها با تغییر برهم کنش میان زنجیره های موسیلاژ، منجر به کاهش ویسکوزیته محلول شدند. افزایش غلظت نمک باعث کاهش τ_0 و ضریب قوام (k) گردید، اما شاخص رفتار جریان (n) افزایش یافت. ساکارز و گلوکز منجر به افزایش ویسکوزیته محلول شدند، به طوری که با افزایش غلظت قندها، ویسکوزیته بالاتری حاصل شد. ارزیابی رفتار جریان وابسته به زمان نشان داد که مدل کنتیکی ساختاری، بهترین توصیف را برای داده ها ارائه می دهد. تمام نمونه ها دارای رفتار تیکسوتروپیک بودند. نتایج میکروسکوپ نیروی اتمی (AFM) نشان دهنده برهم کنش های زنجیره های موسیلاژ و تغییرات رفتار جریان بود. بر اساس نتایج، CaCl_2 و ساکارز بیشترین تاثیر را بر خواص رئولوژیکی موسیلاژ به نشان دادند.