

## Qualitative and Quantitative Changes in the Essential Oil of *Origanum vulgare* L. ssp. *gracile* as Affected by Different Harvesting Times

M. Moradi<sup>1</sup>, A. Hassani<sup>1\*</sup>, F. Sefidkon<sup>2</sup>, and H. Maroofi<sup>3</sup>

### ABSTRACT

Qualitative and quantitative variations in the essential oil of wild growing *Origanum vulgare* L. ssp. *gracile* plants were studied in response to different phenological stages (pre, full and post-flowering). The essential oil of air-dried leaves was isolated by water distillation using a Clevenger-type apparatus and was analyzed by Gas Chromatography (GC) and Gas Chromatography–Mass Spectrometry (GC/MS). The highest (1.87%) and the lowest (1.01%) essential oil content were obtained from post-flowering and pre-flowering stages, respectively. In total, 24 components were identified and quantified in three phenological stages representing 96.75, 97.63, and 98.59% of the oil, respectively. Carvacrol (46.62, 46.5 and 27.6%), *p*-cymene (7.76, 13.54 and 37.08%) and  $\gamma$ -terpinene (21.54, 13.91 and 6.82%) were the main constituents of essential oils in pre, full, and post-flowering stages, respectively. Oxygenated monoterpenes (43.35-61.32%) and monoterpene hydrocarbons (30.81-48.02%) were the main classes of identified compounds in three essential oils. According to the findings of this research, the post-flowering stage can be considered as the most appropriate time for obtaining the highest essential oil content, but to achieve the highest rate of phenolic compounds, the pre-flowering and full-flowering stages can be recommended.

**Keywords:** Carvacrol, Oregano, *p*-Cymene, Phenological stage,  $\gamma$ -Terpinene.

### INTRODUCTION

The genus *Origanum* (Lamiaceae family), which includes about 38 species and 6 subspecies, is native to the Mediterranean Region and it is widely distributed in many parts of the world in West, East and Central Asia, South Europe, and North Africa (Ietswaart, 1980; Vokou *et al.*, 1993). The *Origanum* species, due to richness in essential oils, have been used for centuries as spices and as local medicines in traditional medicine (Fleisher and Fleisher, 1988).

*O. vulgare* L. is one of the most important species of this genus, which is widely distributed in the Mediterranean, Euro-Siberian and Irano-Turanian Regions (Spada and Perrino, 1997). The Iranian flora comprises three subspecies of *O. vulgare* L. (ssp. *viride*, ssp. *vulgare* and ssp. *gracile*), which are distributed in the north and northwest of the country (Mozaffarian, 1996; La Gow, 2004). *O. vulgare* ssp. *gracile* is a perennial herb with white flowers, the leaf is nearly elliptical, covered with glandular hairs, which are more numerous on the lower leaf side. This subspecies is native to Turkey, Afghanistan, Iran, north of Iraq, northwest of

<sup>1</sup>Department of Horticultural Science, Faculty of Agriculture, Urmia University, P. O. Box: 165, Urmia, Islamic Republic of Iran.

<sup>2</sup>Research Institute of Forests and Rangelands, P. O. Box: 13185-116, Tehran, Islamic Republic of Iran.

<sup>3</sup>Research Center of Agriculture and Natural Resources of Kurdistan, P. O. Box: 66169-36311-714, Sanandaj, Islamic Republic of Iran.

\*Corresponding author; e-mail: a.hassani@urmia.ac.ir



Pakistan, and south and center of Russia. In Iran, it grows wild in north and west provinces (Ietswaart, 1982).

Since ancient times, the *Origanum* leaves have been used as a spice for flavoring of food, and it is also used for medicinal purposes, as carminative, diaphoretic, expectorant, sedative, stimulant, stomachic, diuretic, antineuralgic, antitussive, and antirheumatic (Skoula and Kamenopoulos, 1996; Afsharypour et al., 1997; Kintzios, 2002; Dundar et al., 2008). In addition, due to the presence of phenolic compounds (mainly carvacrol and occasionally thymol) in the oil, its antimicrobial, antifungal, insecticidal, and antioxidative effects have received considerable attention in recent years (Kokkini et al., 1997; Kulisic et al., 2004; Bakkali et al., 2008; Hassani et al., 2012).

The biosynthesis of secondary metabolites in plants is not only controlled by interaction between the genotype and environment, but also is strongly influenced by growth stages (harvest time), method of distillation, part of plant utilized, and postharvest drying and storage (Figueiredo et al., 2008; Nejad Ebrahimi et al., 2008; Ozkan et al., 2010; Rostaefar et al., 2017). Like in any other essential oil bearing plants, one of the most important characteristics of oil accumulation in *Origanum* species is its dependence on the developmental (ontogenetic) stage (Ozkan et al., 2010; Toncer et al., 2009). In *Thymus vulgaris*, it was reported that oil qualitative changes induced by ontogenetically variation were more than changes caused by environmental conditions (Ozguven and Tansi, 1998).

The essential oil composition of different species of *Origanum* has been studied in many researches (Sezik et al., 1993; Arnold et al., 1993; Vokou et al., 1993; Baser et al., 1994; Afsharypour et al., 1997; Toncer et al., 2009; Ozkan et al., 2010), but there are only a few reports on the essential oil of *O. vulgare* ssp. *gracile* from France (Chalchat and Bernad, 1999) and Turkey (Kilic and Bagci, 2008). Since there is no bibliographic data reporting the qualitative and quantitative variations in the essential oil of *O.*

*vulgaressp. gracile* plants in response to different growth stages, this study was conducted to investigate the effects of three phenological stages on essential oil content and composition of *O. vulgare* ssp. *gracile* plants growing wild in Iran.

## MATERIALS AND METHODS

### Plant Material

The areal parts of *O. vulgare* ssp. *gracile* were harvested at three phenological stages: pre-flowering or vegetative, just before the flower buds appeared (June 1), full flowering (July 1), and post-flowering or beginning of the seed set (August 10) stages, in 2012, from its wild habitat in the Saral Region (mountains between Sanandaj and Divandareh) (35° 24' to 35° 54' N; 46° 36' to 47° 00' E; 2,300 meter above sea level) in Kurdistan Province, Iran. A voucher specimen (No. 9428) was deposited in the herbarium of the Agriculture and Natural Resources Center of Kurdistan Province.

### Essential Oil Extraction

Plant samples from each harvest time were air dried in shade at room temperature. Essential oils were extracted by water distillation of dried and powdered leaves (25 g) for 3 hours, using Clevenger type apparatus. The essential oil content was expressed as volume per weight (v/w) based on plants material oven-dried weight. The essential oil extraction was repeated three times and the average of three replications was reported as essential oil content for each harvesting time.

### GC and GC/MS Analysis

The analysis of oils was carried out using a Shimadzu 9A gas chromatograph equipped with a Ph-5 column (30 m×0.1 mm, film thickness 0.25 µm). Oven temperature was

held at 60°C for 5 minutes and then programmed to 250°C at a rate of 3°C min<sup>-1</sup> and kept constant at 250°C for 10 minutes. Injector and detector (FID) temperature were 260°C and Helium (with 99.999% purity) was used as carrier gas with a linear velocity of 32 cm s<sup>-1</sup>. Data were calculated by electronic integration of FID peak area without using response correction factor. GC/MS analysis was also carried out on a Varian 3400 GC/MS system equipped with a DB-5 fused silica column (30 m×0.25 mm, film thickness 0.25 µm). Oven temperature program was 50-250°C at a rate of 4°C/min, transfer line temperature was 260°C, carrier gas was Helium (with 99.999% purity) with a linear velocity of 31.5 cm s<sup>-1</sup>, split ratio 1/60, with ionization energy of 70eV.

The essential oil components were identified by comparison of their mass spectra with those

of a computer library or with authentic compounds, and confirmed by comparison of their retention indices either with those of authentic compounds or with data reported in the literature (Davies, 1990; Adams, 2017).

## RESULTS AND DISCUSSION

### Essential Oil Content

The essential oil content of *O. vulgare* ssp. *gracile* differed among the three harvesting times. The highest essential oil content was obtained in post-flowering stage (1.87%) followed by full flowering (1.44%) and pre-flowering (1.01%) stages, respectively (Table 1).

Our results are in agreement with the findings of other researchers who indicated

**Table 1.** Essential oil components of *O. vulgare* ssp. *gracile* leaves in three phenological stages.

No	Compounds	RI <sup>a</sup>	RI <sup>b</sup>	Pre-flowering	Full-flowering	Post-flowering
1	α-Thujene	928	930	0.80	0.72	0.73
2	α-Pinene	937	939	0.70	0.47	0.81
3	Octen-3-ol	983	979	0.49	0.50	0.86
4	3-Octanone	991	984	3.51	3.50	5.37
5	Myrcene	999	991	-	-	0.71
6	α-Phellanderene	1005	1003	0.24	0.24	0.49
7	α-Terpinene	1015	1017	2.06	1.23	0.70
8	ρ-Cymene	1025	1025	7.76	13.54	37.08
9	1,8-Cineole	1033	1031	2.44	2.76	3.75
10	(Z)-β-ocimene	1044	1037	0.43	0.56	0.51
11	γ-Terpinene	1058	1060	21.54	13.91	6.82
12	Terpinolene	1081	1089	0.21	0.14	0.17
13	Cis-ρ-menth-2-en-1-ol	1118	1122	0.51	0.47	0.44
14	Terpinene-4-ol	1161	1177	0.56	0.94	0.86
15	α-Terpineol	1172	1189	0.15	1.07	0.61
16	Thymol methyl ether	1225	1235	0.13	0.15	-
17	Carvacrol methyl ether	1236	1245	4.82	7.19	8.54
18	Thymol	1291	1290	2.61	2.24	1.55
19	Carvacrol	1304	1299	46.62	46.50	27.60
20	E-caryophyllene	1408	1419	0.48	0.85	0.45
21	Germacrene D	1497	1485	0.28	0.24	0.18
22	Germacrene A	1505	1509	0.20	0.19	-
23	Spathulenol	1555	1578	0.21	0.22	0.24
24	Caryophyllene oxide	1560	1583	-	-	0.12
Total				96.75	97.63	98.59
Essential oil content (%)				1.01	1.44	1.87

<sup>a</sup> Retention Indices, <sup>b</sup> RI from Adams (2017).



that the essential oil content of *O. onites* was highest at the end of flowering and beginning of seed formation stage (Yaldiz et al., 2005; Ozkan et al., 2010). Similarly, the highest essential oil content of *O. vulgare* ssp. *Hirtum* (Baranauskiene et al. 2013; Krol et al. 2019), *O. vulgare* ssp. *vulgare* (Baranauskiene et al., 2013), and *Thymus vulgaris* (McGimpsey et al., 1994; Ozguven and Tansi, 1998) plants were obtained at full flowering and post-flowering (after full blooming) stages. According to Sellami et al. (2009), the low rate of volatile compounds biosynthesis during the vegetative stage may be due to partial inactivation of enzymes necessary for the biosynthesis of certain compounds.

The oil content in this study was higher than the values reported in France (0.7%) (Chalchat and Bernad, 1999) and Turkey (0.9-1.2%) (Kilic and Bagci, 2008). These differences may be due to the different environmental and genetic factors, harvesting time, as well as different essential oil extraction procedures.

### Essential Oil Composition

The essential oils of *O. vulgare* ssp. *gracile* were analyzed by GC and GC/MS. The identified essential oil components as well as the main classes and subclasses of oil constituents are shown in Tables 1 and 2, respectively. In total, 24 components were identified and quantified in three

phenological stages representing 96.75, 97.63, and 98.59% of the oil, respectively. The main constituents of leaf oils in pre-flowering (vegetative) and full flowering stages were carvacrol (46.62 and 46.5%),  $\gamma$ -terpinene (21.54 and 13.91%),  $\rho$ -cymene (7.76 and 13.54%), carvacrol methyl ether (4.82 and 7.19%), and 3-octanone (3.51 and 3.5%), respectively. In the post-flowering stage (beginning of seed set),  $\rho$ -cymene (37.08%), carvacrol (27.6%), carvacrol methyl ether (8.54%) and 3-octanone (5.37%) were the main components of essential oil. There are a few published reports on the essential oil analysis of this subspecies from different origins. The main components of *O. vulgare* ssp. *gracile* growing in Turkey were  $\beta$ -ocimene,  $\beta$ -caryophyllene and germacrene (Sezik et al., 1993),  $\beta$ -caryophyllene and thymol (Baser, 2002), and thymol,  $\gamma$ -terpinene,  $\alpha$ -terpinolene, carvacrol,  $\rho$ -cymene, carvacrol methyl ether and thymol methyl ether (Kilic and Bagci, 2008). Investigation on essential oil composition of *O. vulgare* ssp. *gracile* from France showed that oils had high levels of sabinene (Chalchat and Bernad, 1999). In another study, the main components of *O. vulgare* ssp. *gracile* oil from Italy were thymol, carvacrol and  $\rho$ -cymene (De Mastro, 1996). Qualitative differences observed in the chemical composition of the oil samples were not so remarkable. The essential oils from pre-flowering and full flowering stages were found to have a completely similar composition, but in the oil of post-flowering

**Table 2.** Main classes and subclasses of essential oil constituents of *O. vulgare* ssp. *gracile*.

Class and subclass of compounds	(%)		
	Pre-flowering	Full-flowering	Post-flowering
Monoterpenes	91.58	92.13	91.37
Oxygenated monoterpenes	57.84	61.32	43.35
Monoterpene hydrocarbons	33.74	30.81	48.02
Sesquiterpenes	1.17	1.49	0.99
Oxygenated sesquiterpenes	0.21	0.22	0.36
Sesquiterpene hydrocarbons	0.96	1.27	0.63
Others	4.00	4.00	6.23
Total identified	96.75	97.63	98.59

stage, two components (thymol methyl ether and germacrene A) were not identified. In addition, myrcene and caryophyllene oxide were present only in the post-flowering stage oil. However, from a quantitative point of view, remarkable differences existed among the essential oils. The greatest changes occurred in carvacrol,  $\rho$ -cymene and  $\gamma$ -terpinene contents. These components are biosynthetically related (Müller-Riebau *et al.*, 1997). According to a proposed biogenetic pathway,  $\gamma$ -terpinene and  $\rho$ -cymene are the precursors of thymol and carvacrol (Ozguven and Tansi, 1998). The occurrence of the highest amounts of thymol and carvacrol and the lowest values of their precursors ( $\gamma$ -terpinene+ $\rho$ -cymene) in the pre-flowering stage confirms this claim. At the post-flowering stage, the amount of carvacrol+thymol was decreased, but the amount of their precursors increased.

In total, the amounts of octen-3-ol,  $\rho$ -cymene, 1,8-cineole, carvacrol methyl ether and spathulenol increased during plant development to reach a maximum during the post-flowering stage. On the other hand, the highest content of  $\alpha$ -terpinene,  $\gamma$ -terpinene, Cis- $\rho$ -menth-2-en-1-ol, thymol, carvacrol (as major component), and germacrene D was observed in pre-flowering stage and decreased during plant development to reach its minimum percentage during the post-flowering stage.

The main classes of constituents in three oil samples were monoterpenes (91.58, 92.13 and 91.37%) and sesquiterpenes (1.17, 1.49, and 0.99%), respectively. Oxygenated monoterpenes were the dominant subclass of identified compounds in pre-flowering (57.84%) and full-flowering (61.32%) stages, with carvacrol, carvacrol methyl ether, 1,8-cineole and thymol as major components. Monoterpene hydrocarbons were the main subclass (48.2%) at the post-flowering stage due to the presence of  $\rho$ -cymene and  $\gamma$ -terpinene. It was also found that the number and amounts of sesquiterpenes were low in oils (Table 2).

The results of this study showed that a close relationship exists between the

phenological stages and the production of essential oil and phenolic compounds (mainly carvacrol, carvacrol methyl ether, and thymol) in the *Origanum* plants. In agreement with the results of this study, Krol *et al* (2019) reported that the beginning of blooming stage should be chosen when raw material with the highest flavonoids and monoterpenoid phenols (carvacrol and thymol) content is expected.

In the essential oil bearing plants, the content of the essential oil is mostly variable and many factors such as genetic variation, geographical origin, climatic conditions, plant growth phase, and postharvest processes (drying, distillation *etc.*) play an important role in yield and composition of essential oils (Figueiredo *et al.*, 2008; Nejad Ebrahimi *et al.*, 2008; Ozkan *et al.*, 2010; Rostaefar *et al.*, 2017). According to Grevsen *et al.* (2009), the development stage has a significant impact on the content and composition of both terpenes and polyphenols in *O. vulgare* ssp. *hirtum*, and the optimal harvest time depends on the compounds of interest. Hence, optimization of cultivation conditions and harvesting time to obtain higher quantity and quality of essential oils that fit market requirements is crucial. Essential oils rich in phenolic compounds, such as carvacrol and thymol, are widely reported to possess high levels of antimicrobial, antifungal, insecticidal, and antioxidative activity (Kokkini *et al.*, 1997; Kulisic *et al.*, 2004; Bakkali *et al.*, 2008; Nejad Ebrahimi *et al.*, 2008; Hassani *et al.*, 2012).

## CONCLUSIONS

The essential oils obtained from different harvesting times showed considerable variation in content and composition. Post-flowering stage could be considered the best harvesting time for the maximum essential oil content. On the other hand, the oil of this subspecies was found to be rich in the active monoterpene phenols (carvacrol and thymol) and their corresponding monoterpene



hydrocarbon precursors ( $\rho$ -cymene and  $\gamma$ -terpinene). The highest carvacrol content as the main phenolic compound was obtained at pre- and full-flowering stages. Based on the content and biological activity (antimicrobial, antifungal, antioxidant etc.) of essential oil, this subspecies should be harvested in full-flowering stage.

## REFERENCES

1. Adams, R. P. 2017. Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry. 4th Edition, Allured Publishing Corp., Carol Stream, IL, USA, 465 PP.
2. Afsharypour, S., Sajjadi, S. E. and Manesh, M. E. 1997. Volatile Constituents of *Origanum vulgare* ssp. *viride* (Syn.: *O. heracleoticum*) from Iran. *Planta Med.*, **63(2)**: 179-180.
3. Arnold, N., Bellomaria, B., Valentini, G. and Arnold, H. J. 1993. Comparative Study of the Essential Oils from Three Species of *Origanum* Growing Wild in the Eastern Mediterranean Region. *J. Essent. Oil Res.*, **5**: 71-77.
4. Bakkali, F., Averbeck, S., Averbeck, D. and Idaomar, M. 2008. Biological Effects of Essential Oils- A Review. *Food Chem. Toxicol.*, **46(2)**:446-475.
5. Barauskiene, R., Venskutonis, P. R., Dambrauskiene, E. and Viskelis, P. 2013. Harvesting Time Influences the Yield and Oil Composition of *Origanum vulgare* L. ssp. *vulgare* and ssp. *hirtum*. *Ind. Crops Prod.*, **49**: 43-51.
6. Baser, K. H. C. 2002. Aromatic Biodiversity among the Flowering Plant Taxa of Turkey. *Pure Appl. Chem.*, **74(4)**: 527-545.
7. Baser, K. H. C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1994. The Essential Oil of *Origanum vulgare* subsp. *hirtum* of Turkish Origin. *J. Essent. Oil Res.*, **6**: 31-36.
8. Chalchat, J. C. and Bernard, P. 1999. Chemical Studies of *Origanum vulgare* L. ssp. *gracile* (Koh) Ietswaart and *Origanum vulgare* L. ssp. *virens* (Hoffm. et link) Ietswaart. *J. Essent. Oil Res.*, **11**: 143-144.
9. Davies, N. W. 1990. Gas Chromatographic Retention Indices of Monoterpenes and Sesquiterpenes on Methyl Silicone and Carbowax 20M Phases. *J. Chromatogr.*, **503**: 1-24.
10. De Mastro, G. 1996. Crop Domestication and Variability within Accessions of *Origanum* Genus. *Proceedings of the IPGRI International Workshop on Oregano 8-12 May 1996*, CIHEAM, Valenzano (Bari), Italy.
11. Dundar, E., Olgun, E. G., Isiksoy, S., Kurkcuoglu, M., Baser, K. H. C. and Bal, C. 2008. The Effects of Intra-Rectal and Intra-Peritoneal Application of *Origanum monites* L. Essential Oil on 2,4,6-Trinitrobenzenesulfonic Acid-Induced Colitis in the R. *Exp. Toxicol. Pathol.*, **59(6)**: 399-408.
12. Fleisher, A. and Fleisher, Z. 1988. Identification of Biblical Hyssop and Origin of the Traditional Use of Oregano-Group Herbs in the Mediterranean Region. *Econ. Bot.*, **42(2)**: 232-241.
13. Figueiredo, A. C., Barroso, J. G., Pedro, L. G. and Scheffer, J. J. C. 2008. Factors Affecting Secondary Metabolite Production in Plants: Volatile Components and Essential Oils. *Flavour Fragrance J.*, **23(4)**: 213-226.
14. Grevsen, K., Frette, X. C. and Christensen, L. P. 2009. Content and Composition of Volatile Terpenes, Flavonoids and Phenolic Acids in Greek Oregano (*Origanum vulgare* L. ssp. *hirtum*) at Different Development Stages during Cultivation in Cool Temperate Climate. *Europ. J. Hort. Sci.*, **74(5)**: 193-203.
15. Hassani, A., Fathi, Z., Ghosta, Y., Abdollahi, A., Meshkatsadat, M. H. and Jalili Marandi, R. 2012. Evaluation of Plant Essential Oils for Control of Postharvest Brown and Gray Mold Rots on Apricot. *J. Food Saf.*, **32(1)**: 94-101.
16. Ietswaart, J. H. 1980. A taxonomic Revision of the Genus *Origanum* (Labiatae). PhD Thesis, Leiden Botanical Series 4. Leiden University Press, The Hague.
17. Ietswaart, J. H. 1982. *Origanum*. In: "Flora Iranica", (Ed.): Rechinger, K. H. Amsterdam, No. 150, Akademische Druck-u-Verlagsanstalt, Graz, PP: 527-532.
18. Kilic, O. and Bagci, E. 2008. A Study on the Essential Oil Composition of the *Origanum vulgare* L. subsp. *gracile* and the Investigation Probability of Using as Herbal Tea. *Sci. Eng. J. Firat Univ.*, **20(1)**: 83-89.

19. Kintzios, S. E. 2002. Profile of the Multifaceted Prince of the Herbs. In: "Oregano: The Genera *Origanum* and *lippie*", (Ed.): Kintzios S. E. Medicinal and Aromatic Plants, Industrial Profiles 25. Taylor & Francis/CRC Press, USA, PP. 3-9.
20. Kokkini, S., Karousou, R., Dardioti, A., Krigas, N. and Lanaras, T. 1997. Autumn Essential Oils of Greek Oregano. *Phytochemistry*, **44(5)**: 883-886.
21. Król, B., Kołodziej, B., Kędzia, B., Hołderna-Kędzia, E., Sugier, D. and Luchowska, K. 2019. Date of Harvesting Affects Yields and Quality of *Origanum vulgare* ssp. *hirtum* (Link) Ietswaart. *J. Sci. Food Agric.*, **99(12)**: 5432-5443.
22. Kulisic, T., Radoni, A., Katalinic, V. and Milos, M. 2004. Use of Different Methods for Testing Antioxidative Activity of Oregano Essential Oil. *Food Chem.*, **85**: 633-640.
23. LaGow, B. 2004. *PDR for Herbal Medicine*. 3<sup>rd</sup> Edition, Thomson PDR, USA, PP. 609-610, 808-809.
24. McGimpsey, J. A., Douglas, M. H., Van Klink, J. W., Beauregard, D. A. and Perry, N. B. 1994. Seasonal Variation in Essential Oil Yield and Composition from Naturalized *Thymus vulgais* L. in New Zealand. *Flavour. Fragrance. J.*, **9(6)**: 347-352.
25. Mozaffarian, V. 1996. *A Dictionary of Iranian Plant Names*. Farhang-Moaser Publication, Tehran, Iran.
26. Müller-Riebau, F. J., Berger, B. M., Yegen, O. and Cakir, C. 1997. Seasonal Variations in the Chemical Compositions of Essential Oils of Selected Aromatic Plants Growing Wild in Turkey. *J. Agric. Food Chem.*, **45(12)**: 4821-4825.
27. Nejad Ebrahimi, S., Hadian, J., Mirjalili, M. H., Sonboli, A. and Yousefzadi, M. 2008. Essential Oil Composition and Antibacterial Activity of *Thymus caramanicus* at Different Phenological Stages. *Food Chem.*, **110(4)**: 927-931.
28. Ozguven, M. and Tansi, S. 1998. Drug Yield and Essential Oil of *Thymus vulgaris* L. as in Influenced by Ecological and Ontogenetical Variation. *Turk. J. Agric. For.*, **22(6)**: 537-542.
29. Ozkan, G., Baydar, H. and Erbas, S. 2010. The Influence of Harvest Time on Essential Oil Composition, Phenolic Constituents and Antioxidant Properties of Turkish Oregano (*Origanum onites* L.). *J. Sci. Food. Agric.*, **90(2)**: 205-209.
30. Rostaefar, A., Hassani, A. and Sefidkon, F. 2017. Seasonal Variations of Essential Oil Content and Composition in Male and Female Plants of *Juniperus communis* L. ssp. *hemisphaerica* Growing Wild in Iran. *J. Essent. Oil Res.*, **29(4)**: 357-360.
31. Sellami, I. H., Maamouri, E., Chahed, T., Wannes, W. A., Kchouk, M. E. and Marzouk, B. 2009. Effect of Growth Stage on the Content and Composition of the Essential Oil and Phenolic Fraction of Sweet Marjoram (*Origanum majorana* L.). *Ind. Crops Prod.*, **30(3)**: 395-402.
32. Sezik, E., Tümen, G., Kirimer, N., Özek T. and Baser, K. H. C. 1993. Essential Oil Composition in four *Origanum vulgare* Subspecies of Anatolian Origin. *J. Essent. Oil Res.* **5**: 425-431.
33. Skoula, M. and Kamenopoulos, S. 1996. *Origanum dictamnus* L. and *Origanum vulgare* L. subsp. *Hirtum* (Link) Ietswaart: Traditional Uses and Production in Greece. *Proceedings of the IPGRI International Workshop on Oregano*, 8-12 May 1996, CIHEAM, Valenzano (Bari), Italy.
34. Spada, P. and Perrino, P. 1997. Conservation of Oregano Species in National and International Collections: An Assessment. *Proceedings of the IPGRI International Workshop on Oregano*, 8-12 May 1996, CIHEAM, Valenzano (Bari), Italy.
35. Toncer, O., Karaman, S., Kizil S. and Diraz, E. 2009. Changes in Essential Oil Composition of Oregano (*Origanum onites* L.) Due to Diurnal Variations at Different Development Stages. *Not. Bot. Hort. Agrobot. Cluj.*, **37(2)**: 177-181.
36. Vokou, D., Kokkini, S. and Bessiere, J. M. 1993. Geographic Variation of Greek Oregano (*Origanum vulgare* ssp. *Hirtum*) Essential Oils. *Biochem. Syst. Ecol.*, **21(2)**: 287-295.
37. Yaldiz, G., Sekeroglu, N., Ozguven, M. and Kirpik, M. 2005. Seasonal and Diurnal Variability of Essential Oil and Its Components in *Origanum onites* L. Grown in the Ecological Conditions of Cukurova. *Grasas Aceites.*, **56(4)**: 254-258.



تغییرات کمی و کیفی اسانس مرزنجوش بخارایی (*Origanum vulgare* ssp. *gracile*) تحت تاثیر زمان‌های مختلف برداشت

م. مرادی، ع. حسنی، ف. سفیدکن و ح. معروفی

چکیده

تغییرات کمی و کیفی اسانس گیاهان خودروی مرزنجوش بخارایی (*Origanum vulgare* ssp. *gracile*) تحت تاثیر مراحل فنولوژیکی مختلف (قبل از گلدهی، گلدهی کامل و پس از گلدهی) مورد مطالعه قرار گرفت. برای این منظور اسانس برگ‌های خشک شده به روش تقطیر با آب و با استفاده از دستگاه کلونجر استخراج شده و به وسیله‌ی دستگاه‌های کروماتوگرافی گازی (GC) و کروماتوگرافی گازی متصل به طیف‌سنج جرمی (GC/MS) مورد آنالیز قرار گرفتند. بیشترین (۱/۸۷٪) و کمترین (۱/۰۱٪) محتوی اسانس به ترتیب در مراحل پس از گلدهی و قبل از گلدهی بدست آمد. در مجموع، ۲۴ ترکیب که به ترتیب ۹۶/۷۵٪، ۹۷/۶۳٪ و ۹۸/۵۹٪ اسانس سه مرحله فنولوژیکی را تشکیل می‌دادند شناسایی گردید. کارواکرول (۴۶/۶۲٪، ۴۶/۵٪ و ۲۷/۶٪)، پارا-سیمن (۷/۷۶٪، ۱۳/۵۴٪ و ۳۷/۰۸٪) و گاما-ترپنین (۲۱/۵۴٪، ۱۳/۹۱٪ و ۶/۸۲٪) ترکیبات مهم اسانس به ترتیب در مراحل قبل از گلدهی، گلدهی کامل و پس از گلدهی بودند. مونوترپن‌های اکسیژندار (۶۱/۳۲٪ - ۴۳/۳۵٪) و مونوترپن‌های هیدروکربنی (۴۸/۰۲٪ - ۳۰/۸۱٪) گروه‌های عمده ترکیبات شناسایی شده در اسانس سه مرحله بودند. طبق یافته‌های این تحقیق، برای بدست آوردن بالاترین محتوی اسانس، مرحله‌ی پس از گلدهی مناسب‌ترین زمان برداشت است اما برای حصول بالاترین مقادیر ترکیبات فنلی در اسانس، برداشت در مراحل قبل از گلدهی و گلدهی کامل قابل توصیه می‌باشد.