

Successive Rearing on *Sitotroga cerealella* Affects Quality of the Parasitoid Wasp *Trichogramma embryophagum* (Hymenoptera: Trichogrammatidae)

N. Taghikhani¹, A. Sahragard^{1*}, Y. Fathipour^{2*}, and Kh. Madahi¹

ABSTRACT

Demographic parameters of the parasitoid wasp *Trichogramma embryophagum* Hartig reared on *Sitotroga cerealella* (Olivier) were determined for 40 generations (G5-G40). The experiments were done in a growth chamber at $26\pm 2^{\circ}\text{C}$, $65\pm 5\%$ RH and photoperiod of 16:8 h (L:D). The results showed that the female longevity decreased significantly during successive production, ranging from 10.68 (in G5) to 9.64 days (in G40). On the other hand, the male longevity decreased significantly in the 20th generation and no significant difference was found from G20 to G40. The oviposition days and mean total fecundity of *T. embryophagum* decreased as the number of generations increased. Generally, the wasps in earlier generations had longer adult longevity, longer life span, and higher fecundity than later generations. Moreover, sex ratio of *T. embryophagum* was not significantly different in successive generations. Results of paired bootstrap test indicated that all population growth parameters of *T. embryophagum* reared on *S. cerealella* were significantly different in successive generations. The highest and lowest values of the intrinsic rate of increase (r), finite rate of increase (λ), net Reproductive rate (R_0), Gross Reproductive Rate (GRR) and mean generation Time (T) of *T. embryophagum* were observed in G40 and G5, respectively. These results suggest that *T. embryophagum* wasps reared in sequential generations can be used successfully in biological control programs until the 20th generation without any loss of quality or performance; after that, regular rejuvenation of laboratory population by occasional importing of field-collected parasitoids should be done.

Keywords: Different generations, Life table, Population growth parameters, Quality control, Rejuvenating population, *Trichogramma*.

INTRODUCTION

Many species of *Trichogramma* egg parasitoids (Hymenoptera: Trichogrammatidae) are among the most widely distributed biocontrol agents, which have long been utilized successfully in biological control programs against several lepidopteran pests in agricultural and forest areas (Smith, 1996; Razinger *et al.*, 2016). Adult females attack the egg stage of their host and kill the target pest before crop damage by larval feeding start. The

augmentative release of these parasitoids has reduced pest damage by 77-92% in some crops such as sugarcane, wheat, corn, and cabbage in several countries (Li, 1994; Parra, 2010). In Iran, several *Trichogramma* species have been recorded (Ebrahimi *et al.*, 1998; Moghaddassi *et al.*, 2019) and are being mass-reared and released (Chamaani and Poorjavad, 2020). Among them, *T. embryophagum* have been noted to parasitize some important pests including *Ectomyelois ceratoniae* (Zell.) and *Cydia pomonella* L. (Poorjavad *et al.*, 2011) and have potential to

¹ Department of Plant Protection, Faculty of Agricultural Sciences, University of Guilan, Rasht, Islamic Republic of Iran.

² Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Islamic Republic of Iran.

*Corresponding authors; e-mail: sahragard@guilan.ac.ir or e-mail: fathi@modares.ac.ir



be inexpensively produced in large numbers and easily released in fields (Haghani and Fathipour, 2003). *Trichogramma* parasitoids are usually produced on factitious hosts (Li et al., 2019), among them, Angoumois grain moth, *Sitotroga cerealella* (Olivier) (Lepidoptera: Gelechiidae) is considered as a common host of *Trichogramma*, due to its easy and affordable rearing in laboratories (Smith, 1996).

Studying the basic demography of the mass reared biological control agents and determining their efficiency is the first step to conduct successful biological control programs (Ghaemmaghani et al., 2021). However, long-term mass rearing can affect biological attributes, production efficiency and the quality of the mass-reared agents (Sørensen et al., 2012; Khanamani et al., 2017). Thus, an important step in effective mass rearing of natural enemies is to determine the best duration of their production without any unfavorable effect on the biological characteristics (Sørensen et al., 2012). Both inbreeding depression and random genetic drift, which occurred during long-term rearing of natural enemies, led to their lower genetic variability and poor quality, and then resulted in failure of biological control programs (Cônoli et al., 2010). Because of these, the quality control is a crucial issue in mass rearing programs (Pratissoli et al., 2004; Parra, 2010).

Assessment of the quality of artificially reared natural enemies as suggested by the current quality control guidelines are usually done in the laboratory with easily measured life table parameters, such as longevity, fecundity, viability, and sex ratio (Cascone et al., 2015). Life Table studies are effective and comprehensive tools that provide the most detailed information about insect's population dynamics (Khanamani et al., 2017).

A significant number of scientific papers is related to the effects of rearing parasitoids on artificial or natural host eggs only in a single generation (e.g., Grenier and De Clercq, 2003; Cônoli et al., 2010); only few studies have investigated the effects of continuous

production for several generations (Nordlund et al., 1997; Pratissoli et al., 2004; Lü et al., 2015, 2017). In a previous study, adult longevity of *Trichogramma pretiosum* Riley reared on *Anagasta kuehniella* (Zeller) for 23 generations reduced significantly in the last generations (Pratissoli et al., 2004). Also, Ghaemmaghani et al. (2021) showed that mass rearing of *Trichogramma brassicae* (Bezdenko) for 45 generations on *S. cerealella* led to no loss of quality or performance until the 15th generation, but laboratory mass reared colonies declined in quality after G15. The importance of quality control of long-term mass reared natural enemies have been studied in several research works (Bellutti, 2011; Khanamani et al., 2017).

Despite being a species with high economic importance, the biology of *T. embryophagum* is poorly studied (Haghani and Fathipour, 2003; Poorjavad et al., 2011), and no studies have been carried out in the past to determinate its quality control in long-term mass rearing. Thus, the aim of this study was to determine biological characteristic and population parameters of *T. embryophagum* successively reared on *S. cerealella* eggs for 40 generations, in order to understand the effect of sequential mass rearing on its quality.

MATERIALS AND METHODS

Parasitoids

Trichogramma embryophagum was originally collected from the apple orchards located at the suburbs of Karaj (Alborz Province, Iran) by egg traps (pieces of white papers [21×11 cm] containing *S. cerealella* eggs). Egg traps were placed in different parts of the orchards and recollected after 24 hours. Then, the papers were incubated in a growth chamber at 26±2°C, 65±5% RH and 16:8 h (L:D) photoperiod. Stock colonies of *T. embryophagum* were reared on eggs of *S. cerealella* as a factitious host, in the

laboratory under the same conditions mentioned above.

Host

A colony of Angoumois grain moth (*S. cerealella*) was established on barley grains under the same conditions mentioned above. When the flight of the first adults began, they were collected and transferred to funnels (diameter 200 mm) covered with a fine mesh net over the mouth. After mating, the eggs were collected on papers placed under the open side of the funnel.

Assessment of Biological Parameters

Approximately 80 one-day-old parasitized eggs of *S. cerealella* were selected randomly from the colony and kept in a glass tube (10×1.6 cm) until the adult wasps emerged. Pairs of newly emerged male and female *T. embryophagum* were moved into single glass tubes (10×1.6 cm). About 150 eggs of *S. cerealella* stuck to a white paper tape (1×7 cm) were supplied to each tube. Diluted honey (30%) was smeared on the tube wall to feed the parasitoid. If a male died, it was replaced by a newly emerged one (< 24 hours old) from the stock colony. These individuals were omitted from the analyses. When the adults emerged, mortality, survivorship, adult longevity, and the number of parasitized eggs were recorded daily, until the death of the last individual. These procedures were repeated for generations 5, 10, 15, 20, 25, 30, 35, and 40 of the *T. embryophagum* colony. All observations were done in a growth chamber under the same conditions mentioned above.

Statistical Analysis

The life histories data were analyzed by age-stage, two-sex life table theory (Chi and Liu, 1985) and the method described by Chi (1988). All developmental stages and reproductive attributes of different

generations of *T. embryophagum* were subjected to normality testing (Kolmogorov–Smirnov test) and were compared using one-way Analysis Of Variance (ANOVA) and Tukey's test at $P < 0.05$. Statistical analyses were conducted using SAS (v. 9.2) software. The TWOSEX-MSChart program (Chi, 2020) was used to calculate the population parameters including the intrinsic rate of increase (r), net Reproductive rate (R_0), finite rate of increase (λ), Gross Reproductive Rate (GRR), and mean generation Time (T). Bootstrap procedure was used for estimating the standard errors of population parameters with 100,000 resampling (Huang and Chi, 2012). Comparison of bootstrap values of all parameters among different generations of *T. embryophagum* was done using paired bootstrap test procedure (Riahi *et al.*, 2017; Bahari *et al.*, 2018). The sex ratios in different generations were compared based on Chi-square test by SPSS (v. 22). Figures were drawn by Microsoft Excel 2016.

RESULTS

Parasitoid Life Stage Duration, Reproductive Characteristics, and Sex Ratio

The mean duration of different life stages of *T. embryophagum* are shown in Table 1. The egg, larval, and pupal stages occurred inside the host's eggs and all were grouped as "preadult." As the results show, preadult period took approximately 9.95-10 days to be completed, and no significant difference was found among generations. The female longevity varied significantly among sequential generations. The longest female longevity was recorded in G5 and G10; and rearing *T. embryophagum* on *S. cerealella* for G35 and G40 led to the shortest female longevity. On the other hand, the male longevity did not differ significantly between G5, G10, and G15, but hereinafter it decreased significantly and no significant difference was found from G20 to G40. Significant differences were found in the total



life span among different generations of *T. embryophagum* (Table 1). The longest total life span was observed for G5, and during the sequential generations, it was gradually decreased and the lowest total life span was obtained for G40.

The oviposition days of *T. embryophagum* decreased as the number of generations increased, such that the longest oviposition days was recorded in G5 and G10 and the shortest oviposition days was obtained in

G40. The mean total fecundity was highest and lowest in G5 and G40, respectively. Generally, the wasps in earlier generations had longer adult longevity, longer total life span, and higher fecundity than later generations.

Sex ratio (number of females/number of females and males) of *T. embryophagum* was not significantly different among different generations tested (Figure 1).

Table 1. Comparative duration of different developmental stages and reproductive attributes of sequential generations of *Trichogramma embryophagum* reared on *Sitotroga cerealella* eggs.^A

Generation	Number of individuals	Preadult (Day)	Adult longevity (Female) (Day)	Adult longevity (Male) (Day)	Total life span (Day)	Oviposition days (Day)	Fecundity (Eggs)
5	77	9.95 ± 0.03 a	11.68 ± 0.12 a	11.21 ± 0.15 a	19.64 ± 0.49 a	10.29 ± 0.16 a	81.51 ± 2.00 a
10	78	9.98 ± 0.02 a	11.49 ± 0.13 ab	11.09 ± 0.17 a	19.13 ± 0.52 a	10.05 ± 0.16 ab	76.18 ± 1.59 b
15	76	9.97 ± 0.02 a	11.15 ± 0.14 b	11.05 ± 0.15 a	18.89 ± 0.52 abc	9.77 ± 0.15 b	72.92 ± 1.80 b
20	78	9.95 ± 0.02 a	10.64 ± 0.03 c	10.32 ± 0.03 b	18.13 ± 0.49 abcd	9.13 ± 0.16 c	65.00 ± 1.54 c
25	78	10.00 ± 0.02 a	10.24 ± 0.12 d	9.95 ± 0.10 b	17.67 ± 0.50 bcd	8.65 ± 0.14 d	61.65 ± 1.42 cd
30	78	10.00 ± 0.02 a	10.03 ± 0.11 d	9.95 ± 0.12 b	17.56 ± 0.49 cd	8.43 ± 0.12 d	59.30 ± 1.57 de
35	78	10.00 ± 0.02 a	9.94 ± 0.12 de	9.70 ± 0.12 b	17.49 ± 0.49 cd	8.2 ± 0.15 d	56.61 ± 1.26 e
40	78	10.00 ± 0.02 a	9.64 ± 0.10 e	9.68 ± 0.15 b	16.81 ± 0.51 d	7.78 ± 0.14 e	48.03 ± 1.18 f
df		7, 481	7, 305	7, 175	7, 620	7, 305	7, 305
F		1.35	38.66	21.90	3.73	38.16	49.95
P		0.226	< 0.0001	< 0.0001	0.0006	< 0.0001	< 0.0001

^A The means followed by the same letter in each column are not significantly different (Mean±SE) (P < 0.05, Tukey's test).




Figure 1. Sex ratio (females/males+females) of sequential generations of *Trichogramma embryophagum* reared on *Sitotroga cerealella* eggs. The sex ratio in generations with the same letter are not significantly different (P < 0.05, Chi-square test).

For *T. embryophagum*, the age-stage-specific survival rate (s_{xj}), which is the probability of a newborn surviving to age x and stage j , was found to overlap among stages, demonstrating the variable

developmental rates among individuals (Figure 2). The age-specific survivorship (l_x), age-stage-specific fecundity of female (f_{xj}), age-specific fecundity of population (m_x), and maternity ($l_x m_x$) of sequential generations of

T. embryophagum reared on *S. cerealella* eggs are plotted in Figure 3. The age-specific survivorship (l_x) of newly emerged *T. embryophagum* in G5-G40 were obtained as 0.96, 0.97, 0.80, 0.78, 0.75, 0.76, 0.76, and 0.71, respectively. The obtained results showed that f_{xj} of *T. embryophagum* at the first day of oviposition was 21.5, 21, 24, 21.15, 19.76, 19.76, 19.25, and 31.17 eggs, respectively. In addition, the highest value of m_x of *T. embryophagum* in G5-G40 were 14.46, 14.29, 14.11, 13.52, 12.39, 12.39, 11.75, and 11.33 eggs, respectively. According to these graphs, the highest fecundity was obtained in the first two days after the emergence of females.

The age-stage reproductive value of (v_{xj}) of *T. embryophagum* indicates the contribution

of each individual at age x and stage j to population growth. The results revealed that female made the highest contribution to the population growth in days 9, 9, 9, 10, 10, 10, 10 and 10, respectively, as 58.48, 56.77, 58.99, 43.76, 41.81, 40.93, 39.60, and 34.60 (Figure 4).

The value of age-specific life expectancy (e_{xj}) of *T. embryophagum* on *S. cerealella* in successive generations is plotted in Figure 5. Life expectancy of newly emerged females in G5-G40 were 12.63, 12.46, 12.13, 10.59, 10.24, 10.03, 9.94, and 9.64 days, respectively. Also, it was estimated as 12.17, 11.09, 12.00, 11.05, 9.95, 9.95, 9.72, and 9.68 days for adult males.




Figure 2. Age-stage-specific survival rate (s_{xj}) of sequential generations (G5-G40) of *Trichogramma embryophagum* reared on *Sitotroga cerealella* eggs.




Figure 3. Age-specific survivorship (l_x), age-stage specific fecundity of female (f_{xj}) (eggs), age-specific fecundity (m_x), and age-specific maternity ($l_x m_x$) of sequential generations (G5-G40) of *Trichogramma embryophagum* reared on *Sitotroga cerealella* eggs.




Figure 4. Age-stage-specific reproductive value (v_{xj}) of sequential generations (G5-G40) of *Trichogramma embryophagum* reared on *Sitotroga cerealella* eggs.




Figure 5. Age-stage life expectancy (e_{xj}) of sequential generations (G5-G40) of *Trichogramma embryophagum* reared on *Sitotroga cerealella* eggs.

Population Growth Parameters

Results of paired bootstrap test indicated that all population growth parameters of *T. embryophagum* reared on *S. cerealella* were significantly different in successive

generations. The highest and lowest values of the Gross Reproductive Rate (GRR), net Reproductive rate (R_0), intrinsic rate of increase (r), and finite rate of increase (λ) of *T. embryophagum* were observed in G5 and G40, respectively (Table 2).

Table 2. Life table parameters of sequential generations of *Trichogramma embryophagum* reared on *Sitotroga cerealella* eggs.^A

Generation	GRR (Eggs/Individual)	R_0 (Eggs/Individual)	r (d^{-1})	λ (d^{-1})	T (Day)
5	51.60 ± 5.056 a	43.402 ± 4.77 a	0.301 ± 0.009 a	1.351 ± 0.012 a	12.530 ± 0.055 a
10	50.08 ± 4.720 ab	40.397 ± 4.393 ab	0.295 ± 0.009 ab	1.343 ± 0.012 ab	12.546 ± 0.043 ab
15	46.67 ± 4.643 ab	37.420 ± 4.289 abc	0.291 ± 0.009 abc	1.337 ± 0.013 abc	12.468 ± 0.053 abc
20	41.88 ± 4.128 abc	32.500 ± 3.753 abc	0.280 ± 0.009 abc	1.324 ± 0.013 abc	12.411 ± 0.032 bc
25	38.73 ± 3.992 bcd	29.244 ± 3.553 bcd	0.272 ± 0.010 bcd	1.312 ± 0.013 bcd	12.429 ± 0.033 abc
30	37.20 ± 3.87 bcd	28.128 ± 3.429 bcd	0.269 ± 0.010 bcd	1.309 ± 0.013 bcd	12.374 ± 0.039 bcd
35	34.56 ± 3.684 cd	26.128 ± 3.245 cd	0.264 ± 0.010 cd	1.302 ± 0.014 cd	12.352 ± 0.038 cd
40	31.44 ± 3.173 d	22.167 ± 2.762 cd	0.252 ± 0.010 d	1.286 ± 0.013 cd	12.302 ± 0.035 d

^A The means followed by the same letter in each column are not significantly different (Mean±SE) ($P < 0.05$, Paired bootstrap test).



DISCUSSION

Production of biocontrol agents has received much attention in recent years (van Lenteren, 2012; Khanamani et al., 2021). One of the most common natural enemies of agricultural pests are egg parasitoids in the genus *Trichogramma*, which are produced and used in large numbers each year (Cônsooli et al., 2010; Moghaddassi et al., 2019). *T. embryophagum* as one of the most important species of *Trichogramma* has been collected from several regions and reared in several insectaries (Mirkarimi, 2000). In insectaries, long-term mass rearing of parasitoid wasps may decrease their quality, which leads to declining efficiency of integrated pest management programs. Accordingly, frequently assessing the quality of biocontrol agents under successive rearing is vital (Khanamani et al., 2017). In this study, the effect of rearing *T. embryophagum* on *S. cerealella* eggs over 40 sequential generations on its quality were determined using biological characteristics and life table parameters. According to our findings, biological attributes and population parameters of *T. embryophagum* differed significantly among successive generations.

Several factors influence parasitoids' longevity, including species of parasitoid and factitious hosts, population genetic patterns, and rearing conditions. Rearing *T. embryophagum* over 40 generations on *S. cerealella* eggs led to a significant decrease in the male and female longevity. This result is consistent with results obtained by other researchers in different species of *Trichogramma* wasps. For instance, Pratisoli et al. (2004) stated that male and female longevity of *T. pretiosum* decreased significantly over 23 generations. Female longevity of *T. pretiosum* reared on *A. kuehniella* was almost constant in the first 10 generations and then decreased in subsequent generations, up to the 100th generation (Freitas Bueno et al., 2006). This decreasing trend in male and female longevity after a certain number of generations in mass rearing

was also observed in the study of Lü et al. (2017), and Ghaemmaghami et al. (2021).

One of the most important issues in mass rearing of natural enemies is the number of sequential generations that a biological control agent can be mass-reared without a significant reduction in its quality and efficiency (van Lenteren et al., 2002). In the present study, the early generations of *T. embryophagum* reared on *S. cerealella* eggs had the longest female and male longevity, total life span, the number of oviposition days, and fecundity. This shows that parasitoids may demand more energy to complete their life cycle after a certain number of generations without obtaining major benefits (Nordlund et al., 1997; Pratisoli et al., 2004; Lü et al., 2017).

The sex ratio is an economically important trait in assessing the quality of parasitoid wasps (Cerutti and Bigler, 1995; Hassan and Zhang, 2001), which plays an essential role in the dynamics of the insect population, affects the performance of the reared population, and also has a significant effect on the financial profitability of mass rearing (Pipoly et al., 2015). In this research, no significant difference was observed between sex ratio of *T. embryophagum* on *S. cerealella* eggs in different generations ($P= 0.99$). This result differed from the findings of Pratisoli et al. (2004), Lü et al. (2017), and Ghaemmaghami et al. (2021) who showed a fluctuation in the sex ratio of other *Trichogramma* species.

As shown in Figures 2 and 3, the survival of *T. embryophagum* decreased over 40th generations, which is consistent with those results reported in other *Trichogramma* species (Nordlund et al., 1997; Pratisoli et al., 2004; Lü et al., 2015).

The mean generation Time (T) is the period of time it takes for a population to increase to R_0 -folds of its current size at the stable age-stage distribution. Natural logarithm of R_0 divided by r yields this parameter and, therefore, its value is expected to increase with decreasing r . Contrary to expectations, our results showed that T value decreased by decreasing r , which is attributed to the proportion of R_0 and r in successive

generations, such that the value of R_0 decreased more than r . This might be considered one of the study's main findings because it can lead to a larger population size in less time.

The obtained results showed that population parameters of *T. embryophagum* were affected by long-term mass rearing and the highest values of intrinsic rate of increase (r), finite rate of increase (λ), net Reproductive rate (R_0), Gross Reproductive Rate (GRR), and mean generation Time (T) were found in the first generations (up to 20th generation), and all the life table parameters decreased after the 25th generation. The r value is one of the most important criteria used to evaluate the efficiency of natural enemies (Fathipour and Maleknia, 2016). The highest r value was found for parasitoids of G5–G20, indicating their better fitness compared to the previous generations. Similar to the results of this study, a decrease in the quality of natural enemies has been reported by Lü *et al.* (2017) in *Trichogramma dendrolimi* Matsumura and Ghaemmaghani *et al.* (2021) in *T. brassicae*.

Inbreeding, which is the most common form of non-random mating, should be among the main reasons for this decline in fitness over sequential generations (Kazmer and Luck, 1991; van Lenteren and Bigler, 2010). Using small egg cards in the rearing process can increase opportunities for inbreeding, and might gradually decrease the genetic quality of reared parasitoids (Paspati *et al.*, 2019). Contradictory reports have been made about the occurrence of inbreeding in *Trichogramma* wasps. In spite of Sorati *et al.* (1996) who described lack of this phenomenon in the mass rearing of *T. brassicae*, Antolin (1999) reported the negative effect of inbreeding on reproduction and sexual ratio of *T. pretiosum*. According to these contradictory reports, confirmation of the occurrence of inbreeding in long-term mass rearing of *T. embryophagum* requires more detailed studies. Moreover, deformed adults that remained in the colony during the continuous rearing could lead to the reduction

of parasitoids' quality (Suzuki and Hiehata, 1985).

Based on the obtained results, long-term mass rearing on *S. cerealella* eggs up to the 20th generation has no adverse effect on biological characteristics and population parameters of *T. embryophagum*, and the wasps reared in sequential generations up to this generation can be used successfully in biological control programs without any significant loss of quality or performance. Indeed, continuous long-term mass rearing of natural enemies in insectariums may cause the biological control program to fail, and it is necessary to consider strategies to improve the quality of the produced parasitoids. Therefore, to keep the performance fixed, after the 20th generation, the laboratory population should be rejuvenated every few generations by adding field-collected individuals.

REFERENCES

1. Antolin, M. F. 1999. A Genetic Perspective on Mating Systems and Sex Ratios of Parasitoid Wasps. *Res. Popul. Ecol.*, **41**: 29-37.
2. Bahari, F., Fathipour, Y., Talebi, A. A. and Alipour, Z. 2018. Long-Term Feeding on Greenhouse Cucumber Affects Life Table Parameters of Two-Spotted Spider Mite and Its Predator, *Phytoseiulus persimilis*. *Syst. Appl. Acarol.*, **23**: 2304–2316.
3. Bellutti, N. 2011. *Effects of Mass Rearing on Life-History Traits of an Invasive Fruit Moth Species, Grapholita molesta (Busck)*. Institute of Forest Entomology, Forestathology and Forest Protection, BOKU, Vienna, 35 PP.
4. Cascone, P., Carpenito, S., Slotsbo, S., Iodice, L., Sørensen, J. G., Holmstrup, M. and Guerrieri, E. 2015. Improving the Efficiency of *Trichogramma achaeae* to Control *Tuta absoluta*. *BioControl.*, **60**: 761–771.
5. Cerutti, F. and Bigler, F. 1995. Quality Assessment of *Trichogramma brassicae* in the Laboratory. *Entomol. Exp. Appl.*, **75**: 19-26.
6. Chamaani, N. and Poorjavad, N. 2020. Low Efficiency of Four Indigenous



- Trichogramma* Wasp Populations, Collected from Tomato Crops, in Controlling the Invasive Pest *Tuta absoluta* in Iran. *Bull. Insectol.*, **73(2)**: 171-180.
7. Chi, H. and Liu, H. 1985. Two New Methods for the Study of Insect Population Ecology. *Bull. Inst. Zool. Acad. Sin.*, **24**: 225-240.
 8. Chi, H. 1988. Life-Table Analysis Incorporating Both Sexes and Variable Development Rates among Individuals. *Environ. Entomol.*, **17**: 26-34.
 9. Chi, H. 2020. *TWOSEX-MSChart: A Computer Program for the Age-Stage, Two-Sex Life Table Analysis*. Available in: <http://140.120.197.173/Ecology/Download/TWOSEX-MSChart.rar>. (Accessed on January 09, 2020).
 10. Cônsoli, F., Parra, J. R. and Zucchi, R. 2010. *Egg Parasitoids in Agroecosystems with Emphasis on Trichogramma*. Springer, The Netherlands.
 11. Do Thi Khanh, H., Chailleux, A., Tiradon, M., Desneux, N., Colombel, E. and Tabone, E. 2012. Using New Egg Parasitoids (*Trichogramma* spp.) to Improve Integrated Management against *Tuta absoluta*. *EPPO Bull.*, **42**: 249-254.
 12. Ebrahimi, E., Pintureau, B. and Shojai, M. 1998. Morphological and Enzymatic Study of the Genus *Trichogramma* in Iran. *J. Appl. Entomol. Pathol.*, **66(21)**: 39-43.
 13. Fathipour, Y. and Maleknia, B. 2016. Mite Predators. In: "Ecofriendly Pest Management for Food Security", (Ed.): Omkar. Elsevier, San Diego, USA.
 14. Freitas Bueno, R. C. O., Freitas Bueno, A., Pratisoli, D., Vieira, S., Oliveira, L. J., Barros, E. M. and Jakoby, G. L. 2006. Biological Characteristics of *Trichogramma pretiosum* (Riley) Reared on *Anagasta kuehniella* (Zeller) for 100 Generations. *Revista. Ecosistemas.*, **31**: 69-75.
 15. Ghaemmaghami, E., Fathipour, Y., Bagheri, A., Talebi, A. A. and Reddy, G. V. P. 2021. Quality Control of the Parasitoid Wasp *Trichogramma brassicae* (Hymenoptera: Trichogrammatidae) over 45 Generations of Rearing on *Sitotroga cerealella*. *Insect Sci.*, **28**: 180-190.
 16. Grenier, S. and De Clercq, P. 2003. *Quality Control and Production of Biological Control Agents: Theory and Testing Procedures*. CABI Publishing, Wallingford.
 17. Haghani, M. and Fathipour, Y. 2003. The Effect of the Type of Laboratory Host on the Population Growth Parameters of *Trichogramma embryophagum* Hartig (Hym., Trichogrammatidae). *J. Agric. Sci. Natur. Resour.*, **10(2)**: 117-124.
 18. Hassan, S. A. and Zhang, W. Q. 2001. Variability in Quality of *Trichogramma brassicae* (Hymenoptera: Trichogrammatidae) from Commercial Suppliers in Germany. *Biol. Control*, **22**: 115-121.
 19. Huang, Y. B. and Chi, H. 2012. Age-Stage, Two-Sex Life Tables of *Bactrocera cucurbitae* (Coquillett) (Diptera: Tephritidae) with a Discussion on the Problem of Applying Female Age-Specific Life Tables to Insect Populations. *Insect Sci.*, **19**: 263-273.
 20. Kazmer, D. J. and Luck, R. E. 1991. The Genetic-Mating Structure of Natural and Agricultural Populations of *Trichogramma*. *Les Colloques de l'INRA*, **56**: 117-110.
 21. Khanamani, M., Basij, M. and Fathipour, Y. 2021. Effectiveness of Factitious Foods and Artificial Substrate in Mass Rearing and Conservation of *Neoseiulus californicus* (Acari: Phytoseiidae). *Inte. J. Acarol.*, **47(4)**: 273-280.
 22. Khanamani, M., Fathipour, Y., Talebi, A. A. and Mehrabadi, M. 2017. Quantitative Analysis of Long-Term Mass Rearing of *Neoseiulus californicus* (Acari: Phytoseiidae) on Almond Pollen. *J. Econ. Entomol.*, **110**: 1442-1450.
 23. Li, L. L. 1994. *Worldwide Use of Trichogramma for Biological Control in Different Crops*. CAB International, Wallingford, UK.
 24. Li, X. Y., Lei, Q., Hua, H. Q., Song, H. F., Wang, S., Ramirez-Romero, R., et al. 2019. Impact of Host Suitability on Oviposition Preference toward Fertilized and Unfertilized Host Eggs in Two *Trichogramma* Parasitoid Species. *Entomol. Gen.*, **39**: 313-323.
 25. Lü, X., Han, S. and Li, L. 2015. Biochemical Analyses of *Trichogramma dendrolimi* (Hymenoptera: Trichogrammatidae) *in Vitro* and *in Vivo* Rearing for 10 Generations. *Fla. Entomol.*, **98**: 911-915.
 26. Lü, X., Han, S., Li, Z. and Li, L. 2017. Biological Characters of *Trichogramma dendrolimi* (Hymenoptera: Trichogrammatidae) Reared *in Vitro* versus *in Vivo* for Thirty Generations. *Sci. Rep.*, **7**: 17928.

27. Mirkarimi, A. 2000. Biological Control of Carob Moth with Mass Release of *Trichogramma embryophagum* Hartig for Pomegranate Worm Control, the *Ectomyelois (Spectrobates) ceratoniae* Zell. *Iran. J. Agric. Sci.*, **31**: 103-110.
28. Moghaddassi, Y., Ashouri, A., Bandani, A. R., Leppla, N. C. and Shirk, P. D. 2019. Effect of *Ephesia kuehniella* (Lepidoptera: Pyralidae) Larval Diet on Egg Quality and Parasitism by *Trichogramma brassicae* (Hymenoptera: Trichogrammatidae). *J. Insect Sci.*, **19**: 1-7.
29. Nordlund, D. A., Wu, Z. X. and Greenberg, S. M. 1997. *In Vitro* Rearing of *Trichogramma minutum* Riley (Hym: Trichogrammatidae) for Ten Generations, with Quality Assessment Comparisons of *in Vitro* and *in Vivo* Reared Adults. *Biol. Control*, **9**: 201-207.
30. Paspati A., Ferguson, K. B., Verhulst, E. C., Urbaneja, A., González-Cabrera, J. and Pannebakker, B. A. 2019. Effect of Mass Rearing on the Genetic Diversity of the Predatory Mite *Amblyseius swirskii*. *Entomol. Exp. Appl.*, **167**: 670-681.
31. Parra, J. R. P. 2010. Mass Rearing of Egg Parasitoids for Biological Control Programs. In: "Egg Parasitoids in Agroecosystems with Emphasis on *Trichogramma*", (Eds.): Consoli, F. L., Parra, J. R. P. and Zucchi, R. A. Springer, Dordrecht, The Netherlands.
32. Pipoly, I., Bókony, V., Kirkpatrick, M., Donald, P. F., Székely, T. and Liker, A. 2015. The Genetic Sex-Determination System Predicts Adult Sex Ratios in Tetrapods. *Nature*, **527**: 91-94.
33. Poorjavad, N., Goldansaz, H., Hossenineveh, V., Nozari, J., Dehghanian, H. and Enkegaard, A. 2011. Fertility Life Table Parameters of Different Strains of *Trichogramma* spp. Collected from Eggs of the Carob Moth *Ectomyelois ceratoniae*. *Entomol. Sci.*, **14**: 245-253.
34. Pratisoli, D., Oliveira, H. N., Gonçalves, J. R., Zanuncio, J. C. and Holtz, A. M. 2004. Changes in Biological Characteristics of *Trichogramma pretiosum* (Hym.; Trichogrammatidae) Reared on Eggs of *Anagasta kuehniella* (Lep.; Pyralidae) for 23 Generations. *Biocontrol Sci. Tech.*, **14**(3): 313-319.
35. Razinger, J., Vasileiadis, V. P., Giraud, M., van Dijk, W., Modic, Š., Sattin, M. and Urek, G. 2016. On-Farm Evaluation of Inundative Biological Control of *Ostrinia nubilalis* (Lepidoptera: Crambidae) by *Trichogramma brassicae* (Hymenoptera: Trichogrammatidae) in Three European Maize-Producing Regions. *Pest Manag. Sci.*, **72**: 246-25.
36. Riahi, E., Fathipour, Y., Talebi, A. A. and Mehrabadi, M. 2017. Linking Life Table and Consumption Rate of *Amblyseius swirskii* (Acari: Phytoseiidae) in Presence and Absence of Different Pollens. *Ann. Entomol. Soc. Am.*, **110**: 244-253.
37. Smith, S. M. 1996. Biological Control with *Trichogramma*: Advances, Successes, and Potential of Their Use. *Annu. Rev. Entomol.*, **41**: 375-406.
38. Sorati, M., Newman, M. and Hoffmann, A. A. 1996. Inbreeding and Incompatibility in *Trichogramma nr. brassicae*: Evidence and Implications for Quality Control. *Entomol. Exp. Appl.*, **78**: 283-290.
39. Sørensen, J. G., Addison, M. F. and Terblanche, J. S. 2012. Mass-Rearing of Insects for Pest Management: Challenges, Synergies and Advances from Evolutionary Physiology. *Crop Prot.*, **38**: 87-94.
40. Suzuki, Y. and Hiehata, K. 1985. Mating Systems and Sex Ratio in the Egg Parasitoids, *Trichogramma dendrolini* and *T. papilionis* (Hymenoptera: Trichogrammatidae). *Anim. Behav.*, **33**: 1223-1227.
41. van Lenteren, J. C. and Bigler, F. 2010. Quality Control of Mass Reared Egg Parasitoids. In: "Egg Parasitoids in Agroecosystems with Emphasis on *Trichogramma*", (Eds.): Consoli, F. L., Parra, J. R. P. and Zucchi, R. A. Springer Publishing.
42. van Lenteren, J. C. 2012. The State of Commercial Augmentative Biological Control: Plenty of Natural Enemies, But a Frustrating Lack of Uptake. *BioControl.*, **57**: 1-20.
43. van Lenteren, J. C., Hale, A., Klapwijk, J. N., van Schelt, J. and Steinberg, S. 2003. Guidelines for Quality Control of Commercially Produced Natural Enemies. In: "Quality Control and Production of Biological Control Agents: Theory and Testing Procedures", (Ed.): van Lenteren, J. C. CABI Publishing, Wallingford, UK.



پرورش متوالی روی بید غلات، کیفیت زنبور پارازیتوئید *Trichogramma embryophagum* (Hymenoptera: Trichogrammatidae) را تحت تاثیر قرار

می دهد

ن. تقی خانی، ا. صحراگرد، ی. فتحی پور، و خ. مداحی

چکیده

پارامترهای دموگرافی زنبور پارازیتوئید *Trichogramma embryophagum* Hartig پرورش یافته روی بید غلات، *Sitotroga cerealella* (Olivier) به مدت ۴۰ نسل مورد بررسی قرار گرفت. آزمایش ها در اتاقک رشد با شرایط دمایی 26 ± 2 درجه سلسیوس، رطوبت نسبی 65 ± 5 درصد و دوره نوری ۱۶ ساعت روشنایی و ۸ ساعت تاریکی انجام شد. نتایج نشان داد که طول عمر زنبورهای ماده طی پرورش پایایی به طور معنی داری کاهش می یابد و از $10/68$ روز (در نسل پنجم) تا $9/64$ روز (در نسل چهارم) متغیر بود. از سوی دیگر، طول عمر زنبورهای نر در نسل بیستم به طور معنی داری کاهش یافت و هیچ تفاوت معنی داری بین طول عمر زنبورهای نر در نسل بیستم تا نسل چهارم مشاهده نشد. طول دوره تخم‌ریزی و میانگین کل باروری زنبور *T. embryophagum* با افزایش تعداد نسل ها کاهش یافت. عموماً، زنبورهای نسل های ابتدایی داری طول عمر، طول دوره زندگی و باروری بیشتری نسبت به نسل های بعدی بودند. علاوه بر این، نسبت جنسی *T. embryophagum* در نسل های متوالی اختلاف معنی داری نداشت. نتایج آزمون بوت استرپ جفت شده نشان داد که تمام پارامترهای رشد جمعیت زنبور *T. embryophagum* پرورش یافته روی بید غلات در نسل های متوالی به طور معنی داری متفاوت از یکدیگر بودند. بیشترین و کمترین مقادیر نرخ ذاتی افزایش جمعیت (r)، نرخ منتهای افزایش جمعیت (λ)، نرخ خالص تولیدمثل (R_0)، نرخ ناخالص تولیدمثل (GRR) و میانگین طول یک نسل (T) زنبور *T. embryophagum* به ترتیب در نسل چهارم و پنجم مشاهده شد. این نتایج نشان می دهد که زنبورهای *T. embryophagum* پرورش یافته در نسل های متوالی می توانند تا نسل بیستم با موفقیت در برنامه های کنترل بیولوژیک، بدون هیچ گونه افت کیفیت و عملکرد مورد استفاده قرار گیرند و پس از آن، باید جوانسازی منظم جمعیت آزمایشگاهی، با وارد کردن گاه به گاه پارازیتوئیدهای جمع آوری شده از مزرعه صورت گیرد.