Photosynthetic characters of canopy and storage root yield of sweet potato (*Ipomoea batatas* **L.) grow in different soil compaction**

Furong Du¹ , Chengcheng Si² , Yongchen Liu¹ , Xiubo Yin³ , Wenqing Shi³ , Chunyu Shi¹ , Zhe Sun⁴ , Wenjuan Lu⁵ , and Hongjuan Liu¹ *

Abstract

 Soil compaction is a critical constraint that restricts the attainment of high and stable yields of sweet potato. Photosynthesis is a key factor in yield formation. But the mechanism of soil compaction affects the photosynthesis and storage root (SR) yield of sweet potato remains elusive. A field experiments were carried out with two varieties in control, loose, and compacted soil conditions, canopy apparent photosynthesis (CAP), gas exchange parameters and PIabs of the functional leaves, SR yield were measured, and the relationship between yield and photosynthetic characters was studied as well. Compared with the control, the SR yield was significantly increased in loose soil with an average increase of 27.03%~38.74%, but decreased in compacted soil with an average reduction of 17.87%~15.92%. Both loosening and compaction treatments increased the leaf area index (LAI), and the increase in the latter is significantly higher than that in the former. Canopy interception rate in loosening treatment was much higher than that of compaction soil. The CAP showed a similar change in yield, with a strong positive correlation to SR yield and single storage root weight. Loose soil also improved gas exchange parameters, PIabs, the reverse was found in compacted soil. Compared to the control, the loose treatment significantly improved economic coefficient and reduced leaf starch content, while the compaction treatment showed the opposite trend. Path analysis revealed that the net photosynthetic rate (*P*n) had the most total effect and higher direct effect on increasing CAP. Therefore, soil compaction primarily regulates SR yield through CAP, with Pn exerting a significant impact on CAP. Enhanced soil compaction led to reduced photosynthate output in functional leaves, resulting in decreased Pn and increased

 \overline{a}

¹ State Key Laboratory of Crop Biology, Shandong Key Laboratory of Crop Biology, Agricultural College, Shandong Agricultural University, Tai'an, 271018, Shandong Province, China.

² College of Horticulture, Hainan University, Haikou, 5702258, Hainan Province, China.

 Multigrain Department, Shandong Agricultural Technology Extension Station, Jinan, 250100, Shandong Province, China.

Tai'an Academy of Agricultural Sciences, Tai'an 271000, Shandong Province, China.

Weihai Academy of Agricultural Sciences, Weihai 264200, Shandong Province, China.

^{*} Corresponding author; e-mail: [liumei0535@126.com](https://jast.modares.ac.ir/admin_emailer.php?mod=send_form&sid=23&slc_lang=en&em=liumei0535-ATSIGN-126.com&a_ordnum=71905)

LAI. Consequently, an inappropriate canopy structure with low canopy interception is

formed.

 Keywords: canopy apparent photosynthesis, PIabs, gas exchange parameters, storage root yield, soil compaction, sweet potato.

1. Introduction

Sweet potato is a globally significant tuberous crop, with a total production of 8.64×10^7 34 tons on 7.25×10^6 hectares, and an average yield of 11.9 tons/hectare (FAOSTAT, 2023). It demonstrates remarkable adaptability and thrives in diverse terrains, including hilly, mountainous, and plains (Sun et al. 2022). There has been a substantial increase in soil compaction, which is widely recognized as one of the primary challenges to soil fertility, crop productivity, and food safety (Keller et al. 2019). Soil compaction led to a substantial decrease in SRs yield, with reductions ranging from 30% to 90% (Shi et al. 2019). This phenomenon is frequently observed not only in plains (Bogunovic et al. 2018), but also in hilly and mountainous regions (Xoconostlecázares et al. 2010). Hence, it is crucial to make clear the impact of soil compaction on sweet potato yield in order to maximize production potential.

 The bulking rate and single weight of sweet potato SRs decreased with increasing soil compaction, while the root tip number and hypocotyl diameter of soybean showed a significant increase (Li et al., 2024). The soybean root system may change due to limited water and nutrient availability in the soil, caused by restricted gas diffusion between roots and the rhizosphere (Horák et al. 2022). The development of sweet potato SRs is mainly influenced by photosynthesis, as they store photosynthate. However, there are conflicting findings on the effect of soil compaction on photosynthesis. Some studies indicated that soil compaction reduces photosynthesis rates (Mariotti et al. 2020; Huntenburg et al. 2021), which 52 was ascribed to a decrease in stomatal conductance, thus impeding $CO₂$ diffusion to the mesophyll (Philip and Azlin 2005). Net photosynthesis (*P*n), stomatal conductance (gs), and transpiration rates (E) of soybean were reduced up to 50% under compaction (Ferreira et al., 2023). While the study in potato showed that photosynthesis rates did not differ between compaction treatments after ground cover (Huntenburg et al., 2021). The response of photosynthesis to soil compaction may vary across studies, however, a consistent observation

 was the reduction in leaf size and plant carbon assimilation. But sweet potato leaves exhibited an increase in size rather than decrease under soil compaction. This indicated that the response mechanism of sweet potato to soil compaction differed from other crops. The CAP provides a more precise indication of the photosynthesis of field crops (González and Manavella, 2021), while chlorophyll fluorescence parameters have been widely employed to identify disruptions in the photosynthesis apparatus caused by abiotic stress (Grzesiak 2009). So our study aims to investigate the CAP variation at different development stages, gas exchange parameters, chlorophyll fluorescence parameters of functional leaves, and quantify how the CAP, and functional leaf photosynthesis of sweet potato respond to soil compaction and how they affect SR yield.

2 Materials and Methods

2.1 Materials and test design

 The field experiment was conducted at the Agricultural Test Station of Shandong Agricultural University in Taishan District, Tai'an City, Shandong Province (360°09′N, 117°09′E;128 m asl) during 2017 and 2018. The sweet potato cultivars used were Shangshu 19 (SS19) and Jixu 23 (JX23). The tested soil was sandy loam. Three compaction levels were 75 used: (1) compaction (C), where the 0–20-cm soil layer of the treatment was compacted by a 76 vibrating tamper (HS-75R, HANSA, Germany), with a bulk density of $1.40-1.50$ g cm⁻³ and 77 a compaction of > 0.6 MPa and < 1.2 MPa; (2) control (no compaction, CK), where the bulk 78 density of the $0-20$ -cm soil layer was $1.30-1.40$ g cm⁻³ and the compaction was approximately 0.3–0.4 MPa; and (3) loosening (L), the bulk density in the 0–20-cm soil layer 80 was 1.20–1.30 g cm⁻³, and the compaction was approximately 0.1 –0.2 MPa. The soil in this treatment was mixed with organic fertilizer, sand, and common loamy soil. After mixing, the organic matter content of the soil in the loosening was consistent with the other treatments. The nitrogen, phosphorus, and potassium contents in the three treatments were adjusted to similar levels using potassium sulfate and urea. In 2017, the available nitrogen, phosphorus, 85 potassium, and organic matter in the 0–20-cm soil layers were 79.47 mg kg⁻¹, 42.47 mg kg⁻¹, 86 112.33 mg kg⁻¹, and 1.30 %, respectively, and in 2018, it was 88.73 mg kg⁻¹, 35.22 mg kg⁻¹, 90.51 mg kg⁻¹, and 1.13 %, respectively. The physical properties of the soil under the three treatments are shown in Table 1. The field experiment employed a two-factor split-plot experimental design with five replications, using cultivars as the primary plots and 90 compaction as the subplots. Each plot covered an area of 20 m^2 , with row spacing-80 cm and plant spacing-25 cm. Sweet potato was planted on May 10 and May 9, harvested on October 22 and October 20 in 2017 and 2018, respectively.

93 **Table 1.** Physical characteristics of the soil (1 day before planting).

Year	Soil Layer (cm)	Treatme nts	Soil Compactness (kpa)	Soil Soil bulk density (g/cm^3)	Soil Specific Gravity (g/cm^3)	Total Porosity (%)	Capillary porosity (%)	Non-capillary Porosity (%)
2017	$5 - 10$	L	126.49c	1.26c	2.58a	51.35 a	24.86 b	26.49a
		CK	301.16 b	1.33 _b	2.64a	49.81a	24.15 b	25.66 a
		C	541.63 a	1.46a	2.73a	46.38 b	31.78 a	14.60 _b
	$10-15$	L	224.23c	1.30c	2.57 _b	49.35a	25.50c	23.85 a
		CK	464.12 b	1.39 _b	2.73a	49.19a	31.68 b	17.51 b
		C	927.74 a	1.49a	2.75a	45.71 b	38.30 a	7.41 c
2018	$5-10$	L	143.17 c	1.25c	2.57a	50.69a	24.62 b	26.07a
		CK	267.91 b	1.33 _b	2.65a	48.92 a	25.30 _b	23.62 b
		C	826.07 a	1.47a	2.74c	46.66 b	31.21 a	15.45c
	$10-15$	L	174.17 c	1.29c	2.58 _b	49.62a	25.46c	24.16a
		CK	508.06 b	1.38 _b	2.73a	48.65 a	30.14 _b	18.51 _b
		C	1230.6 a	1.49a	2.75a	45.83 b	36.91 a	8.92c

94 Note: Values followed by different lowercase letters within a column in the same year of the same varieties are 95 significantly different among different treatments (P < 0.05). The same as below.

96

97 Soil compaction was measured using a soil compaction metre (CP40 II , Cinstral Exports 98 Pty Ltd T/A Rimik, Australia) at the seedling stage, early, middle and late stage of SR bulking. 99 The soil volumetric moisture content in the 0–20 cm and 20–40 cm layers was measured 100 every five days after transplanting to canopy cover. The moisture content per cubic metre $(m³)$ $101 = H \times 1$ m² × soil volumetric water content (%), where H is the soil depth. The plot with the 102 highest water content served as the standard for adjusting the water content in the remaining 103 plots. From transplanting to canopy cover, irrigated all treatments 1~2 times to ensure the soil 104 relative humidity of the control treatment was above 60%; the irrigation amount was 105 consistent across all treatments. The other management is similiar to that of general field 106 crops, with the climate for two growing seasons detailed in Fig. 1.

Fig. 1 Climate data for the two growing seasons of sweet potato.

2.2 Sampling and measurement method

2.2.1 Canopy apparent photosynthesis

 The CAP was measured in a modified closed gas exchange system using an infrared gas analyser (GXH-305, China) (Hay and Porter, 2006), which was portable and easy to move in 114 the field. The aluminum-framed chamber measured 0.80×0.90 m in area and 0.70 m in height, with the outer cover sealed using a high light transmittance mylene film (about 95% light transmittance). A 25 cm fan was placed at the top to mix the gas and balance the 117 temperature. The $CO₂$ concentrations decreased linearly, usually measured within 60 seconds of the chamber closing. Two rows of sweet potatoes were chosen in the chamber, and measurements were taken every 20 days from 60 to 140 days after planting (DAP), between 120 9:00 and 11:00 A.M. Three sampling site with consistent growth were selected to mearsure for each treatment, and the respiration rates of soil were measured in the open spaces of the adjacent area. The CAP was calculated as follows:

$$
CAP = slope \times n / A \tag{1}
$$

Where the slope presents the reduce in the $CO₂$ concentration per unit time (umol mol⁻¹) 125 s^{-1}), n is moles number of air in the chamber, and A standing for the ground area. The value 126 of n equal to PV/RT, where P is the pressure in kPa, V is the chamber volume in L, T is the 127 temperature in K, and R is the gas constant $(8.314 \text{ k Pa L mol}^{-1} \text{ K}^{-1})$.

2.2.2 Gas exchange measurements

 Gas exchange measurements were conducted at 20-day intervals from 60 to 140 days post-planting. In sunny, windless conditions, the photosynthetic rate of the fifth fully opened leaf (with the highest photosynthetic rate) at the top of the stem was measured between 9:30 and 11:30 A.M. Fifteen leaves with similar growth were selected for each treatment across three replications. The Portable Photosynthesis System (CIRAS-3, PP Systems International, Inc. Amesbury, USA) was used. The measuring head was subject to the following conditions: 137 leaf temperature, 28 °C; reference $CO₂$ content, 380 ppm, and ambient air humidity. The 138 PPFD during the measurements was set to 1000 μ mol (photon) m⁻² s⁻¹ throughout the experiments. The *P*n, E, gs, and Ci in the intercellular spaces were automatically recorded.

2.2.3 Chlorophyll fluorescence parameters of the functional leaves

 The parameters were measured every 20 days from 60 to 140 DAP. The MPEA chlorophyll fluorescence meter (Hansatech Instruments Ltd., King's Lynn, UK) was connected to a computer for precise value determination. Prior to fluorescence signal measurements, the plants were dark-adapted for at least 30 minutes. The fifth unfolded leaf at the top of the stem was measured. 15 leaves with similar growth characteristics were selected for each condition. The following parameters were considered:

148 PIabs = RC/ABS • [φ Po /(1-φ Po)] • [ψ _O /(1- ψ _O)], the performance index of the absorbance basis.

2.2.4 Canopy interception rate

 From 60 to 140 DAP, the canopy interception rate of sweet potato was assessed using a SunScan plant canopy analyzer (Delta-T Devices, UK). Measurements were carried out on clear and calm days between 12:00 and 14:00. Incident and reflectance radiation were measured above the canopy (15 cm above) and below the canopy (5 cm above the ground), with each treatment replicated five times.

157 Canopy transmittance
$$
(\%) = (PAR - PAR_0)/PAR \times 100 \%
$$
 (2)

158 Canopy reflectance
$$
(\%) = PAR'/PAR \times 100\%
$$
 (3)

Canopy interception (%)= 100-Canopy transmittance (%)-Canopy reflectance(%) (4)

160 Where, PAR is the PAR above the canopy, PAR₀ is the PAR at the base of the ridges, and 161 PAR' is the reflected light above the canopy.

162

163 **2.3 Statistical analysis**

 Statistical analysis was conducted with IBM SPSS Statistics 25 (IBM Inc., Chicago, IL, USA). The statistical significance were assessed using ANOVA, followed by Duncan's multiple range test. Figures were generated using SigmaPlot software (SigmaPlot 12.0, Systat Software, San Jose, CA, USA).

168

169 **3 Results**

170 **3.1 Effects of soil compaction on SR yield of sweet potato**

171 Compared with the control, the SR yield was significantly increased in loosening treatment, and the average increase of two years was 27.03%~38.74%, while they were decreased in compaction treatment, and the average reduction was 17.87%~15.92%. The single storage root weight changed similiar to the yield. The change of single SR weight and economic coefficient was similiar to that of yield (Table 2). The result of statistical analysis indicated that treatment, the interaction between year and treatment had significant effect on yield. 177 There were significant differences in the number of SRs, single SR weight and economic coefficients between treatments as well. That is, the soil compaction led to changes in SR yield, primarily through the regulation of individual SR weight.

- 180
-

181 **Table 2.** Effect of soil compaction on SR yield, its economic coefficient of sweet potato.

182 Note: Values followed by different lowercase letters within a column are significantly different among different 183 treatments (*P* < 0.05). The same as below.

184

185 **3.2 Effects of soil compaction on the photosynthetic area and canopy interception rate**

186 *3.2.1 LAI*

187 During SRs bulking, the LAI first increased and then decreased, the peak values were 188 observed at 120 and 100 DAP for SS19 and JX23, respectively. Compared to the control, the LAI was significantly increased in loosening and compaction treatments,with the average increase of 37.54% and 63.81. The highest LAI was obtained in the compaction treatment at 120 and 100 DAP for JX23 and SS19. Subsequently, the LAI in the compaction treatment exhibited a more rapid decline compared to the other treatments after reaching its peak (Fig. 2). In summary, the group structure of loosening treatment appeared to be suitable; however, the group is excessively large within the compact soil.

195

196 **Fig. 2** Effect of soil compaction on LAI (2018).

197 **3.2.2 Canopy interception rate**

198 Compared to the control, the loosening treatment had the highest canopy interception rate, 199 while the compaction treatment showed the lowest despite having a higher LAI. This could 200 be attributed to frequent leaf turnover in the compaction treatment, resulting in numerous

Fig. 3 Effect of soil compaction on canopy interception rate (2018).

3.2 Effects of soil compaction on the CAP rate

 The CAP of sweet potatoes follows a bimodal curve, with peak values at 80 and 120 DAP and a trough at 100 DAP. The occurrence of the trough coincides with the rainy season, during which there was an increase in soil compaction due to sustained precipitation.

 Compared with the control, the loosening treatment increased the CAP rate significantly with an average increase of 29.46% and 38.76%, for the SS19 and JX23, respectively, the greater significant increase appeared at 80 DAP and 140 DAP. And the compaction treatment decreased the CAP rate significantly with the average reduction of 17.31% and 20.21%, the most significant reduction appeared at 60 DAP and 80~100 DAP (Fig. 4). That is, soil compaction affected the CAP greatly in the early and middle growth stages, when the soil compaction was reduced the CAP was increased greatly during the late growth stage.

 [\[Downloaded from jast.modares.ac.ir on 2](https://jast.modares.ac.ir/article-23-71905-en.html)024-11-25]

Downloaded from jast.modares.ac.ir on 2024-11-25

3.4 Effects of soil compaction on the Gas exchange parameters of the functional leaves At 80, 100 and 120 DAP, the loosening treatment significantly improved the *P*n, *Ci*, *gs* and *E* of the functional leaves, whereas the compaction treatment had opposite effects, which was consistent with the 2-year data. In 2017, the gas exchange parameters exhibited significant variation at the trough of CAP rate (100 DAP), whereas in 2018, a noticeable alteration was observed in the second peak of CAP rate (120 DAP) (Fig. 5). The occurrence of periods characterized by significant inter-annual variability is primarily linked to the temporal distribution of intense precipitation.

Fig. 5 The effect of soil compaction on gas exchange parameters in leaves of sweet potato.

3.5 Effect of soil compaction on PIabs

 PIabs is the performance index of the PSⅡ reaction centre, which reflects the overall performance of PSⅡ. The loosening treatment significantly increased the PIabs of the functional leaves by 55.63% and 38.50% in SS19 and JX23, respectively, compared to the control treatment. In contrast, the compaction treatment led to a significant decrease in PIabs by 28.33% and 22.29%, respectively (Fig. 6). Hence, loose treatment is benefit to 233 improvement of the overall performance of PS II.

Fig. 6 The effect of compaction on PIabs of leaves of sweet potato (2018).

3.6 Determining the main influencing factors of the canopy apparent photosynthesis

237 The regression equation Y = $38.38 + 4.67X1 + 1.61X2 + 0.15X3 - 0.34X4 - 0.029 X5 +$ 238 2.03X6 - 18.00X7 - 115.61X8 + 2.24X10 (F = 29.58, R² = 0.938, P=0.0001) was obtained by stepwise regression using the CAP rate as the dependent variable and, LAI (X1), canopy interception (X2), *P*n (X3), *Ci* (X4), *gs* (X5), *E* (X6), Vj (X7), Wk (X8), ψo (X9), and PIabs (X10) as the independent variables. The correlation analysis revealed a significant positive correlation between *P*n, canopy interception, Ci, gs, E, Ψo and PIabs with the CAP. Path analysis revealed that gas exchange parameters and PIabs were the primary contributors to increased CAP. Gas exchange parameters, particularly *P*n and Ci, had the most significant direct effects, and the overall impact of *P*n was mainly driven by its direct effects, while most other factors were primarily influenced by the indirect effects of *P*n (Fig. 7).

247

253

248 **Fig. 7** The path analysis among functional leaf photosynthesis, canopy apparent 249 photosynthesis and storage root yield. **Note:** Dashed and solid lines indicate indirect and 250 direct impacts, respectively. The number above or below the arrow line indicates the direct 251 effects / total effects. R^2 represents the correlation coefficient. The number in the box 252 indicates the correlation coefficient between the item in box with the CAP.

254 **4 Discussion**

255 **4.1 Effects on SR yield of sweet potato**

 This study found a significant increase in SR yield when soil compaction was reduced (Table 2), consistent with previous research (Shi et al. 2019). Improving photosynthesis can increase crop yields (Dong et al. 1993), as more than 90% of crop outputs are directly related to photosynthesis (González and Manavella, 2021; Chen et al. 2022). Within the realm of field crops, the CAP exerts a pronounced impact on yield formation. However, limited research has been conducted on the effect of soil compaction on sweet potato photosynthesis and its correlation with SR yield. This study revealed that variations in single SR weight were the primary contributing factor to differences in SR yields (Table 2). A significant positive correlation was obtained between the CAP and SR yield (*r= 0.99, P<0.05*), as well as the single SR weight (*r= 0.90, P<0.05*). Our previous research has revealed that reducing soil compaction resulted in higher single SR weight by increasing dry matter accumulation in the SRs (Shi et al. 2019). Consequently, soil compaction mainly affects the accumulation of

 photosynthates in SRs by regulating the CAP, thus impacting single SR weight and leading to variations in SR yield.

4.2 Effects on the photosynthetic characteristics of sweet potato

 CAP is closely linked to the canopy architecture and positively correlated with light interception (Bhusal, et al., 2017). The LAI is an important index of canopy architecture. But the correlation between CAP rate and LAI varies among different crop species. The CAP was closely associated with changes in green leaf area in maize (Liu et al. 2015), whereas, the LAI was similar among varieties with distinct CAP rate in wheat (Tang et al. 2017). Moreover, an appropriate LAI can enhance the group's light distribution and interception ability (Maddonni et al 2001). The increase of soil compaction reduced ground cover expansion, decreased plant leaf area, shortened canopy cover duration, and restricted light interception (Assaeed et al. 2008). This study revealed that the loosening treatment appeared the highest CAP rate (Fig. 3), the greater LAI (Fig. 2) and the most canopy interception rate (Fig. 4), compared with the control treatment. The compaction treatment got the highest LAI but the lower canopy interception rate (Fig. 2, 4). This may be because compaction treatment led to excessive growth of stems and leaves, resulting in frequent alternation of new and old leaves. The CAP rate in the compaction treatment was reduced as well (Fig. 3). The previous study has improved that a suitable canopy structure, high chlorophyll content, and prolonged leaf duration can enhance CAP and biomass yield (Tang et al., 2017). Therefore, reducing soil compaction promotes appropriate canopy architecture, improves light penetration, and enhances the CAP.

 Reducing soil compaction has been shown to enhance the net photosynthetic rate of leaves [\[Downloaded from jast.modares.ac.ir on 2](https://jast.modares.ac.ir/article-23-71905-en.html)024-11-25] Downloaded from jast.modares.ac.ir on 2024-11-25] in various crops such as cucumber, strawberry, peanut, ginger, soybean, and potato (Du et al. 292 2010), while its increase has demonstrated an opposite effect. These findings were consistent with previous research on the physiological and agronomic response of soybean cultivars to soil compaction in the Brazilian Cerrado (Maddonni et al. 2001; Ferreira et al., 2023). While the other study in potato showed that *P*n did not differ between compaction treatments after ground cover (Huntenburg et al., 2021). In this study, we found *P*n, Ci, E and gs wers increased in the loosening treatment but decreased in the compation treatment (Fig. 4). The

 significant fluctuations were most pronounced during and after the rainy season (at 80 and 120 DAP). The variation in *P*n for loosening and compaction treatments was less than that of the CAP rate. Furthermore, the PIabs were enhanced in the loosening treatment (Fig. 6), other chlorophyll fluorescence parameters showed relatively small changes (Supplementary Table 1,). And Photosynthesis critically depends on the electron flow through PSII (Hussain et al. 2019), which functions as a fundamental photosynthetic unit within the thylakoid membrane of a chloroplast. The accumulation of starch within chloroplasts may result in the perturbation of thylakoid membranes, leading to a reduction in the photosynthetic efficiency of leaves. When soil compaction was intensified, leaves accumulate a lot of starch, while the loosening treatment leaves have high sucrose and low starch content (Supplementary Table 2). Meanwhile, the economic coefficient of compaction treatment was reduced (Table 2). So the excessive accumulation of starch in functional leaves was the primary factor contributing to the reduction in *P*n. The emphasis of the next research step should be on the output of leaf photosynthetic products.

 Linear regression, correlation analysis, and path analysis were employed to identify the key factors influencing the CAP rate due to soil compaction. The results indicated a highly significant positive correlation between gas exchange measurements (*P*n, Ci, E and gs) and the CAP rate. Gas exchange measurements (*P*n, Ci, and E) had the most significant overall impact on enhancing the CAP. The total effects of *P*n were derived primarily from direct effects, which were the most substantial among the items, while the most items were derived mainly from indirect effects of *P*n (Fig. 7). Overal, the primary determinant of SR yield under soil compaction was the CAP rate. When modulating the CAP rate, it was crucial to consider gas exchange parameters, particularly in controlling the *P*n.

5. Conclusions

 Soil compaction primarily influenced sweet potato SRs yield by modulating their CAP rate. Pn and canopy architecture were the primary determinant of CAP rate. As soil compaction increased, the reducing photosynthate output of leaves leads to starch accumulation, resulting in a marked reduction in photosynthetic rate and a substantial increase in LAI.

Acknowledgments

 Natural Science Foundation of Shandong Province, China (Grant No. ZR2022MC186), Key Research and Development Program of Shandong (2023TZXD024), the Sweetpotato Industry Technology Research System of Shandong Province (SDAIT-16-01), National Natural Science Foundation of China (31701357), Tai'an Science and Technology Commissioner Project.

References

- Anikwe MAN, Ubochi JN (2007) Short-term changes in soil properties under tillage systems and their effect on sweet potato (*Ipomoea batatas* L.) growth and yield in an Ultisol in south-eastern Nigeria. AUST J SOIL RES 45: 351–358
- Assaeed AM, Mcgowan M, Hebblethwaite P D, Brereton J C (2008) Effect of soil compaction on growth, yield and light interception of selected crops. AAB 11: 653–666.
- Bhusal N, Han S, Yoon T (2017). Summer pruning and reflective film enhance fruit quality in excessively tall spindle apple trees. ANN APPL BIOL, 58, 560–567.
- Bogunovic I, Pereira P, Kisic I, Sajko K, Sraka M (2018). Tillage management impacts on soil compaction, erosion and crop yield in stagnosols (Croatia). CATENA 160: 376–384.
- Borgmann C, Secco D, Marins ACD, Junior LAZ, Bassegio D, Souza SNMD, Zhang FN, Silva T RBD (2021) Effect of Soil Compaction and Application of Lime and Gypsum on Soil Properties and Yield of Soybean. COMMUN SOIL SCI PLAN 52: 1434–1447.
- Chen CQ, Tian XY, Li J, Bai S, Zhang ZY, Li Y, Cao HR, Chen ZC (2022) Two central circadian oscillators OsPRR59 and OsPRR95 modulate magnesium homeostasis and carbon fixation in rice. MOL PLANT. 15(10): 1602-1614.
- Dong ST, Hu CH, Gao RQ (1993) Rates of apparent photosynthesis, respiration and dry matter accumulation in maize canopies. BIOL PLANTARUM. 35: 273–277.
- Du GD, Guo XW, Lu DG, Cai M (2010) Effect of soil compaction on photosynthetic characteristics and PSII Photochemistry activities in strawberry leaves. J. FOOD SCI. 27: 542–546.
- Ekwue EI, Stone RJ (1995) Irrigation Scheduling for Sweet Maize Relative to Soil Compaction Conditions. J. AGRIC. ENG. RES. 62: 85–94.

- Ferreira CJB, da Silva AG, Tormena CA, Severiano ED, Tavares RLM, Braz GBP, de Paiva SVD (2023). Physiological and agronomic response of soybean cultivars to soil compaction in the Brazilian Cerrado. SOIL PLANT NUTR 82: e20220160.
- González FG, Manavella PA (2021). Prospects for plant productivity: from the canopy to the nucleus. J. Exp. Bot. 72, 3931–3935.
- Grzesiak MT (2009) Impact of soil compaction on root architecture, leaf water status, gas exchange and growth of maize and triticale seedlings. PLANT ROOT 3: 10–16.
- Hamza MA, Anderson WK (2005) Soil compaction in cropping systems: a review of the nature, causes and possible solutions. SOIL TILL RES 82(2): 121–145.
- Hay RKM, Porter JR (2006) The Physiology of Crop Yield*.* BPL, Oxford.
- Horák J, Šimanský V, Kotuš T, Hnátková T, Trakal L, Lukac M (2022) Mitigation of
- Greenhouse Gas Emissions with Biochar Application in Compacted and Uncompacted Soil. AGRONOMY 12: 546.
- Huntenburg K, Dodd IC, Stalham M (2021) Agronomic and physiological responses of potato subjected to soil compaction and/or drying. ANN APPL BIOL 178: 328–340.
- Hussain S, Iqbal N, Brestic M, Raza MA, Pang T, Langham DR, Safdar ME, Ahmed S, Wen BX, Gao Y, Liu WG (2019) Changes in morphology, chlorophyll fluorescence performance and Rubisco activity of soybean in response to foliar application of ionic titanium under normal light and shade environment. SCI. TOTL. ENVIRON 658: 626–637.
- Keller T, Sandin M, Colombi T, Horn R, Or D (2019) Historical increase in agricultural machinery weights enhanced soil stress levels and adversely affected soil functioning. SOIL TILL RES 194: 104293.
- Liu TN, Gu LM, Don ST, Zhang J, Liu P, Zhao B (2015) Optimum leaf removal increases canopy apparent photosynthesis, 13 C-photosynthate distribution and grain yield of maize crops grown at high density. FIELD CROP RES 170: 32–39.
- Maddonni GA, Otegui ME, Cirilo AG (2001) Plant population density, row spacing and hybrid effects on maize canopy architecture and light attenuation. FIELD CROP RES 71: 183–193.
- Mariotti B, Hoshika Y, Cambi M, Marra E, Marchi E (2020) Vehicle-induced compaction of
- forest soil affects plant morphological and physiological attributes: A meta-analysis. FOREST ECOL MANAG 462: 118004.
- Nedunchezhiyan M, Raya RC (2010) Sweet potato growth, development, production and utilization: Overview. In Sweet Potato: Postharvest Aspects in Food, ed. R.C Raya and K.I Tomkin. NSP Inc pp 1-26.
- Philip E, Azlin YN (2005) Measurement of soil compaction tolerance of *Lagastromia speciosa* (L.) Pers. using chlorophyll fluorescence. URBAN FOR URBAN GREE 3: 203–208.
- Shi WQ, Zhang BB, Liu HJ, Zhao QX, Shi CY, Wang J Si CC (2019) Response mechanism of sweet potato storage root formation and bulking to soil compaction and its relationship with yield. ACTA AGRON SCI 45: 755–763.
- Sun L, Yang Y, Pan H, Zhu J, Zhu M, Xu T, Li Z, Dong T (2022) Molecular Characterization and Target Prediction of Candidate miRNAs Related to Abiotic Stress Responses and/or Storage Root Development in Sweet Potato. GENES (Basel) 13(1): 110.
- Tang YL, Wu XL, Li CS, Yan WY, Huang MB, Ma XL, Li SZ (2017) Yield, growth, canopy traits and photosynthesis in high-yielding, synthetic hexaploid-derived wheats cultivars compared with non-synthetic wheats. CROP PASTURE SCI 68(2): 115–125.
- Wang L, Wang TS, Yao SH, Sun HJ, Zhang B (2024). Soil compaction development facilitated the decadal improvement of the root system architecture and rhizosheath soil traits of soybean in the North China Plain. SOIL TILL RES 237: 105983.
- Xoconostlecázares B, Ramírezortega FA, Floreselenes L, Ruizmedrano R (2010) Drought tolerance in crop plants. J PLANT PHYSIOL 5: 241–256.
- Yang JS, Gao HY, Liu P, Li G, Dong ST, Zhang JW, Wang JF (2010) Effects of planting density and row spacing on canopy apparent photosynthesis of high-yield summer corn. ACTA AGRON SCI 36: 1226–1233.

 [\[Downloaded from jast.modares.ac.ir on 2](https://jast.modares.ac.ir/article-23-71905-en.html)024-11-25] Downloaded from jast.modares.ac.ir on 2024-11-25]