

Changes in some seedling growth parameters, nutrient content and enzyme activity in different melon (*Cucumis melo* L.) genotypes under deficit irrigation conditions

Özlem Cakmakci¹, Selma Kipcak Bitik², Aytekin Ekincialp², Çeknas Erdinc³, Turgay Kabay⁴, Fuat Eser¹, and Suat Sensoy¹

1. Department of Horticulture, Faculty of Agriculture, Van Yuzuncu Yil University, Van, Turkey.
2. Baskale Vocational School, Van Yuzuncu Yil University, Van, Turkey.
3. Department of Agricultural Biotechnology, Agriculture Faculty, Van Yuzuncu Yil University, Van, Turkey.
4. Department of Plant and Animal Production, Erçiş Vocational School, Van Yuzuncu Yil University, Van, Turkey.

*** Corresponding author; e-mail: ozlemguldigen@yyu.edu.tr**

ABSTRACT

Increasing need for drought adaptation measures to conserve water and sustain crop yield in water-scarce regions, driven by severe and recurrent droughts. Achieving sustainable production entails studying deficit irrigation as a means to enhance water productivity and selecting genotypes resilient to soil water deficits. In the present study, 17 different melon (*Cucumis melo* L.) genotypes collected from the Van Lake Basin and 3 hybrids and 1 standard melon cultivar for control purposes were used. The study was carried out under climate room conditions. Two different irrigation levels (I₁₀₀: 100% full irrigation, I₅₀: 50% deficit irrigation-DI) were applied in the study for deficit irrigation. The applications started with the emergence of the second true leaves of the plants and after one-month, different growth, nutrient, and enzyme contents of the seedlings were determined. In general, it was determined that deficit water application negatively affects seedling growth, and root dry matter, stomatal width and density, potassium, APX and SOD enzymes, and MDA content increased, while the other examined parameters decreased. The genotypes of the Van Lake Basin melon were found to vary as a result of the findings.

Keywords: Antioxidative response, Deficit irrigation, Melon, Mineral composition, Seedling growth parameters.

INTRODUCTION

Melon (*Cucumis melo* L.), renowned for its rich nutritional profile, originates from East Africa (Pitrat 2008). Global melon production, totaling around 32 million tons, sees China (PRC) as the leading contributor, accounting for 40%, while Türkiye follows closely with approximately 5.5%

37 (FAO 2019). Türkiye, a recognized gene center for various crops, including melon, stands as a
38 secondary gene center for this species (Sensoy et al. 2007a; Erdinc et al. 2013; Kısaca and Gazioglu
39 Sensoy 2022). The Van province in Eastern Anatolia, Türkiye, holds significance as one of the
40 origins of cantaloupe melon (Sensoy et al. 2007a; Turkmen et al. 2008). Genetic studies by Sensoy
41 and Sahin (2012) revealed a notably high genetic diversity among Sihke melon genotypes in the
42 Lake Van Basin.

43 Drought, a prominent abiotic stressor, significantly jeopardizes global agricultural yield and
44 quality. With the escalation of global warming-induced climate change, arid and semi-arid regions
45 face exacerbated drought challenges (Tan et al. 2006; Pandey et al. 2018). Under drought
46 conditions, plant growth and development are impeded due to slowed cell division, interrupted
47 transpiration, and inhibited nutrient uptake, leading to diminished productivity (Sensoy et al.
48 2007b; Farooq et al. 2009; Cakmakci et al. 2017). In nations heavily reliant on agriculture,
49 optimizing water resource utilization is imperative to alleviate the adverse impacts of climate
50 change.

51 To address future challenges arising from climate change and a growing global population, it is
52 crucial to develop drought-tolerant plant genotypes requiring less water. In Türkiye, insufficient
53 precipitation has led to significant agricultural losses, emphasizing the urgent need to identify and
54 select drought-tolerant genotypes through expanded breeding programs (Kabay and Sensoy 2016).
55 Melon, a globally cultivated fruit with high nutritional and economic value, faces water scarcity
56 issues, particularly in arid regions. Deficit irrigation (DI), a water-saving strategy, seeks to enhance
57 water use efficiency while sustaining plant growth. Numerous studies have explored the impact of
58 DI on melon growth parameters across different genotypes (Sensoy et al. 2007b; Kusvuran et al.
59 2011; Sharma et al. 2014; Kırmak and Dogan 2017; Wang et al. 2017; Barzegar et al. 2018; Lamaoui
60 et al. 2018). Understanding the genetic variability and inheritance of physiological traits under DI
61 is vital. This study focuses on evaluating melon genotypes from the Van Lake Basin for their
62 response to water deficit during the seedling stage, utilizing morphological and physiological
63 parameters to identify tolerant genotypes for future breeding programs.

64 MATERIALS AND METHODS

66 Sihke melon genotypes sourced from the Van Lake Basin, alongside three hybrids and a standard
67 cultivar for control (Table 1), constituted the plant materials for this study. Under climate room
68 conditions (16 hours light, 8 hours dark, % 50-55 humidity and 23-25°C), seeds of the genotypes

69 were sown in 2-liter pots containing a sterile 2:1 peat to perlite ratio. The experiment featured two
 70 irrigation levels: I_{100} (100% full irrigation) and I_{50} (50% deficit irrigation). The study consists of a
 71 two-factor factorial design (melon genotype and irrigation) regime. Employing a randomized
 72 experimental design with three replications, each (total 126 pots) replication housed four plants.
 73 The initiation of applications coincided with the emergence of the second true leaves of the
 74 seedlings.

75 A and B solutions, comprising nutrients (A solutions: 10.03% N, 1.6% NH_4-N , 8.7% NO_3-N , 7.5%
 76 K_2O , 8.6% Ca, 0.3 % Fe and B solution; 2.1% N, 2.1% NO_3-N , 6.4% P_2O_5 , 11.6% K_2O , 1.6% Mg,
 77 0.01% Zn, 0.003% Cu, 0.1% Mn, 0.003% B, 0.004% Mo), were administered (50 ml) to all pots.
 78 Pre-planned irrigations followed, with water applied to reach field capacity before each irrigation
 79 cycle, determined by the pot capacity. The irrigation water volume for each session was computed
 80 using the provided equation.

$$81 \quad I = ((W_i - 1) - W_i) * IR$$

82 Where, I is the amount of irrigation water (ml), W_{i-1} and W_i mass (kg) of the pot at day $i-1$ and i ,
 83 respectively (kg). IR is the irrigation levels (I_{100} : % 100, full irrigation; I_{50} : % 50 deficit).

84 The plants were hand-watered with tap water and the trial was terminated 30 days after sowing
 85 (Kadayifci et al. 2005).

86
 87

Table 1. Melon genotypes employed in the study.

Genotype	Provided Location information	Latitude (N)	Latitude (E)	Genotype	Provided Location information	Latitude (N)	Latitude (E)
YYU-1	Van-Sihke-Kiratli	38° 31' 57.9504"	43° 27' 47.3688"	YYU-21	Van-Unseli	38° 59' 6"	43° 35' 16"
YYU-4	Van-Sihke-Kiratli	38° 31' 57.9504"	43° 27' 47.3688"	YYU-22	Van-Ercis	39° 1' 52"	43° 21' 35"
YYU-6	Van-Sihke-Kiratli	38° 31' 57.9504"	43° 27' 47.3688"	YYU-23	Van-Ercek-Irgatli	38° 36' 38.0628"	43° 36' 52.4766"
YYU-10	Van-Sihke	38° 32' 1"	43° 25' 20"	YYU-25	Van-Ercek-Irgatli	38° 36' 38.0628"	43° 36' 52.4766"
YYU-11	Van-Sihke-Kiratli	38° 31' 57.9504"	43° 27' 47.3688"	YYU-29	Van-Ercek-Irgatli	38° 36' 38.0628"	43° 36' 52.4766"
YYU-12	Van-Sihke-Kiratli	38° 31' 57.9504"	43° 27' 47.3688"	YYU-30	Van-Ercek-Irgatli	38° 36' 38.0628"	43° 36' 52.4766"
YYU-13	Van-Sihke-Kiratli	38° 31' 57.9504"	43° 27' 47.3688"	Galia	Standard		
YYU-14	Van-Sihke-Kiratli	38° 31' 57.9504"	43° 27' 47.3688"	Kirkagac F ₁	Yüksel Tohum		
YYU-15	Van-Sihke-Kiratli	38° 31' 57.9504"	43° 27' 47.3688"	Lokum F ₁	Yüksel Tohum		
YYU-18	Van-Cakirbey	39° 9' 15.2064"	43° 21' 35.6868"	Napolyon F ₁	Yüksel Tohum		
YYU-20	Van-Unseli	38° 59' 6"	43° 35' 16"				

88

89 **Seedling Growth Parameters**

90 Upon completing the experiment, various seedling growth parameters were assessed,
91 encompassing shoot and root lengths, shoot diameter, leaf count, shoot and root fresh weights
92 (SFW and RFW), shoot dry weight (SDW), root dry weight (RDW), shoot and root dry matter
93 (SDM and RDM) quantities, and the root-to-shoot ratio (dry weight %) denoted as R/S. Fresh
94 weights of roots and stems were measured on a precision scale and recorded as SFW AND RFW,
95 and the same samples were kept in an oven at 65 °C for 48 hours and their dry weights were
96 recorded as SDW and RDW. Stem and root dry matter ratios were calculated as percentages and
97 recorded as SDM and RDM. Additionally, the genotypes' responses to deficit irrigation were
98 evaluated on a 0-5 scale, with 0 signifying no effect (akin to control plants) and 5 indicating severe
99 wilting and drying in leaves (Cakmakci et al. 2017).

100

101 **Stomatal Traits**

102 Stomatal traits, including stomatal density (units per mm²), stomatal area (µm²), and stomatal
103 width and length (µm), were determined using the lower epidermis of the 4th leaf of the plants. The
104 epidermis was carefully peeled and mounted on a slide with two water droplets (Kurtar et al. 2016).
105 Stoma quantification was conducted using the LAS EZ 3.0 program, examining tissue samples on
106 the slide at 40x magnification under a light microscope (LEICA DM500). Three randomly selected
107 areas of 0.08 mm² were analyzed for accurate assessment.

108

109 **Mineral matter content**

110 Macro-micro nutrient content in plant leaves was determined through the dry combustion method
111 (Kacar and Inal 2010). Plant leaf samples underwent a 48-hour drying process at 65 °C, followed
112 by crushing with a porcelain mortar. Subsequently, 0.5 grams of the dry samples were ashed at 550
113 °C. The resulting ash was dissolved in 3 N HCl. Potassium (K), calcium (Ca), magnesium (Mg),
114 iron (Fe), zinc (Zn), copper (Cu), and manganese (Mn) levels were quantified using an atomic
115 absorption spectrophotometer, while phosphorus (P) content was determined using a
116 spectrophotometer.

117

118 **Enzyme Activation**

119 Superoxide dismutase (SOD) activity was assessed by inhibiting nitro blue tetrazolium (NBT) at
120 560 nm (Jebara et al., 2005). SOD activity was quantified as the unit reducing 50% of NBT.

121 Catalase (CAT) activity was determined by monitoring H₂O₂ disappearance at 240 nm, following
122 Cakmak and Marschner's method (1992). Ascorbate peroxidase (APX) activity was measured by
123 reducing H₂O₂ bound to ascorbic acid at 290 nm, with APX activity defined as the enzyme amount
124 needed to consume 1 μmol of ascorbate per minute (Cakmak and Marschner 1992).

125
126 **Lipid Peroxidation (MDA)**
127 Lipid peroxidation (MDA) was determined by the method of Heath and Packer (1968). The
128 absorbance value of the mixture was determined at 532 and 600 nm wavelengths and the MDA
129 content was calculated with a molar absorption coefficient of 155 mM cm⁻¹.

130
131 **Statistical evaluation**
132 Data from the study were statistically analyzed using the SPSS program, applying analysis of
133 variance with a significance level of p≤0.05. Significant mean differences were further categorized
134 using Duncan's Multiple Comparison Test. The XLSTAT statistical program, along with Principal
135 Component Analysis (PCA) as a multivariate data analysis method, was employed to discern and
136 emphasize similarities or differences resulting from the study's applications and examined features.
137 The extent to which these differences are explained was also determined.

138
139 **RESULTS**

140 **Seedling Growth**
141 Table 2 summarizes significant variations in leaf number, shoot diameter, and shoot/root lengths
142 among melon genotypes subjected to full and deficit irrigation. Overall, deficit irrigation resulted
143 in reduced leaf numbers across all genotypes, with YYU25 and YYU13 exhibiting the highest
144 (8.25) and lowest (4.38) values under full irrigation. Stem diameter showed considerable diversity,
145 ranging from 6.67 mm (YYU11) to 2.71 mm (YYU30). Full irrigation promoted longer shoot
146 lengths in YYU30 and cv. Napolyon (67.75 cm) and shorter lengths in YYU30 (27.69 cm). Root
147 lengths displayed variability, with certain genotypes displaying resilience to full irrigation.

148 Under deficit irrigation, the number of leaves declined, particularly in YYU25 (5.78) and YYU14
149 (2.97). Stem diameter ranged from 5.67 mm (YYU14) to 2.18 mm (YYU30). Shoot lengths were
150 longest in YYU29 (46.24 cm) and shortest in YYU14 (15.75 cm). Most genotypes experienced
151 reductions in shoot and root lengths, indicating diverse responses to deficit irrigation.

152
153

Table 2. Seedling Growth Parameters in Melon Genotypes under I₁₀₀ and I₅₀ deficit Irrigation.

Genotype	Leaf number		Stem diameter (mm)		Shoot length (cm)		Root length (cm)	
	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀
YYU1	5.50 \pm 0.43 ^{b-f}	4.36 \pm 0.13 ^{b-e}	5.00 \pm 1.92 ^{cf}	3.93 \pm 5.28 ^{d-g}	41.88 \pm 1.92 ^{b-d}	29.51 \pm 5.28 ^{b-e}	21.08 \pm 3.47	17.69 \pm 0.72
YYU4	5.50 \pm 0.43 ^{b-f}	3.50 \pm 0.25 ^{e-g}	5.39 \pm 6.57 ^{be}	4.50 \pm 4.35 ^{b-f}	44.58 \pm 6.57 ^{b-d}	28.54 \pm 4.35 ^{b-e}	18.38 \pm 1.85	14.47 \pm 1.53
YYU6	5.64 \pm 0.13 ^{b-f}	4.08 \pm 0.29 ^{c-f}	5.19 \pm 7.37 ^{cf}	4.40 \pm 2.64 ^{b-f}	38.69 \pm 7.37 ^{b-d}	25.21 \pm 2.64 ^{b-f}	20.75 \pm 2.54	15.75 \pm 3.45
YYU10	5.33 \pm 0.63 ^{c-f}	4.08 \pm 0.80 ^{c-f}	5.45 \pm 5.34 ^{bd}	4.31 \pm 7.06 ^{c-g}	33.51 \pm 5.34 ^{b-d}	25.23 \pm 7.06 ^{b-f}	16.67 \pm 2.80	14.88 \pm 1.44
YYU11	4.83 \pm 0.29 ^{d-f}	3.64 \pm 0.13 ^{d-g}	6.67 \pm 4.21 ^a	5.52 \pm 3.58 ^a	29.79 \pm 4.21 ^{cd}	21.08 \pm 3.58 ^{d-f}	23.11 \pm 3.78	17.65 \pm 0.67
YYU12	4.56 \pm 0.55 ^{ef}	3.00 \pm 0.00 ^{fg}	5.89 \pm 3.52 ^{ac}	4.69 \pm 1.35 ^{a-e}	30.83 \pm 4.07 ^{cd}	17.88 \pm 1.35 ^{ef}	19.00 \pm 0.99	18.04 \pm 1.28
YYU13	4.38 \pm 0.57 ^f	4.39 \pm 1.40 ^{b-e}	6.45 \pm 4.05 ^{ab}	5.19 \pm 2.77 ^{a-c}	30.25 \pm 2.88 ^{cd}	19.29 \pm 2.77 ^{ef}	20.04 \pm 4.33	18.28 \pm 5.00
YYU14	5.47 \pm 1.63 ^{b-f}	2.97 \pm 0.61 ^g	6.28 \pm 8.50 ^{ab}	5.67 \pm 0.75 ^a	27.69 \pm 6.46 ^d	15.7 \pm 0.75 ^f	20.94 \pm 4.16	20.97 \pm 3.26
YYU15	5.50 \pm 1.06 ^{b-f}	4.67 \pm 0.76 ^{a-e}	5.18 \pm 10.34 ^{cf}	5.39 \pm 3.50 ^{ab}	36.63 \pm 9.16 ^{b-d}	23.00 \pm 3.50 ^{c-f}	21.81 \pm 4.51	17.97 \pm 3.39
YYU18	8.17 \pm 2.04 ^a	4.25 \pm 0.25 ^{b-e}	4.95 \pm 10.13 ^f	4.45 \pm 3.86 ^{b-f}	36.00 \pm 0.75 ^{b-d}	24.96 \pm 3.63 ^{b-f}	18.96 \pm 3.71	17.92 \pm 1.23
YYU20	6.67 \pm 0.38 ^{a-e}	4.75 \pm 0.43 ^{a-d}	5.14 \pm 7.56 ^{cf}	4.12 \pm 7.63 ^{d-g}	52.17 \pm 10.13 ^{ac}	24.79 \pm 3.86 ^{b-f}	17.63 \pm 2.13	17.96 \pm 4.12
YYU21	7.31 \pm 2.08 ^a	4.28 \pm 0.05 ^{b-e}	5.12 \pm 17.38 ^f	4.91 \pm 3.06 ^{a-d}	47.08 \pm 7.56 ^{a-d}	34.75 \pm 7.63 ^{bc}	16.79 \pm 3.73	18.51 \pm 3.45
YYU22	7.64 \pm 1.76 ^{ab}	4.53 \pm 0.94 ^{b-e}	4.98 \pm 15.02 ^{cf}	4.52 \pm 8.26 ^{b-f}	55.75 \pm 17.38 ^{ab}	33.13 \pm 3.06 ^{b-d}	21.75 \pm 3.85	18.51 \pm 2.61
YYU23	7.67 \pm 0.29 ^{ab}	4.67 \pm 0.38 ^{a-e}	4.57 \pm 8.72 ^{dg}	3.74 \pm 7.12 ^{e-g}	48.04 \pm 15.02 ^{ad}	27.46 \pm 8.26 ^{b-f}	16.76 \pm 2.56	14.90 \pm 0.57
YYU25	8.25 \pm 1.34 ^a	5.78 \pm 0.46 ^a	4.17 \pm 20.82 ^{fg}	3.61 \pm 16.25 ^{fg}	52.79 \pm 8.72 ^{a-c}	36.79 \pm 7.12 ^{ab}	18.17 \pm 2.89	15.20 \pm 1.92
YYU29	5.75 \pm 1.30 ^{b-f}	4.25 \pm 0.75 ^{b-e}	4.40 \pm 21.41 ^{bg}	3.80 \pm 7.50 ^{e-g}	68.75 \pm 20.82 ^a	46.24 \pm 16.25 ^a	15.29 \pm 1.12	18.42 \pm 0.59
YYU30	5.33 \pm 1.28 ^{c-f}	4.75 \pm 0.90 ^{a-d}	2.71 \pm 10.26 ^h	2.18 \pm 4.88 ^h	39.17 \pm 21.41 ^{bd}	33.58 \pm 7.50 ^{bc}	17.94 \pm 3.77	16.45 \pm 0.51
Galia	6.47 \pm 0.21 ^{a-f}	5.29 \pm 0.25 ^{ab}	4.62 \pm 0.75 ^{dg}	4.77 \pm 3.63 ^{a-e}	31.58 \pm 10.26 ^{cd}	34.82 \pm 4.88 ^{bc}	18.58 \pm 3.00	20.38 \pm 2.26
Kirkagac	6.14 \pm 1.32 ^{a-f}	4.50 \pm 0.43 ^{b-e}	4.30 \pm 24.19 ^{eg}	3.83 \pm 5.03 ^{e-g}	44.64 \pm 24.19 ^{bd}	32.92 \pm 5.03 ^{b-d}	17.21 \pm 3.05	21.40 \pm 8.64
Lokum	6.22 \pm 1.28 ^{a-f}	5.00 \pm 0.66 ^{a-c}	3.63 \pm 14.44 ^{gh}	3.29 \pm 3.31 ^g	32.19 \pm 14.44 ^{cd}	33.25 \pm 3.31 ^{b-d}	13.79 \pm 6.29	15.54 \pm 1.91
Napolyon	7.00 \pm 1.56 ^{ad}	4.50 \pm 0.43 ^{b-e}	4.43 \pm 6.25 ^{dg}	3.95 \pm 8.82 ^{d-g}	68.75 \pm 6.25 ^a	29.78 \pm 8.82 ^{b-e}	17.50 \pm 0.76	16.25 \pm 0.78
*p-value	0.001	0.000	0.001	0.000	0.001	0.000	0.152	0.247

I₁₀₀: Full irrigation, I₅₀: 50% deficit irrigation *: Significant distinctions among groups were observed at the P< 0.05 level, as determined by Duncan's multiple comparison test.

155
156
157

158 Table 3 illustrates significant variations in parameters among genotypes and cultivars under full
159 irrigation, excluding RDW. YYU14 displayed the highest SFW in full irrigation (22.06 g), while
160 YYU30 had the lowest (4.34 g). Under deficit irrigation, YYU15 recorded the highest SFW (13.05
161 g), and YYU30 had the lowest (3.50 g). SDW responded positively to stress in YYU29 and
162 YYU30, while other genotypes showed negative effects. In full irrigation, YYU18 exhibited the
163 highest SDW (1.35 g), and YYU30 showed the lowest (0.39 g). Under deficit irrigation, YYU13
164 and YYU15 displayed the highest SDW (0.88 g and 0.86 g, respectively), while YYU30 had the
165 lowest (0.40 g). For RFW, all genotypes experienced a decrease under stress, while RDW increased
166 in four genotypes and three cultivars. YYU15 demonstrated the highest RFW in full irrigation (1.59
167 g), and YYU30 had the lowest (0.49 g). Under deficit irrigation, YYU14 recorded the highest RFW
168 (0.98 g), with YYU30 displaying the lowest (0.31 g). In deficit irrigation, the highest RDW was in
169 YYU12 (0.083 g), while the lowest was in YYU25 (0.024 g).

170
171
172
173
174
175
176

177 **Table 3. Seedling Growth Traits Variation in Melon Genotypes under I_{100} and I_{50} Deficit Irrigation: Selected Values**
 178 **and Standard Deviations.**

Genotype	Shoot fresh weight (g)		Shoot dry weight (g)		Root fresh weight (g)		Root dry weight (g)	
	I_{100}	I_{50}	I_{100}	I_{50}	I_{100}	I_{50}	I_{100}	I_{50}
YYU1	15.99 ±1.05 ^{b-d}	9.87 ±2.83 ^{b-d}	0.80 ±0.11 ^{b-e}	0.55 ±0.18 ^{b-d}	0.89 ±0.13 ^{b-e}	0.62 ±0.21 ^{a-e}	0.044 ±0.004	0.053 ±0.015 ^{a-f}
YYU4	14.73 ±0.23 ^{b-e}	8.41 ±0.47 ^{b-f}	0.99 ±0.13 ^{a-c}	0.63 ±0.08 ^{a-d}	0.89 ±0.11 ^{b-e}	0.53 ±0.14 ^{b-e}	0.056 ±0.005	0.043 ±0.009 ^{b-f}
YYU6	12.23 ±1.59 ^{c-f}	6.82 ±1.30 ^{d-f}	0.72 ±0.13 ^{c-e}	0.53 ±0.12 ^{b-d}	1.04 ±0.55 ^{a-e}	0.48 ±0.15 ^{c-e}	0.056 ±0.028	0.037 ±0.008 ^{c-f}
YYU10	15.36 ±1.40 ^{b-e}	7.32 ±2.00 ^{a-f}	0.97 ±0.10 ^{a-c}	0.54 ±0.18 ^{b-d}	0.81 ±0.18 ^{b-e}	0.52 ±0.15 ^{b-e}	0.051 ±0.012	0.041 ±0.008 ^{b-f}
YYU11	15.99 ±1.68 ^{b-d}	10.83 ±0.37 ^{ab}	1.12 ±0.12 ^{a-c}	0.74 ±0.09 ^{ab}	1.14 ±0.17 ^{a-e}	0.78 ±0.16 ^{a-d}	0.079 ±0.016	0.073 ±0.005 ^{a-c}
YYU12	16.81 ±1.34 ^{bc}	9.29 ±1.11 ^{b-e}	1.10 ±0.15 ^{a-c}	0.70 ±0.01 ^{a-c}	1.28 ±0.14 ^{a-d}	0.96 ±0.18 ^{ab}	0.080 ±0.016	0.083 ±0.02 ^a
YYU13	16.79 ±0.55 ^{bc}	13.01 ±2.32 ^a	0.98 ±0.12 ^{a-c}	0.88 ±0.10 ^a	1.23 ±0.28 ^{a-d}	0.88 ±0.20 ^{a-c}	0.079 ±0.041	0.072 ±0.025 ^{a-d}
YYU14	22.06 ±6.53 ^a	10.28 ±2.34 ^{a-c}	1.35 ±0.57 ^a	0.72 ±0.14 ^{a-c}	1.42 ±0.69 ^{ab}	0.98 ±0.37 ^a	0.087 ±0.028	0.072 ±0.039 ^{a-c}
YYU15	15.01 ±5.06 ^{b-e}	13.05 ±4.16 ^a	0.88 ±0.27 ^{b-d}	0.86 ±0.25 ^a	1.59 ±0.33 ^a	0.94 ±0.38 ^{ab}	0.062 ±0.040	0.074 ±0.028 ^{ab}
YYU18	18.92 ±4.53 ^{ab}	8.11 ^{b-f} ±0.21	1.35 ±0.26 ^a	0.70 ±0.03 ^{a-c}	1.34 ±0.77 ^{a-c}	0.76 ±0.16 ^{a-d}	0.082 ±0.019	0.061 ±0.008 ^{a-e}
YYU20	15.09 ±2.62 ^{b-e}	9.67 ±1.86 ^{b-d}	1.10 ±0.13 ^{a-c}	0.67 ±0.05 ^{a-c}	1.17 ±0.32 ^{a-d}	0.68 ±0.36 ^{a-e}	0.064 ±0.022	0.045 ±0.005 ^{b-f}
YYU21	15.74 ±1.86 ^{b-e}	7.74 ±0.28 ^{b-f}	1.19 ±0.14 ^{ab}	0.75 ±0.02 ^{ab}	1.01 ±0.18 ^{a-e}	0.60 ±0.08 ^{a-e}	0.085 ±0.013	0.073 ±0.031 ^{ab}
YYU22	12.53 ±1.93 ^{c-f}	7.68 ±1.67 ^{b-f}	0.89 ±0.23 ^{b-d}	0.67 ±0.22 ^{a-c}	0.83 ±0.07 ^{b-e}	0.73 ±0.37 ^{a-e}	0.055 ±0.027	0.065 ±0.032 ^{a-e}
YYU23	11.11 ±2.14 ^{d-g}	6.12 ±0.33 ^{e-g}	0.88 ±0.16 ^{b-d}	0.55 ±0.09 ^{b-d}	0.81 ±0.13 ^{b-e}	0.44 ±0.18 ^{c-e}	0.046 ±0.008	0.036 ±0.008 ^{d-f}
YYU25	13.19 ±3.98 ^{c-f}	6.60 ±1.67 ^{d-g}	0.88 ±0.23 ^{b-d}	0.54 ±0.12 ^{b-d}	0.68 ±0.33 ^{c-e}	0.37 ±0.10 ^{de}	0.040 ±0.022	0.024 ±0.004 ^f
YYU29	9.07 ±2.96 ^{fh}	6.57 ±0.74 ^{d-g}	0.53 ±0.23 ^{de}	0.54 ±0.05 ^{b-d}	0.65 ±0.32 ^{de}	0.40 ±0.15 ^{de}	0.054 ±0.035	0.037 ±0.008 ^{c-f}
YYU30	4.34 ±2.32 ^h	3.50 ±0.81 ^g	0.39 ±0.23 ^e	0.40 ±0.14 ^d	0.49 ±0.34 ^e	0.31 ±0.08 ^e	0.048 ±0.023	0.032 ±0.002 ^{ef}
Galia	11.29 ±1.89 ^{d-g}	7.53 ±1.43 ^{b-f}	0.75 ±0.14 ^{c-e}	0.66 ±0.04 ^{a-c}	0.76 ±0.25 ^{b-e}	0.69 ±0.18 ^{a-e}	0.053 ±0.022	0.054 ±0.008 ^{a-f}
Kirkagac	10.43 ±0.84 ^{e-g}	5.48 ±0.82 ^{fg}	0.99 ±0.22 ^{a-c}	0.55 ±0.07 ^{b-d}	0.87 ±0.38 ^{b-e}	0.55 ±0.21 ^{a-e}	0.068 ±0.049	0.048 ±0.013 ^{a-f}
Lokum	6.77 ±1.65 ^{gh}	6.25 ±0.95 ^{e-g}	0.55 ±0.28 ^{de}	0.47 ±0.12 ^{cd}	0.60 ±0.23 ^{de}	0.48 ±0.23 ^{c-e}	0.038 ±0.040	0.034 ±0.014 ^{ef}
Napolyon	13.14 ±3.42 ^{c-f}	8.28 ±2.20 ^{b-f}	0.87 ±0.16 ^{b-d}	0.67 ±0.23 ^{a-c}	0.65 ±0.23 ^{de}	0.60 ±0.28 ^{a-e}	0.045 ±0.037	0.046 ±0.022 ^{b-f}
*p-value	0.001	0.001	0.001	0.013	0.019	0.008	0.357	0.012

179 I_{100} : Full irrigation, I_{50} : 50% deficit irrigation, *: Significant distinctions among groups were observed at the $P < 0.05$
 180 level, as determined by Duncan's multiple comparison test.

181
 182 Significant variations in SDM content and 0-5 scale values were observed across genotypes and
 183 cultivars, with no notable differences in RDM contents and R/S (Table 4). SDM content generally
 184 increased with deficit irrigation, with exceptions in YYU11, YYU20, and Lokum. In full irrigation,
 185 cv. Kirkagac (9.58%) and genotype YYU30 (8.89%) had the highest SDM, while the lowest was
 186 in genotype YYU1 (4.97%). Similarly, in deficit irrigation, YYU30 (11.29%) and cv. Kirkagac
 187 (10.12%) showed the highest SDM, and genotype YYU1 (5.57%) had the lowest.

188 R/S ratio increased with deficit irrigation, except for YYU6, YYU25, YYU29, and YYU30. In
 189 full irrigation, genotypes YYU30 (0.142) and cv. Lokum (0.049) had the highest and lowest R/S
 190 ratios, respectively. Under deficit irrigation, YYU12 (0.119) and YYU25 (0.050) showed the
 191 highest and lowest R/S ratios, respectively. Genotypes YYU10 and YYU12 shared the highest 0-
 192 5 scale values (3.333), while cv. Lokum had the lowest (1.667).

197 **Table 4.** Dry Matter, Root/Shoot Ratio, and Vigor Assessment in Melon Genotypes under Full and 50% Deficit
 198 **Irrigation: Mean Values and Standard Deviations.**

Genotype	Shoot dry matter content (%)		Root dry matter content (%)		Root: Shoot ratio		0-5 scale
	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀	I ₅₀
YYU1	4.97±0.40 ^d	5.57±0.38 ^g	5.11±1.20	8.88±2.89	0.056±0.003	0.102±0.047	2.333 ±1.15 ^{a-c}
YYU4	6.75 ±0.97 ^{b-d}	7.53±0.78 ^{d-f}	6.42±1.33	8.25±1.16	0.057±0.002	0.067±0.007	2.667 ±0.58 ^{a-c}
YYU6	5.87±0.65 ^{cd}	7.69±0.61 ^{d-f}	5.55±1.48	8.00±2.24	0.077±0.034	0.072±0.021	3.000±0.00 ^{ab}
YYU10	6.33 ±0.12 ^{b-d}	7.36±0.46 ^{d-f}	6.27±0.60	8.02±0.93	0.052±0.007	0.078±0.016	3.333±0.58 ^a
YYU11	7.00 ±0.70 ^{a-d}	6.80±0.75 ^{fg}	6.93±0.92	9.68±2.73	0.071±0.013	0.099±0.008	2.333 ±0.58 ^{a-c}
YYU12	6.54 ±0.48 ^{b-d}	7.58±0.98 ^{d-f}	6.22±0.76	8.75±2.31	0.069±0.016	0.119±0.033	3.333±0.58 ^a
YYU13	5.83±0.88 ^{cd}	6.79±0.46 ^{fg}	6.09±1.86	8.08±1.23	0.068±0.022	0.083±0.031	2.667 ±0.58 ^{a-c}
YYU14	5.99±0.93 ^{cd}	7.03 ±0.31 ^{e-g}	6.48±2.03	7.10±1.07	0.090±0.026	0.097±0.033	2.333 ±0.58 ^{a-c}
YYU15	5.90±0.20 ^{cd}	6.66±0.71 ^{fg}	3.69±1.77	7.97±0.96	0.070±0.053	0.084±0.010	2.000±0.00 ^{bc}
YYU18	7.19 ±0.93 ^{a-d}	8.63 ±0.19 ^{b-e}	7.06±2.92	8.26±2.00	0.061±0.003	0.087±0.012	1.667±0.58 ^c
YYU20	7.37 ±1.12 ^{a-d}	7.06 ±1.38 ^{e-g}	5.89±2.79	9.55±0.45	0.058±0.013	0.066±0.004	2.667 ±0.58 ^{a-c}
YYU21	7.58 ±1.06 ^{a-d}	9.74 ±0.40 ^{a-c}	8.62±2.06	11.81±3.57	0.073±0.021	0.096±0.039	2.667 ±0.58 ^{a-c}
YYU22	7.01 ±0.83 ^{a-d}	8.64 ±0.88 ^{b-e}	6.60±2.99	8.98±2.15	0.060±0.014	0.094±0.029	3.000±0.00 ^{ab}
YYU23	7.98 ±0.72 ^{a-c}	8.96 ±1.02 ^{b-d}	5.74±0.75	8.90±2.11	0.052±0.003	0.067±0.021	2.667 ±0.58 ^{a-c}
YYU25	6.73 ±0.33 ^{b-d}	8.31 ±0.85 ^{c-f}	5.61±0.68	7.26±0.74	0.050±0.037	0.050±0.010	3.000±1.00 ^{ab}
YYU29	6.13 ±3.19 ^{b-d}	8.25 ±0.57 ^{c-f}	7.80±3.08	9.67±2.83	0.122±0.090	0.069±0.014	2.000±0.00 ^{bc}
YYU30	8.89±0.87 ^{ab}	11.29±1.64 ^a	12.72±7.56	9.44±1.94	0.142±0.066	0.092±0.040	3.000±1.00 ^{ab}
Galia	6.81 ±1.88 ^{b-d}	9.02 ±2.11 ^{b-d}	7.72±4.60	8.07±1.19	0.071±0.031	0.082±0.072	2.333 ±0.58 ^{a-c}
Kirkagac	9.58±2.42 ^a	10.12 ±0.25 ^{a-b}	7.23±3.33	9.02±1.67	0.067±0.022	0.085±0.011	3.000±0.00 ^{ab}
Lokum	7.83 ±2.61 ^{a-c}	7.50±1.08 ^{df}	5.27±3.86	7.46±2.05	0.049±0.006	0.068±0.005	1.667±0.58 ^c
Napolyon	6.88 ±2.15 ^{b-d}	7.91±0.92 ^{df}	6.28±2.97	7.75±0.35	0.057±0.012	0.069±0.006	2.667 ±0.58 ^{a-c}
*p-value	0.061	0.000	0.398	0.581	0.227	0.253	0.052

199 I₁₀₀: Full irrigation, I₅₀: 50% deficit irrigation, *: Significant distinctions among groups were observed at the P < 0.05
 200 level, as determined by Duncan's multiple comparison test.

202 Stomatal Traits

203 Stomatal characteristics in melon genotypes and cultivars significantly differed under full and
 204 deficit irrigation (Table 5). Stomatal length and area decreased in 52.38% of cases, while width
 205 and density increased by 71.43% and 61.91%, respectively. In full irrigation, genotype YYU22 had
 206 the tallest stoma (23.73 μm), and genotype YYU6 had the shortest (8.47 μm). Under deficit
 207 irrigation, genotypes YYU25 and YYU22 showed the tallest stomata (21.80 μm and 21.47 μm),
 208 and genotype YYU6 had the shortest (14.47 μm). For stoma width, genotype YYU13 had the
 209 widest (15.58 μm) in full irrigation, and genotype YYU6 had the narrowest (7.97 μm). In deficit
 210 irrigation, genotype YYU21 had the widest (17.20 μm), and genotype YYU6 had the narrowest
 211 (10.78 μm). Regarding stoma area, genotype YYU22 had the widest (252.94 μm²) in full irrigation,
 212 and genotype YYU6 had the narrowest (53.03 μm²). In deficit irrigation, genotype YYU21 had the
 213 widest (288.05 μm²), and genotype YYU6 had the narrowest (124.28 μm²). In full irrigation, cv.
 214 Kirkagac displayed the highest stoma intensity (362.50 units per mm²), and genotype YYU15 had
 215 the lowest (56.25 units per mm²). Under deficit irrigation, genotype YYU6 showed the highest
 216 stoma intensity (516.67 units per mm²), with genotype YYU15 displaying the lowest (108.33 units
 217 per mm²).

218 **Table 5.** Stomatal Traits of Melon Genotypes under Full and 50% Deficit Irrigation; Mean Values and Standard
 219 Deviations.

Genotype	Stoma height (μm)		Stoma width (μm)		Stoma area (μm^2)		Stoma intensity (unit per mm^2)	
	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀
YYU1	19.04 ±1.98 ^{b-f}	17.93 ±1.66 ^{a-f}	12.76 ±1.37 ^{c-h}	13.82 ±1.37 ^{a-f}	189.56 ±12.55 ^{b-g}	194.09 ±20.41 ^{b-g}	108.33 ±19.09 ^{b-i}	137.50 ±21.65 ^{gh}
YYU4	12.61 ±1.63 ^h	15.90 ±0.80 ^{d-f}	9.49 ±0.78 ⁱⁱ	13.16 ±2.13 ^{b-f}	94.42 ±18.34 ^{hi}	164.91 ±33.80 ^{d-f}	250.00 ±12.50 ^{b-d}	350.00 ±33.07 ^b
YYU6	8.47 ±0.71 ^l	14.47 ±0.2 ^f	7.97 ±0.15 ^l	10.95 ±0.98 ^{ef}	53.03 ±4.76 ^l	124.28 ±9.46 ^f	162.50 ±12.50 ^{c-h}	516.67 ±7.22 ^a
YYU10	17.32 ±0.64 ^{c-g}	16.97 ±2.32 ^{b-f}	13.43 ±0.80 ^{a-g}	13.50 ±2.10 ^{b-f}	182.38 ±4.70 ^{c-g}	179.11 ±33.72 ^{b-f}	129.17 ±28.87 ^{g-i}	320.83 ±14.43 ^{bc}
YYU11	19.02 ±0.98 ^{b-f}	17.10 ±4.10 ^{b-f}	14.33 ±1.61 ^{a-d}	14.25 ±3.97 ^{a-f}	214.76 ±34.44 ^{a-e}	198.75 ±102.02 ^{b-f}	216.67 ±14.43 ^{c-e}	316.67 ±19.09 ^{bc}
YYU12	18.82 ±1.50 ^{b-f}	19.33 ±0.91 ^{a-e}	14.00 ±1.8 ^{a-e}	16.69 ±2.24 ^{ab}	206.30 ±28.21 ^{a-f}	253.59 ±38.79 ^{a-c}	278.13 ±57.17 ^{bc}	270.83 ±52.04 ^{cd}
YYU13	16.65 ±1.54 ^{c-g}	20.37 ±1.01 ^{a-c}	15.58 ±0.70 ^a	13.94 ±0.85 ^{a-f}	204.03 ±25.93 ^{a-f}	222.66 ±11.80 ^{a-e}	312.50 ±54.49 ^{ab}	133.33 ±7.22 ^{gh}
YYU14	17.08 ±2.48 ^{b-g}	15.40 ±2.46 ^{def}	11.64 ±0.73 ^{e-i}	10.78 ±0.85 ^f	156.35 ±26.69 ^{fg}	130.05 ±19.67 ^f	220.83 ±127.68 ^{ce}	212.50 ±12.50 ^{ef}
YYU15	18.00 ±1.60 ^{b-g}	18.60 ±0.56 ^{a-f}	12.12 ±0.04 ^{d-h}	13.02 ±1.97 ^{b-f}	171.23 ±14.60 ^{e-g}	189.50 ±22.77 ^{b-f}	56.25 ±8.84 ⁱ	108.33 ±14.43 ^h
YYU18	18.09 ±2.75 ^{b-g}	16.07 ±0.40 ^{d-f}	10.96 ±1.35 ^{g-i}	11.54 ±0.88 ^{d-f}	155.50 ±29.53 ^{fg}	145.77 ±14.85 ^{ef}	116.67 ±7.22 ^{h-i}	237.50 ±45.07 ^{de}
YYU20	20.52 ±1.12 ^{bc}	17.17 ±2.16 ^{b-f}	14.56 ±1.79 ^{a-d}	14.69 ±1.23 ^{a-d}	235.00 ±37.37 ^{a-d}	198.37 ±33.77 ^{b-f}	87.50 ±12.50 ⁱⁱ	200.00 ±45.07 ^{ef}
YYU21	16.80 ±1.27 ^{d-g}	20.97 ±2.65 ^{ab}	12.98 ±1.40 ^{b-h}	17.20 ±4.23 ^a	170.25 ±6.95 ^{e-g}	288.05 ±102.96 ^a	179.17 ±7.22 ^{e-h}	133.33 ±7.22 ^{gh}
YYU22	23.73 ±2.34 ^a	21.47 ±2.17 ^a	13.56 ±0.61 ^{a-f}	14.66 ±1.00 ^{a-e}	252.94 ±31.48 ^a	248.11 ±42.36 ^{a-d}	75.00 ±21.65 ^{l-i}	166.67 ±7.22 ^{fg}
YYU23	15.27 ±2.20 ^{gh}	19.97 ±3.09 ^{a-d}	11.51 ±0.88 ^{e-i}	12.77 ±1.98 ^{c-f}	137.74 ±20.67 ^{fg}	203.29 ±64.20 ^{b-f}	129.17 ±7.22 ^{g-i}	204.17 ±28.87 ^{ef}
YYU25	16.50 ±1.84 ^{fg}	21.80 ±2.84 ^a	10.74 ±1.02 ^{hi}	15.47 ±1.27 ^{a-c}	138.16 ±6.71 ^{fg}	264.78 ±42.66 ^{ab}	137.50 ±21.65 ^{g-i}	170.83 ±14.43 ^{fg}
YYU29	20.06 ±1.79 ^{b-e}	15.90 ±1.22 ^{d-f}	15.40 ±1.18 ^{ab}	15.22 ±0.97 ^{a-d}	243.55 ±39.48 ^{ab}	189.34 ±4.10 ^{b-f}	141.67 ±19.09 ^{f-i}	170.83 ±7.22 ^{fg}
YYU30	19.71 ±2.72 ^{b-f}	20.50 ±2.52 ^{a-c}	15.21 ±2.34 ^{a-c}	12.95 ±1.19 ^{c-f}	238.61 ±70.97 ^{a-c}	208.50 ±31.91 ^{a-f}	195.83 ±7.22 ^{d-g}	166.67 ±47.32 ^{fg}
Galia	21.29 ±1.35 ^{ab}	19.50 ±2.65 ^{a-e}	13.02 ±1.76 ^{b-h}	13.39 ±1.22 ^{b-f}	218.30 ±39.26 ^{a-e}	206.52 ±46.52 ^{a-f}	87.50 ±33.07 ⁱⁱ	133.33 ±14.43 ^{gh}
Kirkagac	18.56 ±1.94 ^{b-g}	16.63 ±0.75 ^{c-f}	12.42 ±0.82 ^{d-h}	13.72 ±1.65 ^{a-f}	180.46 ±13.41 ^{d-g}	179.66 ±29.11 ^{b-f}	362.50 ±12.50 ^a	279.17 ±40.18 ^{cd}
Lokum	20.27 ±1.11 ^{b-d}	19.60 ±3.55 ^{a-e}	14.47 ±1.42 ^{a-d}	11.59 ±1.65 ^{d-f}	230.48 ±28.60 ^{a-d}	175.43 ±9.58 ^{c-f}	208.33 ±7.22 ^{c-e}	200.00 ±54.49 ^{ef}
Napolyon	16.41 ±1.91 ^{fg}	19.67 ±0.35 ^{a-d}	11.31 ±0.95 ^{f-i}	12.67 ±0.29 ^{c-f}	146.46 ±27.17 ^{gh}	195.61 ±1.91 ^{b-f}	216.67 ±19.09 ^{c-e}	137.50 ±12.50 ^{gh}
*p-value	0.001	0.001	0.001	0.007	0.001	0.003	0.001	0.001

220 I₁₀₀: Full irrigation, I₅₀: 50% deficit irrigation, *: Significant distinctions among groups were observed at the P < 0.05
 221 level, as determined by Duncan's multiple comparison test.

223 Mineral Content

224 Differences in mean K, Ca, and Mg contents among melon genotypes and cultivars were
 225 significant under both full and deficit irrigation conditions, with P content showing significance
 226 exclusively in deficit irrigation (Table 6). Deficit irrigation led to a decrease in P and Mg uptake
 227 in 66% of genotypes and cultivars, while 90% experienced reduced Ca intake. In full irrigation, the
 228 highest K content occurred in genotypes YYU6 (6.28%) and YYU21 (6.25%), while the lowest
 229 was in genotype YYU20 (2.91%). Under deficit irrigation, the highest K content was in genotype
 230 YYU29 (6.58%), and the lowest was in genotype YYU12 (4.56%) and cv. Lokum (4.44%). For P
 231 content in deficit irrigation, the highest was in genotype YYU29 (0.92%), and the lowest was in
 232 genotype YYU10 (0.46%). In both full and deficit irrigation, the highest Ca content was in
 233 genotypes YYU30 (7.72% and 5.65%, respectively), and the lowest was in genotype YYU12
 234 (3.83% and 4.22%, respectively). In full irrigation, the highest Mg content was in genotype YYU30
 235 (7.72%), and the lowest was in genotype YYU20 (0.49%). Under deficit irrigation, the highest Mg
 236 content was in genotype YYU30 (0.73%), and the lowest was in cv. Lokum (0.48%).

Table 6. Macro mineral contents in melon genotypes under full and 50% deficit irrigation.

Genotype	K (%)		P (%)		Ca (%)		Mg (%)	
	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀
YYU1	6.00 ± 0.57 ^{ac}	5.78 ± 0.51 ^{ac}	1.06 ± 0.02	0.76 ± 0.07 ^{ab}	5.73 ± 0.52 ^{b-e}	4.34 ± 0.60 ^{cd}	0.64 ± 0.05 ^{d-g}	0.57 ± 0.06 ^{b-e}
YYU4	4.87 ± 0.14 ^{af}	5.55 ± 0.61 ^{ad}	0.71 ± 0.28	0.85 ± 0.10 ^{ab}	4.39 ± 1.48 ^{ef}	4.28 ± 0.52 ^{cd}	0.71 ± 0.09 ^{c-g}	0.52 ± 0.10 ^{de}
YYU6	6.28 ± 0.61 ^a	6.09 ± 0.91 ^{ab}	0.86 ± 0.33	0.79 ± 0.15 ^{ab}	5.93 ± 1.06 ^{b-e}	4.31 ± 0.75 ^{cd}	0.69 ± 0.20 ^{d-g}	0.51 ± 0.06 ^{de}
YYU10	5.81 ± 0.42 ^{a-d}	5.92 ± 0.58 ^{ac}	0.80 ± 0.09	0.46 ± 0.07 ^c	5.75 ± 0.65 ^{b-e}	4.32 ± 0.43 ^{cd}	0.68 ± 0.10 ^{d-g}	0.52 ± 0.03 ^{de}
YYU11	4.36 ± 0.43 ^{cg}	4.97 ± 0.25 ^{bd}	0.73 ± 0.19	0.62 ± 0.12 ^{bc}	5.26 ± 0.33 ^{c-f}	4.29 ± 0.39 ^{cd}	0.75 ± 0.10 ^{c-f}	0.56 ± 0.02 ^{be}
YYU12	4.52 ± 0.59 ^{bf}	4.56 ± 0.64 ^d	0.87 ± 0.12	0.59 ± 0.10 ^{bc}	6.19 ± 0.64 ^{a-e}	4.22 ± 0.21 ^d	0.92 ± 0.15 ^{bc}	0.60 ± 0.04 ^{a-e}
YYU13	5.48 ± 0.27 ^{ae}	5.52 ± 0.64 ^{ad}	0.92 ± 0.01	0.82 ± 0.17 ^{ab}	7.22 ± 0.42 ^{ab}	5.04 ± 0.45 ^{a-d}	1.01 ± 0.13 ^b	0.66 ± 0.03 ^{a-c}
YYU14	4.76 ± 0.22 ^{af}	5.12 ± 0.72 ^{bd}	0.78 ± 0.14	0.69 ± 0.20 ^{ac}	5.83 ± 0.73 ^{b-e}	5.30 ± 0.31 ^{a-d}	0.71 ± 0.17 ^{c-g}	0.69 ± 0.06 ^{ab}
YYU15	3.42 ± 0.68 ^{fg}	5.58 ± 0.09 ^{ad}	0.87 ± 0.14	0.74 ± 0.03 ^{ab}	5.57 ± 2.99 ^{b-f}	4.59 ± 0.38 ^{a-d}	0.65 ± 0.31 ^{d-g}	0.57 ± 0.02 ^{b-e}
YYU18	3.95 ± 0.44 ^{eg}	5.48 ± 0.79 ^{ad}	0.69 ± 0.00	0.63 ± 0.16 ^{bc}	4.61 ± 0.18 ^{ef}	5.11 ± 0.79 ^{a-d}	0.58 ± 0.09 ^{e-g}	0.66 ± 0.10 ^{a-c}
YYU20	2.91 ± 0.53 ^g	5.19 ± 1.17 ^{bd}	0.78 ± 0.03	0.73 ± 0.14 ^{ab}	3.83 ± 0.53 ^f	4.43 ± 0.47 ^{cd}	0.49 ± 0.09 ^g	0.57 ± 0.06 ^{b-e}
YYU21	6.25 ± 1.69 ^a	5.34 ± 0.71 ^{bd}	0.79 ± 0.10	0.75 ± 0.11 ^{ab}	6.45 ± 0.51 ^{a-d}	4.87 ± 0.30 ^{a-d}	0.85 ± 0.18 ^{b-d}	0.69 ± 0.05 ^{a-b}
YYU22	5.29 ± 1.19 ^{ae}	5.33 ± 0.46 ^{bd}	0.98 ± 0.11	0.76 ± 0.06 ^{ab}	5.55 ± 0.98 ^{b-f}	5.10 ± 0.60 ^{a-d}	0.65 ± 0.10 ^{d-g}	0.64 ± 0.14 ^{a-d}
YYU23	5.32 ± 0.29 ^{ae}	5.33 ± 0.16 ^{bd}	0.93 ± 0.00	0.69 ± 0.35 ^{ac}	6.83 ± 0.17 ^{a-c}	4.53 ± 1.17 ^{b-d}	0.78 ± 0.06 ^{c-e}	0.58 ± 0.04 ^{b-e}
YYU25	4.52 ± 0.04 ^{bf}	4.71 ± 0.80 ^{cd}	0.71 ± 0.36	0.85 ± 0.07 ^{ab}	6.06 ± 0.54 ^{a-e}	4.97 ± 0.62 ^{a-d}	0.70 ± 0.08 ^{c-g}	0.61 ± 0.09 ^{a-d}
YYU29	4.64 ± 1.47 ^{af}	6.58 ± 0.69 ^a	0.85 ± 0.21	0.92 ± 0.12 ^a	5.14 ± 1.32 ^{c-f}	5.53 ± 0.36 ^{ab}	0.60 ± 0.11 ^{e-g}	0.67 ± 0.02 ^{ab}
YYU30	6.16 ± 1.45 ^{ab}	5.01 ± 0.19 ^{bd}	0.85 ± 0.17	0.65 ± 0.04 ^{bc}	7.72 ± 1.25 ^a	5.65 ± 0.34 ^a	1.41 ± 0.01 ^a	0.73 ± 0.05 ^a
Galia	5.14 ± 0.37 ^{ae}	4.87 ± 0.18 ^{cd}	0.93 ± 0.12	0.74 ± 0.08 ^{ab}	5.17 ± 0.61 ^{a-e}	5.37 ± 0.56 ^{a-c}	0.67 ± 0.09 ^{d-g}	0.64 ± 0.07 ^{a-d}
Kirkagac	4.32 ± 1.55 ^{dg}	5.23 ± 0.58 ^{bd}	0.78 ± 0.26	0.69 ± 0.07 ^{ac}	6.12 ± 0.74 ^{c-f}	4.69 ± 0.37 ^{a-d}	0.55 ± 0.02 ^{e-g}	0.53 ± 0.04 ^{c-e}
Lokum	5.09 ± 1.03 ^{ae}	4.44 ± 0.53 ^d	0.85 ± 0.33	0.80 ± 0.10 ^{ab}	5.00 ± 0.36 ^{d-f}	4.78 ± 0.25 ^{a-d}	0.52 ± 0.08 ^{f-g}	0.48 ± 0.03 ^e
Napolyon	5.73 ± 0.03 ^{ad}	4.97 ± 0.27 ^{b-d}	0.83 ± 0.01	0.77 ± 0.01 ^{ab}	5.34 ± 1.25 ^{c-f}	5.08 ± 0.74 ^{a-d}	0.61 ± 0.10 ^{d-g}	0.60 ± 0.11 ^{a-e}
*p-value	0.000	0.020	0.709	0.051	0.001	0.031	0.000	0.001

I₁₀₀: Full irrigation, I₅₀: 50% deficit irrigation, *: Significant distinctions among groups were observed at the P < 0.05 level, as determined by Duncan's multiple comparison test.

240
241
242
243 Significant differences in Zn and Cu contents among melon genotypes and cultivars were
244 observed under both full and deficit irrigation conditions. Meanwhile, Fe content showed
245 significant differences only under full irrigation, and Mn content exhibited significance solely
246 under limited irrigation conditions (Table 7). Approximately 76% of all genotypes and cultivars
247 were adversely affected by deficit irrigation for Fe uptake, and all genotypes showed negative
248 effects on Mn uptake under deficit irrigation. In full irrigation, the highest Zn content was in
249 genotype YYU29 (238.35 ppm), and the lowest was in genotype YYU4 (47.27 ppm). Under deficit
250 irrigation, the highest Zn content was in genotype YYU29 (193.78 ppm), and the lowest was in
251 genotype cv. Napolyon (46.22 ppm). Regarding Cu content, in full irrigation, the highest was in
252 genotype YYU23 (25.82 ppm), and the lowest was in genotypes YYU25 (10.39 ppm) and YYU11
253 (10.71 ppm). Under deficit irrigation, the highest Cu content was in genotype YYU25 (14.56 ppm),
254 and the lowest was in genotype YYU10 (6.39 ppm). For Fe content, in full irrigation, the highest
255 was in genotype YYU6 (232.57 ppm), and the lowest was in genotype YYU25 (111.70 ppm). In
256 deficit irrigation, the highest Mn content was in cv. Galia (64.41 ppm), and the lowest was in cv.
257 Kirkagac (32.67 ppm).

258

259

Table 7. Micro mineral contents in melon genotypes under full and 50% deficit irrigation.

Genotype	Zn (ppm)		Cu (ppm)		Fe (ppm)		Mn (ppm)	
	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀
YYU1	130.26 \pm 106.20 ^{b-g}	67.01 \pm 43.97 ^{e-g}	13.63 \pm 1.06 ^{c-f}	6.81 \pm 0.89 ^{ef}	162.10 \pm 37.56 ^{b-f}	134.56 \pm 20.15	67.29 \pm 12.60	43.21 \pm 8.46
YYU4	47.27 \pm 10.28 ^g	164.38 \pm 12.30 ^{a-c}	18.63 \pm 2.39 ^b	6.61 \pm 0.04 ^{ef}	155.20 \pm 10.19 ^{c-f}	136.38 \pm 7.82	60.74 \pm 20.91	37.87 \pm 9.79
YYU6	147.04 \pm 104.75 ^{a-f}	157.13 \pm 23.60 ^{a-d}	11.88 \pm 1.87 ^{df}	8.35 \pm 1.77 ^{c-f}	232.57 \pm 69.14 ^a	162.69 \pm 5.29	68.66 \pm 22.35	50.87 \pm 6.39
YYU10	183.23 \pm 35.97 ^{a-e}	50.91 \pm 11.64 ^{f-g}	11.18 \pm 0.65 ^{ef}	6.39 \pm 0.42 ^f	149.46 \pm 8.51 ^{c-f}	123.35 \pm 4.33	63.18 \pm 20.97	39.95 \pm 2.19
YYU11	140.05 \pm 52.77 ^{a-g}	178.57 \pm 7.78 ^{ab}	10.71 \pm 1.13 ^f	7.09 \pm 1.19 ^{e-f}	160.95 \pm 9.29 ^{b-f}	153.55 \pm 21.76	75.77 \pm 4.99	40.67 \pm 6.39
YYU12	121.00 \pm 21.96 ^{c-g}	129.88 \pm 6.49 ^{a-f}	12.50 \pm 1.64 ^{c-f}	8.10 \pm 0.65 ^{c-f}	165.81 \pm 14.91 ^{b-f}	139.63 \pm 8.69	73.66 \pm 23.93	52.88 \pm 4.89
YYU13	89.32 \pm 52.03 ^{e-g}	62.54 \pm 18.39 ^{fg}	14.72 \pm 2.76 ^{b-f}	6.50 \pm 2.14 ^f	185.54 \pm 4.53 ^{a-e}	141.08 \pm 7.42	92.86 \pm 15.06	55.40 \pm 8.37
YYU14	172.39 \pm 54.82 ^{a-e}	69.13 \pm 48.34 ^{e-g}	12.00 \pm 1.45 ^{d-f}	7.48 \pm 0.93 ^{d-f}	191.18 \pm 18.82 ^{a-d}	145.41 \pm 17.35	89.48 \pm 26.99	47.9 \pm 8.61
YYU15	219.47 \pm 31.76 ^{a-c}	114.18 \pm 64.39 ^{b-g}	11.60 \pm 0.18 ^{ef}	11.10 \pm 3.17 ^{bc}	169.04 \pm 1.20 ^{b-f}	177.98 \pm 40.13	49.98 \pm 23.99	34.20 \pm 1.05
YYU18	150.62 \pm 25.86 ^{a-f}	57.85 \pm 62.04 ^{f-g}	13.00 \pm 1.26 ^{c-f}	9.33 \pm 0.35 ^{c-f}	215.94 \pm 60.01 ^{ab}	130.42 \pm 8.76	74.21 \pm 5.87	40.13 \pm 4.75
YYU20	173.19 \pm 73.28 ^{a-e}	119.20 \pm 18.38 ^{a-g}	14.24 \pm 0.90 ^{b-f}	9.11 \pm 1.63 ^{c-f}	147.46 \pm 7.77 ^{c-f}	168.24 \pm 88.29	65.51 \pm 18.79	39.96 \pm 9.89
YYU21	156.17 \pm 75.16 ^{a-f}	61.42 \pm 27.11 ^{fg}	12.70 \pm 1.44 ^{c-f}	9.40 \pm 0.14 ^{c-f}	155.61 \pm 21.06 ^{c-f}	146.78 \pm 26.75	68.18 \pm 21.50	46.23 \pm 5.28
YYU22	226.33 \pm 43.38 ^{ab}	99.74 \pm 57.25 ^{b-g}	12.92 \pm 0.41 ^{c-f}	9.78 \pm 0.26 ^{c-f}	149.75 \pm 6.44 ^{c-f}	128.18 \pm 7.89	59.81 \pm 5.89	34.77 \pm 3.16
YYU23	57.31 \pm 1.51 ^{fg}	99.26 \pm 18.13 ^{b-g}	25.82 \pm 7.49 ^a	7.68 \pm 1.95 ^{c-f}	128.97 \pm 5.14 ^{ef}	101.85 \pm 39.90	81.25 \pm 17.10	44.03 \pm 18.17
YYU25	96.00 \pm 24.49 ^{d-g}	86.00 \pm 83.82 ^{c-g}	10.39 \pm 2.05 ^f	14.56 \pm 4.35 ^a	111.70 \pm 39.83 ^f	134.48 \pm 17.05	57.89 \pm 33.99	44.84 \pm 8.36
YYU29	238.35 \pm 23.15 ^a	193.78 \pm 1.36 ^a	12.84 \pm 3.13 ^{c-f}	10.72 \pm 2.58 ^{b-d}	142.19 \pm 8.77 ^{d-f}	117.36 \pm 4.07	88.90 \pm 11.45	41.48 \pm 9.41
YYU30	217.16 \pm 15.01 ^{a-c}	78.90 \pm 55.14 ^{d-g}	16.02 \pm 0.58 ^{b-e}	9.25 \pm 0.98 ^{c-f}	206.30 \pm 55.75 ^{a-c}	139.05 \pm 48.08	95.26 \pm 30.12	42.74 \pm 6.74
Galia	95.83 \pm 23.83 ^{d-g}	144.02 \pm 29.03 ^{a-e}	14.25 \pm 1.13 ^{b-f}	11.09 \pm 1.86 ^{bc}	181.86 \pm 21.21 ^{a-e}	219.30 \pm 70.51	98.72 \pm 22.67	64.41 \pm 16.84
Kirkagac	148.50 \pm 72.15 ^{a-f}	52.6 \pm 17.37 ^{f-g}	12.62 \pm 2.11 ^{c-f}	8.43 \pm 0.62 ^{c-f}	137.45 \pm 25.57 ^{d-f}	150.76 \pm 35.76	59.58 \pm 5.41	32.67 \pm 1.86
Lokum	185.73 \pm 9.31 ^{a-e}	146.74 \pm 79.52 ^{a-e}	17.03 \pm 2.03 ^{bc}	10.17 \pm 1.11 ^{c-e}	169.38 \pm 35.88 ^{b-f}	136.44 \pm 14.47	54.66 \pm 26.67	39.72 \pm 11.90
Napolyon	194.65 \pm 3.52 ^{a-d}	46.22 \pm 7.29 ^g	16.69 \pm 4.49 ^{b-d}	13.96 \pm 3.15 ^{ab}	159.86 \pm 13.59 ^{b-f}	143.42 \pm 16.24	58.99 \pm 15.33	35.18 \pm 2.30
*p-value	0.001	0.000	0.000	0.000	0.002	0.094	0.144	0.008

I₁₀₀: Full irrigation, I₅₀: 50% deficit irrigation, *: Significant distinctions among groups were observed at the P < 0.05 level, as determined by Duncan's multiple comparison test.

Enzyme Activation and MDA Content

CAT activity significantly increased under deficit irrigation, indicating a response to water stress. APX activity increased across all melon genotypes and cultivars in deficit irrigation, with significant differences found only in the full irrigation group (Table 8). SOD enzyme activity increased with deficit irrigation, and MDA content increased in 76% of genotypes, yet no significant differences were observed among genotypes and cultivars in full and restricted irrigation applications. CAT activity in melon genotypes showed a 57.14% decrease under deficit irrigation-induced water stress. The genotype YYU25 exhibited the highest CAT content (0.104 mmol g⁻¹ FW), followed by cv. Kirkagac (0.086 mmol g⁻¹ FW), and then genotypes YYU14, YYU23, YYU29, YYU30, and cv. Lokum (0.040-0.040 mmol g⁻¹ FW). Other melon genotypes and cultivars displayed lower CAT content. In full irrigation, the highest APX content was in genotype YYU4 (0.714 mmol g⁻¹ FW), while the lowest Fe content was in genotype YYU29 (0.152 mmol g⁻¹ FW).

Table 8. Enzyme Activation and MDA Content in Melon Genotypes under Full and 50% Deficit Irrigation.

Genotip	CAT (mmol g ⁻¹ FW)		SOD (U mg ⁻¹ FW)		APX (mmol g ⁻¹ FW)		MDA (μmol g ⁻¹ FW)	
	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀	I ₁₀₀	I ₅₀
	YYU1	0.052 _{±0.023}	0.004 _{±0.002} ^c	82.68 _{±45.60}	274.48 _{±223.88}	0.393 _{±0.287} ^{a-f}	1.101 _{±0.352}	2.624 _{±0.711}
YYU4	0.013 _{±0.010}	0.015 _{±0.007} ^c	83.86 _{±27.82}	312.28 _{±69.96}	0.714 _{±0.245} ^a	1.548 _{±0.267}	3.570 _{±1.273}	3.355 _{±1.419}
YYU6	0.110 _{±0.006}	0.017 _{±0.017} ^c	104.33 _{±16.83}	267.38 _{±141.04}	0.571 _{±0.161} ^{a-d}	1.313 _{±0.180}	3.140 _{±0.521}	2.968 _{±1.677}
YYU10	0.023 _{±0.011}	0.005 _{±0.005} ^c	103.33 _{±13.33}	252.12 _{±85.92}	0.506 _{±0.037} ^{a-e}	1.077 _{±0.172}	4.000 _{±0.785}	6.194 _{±1.707}
YYU11	0.069 _{±0.006}	0.030 _{±0.025} ^c	98.72 _{±17.95}	167.02 _{±40.83}	0.473 _{±0.045} ^{a-f}	1.393 _{±0.179}	4.430 _{±0.521}	5.548 _{±1.359}
YYU12	0.076 _{±0.069}	0.024 _{±0.033} ^c	103.28 _{±32.44}	266.15 _{±23.56}	0.670 _{±0.143} ^{ab}	1.411 _{±0.240}	5.129 _{±0.581}	6.409 _{±1.171}
YYU13	0.076 _{±0.019}	0.015 _{±0.025} ^c	89.35 _{±8.10}	393.65 _{±240.29}	0.586 _{±0.163} ^{a-c}	1.113 _{±0.552}	4.710 _{±0.281}	4.903 _{±1.617}
YYU14	0.027 _{±0.015}	0.048 _{±0.017} ^{bc}	93.61 _{±11.23}	156.78 _{±111.66}	0.571 _{±0.093} ^{a-d}	1.060 _{±0.435}	3.828 _{±1.659}	3.484 _{±0.930}
YYU15	0.035 _{±0.039}	0.036 _{±0.009} ^c	98.53 _{±4.35}	336.39 _{±123.65}	0.429 _{±0.227} ^{a-f}	0.887 _{±0.627}	5.355 _{±0.091}	3.484 _{±1.466}
YYU18	0.037 _{±0.018}	0.024 _{±0.012} ^c	99.85 _{±66.69}	180.90 _{±80.44}	0.458 _{±0.135} ^{a-f}	1.083 _{±0.318}	5.376 _{±1.624}	7.312 _{±5.099}
YYU20	0.095 _{±0.071}	0.013 _{±0.015} ^c	91.81 _{±39.10}	222.62 _{±132.01}	0.345 _{±0.220} ^{b-f}	1.131 _{±0.530}	4.301 _{±0.878}	5.118 _{±1.496}
YYU21	0.050 _{±0.022}	0.017 _{±0.027} ^c	100.39 _{±16.92}	273.18 _{±162.02}	0.232 _{±0.107} ^{ef}	1.113 _{±0.072}	4.086 _{±0.269}	6.323 _{±2.323}
YYU22	0.014 _{±0.015}	0.026 _{±0.029} ^c	118.53 _{±49.18}	272.22 _{±39.15}	0.440 _{±0.213} ^{a-f}	1.238 _{±0.438}	5.505 _{±2.168}	7.785 _{±7.254}
YYU23	0.052 _{±0.051}	0.040 _{±0.019} ^{bc}	116.63 _{±85.74}	294.33 _{±121.03}	0.369 _{±0.021} ^{b-f}	1.298 _{±0.254}	5.591 _{±1.871}	4.817 _{±2.831}
YYU25	0.053 _{±0.002}	0.104 _{±0.084} ^a	118.31 _{±38.39}	339.11 _{±286.34}	0.268 _{±0.263} ^{c-f}	0.839 _{±0.568}	3.656 _{±0.197}	4.430 _{±0.711}
YYU29	0.042 _{±0.059}	0.047 _{±0.005} ^{bc}	105.19 _{±33.61}	360.50 _{±64.62}	0.152 _{±0.116} ^f	1.262 _{±0.332}	3.613 _{±1.359}	4.860 _{±0.662}
YYU30	0.055 _{±0.048}	0.044 _{±0.004} ^{bc}	129.06 _{±70.78}	398.75 _{±163.67}	0.304 _{±0.000} ^{c-f}	0.702 _{±0.320}	4.344 _{±0.537}	4.473 _{±1.844}
Galia	0.033 _{±0.018}	0.015 _{±0.020} ^c	118.38 _{±37.70}	471.32 _{±347.81}	0.423 _{±0.297} ^{a-f}	0.857 _{±0.295}	3.269 _{±0.649}	3.699 _{±1.171}
Kirkagac	0.045 _{±0.016}	0.086 _{±0.029} ^{ab}	82.94 _{±43.06}	135.11 _{±38.43}	0.232 _{±0.125} ^{ef}	0.774 _{±0.260}	4.645 _{±0.683}	5.677 _{±3.307}
Lokum	0.093 _{±0.127}	0.050 _{±0.030} ^{bc}	101.74 _{±22.80}	230.04 _{±200.48}	0.411 _{±0.054} ^{a-f}	0.792 _{±0.115}	3.441 _{±0.778}	5.204 _{±3.492}
Napolyon	0.040 _{±0.054}	0.022 _{±0.013} ^c	86.56 _{±13.04}	412.12 _{±243.99}	0.244 _{±0.108} ^{d-f}	1.268 _{±0.189}	3.656 _{±2.086}	6.538 _{±3.662}
*p-value	0.472	0.003	0.995	0.606	0.005	0.481	0.074	0.658

I₁₀₀: Full irrigation, I₅₀: 50% deficit irrigation. *: Significant distinctions among groups were observed at the P < 0.05 level, as determined by Duncan's multiple comparison test.

Principle Component (PCA) and Cluster Analysis


Eigenvalues and variances resulting from PCA elucidated the contributions of traits (PCA loads) causing distinctions in deficit irrigation. In the analysis encompassing 28 traits, the first six components with Eigenvalues exceeding 1.00 collectively explicated 77.91% of the total variation (Table 9). PC1 (32.36%) was primarily influenced by APX, SFW, RFW, SDW, 0-5 scale, and Mn content. PC2 (15.39%) was characterized by leaf number, shoot diameter, shoot length, RDW, R/S, and Cu content. PC3 (11.32%) featured stoma length, stoma width, and stoma area as prominent contributors. PC4 (8.77%) revealed the significance of RDM, K, Ca, Mg, and Fe contents. PC5 (5.73%) and PC6 (4.34%) portrayed the importance of MDA, SDM, and P content in the former, while CAT, SOD, stoma density, root length, and Zn content were crucial in the latter (Table 9).

Table 9. PCA loads of the investigated properties in deficit irrigation.

	PC1	PC2	PC3	PC4	PC5	PC6
Eigenvalues	9.06	4.309	3.17	2.46	1.61	1.21
Explained proportion of variation (%)	32.36	15.39	11.32	8.77	5.73	4.34
Cumulative proportion of variation (%)	32.36	47.75	59.07	67.84	73.57	77.91
Factors (Eigen Vectors)						
CAT	-0.105	-0.174	0.012	0.010	0.066	0.403
SOD	0.255	0.055	0.051	0.027	-0.089	-0.314
APX	0.249	0.249	-0.063	0.070	-0.067	-0.091
MDA	0.109	0.116	0.206	-0.128	0.450	-0.048
Stoma length	0.069	-0.075	0.452	-0.235	-0.124	-0.078
Stoma width	0.109	0.029	0.460	0.018	-0.138	0.161
Stoma area	0.097	-0.029	0.507	-0.107	-0.142	0.054
Stoma density	0.095	0.114	-0.202	0.186	0.085	0.567
Leaf number	-0.196	-0.282	-0.006	-0.234	0.122	-0.137
Shoot diameter	-0.216	0.317	0.004	0.005	-0.058	-0.018
Shoot length	-0.147	-0.295	-0.095	-0.234	0.156	-0.095
Root length	-0.170	0.176	0.085	0.106	0.072	-0.212
SFW	-0.287	0.176	-0.016	-0.093	-0.031	-0.106
RFW	-0.267	0.242	0.051	-0.071	0.057	0.034
SDW	-0.262	0.170	0.026	-0.158	0.170	-0.051
RDW	-0.167	0.337	0.179	0.039	0.171	0.062
SDM	0.200	-0.133	0.115	-0.010	0.420	0.141
RDM	0.214	0.022	0.203	0.249	0.142	0.011
R/S	-0.219	0.321	0.096	0.064	-0.055	-0.025
0-5 Scale	0.299	0.152	-0.011	0.006	0.010	-0.085
K	0.085	-0.069	-0.131	0.445	-0.105	-0.261
Ca	-0.163	-0.193	0.138	0.368	0.171	-0.152
Mg	-0.113	-0.095	0.209	0.447	0.237	-0.027
P	-0.153	-0.216	0.093	0.119	-0.352	-0.175
Cu	-0.178	-0.277	0.025	-0.004	0.160	-0.042
Fe	-0.172	0.037	0.018	0.251	-0.079	0.074
Mn	-0.247	-0.103	0.146	0.199	-0.006	0.104
Zn	-0.115	-0.102	0.102	0.000	-0.396	0.335


304 SFW: Shoot fresh weights, RFW: Root fresh weights, SDW: Shoot dry weight, RDW: Root dry weight, SDM: Shoot
305 dry matter, RDM: Root dry matter, R/S: Root-to-shoot ratio (dry weight %).

306
307 A loading plot, derived from the initial two components (PC1 and PC2), elucidated the intricate
308 relationships among the 28 examined traits (Figure 1). A corresponding score plot, integrating PC1
309 and PC2 components, effectively depicted the impact of deficit irrigation (Figure 2). Notably, a
310 clear demarcation was observed between full and deficit irrigation applications, with a propensity
311 for close proximity. Additionally, in the deficit irrigation application, genotypes YYU25 and
312 YYU29 were discernibly positioned in the positive regions of both PC1 and PC2. These findings
313 underscore the nuanced interplay of traits under deficit irrigation conditions, shedding light on the
314 pivotal role of certain genotypes in this context.



316
317
318

Figure 1. PCA Loading plot of study traits using first two principal components.



319
320
321
322
323
324
325

Figure 2. Melon Genotypes/Cultivars under Full and 50% Deficit Irrigation Conditions (-WD Suffix) Mapped on PCA Score Plot.

DISCUSSION

Water scarcity poses a significant challenge to agriculture, impacting crop productivity and yield. Deficit irrigation (DI), a water conservation strategy, influences physiological and yield traits in

326 crop genotypes. The response to DI varies across species and cultivars, emphasizing the importance
327 of understanding this variability for effective water management. Studies on upland cotton
328 (Matniyazova et al., 2022) and muskmelon (Pandey et al., 2018) underscored the significance of
329 genetic variability in drought tolerance traits, suggesting the potential for developing drought-
330 tolerant cultivars. Similarly, the present study on melon revealed varying responses among
331 genotypes and cultivars to deficit irrigation.

332 The observed fluctuations in parameters indicate both positive and negative impacts of stress.
333 Notably, SDM and RDM, stomatal width, stomatal density, K, APX, SOD enzymes, and MDA
334 increased, while other parameters (shoot and root length, stem diameter, leaf number, shoot fresh
335 and drsy weight, root fresh and dry weight, stoma height and area, Mg, Ca, P, Zn, Cu, Fe, Mn and
336 CAT) decreased. Overall, deficit irrigation adversely affected plant growth, aligning with the
337 common response of decreased growth rate and visible stress symptoms (Dasgan et al., 2002;
338 Cakmakci et al., 2022b). Genotypic variations were evident in the reduction of leaves, shoot length,
339 shoot diameter, root length, SDW, RFW, and RDW under stress. The root system, crucial for water
340 uptake, suffered significant decreases, likely impacting water retention and, consequently, plant
341 survival.

342 Root length, a key indicator of drought avoidance, was negatively affected, aligning with studies
343 emphasizing the importance of a long root system in drought tolerance (Serraj et al., 2004). The
344 study implies that decreased root length results from stress-induced damage to cell growth and
345 division. The effects of DI on root growth have been reported, suggesting stimulation of root
346 growth and improved water use efficiency (Costa et al., 2007).

347 A notable finding is the increase in SDM and RDM ratio under stress, indicating better water-
348 holding capacity in more stress-tolerant plants. This aligns with the notion that higher water-
349 holding capacity correlates with better drought tolerance. However, such changes in dry matter
350 ratios may also be indicative of osmotic stress (Kravić et al., 2013). The negative impact of stress
351 on the nutritional status of plants was evident, with decreases in P, Ca, Mg, Zn, Cu, Fe, and Mn
352 uptake, except for a 67% increase in K uptake. Potassium, vital for osmoregulation and enzyme
353 activation, demonstrated an increased role under stress, potentially maintaining osmotic balance.
354 The observed decrease in Ca uptake aligns with reports of limited mobility in the phloem,
355 restricting Ca transport under water limitation (Hessini et al., 2009; Kiegle et al., 2000).

356 Enzyme activities showed a nuanced response, with a 57.14% decrease in CAT activity and
357 increased SOD and APX activities in genotypes under water stress. Antioxidative enzymes, crucial
358 during stress, are stimulated to defend against stress. The elevated MDA content in 76% of
359 genotypes suggests cell membrane damage under stress conditions. One of the most important
360 effects of water shortage stress is the decrease in plant nutritional elements and the other is the
361 damage to the cell walls. Malondialdehyde (MDA), as an indicator of oxidative damage caused by
362 water shortage stress, increased in the leaf tissues of all melon genotypes in the experiment after
363 the stress application (Kırathı et al. 2015). In melon, it was determined that the amount of MDA in
364 the leaves of drought-stressed plants was significantly higher than in control plants (Kusvuran
365 2010). SOD, CAT, GR, APX enzyme activities are stimulated under drought stress conditions
366 (Mohammadkhani and Heidari 2007; Bahadur et al., 2011; Fghire et al., 2013). In this study, the
367 results of which were given, increased enzyme activities such as SOD and APX occurred due to
368 stress factor and injury. It has been determined that these increases are at varying rates. Sources
369 point out that increases in enzyme activation may have an effective role in establishing drought
370 tolerance of genotypes (Kıran et al. 2015). MDA levels have been linked to plant stress responses,
371 varying across species and varieties. The findings align with other studies reporting increased MDA
372 content under drought stress (Sevengor et al., 2011; Sánchez-Rodríguez et al., 2010; Keling et al.,
373 2013), highlighting the variability in stress responses.

374 PCA analysis effectively elucidated stress-induced variations, explaining 47.75% of the total
375 variation. The differentiation among irrigation regimes and mycorrhizal inoculum in melon plants
376 further emphasizes the importance of selecting appropriate parameters for discriminating among
377 treatments. The relationship between vectors, as illustrated in Figure 1, provides insights into the
378 positive correlation within certain growth and physiological parameters. The study concludes by
379 emphasizing the complexity of plant responses to deficit irrigation and the need for tailored
380 approaches to mitigate the impact of water stress on crop productivity.

381 382 **CONCLUSIONS**

383 Global challenges like population growth and environmental issues demand sustainable solutions
384 in agriculture. Deficit irrigation (DI) stands out as an effective water-saving strategy for melon
385 production, but its impact varies among plants. Genetic diversity plays a crucial role in developing
386 drought-tolerant cultivars, a key focus in breeding programs. The study, conducted in the Lake Van

387 Basin, highlights variations in melon genotypes' responses to deficit irrigation. Promising
388 genotypes, selected for traits like seedling development and ion balance, show potential for future
389 breeding programs, enhancing fruit quality and sustainability. In summary, addressing water
390 scarcity requires leveraging genetic diversity and tailored breeding efforts. The identified
391 genotypes offer promising prospects for sustainable agriculture in water-scarce regions like the
392 Lake Van Basin.

393

394 REFERENCES

395 1. Bahadur, A, Chatterjee, A., Kumar, R., Singh, M., Naik, Ps. 2011. Physiological and
396 biochemical basis of drought tolerance in vegetables. *Vegetable Science* 38 (1): 1-16.

397 2. Barzegar T., Heidaryan N., Lofti H., Ghahremani Z. (2018) Yield, fruit quality and
398 physiological responses of melon cv. Khatooni under deficit irrigation. *Advances in Horticultural
399 Science*, 32(4), 451-458.

400 3. Cakmak I., Marschner H. (1992) Magnesium deficiency and high light intensity enhance
401 activities of superoxide dismutase, ascorbate peroxidase, and glutathione reductase in bean leaves.
402 *Plant Physiol*, 98: 1222-1227. [DOI: 10.1104/pp.98.4.1222](https://doi.org/10.1104/pp.98.4.1222)

403 4. Cakmakci O., Cakmakci T., Durak E.D., Demir S., Sensoy S. (2017) Effects of arbuscular
404 mycorrhizal fungi in melon (*Cucumis melo* L.) seedling under deficit irrigation. *Fresenius
405 Environmental Bulletin*, 26(12): 7513-7520.

406 5. Cakmakci T., Cakmakci O., Sahin U. (2022a) The effect of biochar amendment on
407 physiological and biochemical properties and nutrient content of lettuce in saline water irrigation
408 conditions. *Turkish Journal of Agriculture-Food Science and Technology*, 10(12): 2560-2570.
409 [DOI: 10.24925/turjaf.v10i12.2560-2570.5653](https://doi.org/10.24925/turjaf.v10i12.2560-2570.5653)

410 6. Cakmakci Ö., Cakmakci T., Şensoy S. (2022b) Effects of silver nanoparticles on growth
411 parameters of radish (*Raphanus sativus* l. var. *radicula*) grown under deficit irrigation. *Current
412 Trends in Natural Sciences*, 11(21): 37-44. [DOI: 10.47068/ctns.2022.v11i21.004](https://doi.org/10.47068/ctns.2022.v11i21.004)

413 7. Erdinc C., Ekinialp A., Yıldız M., Kabay T., Turkmen O., Sensoy S. (2013) Molecular
414 genetic diversity in Lake Van Basin melons (*Cucumis melo* L.) based on RAPD and ISSR markers.
415 *YYU J. Agr Sci.*, 23(3): 264-270.

416 8. FAO (2019) Food and Agriculture Organization of the United Nations. [Link](#). Accessed: 13
417 November 2022

- 418 9. Farooq M., Wahid A., Kobayashi N., Fujita D., Basra S.M.A. (2009) Plant drought stress:
419 effects mechanisms and management. *Sustainable Agriculture*, 153-188. DOI: [10.1007/978-90-](https://doi.org/10.1007/978-90-481-2666-8_12)
420 [481-2666-8_12](https://doi.org/10.1007/978-90-481-2666-8_12)
- 421 10. Fghire, R., Issaali, O., Anaya, F., Benhabib, O., Jacobsen, S.E., Wahbi, S. 2013. Protective
422 antioxidant enzyme activities are affected by drought in quinoa (*Chenopodium quinoa* Willd).
423 *Journal of Biology, Agriculture and Healthcare* 3 (4): 62-68.
- 424 11. Heath R.L., Packer L. (1968) Photoperoxidation in isolated chloroplasts: I. Kinetics and
425 stoichiometry of fatty acid peroxidation. *Arch Biochem Biophys.*, 125: 189-198. DOI:
426 [10.1016/0003-9861\(68\)90654-1](https://doi.org/10.1016/0003-9861(68)90654-1)
- 427 12. Hessini K., Martínez J.P., Gandour M., Albouchi A., Soltani A., Abdelly C. (2009) Effect
428 of water stress on growth osmotic adjustment cell wall elasticity and water-use efficiency in
429 *Spartina alterniflora*. *Environmental and Experimental Botany*, 67(2): 312-319. DOI:
430 [10.1016/j.envexpbot.2009.06.010](https://doi.org/10.1016/j.envexpbot.2009.06.010)
- 431 13. Jebara S., Jebara M., Limam F., Aouani M.E. (2005) Changes in ascorbate peroxidase
432 catalase guaiacol peroxidase and superoxide dismutase activities in common bean (*Phaseolus*
433 *vulgaris*) nodules under salt stress. *J Plant Physiol.*, 162: 929-936. DOI:
434 [10.1016/j.jplph.2004.10.005](https://doi.org/10.1016/j.jplph.2004.10.005)
- 435 14. Kabay T., Şensoy S. (2016) Enzyme, chlorophyll, and ion changes in some common bean
436 genotypes by drought stress. *YYU J of Agri Sci.*, 26(3): 380-395.
- 437 15. Kacar B., İnal A. (2010) Bitki analizleri. Nobel Yayın Dağıtım.
- 438 16. Kadayifci A., Tuylu G.İ., Ucar Y., Cakmak B. (2005) Crop water use of onion (*Allium cepa*
439 L.) in Turkey. *Agricultural Water Management*, 72(1): 59-68. DOI: [10.1016/j.agwat.2004.08.002](https://doi.org/10.1016/j.agwat.2004.08.002)
- 440 17. Keling H., Ling Z., Jitao W., Yang Y. (2013) Influence of selenium on growth, lipid
441 peroxidation, and antioxidative enzyme activity in melon (*Cucumis melo* L.) seedlings under salt
442 stress. *Acta Soc Bot Pol.*, 82: 193-197. DOI: [10.5586/asbp.2013.023](https://doi.org/10.5586/asbp.2013.023)
- 443 18. Kırmak H., Doğan E. (2017). The effects of deficit irrigation on some quantitative
444 parameters of muskmelon with subsurface and surface drip irrigation systems. *Gaziosmanpaşa*
445 *Üniversitesi Ziraat Fakültesi Dergisi*, 34(Ek Sayı), 80-86.
- 446 19. Kiegle E., Moore C.A., Haselof J., Tester M.A., Knight M.R. (2000) Cell type-specific
447 calcium response to drought salt and cold in *Arabidopsis* root. *The Plant Journal*, 23(2): 267-278.
448 DOI: [10.1046/j.1365-313x.2000.00786.x](https://doi.org/10.1046/j.1365-313x.2000.00786.x)

- 449 20. Kısaca G., Gazioglu Sensoy R.I. (2022) Phenolic contents, organic acids, and antioxidant
450 capacities of twenty grape (*Vitis vinifera* L.) cultivars having different berry colors. *J of Food*
451 *Measurement and Characterization*, 1-17. [DOI: 10.1007/s11694-022-01698-3](https://doi.org/10.1007/s11694-022-01698-3)
- 452 21. Kravić N., Marković K., Anđelković V., Hadži-Tašković Šukalović V., Babić V., Vuletić
453 M. (2013) Growth proline accumulation and peroxidase activity in maize seedlings under osmotic
454 stress. *Acta Physiologiae Plantarum*, 35(1): 233-239. [DOI: 10.1007/s11738-012-1068-x](https://doi.org/10.1007/s11738-012-1068-x)
- 455 22. Kurtar E.S., Balkaya A., Kandemir D. (2016) Screening for salinity tolerance in developed
456 winter squash (*Cucurbita moschata*) lines. *YYU J Agr Sci.*, 26(2): 183-195.
- 457 23. Kuşvuran, Ş. 2010. Kavunlarda Kuraklık ve Tuzluluğa Toleransın Fizyolojik
458 Mekanizmaları Arasındaki Bağlantılar. Çukurova Üniversitesi Fen Bilimleri Enst., Doktora Tezi,
459 355 s., Adana
- 460 24. Kuşvuran Ş., Daşgan H.Y., Abak K. (2011) Farklı Kavun Genotiplerinin Kuraklık Stresine
461 Tepkileri. *YYÜ Tarım Bilimleri Dergisi*, 21 (3): 209-219.
- 462 25. Lamaoui M., Chakhchar A., Kharrassi Y.E., Wahbi S., El Modafar C. (2018)
463 Morphological, Physiological, and Biochemical Responses to Water Stress in Melon (*Cucumis*
464 *melo*) Subjected to Regulated Deficit Irrigation (RDI) and Partial Rootzone Drying (PRD). *Journal*
465 *of Crop Science and Biotechnology*, 21(4), 407-416
- 466 26. Mohammadkhani, N., Heidari, R. 2007. Effects of drought stresses on protective enzyme
467 activities and lipid peroxidation in two maize cultivars. *Pakistan Journal of Biological Sciences* 10
468 (2): 3835-3840.
- 469 27. Pandey S., Ansari W.A., Atri N., Singh B., Gupta S., Bhat K.V. (2018) Standardization of
470 screening technique and evaluation of muskmelon genotypes for drought tolerance. *Plant Genetic*
471 *Resources*, 16(1): 1-8. [DOI: 10.1017/S1479262116000253](https://doi.org/10.1017/S1479262116000253)
- 472 28. Pitrat M. (2008) Melon in vegetables I. Asteraceae. Brassicaceae. and Cucurbitaceae (Ed:
473 Jaime Prohens. Fernando Nuez). Springer Science. Business Media. LLC. p: 283-316.
- 474 29. Sánchez-Rodríguez E., Rubio-Wilhelmi M., Cervilla L.M., Blasco B., Rios J.J., Rosales
475 M.A., Ruiz J.M. (2010) Genotypic differences in some physiological parameters symptomatic for
476 oxidative stress under moderate drought in tomato plants. *Plant Sci.*, 178(1): 30-40. [DOI:](https://doi.org/10.1080/01904169609365104)
477 [10.1080/01904169609365104](https://doi.org/10.1080/01904169609365104)
- 478 30. Sensoy S., Sahin U. (2012) Genetic relationships among various Sıhke melon landraces.
479 *YYU JI of Agri Sci*, 22(3): 147-154.

- 480 31. Sensoy S., Büyükalaca S., Abak K. (2007a) Evaluation of genetic diversity in Turkish
481 melons (*Cucumis melo* L) based on phenotypic characters and RAPD markers. *Genet Resour Crop*
482 *Ev*, 54: 1351-1365. [DOI: 10.1007/s10722-006-9120-6](https://doi.org/10.1007/s10722-006-9120-6)
- 483 32. Sensoy S., Ertek A., Gedik I., Kucukyumuk C. (2007b) Irrigation frequency and amount
484 affect yield and quality of field-grown melon (*Cucumis melo* L). *Agriculture Water Management*,
485 88 (1-3): 269-274. [DOI: 10.1016/j.agwat.2006.10.015](https://doi.org/10.1016/j.agwat.2006.10.015)
- 486 33. Serraj R., Krishnamurthy L., Kashiwagi J., Kumar J., Chandra S., Crouch J.H. (2004)
487 Variation in root traits of chickpea (*Cicer arietinum* L.) grown under terminal drought. *Field Crops*
488 *Research*, 88(2-3): 115-127. [DOI: 10.1016/j.fcr.2003.12.001](https://doi.org/10.1016/j.fcr.2003.12.001)
- 489 34. Sevengor S., Yasar F., Kusvuran S., Ellialtioglu S. (2011) The effect of salt stress on
490 growth. chlorophyll content lipid peroxidation and antioxidative enzymes of pumpkin seedling.
491 *Afr J Agr Res.*, 6: 4920-4924.
- 492 35. Sharma S. P., Leskovar D. I., Crosby K. M., Volder A., Ibrahim A.M.H. (2014) Root
493 growth, yield, and fruit quality responses of *reticulatus* and *inodorus* melons (*Cucumis melo* L.) to
494 deficit subsurface drip irrigation. *Agricultural water management*, 136, 75-85.
- 495 36. Tan Y., Liang Z., Shao H., Du F. (2006) Effect of water deficits on the activity of
496 antioxidative enzymes and osmoregulation among three different genotypes of *Radix Astragali* at
497 seeding stage. *Colloids and surfaces B: Biointerfaces*, 49(1): 60-65. [DOI:](https://doi.org/10.1016/j.colsurfb.2006.02.014)
498 [10.1016/j.colsurfb.2006.02.014](https://doi.org/10.1016/j.colsurfb.2006.02.014)
- 499 37. Turkmen O., Sensoy S., Demir S., Erdinc C. (2008) Effects of two different AMF species
500 on growth and nutrient content of pepper seedlings grown under moderate salt stress. *Afr J of*
501 *Biotechnology*, 7(4): 392-396.
- 502 38. Wang J., Huang G., Li J., Zheng J., Huang Q., Liu H. (2017) Effect of soil moisturebased
503 furrow irrigation scheduling on melon (*Cucumis melo* L.) yield and quality in an arid region of
504 NorthwestChina. *Agric.l Water Manage.* 179, 167–176.
- 505