

Fruit Biochemical and nutritional properties of some Asian and European pears (*Pyrus* spp.) grown under Tehran environmental conditions

Somayeh Kadkhodaei¹, and Kazem Arzani^{1*}

Abstract

Pear is one of the most important pome fruits in the world fruit market with a high nutritional value. This study was performed to determine the phenolic compounds and some chemical properties of the flesh and peel of 12 Asian and European pears. Chlorogenic acid and rutin were found the important phenolic compounds in the peel which were measured using HPLC. Results showed fruit titratable acidity (TA, 0.17-0.53%), total soluble solids (TSS, 13.33-17.33 °Brix), firmness (1.7-2.75, kg/cm²), and color parameters. The highest L* value was observed in KS7 (40.55), while the lowest was in KS12 (14.26) and KS13 (14.78). Additionally, the study assessed the nutrient and total phenol content of fruit samples. The 'Shahmiveh' cultivar displayed the highest total phenol content (638.01 mg/100 g FW), while the KS7 cultivar had the lowest (420.02 mg/100 g FW). Potassium was the most abundant nutrient (1.16 mg/100g DW), followed by nitrogen and calcium contents. As the total phenol increased, so did the amount of rutin. Principal components analysis (PCA) of all data showed that European and Asian pear cultivars were categorized and placed into two distinct groups. In conclusion, the different European and Asian pear studied cultivars and genotypes were different in terms of most of the studied biochemical traits, and significant relationships were observed between some traits. Besides, the obtained results help in the selection of the best pear cultivars or genotypes in terms of the highest phenolic content and nutrients, both for fresh consumption and in the juice industry.

Keywords: Pear. Rutin. Chlorogenic acid. Macronutrients. Micronutrients. Total phenol.

Introduction

Pyrus (*Pyrus* spp.) is the second most important crop following apple in the *Rosaceae* family and can be divided into two major groups of Asian and European type pears (Arzani, 2019; Wang & Arzani, 2019). European species (*Pyrus. communis* L.) have more than 5000 cultivars (Kadkhodaei et al., 2021; Monte-Corvo et al., 2001). Asian pears (*Pyrus pyrifolia*) are mainly cultivated in

¹ Department of Horticultural Science, College of Agriculture, Tarbiat Modares University, Tehran, Islamic Republic of Iran.

* Corresponding author; e-mail: arzani_k@modares.ac.ir

countries such as Korea, China, and Japan in East Asia. They have been cultivated in various parts of Asia for over 3,000 years. Currently, this species is grown commercially in more than 50 countries under temperate climate regions. At least 22 early *Pyrus* species have been identified, all of which are native to Asia, Europe, and the mountainous regions of North America (Bell et al.; 1996). This crop is cultivated mainly due to its commercial value and desirable fruit taste (Arzani, 2002; Wang & Arzani, 2019; Arzani, 2019; Arzani, 2021).

Today, plants can be used to prevent and treat diseases (Jimenez-Garcia et al., 2021; Fattahi et al., 2021). Pears contain phenolic substances and have antioxidant and antimicrobial properties (Jennings et al., 2017; Tiwari et al., 2023). Antioxidant supplements can be mentioned as an important non-pharmacological strategy against oxidative stress (Mota et al., 2022). A self-incompatible, high heterogeneity and allelic diversity have been reported in this genus (Monte-Corvo et al., 2001). In recent decades, numerous studies have been devoted to assessing the genetic diversity in various *Pyrus* species including morphological, biochemical, and DNA markers among European, Asian, and other species. Polyphenols are secondary metabolites (de Paulo Farias et al., 2020). Plant secondary metabolites are a group of chemicals that play a major role in plant growth and survival (Singh et al., 2021; Fattahi et al., 2021). Polyphenols are compounds that occur naturally in fruits and vegetables and are important because of their healing properties and application in technology (de Araújo et al., 2021), and as such, they have received much attention in recent years (de Paulo Farias et al., 2020). Consumption of plants helps the supply of macro and micronutrient elements and reduces the incidence of functional disorders in the body and human health (Li et al., 2017). The role of nutrients in improving the quality of fruit and its other effects- in particular, respiratory failure- causes delay in ripening, increased fruit firmness, and improved fruit storage. Meanwhile, over-consumption of nutrients also impedes the production of quality fruit. Besides, nutrient imbalances cause numerous disorders that affect the quality and performance of pears (Dar et al., 2015; Wang & Arzani, 2019; Arzani, 2019).

Phenolic compounds are chemicals found in most plant tissues of fruits and vegetables. Also, chlorogenic acid (5-Ocaffeoylquinic acid) is a secondary metabolite of phenolic acids and is found in many plants it has an obese anti-obesity mechanism (He et al., 2021). Chlorogenic acid (CGA) is a natural product that has medicinal properties such as anti-cancer, light protection, antioxidant, anti-inflammatory, hypoglycemic, and hypoglycemic effects. After absorption, CGA is further metabolized into sulfate metabolites, glucuronic acid, and glycosides (Sanchez et al., 2017;

Nwafor et al.,2022). Rutin is an important flavonoid also known as vitamin P and quercetin-3-o-rutinoside and has a protective role against liver and gastrointestinal problems (Hosseinzadeh & Nassiri-Asl, 2014), with anti-inflammatory, anti-tumor, antioxidant, and neuroprotective effects (Muvhulawa et al., 2022; Song et al., 2014). Recently, eight phenolic compounds have been identified in the fruit peel of Asian pear (Lee et al., 2011). Pear fruit peel has far higher and more varied phenolic contents than its flesh (Chen et al., 2006). In addition, it is highly recommended that natural products are eaten with their peels because if they reduce oxidation, they will be useful for well-being and disease reduction (Nazir et al., 2020). Previous studies on pears have shown that they contain minerals (Brunetto et al., 2015; Ozturk et al., 2009), and pear fruit is rich in macro and micro-nutrients (Nazir et al., 2020). Mineral nutrients play an important role in plant growth and metabolic functions and are heavily involved in maintaining the health and proper functioning of an organism (Tewari et al., 2021). There are human, plant, and animal diseases associated with micronutrient deficiencies. Also, efforts should be made to produce aggregating microelements of genotypes with an overexpression approach (bio-genetic enhancement) (Izydorczyk et al., 2021). 'Shahmiveh' and 'Sebri' are native commercial European pears cultivars to Iran, in addition A95 promising chance seedling genotype showed superiority in some qualitative fruit characteristics (Wang & Arzani, 2019; Kadkhodaei et al., 2021; Yadegari & Arzani, 2023). The objective of this research was to explore the phenolic and biochemical compounds of 'Shahmiveh', 'Sebri' and A95 promising genotype in compare with some commercial Asian pear cultivars that grown under Tehran, Iran environmental conditions.

Materials and methods

In this experiment, 9 Asian pear cultivars including KS6, KS7, KS8, KS9, KS10, KS11, KS12, KS13, and KS14 (Arzani, 2002), as well as 3 European pear cultivars, 'Shahmiveh', 'Sebri', and A95 promising genotypes (Najafzadeh, 2015; Wang and Arzani, 2019), were used. Trees were planted under Tehran environmental conditions at Tarbiat Modares University (TMU) Asian pear collection orchard with the geographic locations of latitude 35° 41' 39.80" N, longitude 51° 25' 17.44" E. Besides, fruit were harvested at the commercial maturity harvest index (Arzani, 2019) mainly based on the fruit background color, flesh firmness, and total soluble solids (TSS) for further assessments.

Determination of total phenolic content

Fruit samples (fruit with the peel) were freeze-dried for 48 hours and then powdered. For the extraction of phenolic compounds, 2 g. of powdered pulp was used according to the method described previously by Lister et al (1994) with a slight modification. Then, 5 ml of extraction solvent consisting of 85% methanol and 15% acetic acid was added. The samples were placed at 4° C for 24 h and then centrifuged at 10,000 rpm for 10 min. About 1 ml of the supernatant of each sample was filtered using a 0.45 µm syringe filter.

The total phenol content of the extracts was measured by the Folin-ciocalteu method. The absorbance was measured using a spectrophotometer at 765 nm. Total phenol contents were expressed in terms of a milligram of gallic acid content per 100 g fruit fresh weight.

Determination of the phenolic content and components

A water liquid chromatography apparatus consisting of a Separations module: waters 2695 (USA) and a PDA Detector waters 996 (USA) was used for the HPLC analysis. Data acquisition and integration were performed via Millennium32 software. The injection was performed by an autosampler injector equipped with chromatographic assay performed on a 15 cm×4.6 mm with pre-column, Eurospher 100-5 C18 analytical column provided by waters (Sunfire) reversed-phase matrix (3.5 µm) (Waters). The elution was carried out in a gradient system with methanol as the organic phase and distilled water with a flow rate of 1 mL/min. Peaks were monitored at 195-400 nm wavelength. The injection volume was 20 µL and the temperature was maintained at 25°C.

Fruit sample preparation for TSS, TA, pH, firmness, and color

To determine total soluble solids (TSS), a few drops of fruit flesh extract were poured onto the refractometer. For this purpose, Japan's portable refractometer Model 9703 was used (Bexiga et al., 2017). For titratable acidity (TA) and pH of fruit extract, 10 g of smashed fruit flesh was used, with the addition of about 30 ml of distilled water. The extract was centrifuged at 50 °C for 30 min at 4000 rpm. To determine the TA with 0.1 normal solutions, it was titrated to reach pH 8.3 (pH meter Consort- model C860), after which the acidity was calculated based on a milligram of malic acid per 100 g of fruit tissue. To measure the flesh firmness of the fruit tissue, after removing a thin layer of fruit peel, an 8 mm diameter probe by penetrometer (Wagner) was used and fruit

firmness was measured in kg/cm². The pear fruit peel color was measured via the Lutron RGB-1002 color analyzer and converted to L* (lightness), a* (green to red), and, b* (blue to yellow).

Extraction to measure nutrients

Initially, the fruit samples (fruit with the peel) were washed with tap water, followed by 0.1 M hydrochloric acid (HCl), and then rinsed again with distilled water. The sample was dried in an oven at 70 °C and then powdered. The 0.5 mm sieve mesh was used for collecting the clean powdered samples. Extraction steps were performed as follows. Briefly, 2 g of the dried sample was heated in an oven at 550 °C for 4 hours; the ash was slightly moistened with distilled water, and 10 ml of 2 M HCl was added. The final extract was delivered in a volume of 100 ml (Waling et al., 1989). Distillation and sample titration were used to measure the percentage of plant nitrogen (Waling et al., 1989). Phosphorus was measured using a colorimetric (Vanadate-Molybdate) method. The absorbance was measured via a spectrophotometer at 470 nm (Chapman & Pratt, 1962). Potassium and sodium were measured by atomic emission spectrometry.

The absorbance was read by a flame photometer with 766.5 nm for K and 589 nm for N (Waling et al., 1989). The Azomethine colorimetric method was employed to measure the amount of boron and the spectrophotometer device was used at 430 nm (31). An atomic absorption device was utilized to measure the percentage of magnesium (285.2 nm) and calcium (422.7 nm) in the plant (Waling et al., 1989). Measurement of microminerals (iron-manganese-zinc and copper) was performed by atomic absorption spectrometry (A.A.S) method. Measurement of the resulting extraction was carried out by dry burning and the use of HCl. The absorption rate of Fe, Mg, Zn, and Cu was measured at 248.3, 289.5, 213.9, and 324.7 nm, respectively (Elmer & Conn, 1982).

Statistical Analysis

The obtained data were initially checked for normality and analyzed using SAS (Ver. 9.3, SAS Institute, Cary, NC). The results were statistically evaluated by analysis of variance (ANOVA) and expressed as mean ± standard error (SE). Biochemical data for principal component analysis (PCA) and cluster analysis were used. PCA and cluster analysis were performed using Minitab software (Ver. 17). For the Heat map, R 3.5.3 software was used.

Results and discussion

Total Phenol

The highest total phenol amount was observed in the 'Shahmiveh' cultivar and the lowest in KS7 which were 638.01 and 420.02 (mg/100 g FW), respectively. Based on the results, the total phenol amount in different cultivars and the studied genotype showed significant differences (Table 1). The differences in the phenolic composition of different cultivars confirm the genetic role in the synthesis of phenolic compounds since the amount of polyphenols is affected by genotype, rootstock, and climatic conditions (Lin & Harnly, 2008; Manila et al., 2011; Maleki Asayesh et al., 2023). Phenolic compounds may impair callus formation by affecting cell division, development, and differentiation (Bennett & Wallsgrove, 1994). The total phenolic compound in the fruit tissue of the KS13 was higher than KS6 and KS9 Asian pear cultivars (Maghdori et al., 2015). In another study performed on several Australian pear cultivars, the highest amount of total phenol was observed in 'Beurre Bosc' (3.14 ± 0.02 a mg GAE/g) and the lowest in 'Winter Nelis' European pear (1.89 ± 0.03 a mg GAE/g) (Wang et al., 2021).

Phenolic compounds

The minimum amount of chlorogenic acid was observed in 'Sebri' and KS8 which were 3.48 and 3.55 (mg/g FW), respectively, while its maximum was obtained in KS14 and was 9.48 (mg/g FW). The amount of Rutin was between 0.04 (KS8, KS9) and 0.66 (mg/g FW) (A95) in different cultivars (Table 1).

Maghdori et al., 2015 reported a higher amount of chlorogenic acid and catechin in KS6 Asian pear fruit tissues that were grown under Tehran environment conditions. The highest amount of phenolic compounds among the two measured phenolic compounds belonged to chlorogenic acid. It is the most crucial derivative of cinnamic acid in fruits and is known as a disinfectant and radical modifier. These antifungal properties were also evaluated in vitro and the results were satisfying (Martínez et al., 2017).

Pyrus pashia and *Pyrus pyrifolia* are two important sources of chlorogenic acid and rutin (Tiwari et al., 2023). Rutin is a phenolic compound found in other plants, including peaches (Chang et al., 2000). In this study, the amount of rutin in the A95 promising genotype, which is one of the European pears, was higher than in the others. Due to the role of phenolic compounds in human health, cultivars and genotypes with higher amounts of these compounds are important. Also, in

another experiment, several phenolic compounds such as chlorogenic acid and rutin were measured in some popular pear cultivars. The highest and lowest values of chlorogenic acid were 0.69 ± 0.033 mg/g as well as 0.32 ± 0.005 mg/g in 'Graboid' and 'Grabova' cultivars, respectively. Also, the highest and lowest routine values of 0.09 ± 0.001 and 0.01 ± 0.001 were reported in 'Patten' and 'Conference' cultivars (Liaudanskas et al., 2017). It has been reported that the amount of phenolic compounds may vary among Asian and European pears (Lin & Harnly, 2008).

According to Fig 3, for every 484.34 (mg/100 g FW) of total phenol, the amount of Rutin increased by 1 (mg/g FW), which showed a linear relationship.

PH, TSS, TA, color, and firmness

The results showed that the amount of the total soluble solids (TSS) was significant among the studied cultivars and the highest amount was observed in cultivar KS8 (17.33 °Brix) while the lowest was observed in KS10 (13.67 °Brix), KS11 (13.34 °Brix), and KS14 (13.33 °Brix) cultivars (Table 1). TA was 0.53 % in 'Sebri' and 0.17% in the KS13 cultivar. Also, according to the results, the amount of TSS/TA ratio was higher in KS13 (84.95 %) and less in 'Sebri' (29.32%) than in other cultivars (Table 1). Three European pear cultivars had lower pH levels than Asian cultivars. KS13 (5.72) had the highest pH among other cultivars. In general, firmness was higher in Asian pear cultivars than in European ones (Table 1). The highest amount of L^* was in KS7 (40.55) and the lowest was in KS12 (14.26) as well as KS13 (14.78). The a^* value was higher in KS9 (11.65), KS12 (11.70), and KS13 (10.39) than in the other studied cultivars, and b^* was significantly higher in 'Sebri' (29.25) cultivars than in the other cultivars (Table 2).

The acidity concentration and pH extraction were among the studied traits and have a great influence on the aroma, taste, as well as quality of edibility, and fruit storage. It was shown in an experiment that the amount of pH pear fruit is within the range of 3.94 -4.28. Also, the pH in fruit depends on the cultivar and the condition of the planting location (Ozturk et al., 2009). In this experiment, the pH was between 3.9 and 5.72. It has been reported that the titratable acidity (TA) of pear fruit was in the range of 0.5-0.21% in different parts of Turkey, which was consistent with the results of the present research (Ozturk et al., 2009). In another published report, it was mentioned the range of 0.1 to 46% (Chen et al., 2007).

The aroma and taste of fruit are a combination of the amount and type of sugars, organic acids, and aromatic substances. The standard titratable acidity varies depending on the cultivar and

season. Malic acid is the main acid in most pear cultivars at maturity (Colaric et al., 2007). Fruit skin color is one of the most important indicators of determining quality and maturity in pears. Previous experiments have shown a link between fruit ripening, L^* , a^* , and b^* (Kawamura, 2000). Also, L^* , a^* , and b^* are different in pear cultivars (Feng et al., 2023). In addition, fruit tissue firmness is one of the most important traits of quality and physiology that directly affects the texture of the fruit. In many fruits, softening is a programmed process to ripen fruit. Much of this process is a consequence of the chemical alteration of the cell wall, which eventually results in variations in the fruit tissue at the time of maturity (Chen et al., 2006; Ozturk et al., 2009; Wang & Arzani, 2019; Arzani, 2019). Our results showed that pear firmness in the commercial maturity stage was 8.66 to 4.06 (kg/cm²). Overall, Asian pear fruits showed more firmness than European pear cultivars (Arzani, 2019). In the present research, 'Sebri' as one of the European pear cultivars showed higher firmness within the studied Asian and European pear cultivars.

In an experiment, the degree of fruit firmness in different pear genotypes was reported at 3.4 to 8 kg/cm² (Tatari et al., 2020). In this research, we found that by increasing 484.34 mg/100 g FW total Phenol of a pear, and rutin increased by 1 mg/g FW.

Also, by increasing the total phenol by 125.69 mg/100 g FW, one °Brix was increased in fruit TSS. It has been reported that the fruit TSS is an important indicator that is used for the proper time of harvest (Arzani et al., 2008). According to the reported results of the experiments, the total soluble solids (TSS) in different cultivars of pears was 12.5-14 (Bexiga et al., 2017). Besides, TSS is one of the ways to control the quality of fruit, which is important in the grading of fruits in the agricultural industry. Also, the fruit TSS monitoring is a fast and easy as well as cheap record used by the orchardist for considering as one of the good indicators for proper pear fruit harvest (Bexiga et al., 2017; Wang & Arzani 2019; Arzani, 2019).

In research, TSS was measured in 9 pear genotypes and was reported to be 7-13% (Tatari et al., 2020). The ratio of TSS/TA is an indicator of fruit flavor. Natanz and Arbakhøj genotypes had the highest and lowest TSS/TA ratios with averages of 97.54 and 94.11 respectively (Rezaeirad et al., 2013).

Fruit nutrients

Nitrogen content in KS8 and 'Sebri' was 0.44 and 0.5 (mg/100g DW), respectively, and significantly higher than in other cultivars. The KS13 (0.18 mg/100g DW) cultivar had less N than

other cultivars (Table 3). The highest phosphorus levels were observed in KS14 (0.24 mg/100g DW) and the lowest levels were in 'Shahmiveh' as well as the A95 promising genotype (0.14 and 0.16 mg/100g DW). Potassium in KS14 (1.15 mg/100 g DW) and KS9 (1.16 mg/100g DW) was higher than in other cultivars and had the highest potassium among pears cultivars (Table 3). KS8 (0.42 mg/100g DW) showed the higher amount and 'Shahmiveh' (0.04 mg/100g DW) had the minimum amount of calcium. In KS7 (0.15 mg/100 g DW) and KS13 (0.13 mg/100 g DW) Asian pears, the amount of magnesium was higher than in the other studied cultivars (Table 3). In the KS9 (100 mg/kg DW) cultivar, Fe was higher than in the other cultivars. 'Shahmiveh' (29 mg/kg DW) and KS9 (25.67 mg/kg DW) had also less iron than other cultivars. The highest amount of Mn was found in cultivar KS9 (3.46 mg/kg DW). KS12 (11.27 mg/kg DW) had the highest amount of Zn while KS6 (7.26 mg/kg DW) had the lowest amount of Zn compared to other cultivars. The Cu was also higher in KS12 (18.5 mg/kg DW) and lower in KS8 (8.5 mg/kg DW), while the highest B was observed in the KS12 cultivar (96.53 mg/kg DW) (Table 4).

Principal component analysis (PCA) and heat map

PCA showed two groups in the studied pears in this research: Asian pear (*Pyrus serotina* Rehd.) cultivars in one group and European pear (*Pyrus communis* L.) cultivars and A95 promising pear genotype in another group (Fig. 1). The heat map shows a positive (light color) and negative (dark color) relationship between all the measured traits in the study of two group types. There is a positive relationship between total phenol and TSS, firmness, calcium, TA and Na, Ph and TSS/TA, chlorogenic acid, K, and Ca (Fig. 2).

Nutrients play an important role in increasing the quantity, quality, and shelf life and reducing fruit physiological disorders in European as well as Asian pears (Wang & Arzani, 2019; Arzani, 2019). The proper concentration of nitrogen can improve the color, taste, and size of the fruit (Brunetto et al., 2015). Potassium affects fruit size, firmness, color, acidity, and TSS of fruit juice and its aroma. The imbalance in the K to Ca ratio in the plant could cause the cork spot in the 'D Anjou' cultivar (Brunetto et al., 2015; Wang & Arzani, 2019). Many physiological disorders in pear fruits are closely related to calcium deficiency whose prevention requires adequate application of calcium fertilizers (Duan et al., 2019; Wang & Arzani, 2019; Arzani, 2019). In this study, K was found to be the most abundant nutrient (Table 3). Also, previously published research results showed the amount of this element was higher than other nutrients (Chen et al., 2007). As

reported the presence of K under stress enhanced the function of photosystem II, the biosynthesis of chlorophyll, and the antioxidant enzyme activity (Shahid et al., 2019). Besides, Fe deficiency causes disturbances in the photosynthetic system, a decrease in chlorophyll content, and leads to the reduction of crop yield. Also, iron deficiency causes photosynthetic system abnormalities and diminished chlorophyll content (Pestana et al., 2001). The amount of iron in pear fruit is low and has been reported between 20 and 35 mg Fe /kg DW (Brunetto et al., 2015). In our study, the amount of iron varied from 29 to 100 mg/kg DW (Table 4). Manganese deficiency, which is more likely to occur under alkaline conditions, can markedly reduce the pear yield (Brunetto et al., 2015). Zinc deficiency in the soil disrupts the growth of the plant and the fruit set conditions, accordingly, the fruits get miniaturized, and the yield decreases (Wójcik & Popińska, 2009). In another experiment, the amount of zinc among pears was reported from 14 to 27 (mg. kg⁻¹ dry weight) (Arzani et al., 2008). Also, Boron plays an important role in pollen tube growth, pollen germination, fruit size, sourness, and early ripening (Wojcik & Wojcik, 2003).

In this research, we found a linear relationship between total phenol and rutin, total phenol and TSS, firmness, and Ca, and TSS and rutin (Fig. 3). Due to the importance of Ca in the cell wall structure which led to the increase in fruit storage and shelf life, it was observed that for the increase of 5.9249 (kg/cm²) of fruit firmness, one (mg/100g DW) of Ca increased (Fig. 3). It has been reported that (Wang & Arzani, 2019) Ca deficiency in the pear fruit also causes disorders such as cork spot (bitter pit). Also, calcium treatment decreased the peak of ethylene production and decreased respiration rate during fruit storage (Han et al., 2021).

We found in another linear relationship that by increasing one (mg/100 g FW) of total phenol in the pear, 125.69 TSS (°Brix) increased. It was also observed that for every increase of 14.61 TSS, the amount of rutin increased by 1 (mg/g FW) (Fig.3).

Conclusions

In general, the results showed that the amounts of nutrients and biochemistry measurements in this study were different across pear cultivars. Important factors including mineral compounds and sugars affect the quality of pears. Identifying promising genotypes of native pears in the world is important. Recently, A95 as a promising pear genotype has been identified through the breeding program at Tarbiat Modares University, Iran. Also, 'Shahmiveh' and 'Sebri' are native to Iran belonging to the European pear group (*Pyrus communis* L.), and have been compared to Asian

pear cultivars in terms of chemical characteristics. It is necessary to examine indigenous cultivars that also have commercial properties. The genotype of A95 could be one of them. In addition to genetics, factors such as climate, region, and orchard management also affect fruit quality. The highest amount of phenol was found in the 'Shahmiveh' cultivar, which is one of the important native cultivars of Iran. Chlorogenic acid and rutin were relatively high in KS14 and A95. However, in previous experiments, the highest amounts of TA, TSS, and PH were observed in KS8, 'Sebri', and KS13, respectively. Finally, pears can be considered important fruits in terms of minerals and phenolic compounds, and their role in human health is important. The above experiment, which was conducted among 12 European and Asian pear genotypes, will help us choose the best pear genotype in terms of the highest phenolic content and nutrients, both for fresh consumption and in the juice industry.

Acknowledgements

We would like to thank Tarbiat Modares University (TMU) for its financial support. This work was supported by Ph.D. Student Grant Program by TMU. In addition, greenhouse and laboratory facilities provided by Pomology Lab., Department of Horticultural Science at TMU are acknowledged.

References

- Arzani, K. (2002). The position of pear breeding and culture in Iran: Introduction of some Asian pear (*Pyrus serotina* Rehd.) cultivars. *Acta Horticulturae*, 587(October 2002), 167–173. <https://doi.org/10.17660/ActaHortic.2002.587.18>
- Arzani, K., Khoshghalb, H., Malakouti, M.-J., & Barzegar, M. (2008). Postharvest fruit physicochemical changes and properties of Asian (*Pyrus serotina* Rehd.) and European (*Pyrus communis* L.) pear cultivars. *Horticulture Environment and Biotechnology*, 49 (1), 244–252.
- Arzani, K. (2019). Asian Pear. *Postharvest Physiological Disorders in Fruits and Vegetables*. First Edition. CRC Press, Taylor & Francis Group, London, United Kingdom & New York, USA, pp. 329-345.
- Arzani, K. (2021). The National Asian Pear (*Pyrus serotina* Rehd) project in Iran: compatibility and commercial studies of introduced cultivars. *Acta Hortic.* 1315, 91-98 <https://doi.org/10.17660/ActaHortic.2021.1315.14>

- 339 Bell, R. L., Quamme, H. A., Layne, R. E. C., & Skirvin, R. M. (1996). *Pears In: Janick J, Moore*
 340 *JN (eds) Fruit breeding, volume I: tree and tropical fruit*. Wiley, New York.
- 341 Bennett, R. N., & Wallsgrove, R. M. (1994). Secondary metabolites in plant defense mechanisms.
 342 *New Phytologist*, 127(4), 617–633. <https://doi.org/10.1111/j.1469-8137.1994.tb02968.x>
- 343 Bexiga, F., Rodrigues, D., Guerra, R., Brázio, A., Balegas, T., Cavaco, A. M., Antunes, M. D.,
 344 & Valente de Oliveira, J. (2017). A TSS classification study of ‘Rocha’ pear (*Pyrus communis* L.)
 345 based on non-invasive visible/near infra-red reflectance spectra. *Postharvest Biology and*
 346 *Technology*, 132(May), 23–30. <https://doi.org/10.1016/j.postharvbio.2017.05.014>
- 347 Brunetto, G., Melo, Gewellington Bastosde Melo, M., Quartieri, M., & Tagliavini, M. (2015).
 348 The Role of Mineral Nutrition on Yields and Fruit Quality in Grapevine, Pear, and Apple. *Revista*
 349 *Brasileira de Fruticultura*, 37(4), 1089–1104. <https://doi.org/10.1590/0100-2945-103/15>
- 350 Chang, S., Tan, C., Frankel, E. N., & Barrett, D. M. (2000). Low-density lipoprotein antioxidant
 351 activity of phenolic compounds and polyphenol oxidase activity in selected clingstone peach
 352 cultivars. *Journal of Agricultural and Food Chemistry*, 48(2), 147–151.
- 353 Chapman, H. D., & Pratt, P. F. (1962). Methods of analysis for soils, plants, and waters. *Soil*
 354 *Science*, 93(1), 68.
- 355 Chen, J. L., Yan, S., Feng, Z., Xiao, L., & Hu, X. S. (2006). Changes in the volatile compounds
 356 and chemical and physical properties of Yali pear (*Pyrus bertschneideri* Reld) during storage.
 357 *Food Chemistry*, 97(2), 248–255. <https://doi.org/10.1016/j.foodchem.2005.03.044>
- 358 Chen, J., Wang, Z., Wu, J., Wang, Q., & Hu, X. (2007). Chemical compositional characterization
 359 of eight pear cultivars grown in China. *Food Chemistry*, 104(1), 268–275.
 360 <https://doi.org/10.1016/j.foodchem.2006.11.038>
- 361 Chinnici, F., Spinabelli, U., Riponi, C., & Amati, A. (2005). Optimization of the determination
 362 of organic acids and sugars in fruit juices by ion-exclusion liquid chromatography. *Journal of Food*
 363 *Composition and Analysis*, 18(2–3), 121–130. <https://doi.org/10.1016/j.jfca.2004.01.005>
- 364 Colaric, M., Stampar, F., & Hudina, M. (2007). Content levels of various fruit metabolites in the
 365 “Conference” pear response to branch bending. *Scientia Horticulturae*, 113(3), 261–266.
 366 <https://doi.org/10.1016/j.scienta.2007.03.016>
- 367 Dar, M. A., Wani, J. A., Raina, S. K., Bhat, M. Y., & Malik, M. A. (2015). Relationship of leaf
 368 nutrient content with fruit yield and quality of pear. *Journal of Environmental Biology*, 36(3), 649–
 369 653.

- 370 De Araújo, F. F., de Paulo Farias, D., Neri-Numa, I. A., & Pastore, G. M. (2021). Polyphenols
371 and their applications: An approach in food chemistry and innovation potential. *Food Chemistry*,
372 338, 127535. <https://doi.org/10.1016/j.foodchem.2020.127535>
- 373 De Paulo Farias, D., Neri-Numa, I. A., de Araújo, F. F., & Pastore, G. M. (2020). A critical
374 review of some fruit trees from the Myrtaceae family as promising sources for food applications
375 with functional claims. *Food Chemistry*, 306(October 2019), 125630.
376 <https://doi.org/10.1016/j.foodchem.2019.125630>
- 377 Dong, Y., Guan, J. F., Ma, S. J., Liu, L. L., Feng, Y. X., & Cheng, Y. D. (2014). Calcium content
378 and its correlated distribution with skin browning spot in bagged Huangguan pear. *Protoplasma*,
379 252(1), 165–171. <https://doi.org/10.1007/s00709-014-0665-5>
- 380 Duan, Y. X., Xu, Y., Wang, R., & Ma, C. H. (2019). Investigation and prevention of cork spot
381 disorder in ‘Akizuki’ pear (*Pyrus pyrifolia* Nakai). *HortScience*, 54(3), 480–486.
382 <https://doi.org/10.21273/HORTSCI13775-18>
- 383 Elmer, P., & Conn, N. (1982). Analytical methods for atomic absorption spectrophotometry.
384 *Perkin Elmer, Norwalk, CT*.
- 385 Fattahi, B., Arzani, K., Souri, M.K., & Barzegar, M. (2021). Morphological and phytochemical
386 responses to cadmium and lead stress in coriander (*Coriandrum sativum* L.). *Industrial Crops and*
387 *Products* 171(1) : 113979 ISSN 0926-6690, <https://doi.org/10.1016/j.indcrop.2021.113979>
- 388 Feng, Y., Cheng, H., Cheng, Y., Zhao, J., He, J., Li, N., ... & Guan, J. (2023). Chinese Traditional
389 Pear Paste: Physicochemical Properties, Antioxidant Activities, and Quality
390 Evaluation. *Foods*, 12(1), 187.
- 391 Han, S., Liu, H., Han, Y., He, Y., Nan, Y., Qu, W., & Rao, J. (2021). Effects of calcium treatment
392 on malate metabolism and γ -aminobutyric acid (GABA) pathway in postharvest apple fruit. *Food*
393 *Chemistry*, 334, 127479.
- 394 He, X., Zheng, S., Sheng, Y., Miao, T., Xu, J., Xu, W., Huang, K., & Zhao, C. (2021).
395 Chlorogenic acid ameliorates obesity by preventing energy balance shifts in high-fat diet-induced
396 obese mice. *Journal of the Science of Food and Agriculture*, 101(2), 631–637.
397 <https://doi.org/10.1002/jsfa.10675>
- 398 Hosseinzadeh, H., & Nassiri-Asl, M. (2014). Review of the protective effects of rutin on the
399 metabolic function as an important dietary flavonoid. *Journal of Endocrinological Investigation*,
400 37(9), 783–788. <https://doi.org/10.1007/s40618-014-0096-3>

- 401 Izydorczyk, G., Ligas, B., Mikula, K., Witek-Krowiak, A., Moustakas, K., & Chojnacka, K.
 402 (2021). Biofortification of edible plants with selenium and iodine – A systematic literature review.
 403 *Science of the Total Environment*, 754, 141983. <https://doi.org/10.1016/j.scitotenv.2020.141983>
- 404 Jennings, A., MacGregor, A., Spector, T., & Cassidy, A. (2017). Higher dietary flavonoid intakes
 405 are associated with lower objectively measured body composition in women: Evidence from
 406 discordant monozygotic twins. *American Journal of Clinical Nutrition*, 105(3), 626–634.
 407 <https://doi.org/10.3945/ajcn.116.144394>
- 408 Jimenez-Garcia, S. N., Garcia-Mier, L., Vazquez-Cruz, M. A., Ramirez-Gomez, X. S., Guevara-
 409 Gonzalez, R. G., Garcia-Trejo, J. F., & Feregrino-Perez, A. A. (2021). Role of Natural Bio-active
 410 Compounds as Antidiabetic Agents. In *Bioactive Natural Products for Pharmaceutical*
 411 *Applications* (pp. 535–561). Springer.
- 412 Kadkhodaei, S., Arzani, K., Yadollahi, A., Karimzadeh, G., & Abdollahi, H. (2021). Genetic
 413 Diversity and Similarity of Asian and European Pears (*Pyrus* Spp .) Revealed by Genome Size
 414 and Morphological Traits Prediction Genetic Diversity and Similarity of Asian and European Pears
 415 (*Pyrus*. *International Journal of Fruit Science*, 21(1), 619–633.
 416 <https://doi.org/10.1080/15538362.2021.1908201>
- 417 Kawamura, T. (2000). Relationship Between Skin Color and Maturity of Japanese Pear 'Housui'.
 418 *Japanese Journal of Farm Work Research*, 35(1), 33–38.
- 419 Lee, K. H., Cho, J. Y., Lee, H. J., Park, K. Y., Ma, Y. K., Lee, S. H., Cho, J. A., Kim, W. S.,
 420 Park, K. H., & Moon, J. H. (2011). Isolation and identification of phenolic compounds from an
 421 Asian pear (*Pyrus pyrifolia* Nakai) fruit peel. *Food Science and Biotechnology*, 20(6), 1539–1545.
 422 <https://doi.org/10.1007/s10068-011-0213-4>
- 423 Li, X., Liao, W., Yu, H., Liu, M., Yuan, S., Tang, B., Yang, X., Song, Y., Huang, Y., & Cheng,
 424 S. (2017). Combined effects of fruit and vegetable intake and physical activity on the risk of
 425 metabolic syndrome among Chinese adults. *PloS One*, 12(11), e0188533.
- 426 Liaudanskas, M., Zymone, K., Viškelis, J., Klevinskas, A., & Janulis, V. (2017). Determination
 427 of the phenolic composition and antioxidant activity of pear extracts. *Journal of Chemistry*, 2017.
 428 <https://doi.org/10.1155/2017/7856521>
- 429 Lin, L. Z., & Harnly, J. H. (2008). Phenolic compounds and chromatographic profiles of pear
 430 skins (*Pyrus* spp.). *Journal of Agricultural and Food Chemistry* 56: 9094-9101.
- 431 Lister, C. E., Lancaster, J. E., Sutton, K. H., & Walker, J. R. L. (1994). Developmental changes

in the concentration and composition of flavonoids in the skin of a red and a green apple cultivar. *Journal of the Science of Food and Agriculture*, 64(2), 155–161.

Maghdouri, M., Arzani, K. & Bakhshi, D. (2015). Phenolic compounds and antioxidant activity of some Asian pears (*Pyrus serotina* Rehd.) cultivars under Tehran climatic conditions. *Seed & Plant Production Journal* 30(3):315-326 (In Persian).

Mainla, L., Moor, U., Karp, K., & Puessa, T. (2011). The effect of genotype and rootstock on polyphenol composition of selected apple cultivars in Estonia. *Žemdirbystė= Agriculture*, 98(1), 63–70.

Maleki Asayesh, Z., Arzani, K., Mokhtassi-Bidgoli, A. & Abdollahi, H. 2023. Enzymatic and non-enzymatic response of grafted and ungrafted young European pear (*Pyrus communis* L.) trees to drought stress. *Scientia Horticulturae* 310: 111745.

<https://doi.org/10.1016/j.scienta.2022.111745>

Martínez, G., Regente, M., Jacobi, S., Del Rio, M., Pinedo, M., & de la Canal, L. (2017). Chlorogenic acid is a fungicide active against phytopathogenic fungi. *Pesticide Biochemistry and Physiology*, 140, 30–35. <https://doi.org/10.1016/j.pestbp.2017.05.012>.

Mota, J. C., Almeida, P. P., Freitas, M. Q., Stockler-Pinto, M. B., & Guimarães, J. T. (2022). Far from being a simple question: The complexity between in vitro and in vivo responses from nutrients and bioactive compounds with antioxidant potential. *Food Chemistry*, 134351.

Monte-Corvo, L., Goulão, L., & Oliveira, C. (2001). ISSR analysis of cultivars of pear and suitability of molecular markers for clone discrimination. *Journal of the American Society for Horticultural Science*, 126(5), 517–522. <https://doi.org/10.21273/jashs.126.5.517>.

Muvhulawa, N., Dlodla, P. V., Ziqubu, K., Mthembu, S. X., Mthiyane, F., Nkambule, B. B., & Mazibuko-Mbeje, S. E. (2022). Rutin ameliorates inflammation and improves metabolic function: A comprehensive analysis of scientific literature. *Pharmacological Research*, 106163.

Najafzadeh, N., & Arzani, K. 2015. Superior growth characteristics, yield, and fruit quality in promising European pear (*Pyrus communis* L.) chance seedlings in Iran. *J. Agr. Sci. Tech. (JAST)* 17(2): 427-442.

Nazir, N., Nisar, S., Mubarak, S., Khalil, A., Javeed, K., Banerjee, S., Kour, J., & Nayak, G. A. (2020). Pear. In *Antioxidants in Fruits: Properties and Health Benefits* (pp. 435–447). Springer.

Nwafor, E. O., Lu, P., Zhang, Y., Liu, R., Peng, H., Xing, B., ... & Liu, Z. (2022). Chlorogenic acid: Potential source of natural drugs for the therapeutics of fibrosis and cancer. *Translational*

- 463 Oncology, 15(1), 101294.
- 464 Ozturk, I., Ercisli, S., Kalkan, F., & Demir, B. (2009). Some chemical and physicochemical
465 properties of pear cultivars. *Journal of Biotechnology*, 8(4), 687–693.
- 466 Pestana, M., Correia, P. J., de Varennes, A., Abadía, J., & Faria, E. A. (2001). Effectiveness of
467 different foliar iron applications to control iron in orange trees grown on calcareous soil. *Journal*
468 *of Plant Nutrition*, 24(4–5), 613–622. <https://doi.org/10.1081/PLN-100103656>
- 469 Rezaeirad, D., Bakhshi, D., Ghasemnezhad, M., & Lahiji, H. S. (2013). Evaluation of some
470 quantitative and qualitative characteristics of local pears (*Pyrus* sp.) in the north of Iran.
471 *International Journal of Agriculture and Crop Sciences (IJACS)*, 5(8), 882–887.
- 472 Sanchez, M. B., Miranda-Perez, E., Verjan, J. C. G., de los Angeles Fortis Barrera, M., Perez-
473 Ramos, J., & Alarcon-Aguilar, F. J. (2017). The potential of the chlorogenic acid as a multitarget
474 agent: Insulin-secretagogue and PPAR α/γ dual agonist. *Biomedicine and Pharmacotherapy*, 94,
475 169–175. <https://doi.org/10.1016/j.biopha.2017.07.086>
- 476 Shahid, M., Saleem, M. F., Saleem, A., Raza, M. A. S., Kashif, M., Shakoor, A., & Sarwar, M.
477 (2019). Exogenous Potassium–Instigated Biochemical Regulations Confer Terminal Heat
478 Tolerance in Wheat. *Journal of Soil Science and Plant Nutrition*, 19(1), 137–147.
479 <https://doi.org/10.1007/s42729-019-00020-3>
- 480 Singh, A., Chaubey, R., Srivastava, S., Kushwaha, S., & Pandey, R. (2021). Beneficial Root
481 Microbiota: Transmogrifiers of Secondary Metabolism in Plants. In *Emerging Trends in Plant*
482 *Pathology* (pp. 343–365). Springer.
- 483 Song, K., Kim, S., Na, J. Y., Park, J. H., Kim, J. K., Kim, J. H., & Kwon, J. (2014). Rutin
484 attenuates ethanol-induced neurotoxicity in hippocampal neuronal cells by increasing aldehyde
485 dehydrogenase 2. *Food and Chemical Toxicology*, 72, 228–233.
486 <https://doi.org/10.1016/j.fct.2014.07.028>
- 487 Tatari, M., Ghasemi, A., & Mousavi, A. (2020). Diversity of Local and Wild Pear Germplasm in
488 Central Regions of Iran. *International Journal of Fruit Science*, 20(S2), S432–S447.
489 <https://doi.org/10.1080/15538362.2020.1738974>
- 490 Tewari, R. K., Yadav, N., Gupta, R., & Kumar, P. (2021). Oxidative Stress Under Macronutrient
491 Deficiency in Plants. *Journal of Soil Science and Plant Nutrition*, 21(1), 832–859.
492 <https://doi.org/10.1007/s42729-020-00405-9>
- 493 Tiwari, D. C., Bahukhandi, A., Durgapal, M., & Bhatt, I. D. (2023). *Pyrus* spp.(*Pyrus pashia*

Buch.-Ham. ex D. Don, *Pyrus pyrifolia* (Burm. f) Nakai). In *Himalayan Fruits and Berries* (pp. 331-341). Academic Press.

Waling, I., Van Vark, W., Houba, V. J. G., & Van der Lee, J. J. (1989). *Soil and plant analysis, a series of syllabi: Part 7. Plant Analysis Procedures Wageningen Agriculture University*.

Wang, Y., & Arzani, K. (2019). *European Pear. Postharvest Physiological Disorders in Fruits and Vegetables*. First Edition. CRC Press, Taylor & Francis Group, London, United Kingdom & New York, USA, pp. 305-328.

Wang, Z., Barrow, C. J., Dunshea, F. R., & Suleria, H. A. R. (2021). A comparative investigation on phenolic composition, characterization, and antioxidant potentials of five different Australian-grown pear varieties. *Antioxidants*, 10(2), 1–22. <https://doi.org/10.3390/antiox10020151>

Wójcik, P., & Popińska, W. (2009). Response of Lukasovka pear trees to foliar zinc sprays. *Journal of Elementology*, 14(1/2009), 181–188. <https://doi.org/10.5601/jelem.2009.14.1.19>

Wojcik, P., & Wojcik, M. (2003). Effects of boron fertilization on “Conference” pear tree vigor, nutrition, and fruit yield and storability. *Plant and Soil*, 256(2), 413–421. <https://doi.org/10.1023/A:1026126724095>

Yadegari, P., and Arzani, K. (2023). The importance, conservation, maintenance, and propagation of European pear (*Pyrus communis* L.) A95 promising genotype. Thirteen *National Horticultural Science Congress of Iran (IrHC2023)*, September 18-21, Gorgan University of Agricultural Science and Natural Resources, Gorgan, Iran, Congress Proceeding, Pages: 3186-3189.

Table 1. Total Phenole, Chlorogenic acid, Rutin, TSS, TA, and TSS/TA of pear cultivars.

Cultivar	Total Phenol (mg/100 g FW)	Chlorogenic acid (mg/g FW)	Rutin (mg/g FW)	TSS °Brix	TA (%)	TSS/TA
KS6	466.93±9.19efg	7.8±0.0274b	0.08±0.000cde	14.67±0.033cde	0.24±0.003g	60.33±2.166b
KS7	420.02±14.6g	5.56±0.140gh	0.05±0.001fg	14.33±0.033de	0.38±0.015bc	37.82±1.478f
KS8	586.68±32.6ab	3.55±0.177i	0.04±0.000g	17.33±0.033a	0.32±0.011def	54.25±1.212bc
KS9	450.92±3.45fg	7.67±0.328bc	0.04±0.000g	16±0.577ab	0.33±0.003de	48.96±1.360cd
KS10	443.75±16.1fg	7.2±0.208cd	0.07±0.000def	13.67±0.333e	0.23±0.003g	58.57±1.260b
KS11	489.55±7.42def	7.71±0.158bc	0.11±0.006c	13.34±0.333e	0.29±0.005f	46.01±1.441de
KS12	487.35±7.82def	6.54±0.144ef	0.10±0.008cd	15.33±0.333bcd	0.33±0.005de	46.5±1.275cde
KS13	560.75±7.05bc	6.89±0.058de	0.06±0.002efg	14.66±0.333cde	0.17±0.006h	84.95±4.681a
KS14	473.55±8.88efg	9.46±0.088a	0.06±0.003efg	13.33±0.333e	0.33±0.012de	41.43±2.638def
A95	520.46±11.6cd	5.03±0.088h	0.66±0.003a	16.67±0.333ab	0.41±0.008b	40.39±1.618ef
Shahmiveh	638.01±18.2a	5.99±0.106fg	0.47±0.020b	16.33±0.333ab	0.35±0.003de	45.79±0.802de
Sebri	530.40±1.91cd	3.48±0.061i	0.06±0.008efg	15.33±0.333bcd	0.53±0.012a	29.32±0.774g

Values represent the mean ± standard errors (SE). Different letters in the same column indicate significant differences between treatments at $P \leq 0.05$.

Table 2. Fruit color L*, a*, b*, firmness, and pH of the studied pear cultivars.

Cultivar	L*	a*	b*	Firmness (kg/cm ²)	pH
KS6	21.16±0.88g	7.99±0.39bc	20.68±0.99d	5.5±0.288de	5.17±0.088b
KS7	40.55±0.44a	5.75±0.53c	25.97±0.52bc	6.16±0.166d	4.77±0.033bc
KS8	17.00±0.38h	6.36±0.16c	14.26±0.07e	8.66±0.333a	4.63±0.033c
KS9	24.16±0.13f	11.65±1.28a	20.98±0.17d	5.26±0.145e	4.83±0.033bc
KS10	40.32±0.07ab	7.02±0.28bc	26.72±0.49ab	7.4±0.264bc	5±0.057bc
KS11	38.19±0.35c	6.39±0.21c	23.78±0.68c	7.9±0.208abc	5.17±0.033b
KS12	14.26±0.52i	11.70±2.17a	13.10±0.46e	7.33±0.202c	4.58±0.346c
KS13	14.87±0.58i	10.39±0.63a	14.04±0.51e	8.23±0.145ab	5.72±0.044a
KS14	31.67±0.27e	7.59±0.82bc	25.94±0.57bc	7.43±0.296bc	4.92±0.044bc
A95	36.36±0.11d	6.96±0.26bc	26.31±0.27bc	4.06±0.166f	3.9±0.054d
Shahmiveh	38.19±0.62c	7.41±0.59bc	28.28±1.35ab	4.76±0.066ef	4.1±0.057d
Sebri	35.47±0.44d	6.60±0.72c	29.25±0.46a	7.33±0.145c	3.93±0.033d

Values represent the mean ± standard errors (SE). Different letters in the same column indicate significant differences between treatments at $P \leq 0.05$.

Table 3. Fruit macronutrient composition of different pear cultivars (mg/100g DW).

Cultivar	Mg	Ca	K	P	N
KS6	0.06±0.01bc	0.21±0.02bcd	1.13±0.008ab	0.23±0.02ab	0.28±0.008bc
KS7	0.15±0.01a	0.19±0.03bcde	1.07±0.008c	0.19±0.01abc	0.34±0.02b
KS8	0.12±0.02ab	0.42±0.04a	1.13±0.01ab	0.19±0.01abc	0.44±0.03a
KS9	0.06±0.005c	0.22±0.04bc	1.16±0.01a	0.22±0.01ab	0.26±0.02c
KS10	0.06±0.01bc	0.16±0.03bcdef	1.09±0.005bc	0.19±0.02abc	0.29±0.008bc
KS11	0.04±0.008c	0.21±0.05bcd	1.13±0.01ab	0.19±0.01abc	0.24±0.02cd
KS12	0.03±0.007c	0.07±0.01def	1.14±0.008ab	0.21±0.02abc	0.3±0.01bc
KS13	0.13±0.01a	0.25±0.06b	1.13±0.008ab	0.23±0.02ab	0.18±0.008d
KS14	0.04±0.01c	0.06±0.01def	1.15±0.01a	0.24±0.01a	0.30±0.01bc
A95	0.03±0.006c	0.08±0.01cdef	1.06±0.03c	0.16±0.01bc	0.30±0.01bc
Shahmiveh	0.03±0.008c	0.04±0.01f	1.07±0.008c	0.14±0.008c	0.29±0.02bc
Sebri	0.03±0.008c	0.05±0.01ef	1.03±0.01c	0.22±0.02ab	0.5±0.02a

Values represent the mean ± standard errors (SE). Different letters in the same column indicate significant differences between treatments at $P \leq 0.05$.

Table 4. Fruit micronutrient composition of different pear cultivars (mg/kg DW)..

Cultivar	B	Cu	Zn	Fe	Mn
KS6	74.67±1.76cd	12±0.28cd	7.26±0.21f	35.57±1.68g	2.8±0.11b
KS7	56.67±0.88g	17.83±0.72a	10.06±0.47cd	51.77±1.47cd	1.67±0.17c
KS8	78±1.15bc	8.5±0.50f	8.33±0.32ef	43.97±0.84f	2.8±0.20b
KS9	52.7±1.56g	11.67±0.66cd	8.43±0.12e	100±1.53a	3.46±0.27a
KS10	73.4±0.83de	10±0.57def	8.9±0.11e	48.9±1.16def	1.7±0.15c
KS11	73.33±0.88de	16.5±0.28ab	8.13±0.34ef	56±1.53c	1.83±0.12c
KS12	96.53±0.74a	18.5±0.28a	11.27±0.21a	56.33±0.88c	2.26±0.23bc
KS13	68.33±1.36f	15.67±0.88b	11.13±0.13abc	50.66±1.20de	2.2±0.05bc
KS14	70.33±0.88ef	12.43±0.34c	10.5±0.50abcd	46.33±1.45ef	2.23±0.14bc
A95	56.67±0.66g	9.1±0.30ef	11.2±0.26ab	25.67±1.76h	1.7±0.15c
Shahmiveh	78.5±0.76bc	10.33±0.33def	10.13±0.08bcd	29±0.57h	2.3±0.05bc
Sebri	79±0.57b	11±0.57cde	10.03±0.08d	94.67±1.20b	2.2±0.15bc

Values represent the mean ± standard errors (SE). Different letters in the same column indicate significant differences between treatments at $P \leq 0.05$.




Figure 1. Heat map of Spearman's correlations between chemical traits of the studied Asian and European pear cultivars.





Figure 2. Principal component analysis of the studied Asian and European pear cultivars based on chemical traits. Each point represents one genotype and the surrounding green and blue lines are drawn to indicate the division of cultivars and genotypes into two large groups of the Asian and European pears, respectively.

568



569

570 **Figure 3.** Linear regression between total Phenol and rutin, total Phenol and TSS, firmness and
571 Ca, and TSS and rutin of the studied European, and Asian pear genotypes. The Pearson correlation
572 coefficient between total Phenol and rutin was 0.413*, total Phenol and TSS 0.532*, firmness and
573 Ca 0.369*, and TSS and rutin 0.426**. In correlations, * and ** were used to show the significance
574 at $P \leq 0.05$ and $P \leq 0.01$ respectively.