

Evaluation of red currant cultivars' technological effectiveness for mechanized harvesting

Olga Panfilova^{1*}, Ibrahim Kahramanoğlu², Gabrijel Ondrasek³, Olga Golyaeva¹, Volkan Okatan⁴, Sebastián Meier^{5, 6}, Nelli Ryago¹, Mikhail Tsoy¹, Alex Seguel⁷, Serhat Usanmaz², Murat Helvacı², Viktor Kukhar⁸, and Mikhail Karpukhin⁸

ABSTRACT

This study aimed to identify the optimal red currant cultivars for mechanized harvesting by evaluating the bush's morphological structure and the berries' mechanical parameters. Fourteen red currant cultivars were assessed during the 2021-2022 season, and their berry quality indicators and mechanical parameters, including separation force (Fs) and crushing force (Fc), were measured. The results showed that the cultivar plays an important role on the bush's morphological structure, while the berries' mechanical parameters impact the efficiency of harvesting process. The high correlation between Fs and Fc allowed for predicting the optimal harvesting periods of the cultivars. Among the cultivars, 'Rolan', 'Jonkheer Van Tets', 'Rovada', 'Red Lake', 'Asya', 'Vika', and 'Niva' were considered as technological cultivars and suitable for mechanized harvesting. This study emphasizes the importance of considering genetic and morphological factors when selecting red currant cultivars for mechanized harvesting and provides valuable insights for breeding and developing new cultivars adapted to mechanized harvesting.

Keywords: bush habit, optimal harvesting, qualitative characteristics, technology, yield.

INTRODUCTION

Small fruits are highly valued for their biological and nutritional benefits to human's health (Kirina *et al.*, 2020; Kahramanoğlu *et al.*, 2022) and are widely cultivated for their commercial potential (Asănică, 2019; FAO, 2021). Among the berry crops, red currants are particularly valuable due to their numerous biological and pharmacological properties (Bilici, 2021). Many European countries (*i.e.* Poland, Germany, Hungary, United Kingdom, etc.) grow certain red currant cultivars in industrial volumes. One of the most important factors that negatively affect red currant production

¹ Russian Research Institute of Fruit Crop Breeding (VNIISPK), Zhilina, Orel District, Orel Region 302530, Russia.

² Department of Horticulture, Faculty of Agricultural Sciences and Technologies, European University of Lefke, Gemikonagi, via Mersin 10, Ankara 99780, Northern Cyprus, Turkiye.

³ Faculty of Agriculture, University of Zagreb, 10000 Zagreb, Croatia.

⁴ Department of Horticulture, Faculty of Agriculture, Eskisehir Osmangazi University, Eskisehir 26160, Turkiye.

⁵ Agricultural Research Institute. INIA Carillanca, P. O. Box: 929, Temuco, Chile.

⁶ School of Veterinary Medicine, Faculty of Medicine and Health Sciences, Universidad Mayor, Temuco, Chile

⁷ Department of Agricultural Sciences and Natural Resources, Faculty of Agricultural Sciences and Environment, University of La Frontera, Temuco, Chile

⁸ Ural State Agrarian University, ul. Karl Liebknecht, 42, Yekaterinburg 620075, Russia.

*Corresponding author; e-mail: panfilova@orel.vniispk.ru or 302530olga@gmail.com

1 is harvesting costs, which can reach up to 60-70% (Wróblewska *et al.*, 2019; Brondino *et al.*, 2021;
2 FAO, 2021). Many modern cultivars of currants do not meet the requirements for intensive
3 cultivation (Wang *et al.*, 2009; Sarig, 2012; Čejka *et al.*, 2018; Szmajda and Nowakowski, 2020;
4 Perekopskiy *et al.*, 2021). The main parameters of suitability for mechanized harvesting are the
5 architectonics of the bush, high yield, resistance to various stressors, simultaneous ripening of
6 berries in the raceme, and the rapid recovery characteristics of shoots after the combine (Sava and
7 Bodi, 2012; Djordjević *et al.*, 2014; Rakonjac *et al.*, 2015). The physico-mechanical criteria of
8 currant berries have also been developed to reduce the percentage of berry loss during mechanized
9 harvesting (Gurin, 2000). As a result, adjustments are being made to the breeding programs for
10 currants and an important criterion is the mechanization of harvesting (Wang *et al.*, 2009;
11 Rakonjac, 2015; Sasnauskas *et al.*, 2019). One of the main criteria is the placement of berries on
12 bushes, simultaneous ripening of berries, the strength of the skin and the separation of berries from
13 the raceme. Secondary features include the habitus of the bush (height, width and compactness of
14 the bush, flexibility of branches), as well as a long harvest period (Masny *et al.*, 2018).
15 The objective of this study is to evaluate the technological qualities of some cultivars based on the
16 architectonics of the bush and the mechanical properties of the berries. In particular, the research
17 is aimed at identifying promising technological cultivars for inclusion in breeding programs, as
18 well as for industrial cultivation.

19

20 MATERIALS AND METHODS

21 The study was conducted over two seasons (2021-2022), at the growing and testing site of the
22 Russian Research Institute of Fruit Crop Breeding (VNIISPK), located at 52°96' north latitude and
23 36°07' east longitude. The two years-old bare rooted plants were planted in 2017 to evaluate the
24 suitability of cultivars for machine harvesting using a planting scheme of 3.5×0.5 m. The
25 experimental design included 3 replications for each cultivar, with 10 plants in each repetition.
26 The experimental site was on loamy haplic luvisol, and the organic horizon was 50-55 cm thick.
27 The soil of the experimental site was medium acidic (pH=4.82). The content of Potassium and
28 Phosphorus varied depending on the depth of the soil profile and was low for berry crops. The
29 content of Potassium at a depth of 0-0.2 m was 41.75mg·kg⁻¹, Phosphorus - 66.50 mg·kg⁻¹. The
30 content of Potassium at a depth of 0.2-0.4 m was 12.00 mg·kg⁻¹, Phosphorus - 45.00 mg·kg⁻¹.

31 The weather during the growing seasons of 2021 and 2022 was contrasting. While the weather
32 conditions in 2021 were close to the average annual values for the Central region in terms of
33 temperature and precipitation, 2022 was characterized by a 10-12-day delay in spring, and the
34 average temperature in May was 2.5°C lower than usual. The temperature during the summer

1 months (June-August) was also lower by 3 °C compared to the climatic norm for the region. These
2 variations in temperature and precipitation over the years allowed us to calculate the maximum
3 possible duration of the currant harvest.

4 Red currant cultivars of Russian and foreign breeding were selected as the plant materials of the
5 research: 'Red Lake' (USA), 'Lozan' (Slovakia), 'Viksne' (Latvia), 'Jonkheer Van Tets'
6 (Netherlands), 'Rondom' (Netherlands), 'Englische Grosse Weisse' (Netherlands), 'Rovada'
7 (Netherlands), 'Hollandische Rote' (Netherlands), 'Rolan' (Netherlands), 'Natali' (Russia), 'Vika'
8 (Russia), 'Niva' (Russia), 'Osipovskaya' (Russia), and 'Asya' (Russia). Full fruiting of the studied
9 cultivars, when berry harvesting equipment can be used, occurred four years after planting due to
10 the peculiarities of generative bud formation.

11

12 Parameters of the study

13 The morphological features of each cultivar were assessed in triplicate. Biometric indicators of
14 plants such as the height of the bush, the width of the bush, the height of the laying of berries from
15 the ground surface were determined using a technical ruler with an accuracy of 0.1cm. The
16 determination of the volume of the bush was defined as the product of the height of the bush, the
17 width of the bush along and across the row. The compactness of the bush (C) was calculated as
18 the ratio of the height of the bush to the width of the bush across the row.

19 The bush form is compact if the angle between the main fruiting branches and soil surface is 60-
20 75° (coefficient 0.7-0.9) (Sava *et al.*, 2012; Utkov, 2015).

21 The percentage of broken shoots after mechanized harvesting was calculated as the number of
22 broken branches to the total number of shoots of the cultivar on the production plot. Harvesting of
23 berries by the combine harvester was performed at the stage of biological maturity, considering
24 the berries' mechanical characteristics for each cultivar. The weight method determined crop loss
25 from the bush using electronic scales CAS SWN-6 (South Korea). Crop losses refer to undamaged
26 berries left on the bushes after machine harvesting. Repeated mechanized harvesting of the
27 remaining berries on the bushes leads to additional damage to the bush, breakage of annual and
28 perennial shoots, and can result in a prolonged recovery period for the bushes and a significant
29 reduction in yield for the following year. After the combine harvester, the remaining berries on the
30 bushes harvested by hand from each bush, using plastic containers, and weighed on electronic
31 scales in the field. Accounting was conducted on three bushes of the same cultivar, and the average
32 value was calculated. Mechanized harvesting was carried out by the Joonas-2000 combine
33 harvester (Finland), which has no technical design for lifting shoots from the soil.

1 The degree of illumination of the bushes was measured using a luxmeter Light Meter H.S. 1010A
2 (China). The mechanical parameters of the berries were evaluated by separation force (F_s) and
3 crushing force (F_c) during the period of biological ripening of the berries. F_s was determined using
4 the "Dina-2" device (Russia); F_c - using the "Plodtest-1" device (Russia). The measurement
5 interval was 3 days. The measurement was carried out on 5 racemes of each cultivar (an average
6 of 40 berries) in threefold repetition. The strength of the skin (σ) was determined as the F_c ratio to
7 the cross-sectional area of the crushing plunger (Mikhailova, 2014). The strength coefficient (C)
8 of berry depended from the crushing force and crushing separation is given by Equation by
9 Equation (1) (Aleynikov and Mineyev, 2016).

$$10 \quad C = \frac{F_c - F_s}{F_s} \quad (1)$$

11 The cultivar is suitable for mechanized harvesting if $C \geq 0.8$. The multiple regression equation was
12 used to determine the duration of the mechanized harvesting period (Draper and Smith, 1998).
13 Microsoft Excel 2017 was used to calculate the average and standard deviations for the studied
14 indicators and to present the figures. Statistical analysis was carried out using the software package
15 SPSS 22.0 (ANOVA) and Tukey's HSD test ($p < 0.05$). To analyze the similarities and differences
16 between the analyzed groups and build a correlation, the program R 4.2.2 was used; for cluster
17 analysis, the program FactoMineR R was used; the RSA analysis was carried out using the
18 program factoextra R.

19 20 RESULTS

21 The volume of the bush of the studied cultivars varied in the range from 0.37 ('Red Lake') to 5.16
22 ('Osipovskaya') m^3 (Table 1). The optimal area for placing the bush harvest ranges from 0.30 m to
23 1.10 m from the soil surface (Korovin, 2011). The cultivars 'Viksne', 'Niva', and 'Osipovskaya'
24 exhibited an optimal berry placement zone in their large and tall bushes. However, the bush shape
25 of 'Viksne' and 'Osipovskaya' is spreading, which may result in difficulties during mechanized
26 harvesting. On the other hand, 'Niva' has the highest bushes and an erect habit, but the combined
27 harvester can cause damage to the upper part of the shoots. The problem above results in need for
28 additional technical operations such as contour pruning of bushes. This issue is no observed in
29 cultivars such as 'Englische Grosse Weisse', 'Red Lak'e, and 'Lozan', where despite the smaller
30 volume of the bush, the majority of the yield is concentrated in a zone closer to the base of the
31 bush, as shown in Table 1.

32 In this experiment, the morphological characteristics of the bushes were found to be relatively
33 stable with low variation coefficient (CV) values not exceeding 30%: CV (h) was up to 22.98%,

1 CV(B1) up to 19.74%, and CV(V) up to 22.85%. The phenotypic variability of the red currant
2 bush habit was insignificant. In most of the studied cultivars, the percentage of broken branches
3 after the mechanized harvesting did not exceed 10%. A low percentage of damaged branches was
4 not critical for reducing the duration of plantings exploitation. The high flexibility of the
5 'Osipovskaya' branches led to significant damage to the bush. The percentage of broken shoots
6 ranged from 24.32 to 27.39%. The sanitary pruning carried out to remove damaged branches has
7 shown a high regenerative ability of bushes for most cultivars. Nevertheless, 'Rovada' and
8 'Englische Grosse Weisse' demonstrated a weak regenerative ability of the bush, and the number
9 of new annual shoots did not exceed 2-3 pieces per bush for 6 months. The complete restoration
10 of these cultivars took place after 24 months.


Table 1. Morphometric parameters of red currants.

Cultivar	H (m)	B (m)	A (m)	V (m ³)	K	H (m)	B (m)	A (m)	V (m ³)	K	
			'Asya'						'Viksne'		
2021	1.45±0.66 ^{ab}	1.03±0.26 ^{bc}	1.32±0.19 ^b	1.97±0.90 ^c	1.10±0.31 ^{bcd}	1.62±0.12 ^{ab}	1.47±0.50 ^{ab}	1.79±0.07 ^a	5.51±1.82 ^a	0.70±0.07 ^{de}	
2022	1.15±0.11 ^{cd}	1.47±0.80 ^{ab}	1.35±0.76 ^{bc}	2.28±0.88 ^d	0.85±0.23 ^{ab}	1.58±0.06 ^{ab}	1.33±0.38 ^{ab}	2.13±0.14 ^a	4.48±0.83 ^a	0.74±0.02 ^b	
Mean	1.30±0.38 ^{cd}	1.25±0.53 ^c	1.33±0.47 ^{abc}	2.13±0.89 ^c	0.98±0.27 ^{de}	1.60±0.09 ^{ab}	1.40±0.44 ^{ab}	1.80±0.11 ^a	5.00±1.33 ^{ab}	0.72±0.05 ^{de}	
			'Niva'						'Rondom'		
2021	1.72±0.19 ^a	1.57±0.63 ^a	1.40±0.25 ^b	3.78±1.67 ^b	1.23±0.39 ^{bc}	0.98±0.19 ^e	0.73±0.50 ^{fg}	0.75±0.16 ^{fg}	0.54±0.19 ^{ef}	1.31±0.28 ^{ab}	
2022	1.68±0.19 ^a	1.37±0.81 ^{ab}	1.80±0.25 ^{ab}	4.14±2.11 ^a	0.94±0.20 ^{ab}	1.07±0.19 ^{de}	0.77±0.38 ^{de}	0.92±0.26 ^{dc}	0.75±0.02 ^{cd}	1.16±0.18 ^a	
Mean	1.70±0.19 ^a	1.47±0.72 ^a	1.60±0.25 ^b	3.96±1.89 ^b	1.08±0.30 ^{de}	1.03±0.19 ^{df}	0.75±0.44 ^{de}	0.84±0.21 ^{fg}	0.65±0.11 ^{fg}	1.23±0.23 ^a	
			'Vika'					'Englische Grosse Weisse'			
2021	1.37±0.14 ^{bc}	0.88±0.18 ^{de}	1.03±0.29 ^{cd}	1.24±0.58 ^{de}	1.32±0.39 ^{ab}	1.28±0.19 ^{cd}	0.89±0.05 ^{fg}	0.92±0.07 ^{fg}	1.05±0.24 ^{de}	1.40±0.22 ^{ab}	
2022	1.35±0.37 ^{bc}	1.10±0.35 ^{de}	1.33±0.29 ^{bc}	1.98±0.39 ^{cd}	1.01±0.40 ^{ab}	0.86±0.36 ^e	0.65±0.12 ^{de}	0.92±0.26 ^{dc}	0.51±0.11 ^{cd}	0.94±0.30 ^{ab}	
Mean	1.36±0.25 ^{bc}	0.99±0.26 ^{de}	1.18±0.29 ^{de}	1.61±0.49 ^{de}	1.17±0.40 ^{dc}	1.07±0.28 ^{df}	0.77±0.08 ^{de}	0.92±0.17 ^{fg}	0.78±0.18 ^{fg}	1.17±0.26 ^{bc}	
			'Lozan'					'Red Lake'			
2021	1.02±0.19 ^{de}	0.60±0.04 ^{fg}	0.64±0.13 ^g	0.39±0.07 ^{ef}	1.58±0.51 ^a	0.91±0.07 ^e	0.50±0.38 ^g	0.70±0.24 ^{fg}	0.32±0.10 ^f	1.29±0.35 ^{ab}	
2022	0.95±0.22 ^{de}	0.75±0.12 ^{de}	0.95±0.33 ^{dc}	0.68±0.18 ^{cd}	1.00±0.13 ^{ab}	0.98±0.10 ^{de}	0.58±0.07 ^e	0.72±0.26 ^d	0.41±0.11 ^d	1.36±0.37 ^a	
Mean	0.98±0.20 ^f	0.67±0.08 ^{de}	0.80±0.23 ^g	0.53±0.13 ^g	1.29±0.32 ^a	0.94±0.09 ^f	0.54±0.23 ^e	0.71±0.25 ^g	0.37±0.11 ^g	1.33±0.36 ^f	
			'Rovada'					'Rolan'			
2021	1.02±0.21 ^{de}	0.79±0.31 ^{de}	0.93±0.14 ^{ef}	0.75±0.15 ^{ef}	1.09±0.37 ^{bcd}	1.37±0.01 ^{bc}	0.75±0.39 ^{df}	1.03±0.16 ^{de}	1.06±0.28 ^{de}	1.34±0.21 ^{ab}	
2022	0.97±0.26 ^{de}	1.18±0.19 ^{cd}	1.10±0.12 ^{dc}	1.26±0.42 ^{cd}	0.88±0.03 ^{ab}	1.13±0.19 ^{cd}	0.87±0.19 ^{de}	1.03±0.52 ^{dc}	1.01±0.36 ^{cd}	1.10±0.12 ^{ab}	
Mean	0.99±0.23 ^f	0.99±0.25 ^{de}	1.02±0.13 ^{fg}	1.00±0.29 ^{ef}	0.99±0.20 ^{de}	1.25±0.10 ^{cd}	0.81±0.29 ^{de}	1.03±0.34 ^{fg}	1.04±0.32 ^{ef}	1.22±0.17 ^{ab}	
			'Natali'					'Hollandische Rote'			
2021	1.05±0.17 ^{de}	1.20±0.25 ^{ab}	1.23±0.35 ^{bc}	1.55±0.76 ^{de}	0.85±0.30 ^{de}	1.07±0.09 ^{de}	0.83±0.28 ^{de}	0.77±0.15 ^{fg}	0.69±0.26 ^{ef}	1.39±0.37 ^{ab}	
2022	1.02±0.07 ^{de}	1.43±0.34 ^{ab}	1.33±0.17 ^{bc}	1.94±0.72 ^{cd}	0.76±0.16 ^b	1.05±0.22 ^{de}	0.93±0.14 ^{de}	1.27±0.14 ^{dc}	1.24±0.16 ^{cd}	0.83±0.27 ^b	
Mean	1.03±0.12 ^{df}	1.32±0.29 ^{bc}	1.28±0.35 ^{de}	1.75±0.74 ^{de}	0.81±0.23 ^{de}	1.06±0.16 ^{df}	0.88±0.21 ^{de}	1.02±0.15 ^{ad}	0.96±0.21 ^{fg}	1.11±0.32 ^{bc}	
			'Jonkheer Van Tets'					'Osipovskaya'			
2021	1.19±0.08 ^{de}	1.02±0.19 ^{bc}	1.41±0.03 ^b	1.70±0.18 ^{cd}	0.84±0.05 ^{de}	1.66±0.15 ^a	1.20±0.43 ^a	2.43±0.29 ^a	4.85±0.91 ^{ab}	0.68±0.07 ^e	
2022	1.08±0.07 ^{de}	1.38±0.19 ^{ab}	1.48±0.19 ^{bc}	2.22±0.49 ^{bc}	0.73±0.07 ^b	1.50±0.25 ^{ab}	1.63±0.29 ^a	2.23±0.52 ^a	5.47±1.41 ^a	0.67±0.06 ^b	
Mean	1.14±0.07 ^{df}	1.20±0.19 ^{cd}	1.45±0.11 ^{bc}	1.96±0.34 ^{cd}	0.79±0.06 ^{de}	1.58±0.20 ^{ab}	1.42±0.36 ^{bc}	2.33±0.41 ^a	5.16±1.16 ^a	0.68±0.07 ^e	

Note: H – the height of the bush (m); A – length of the bush diagonally along the row (m); B – width of the bush across the row (m); V – volume of the bush (m³); and K – compactness of the bush. The data represent the average values for three repetitions (n = 10) ± standard error (S.E.). Different letters show significant differences between the parameters of the bush according to Tukey's HSD test (p < 0.05).

1 One of the critical factors determining the suitability of a cultivar for combine harvesting is its
 2 ability to produce a stable yield with a minimum density of 2-3 kg of the bush. To assess the
 3 performance of various cultivars under mechanized harvesting, we measured their yield and crop
 4 losses, as shown in Figure 1. It should be noted that the reduction in yield during the second year
 5 of the experiment can be attributed to damage caused by the combine harvester to some of the
 6 perennial shoots, which are responsible for producing generative buds.

7



8

9 **Figure 1.** The total yield of cultivars and yield loss after the mechanized harvesting. The letters
 10 above the columns are used to compare yield data and its loss for each cultivar. Different letters
 11 indicate significant differences between the values according to Tukey's HSD test ($p < 0.05$).

12

13 The berries on the bushes should be positioned in such a way as to ensure maximum efficiency of
 14 berry harvesting by a combine harvester and the berries should have a simultaneous ripening
 15 period. The simultaneity of maturation is determined by varietal differences and bush lighting. The
 16 sparse architectonics of the bush in 'Jonkheer Van Tets', 'Asya', 'Vika', 'Natali', 'Red Lake' and
 17 'Rovada' provided uniform illumination of berries on the bushes (687.2 Lx). By the beginning of
 18 the mechanized harvesting, more than 90% of the berries at the stage of biological maturation had
 19 a uniform color and met the quality standard of berry products. In 'Osipovskaya', the strong
 20 thickening of the bushes reduced the intake of sunlight (425 Lx). During mechanized harvesting,
 21 about 30% of the yield had unripe berries, which were poorly separated from the raceme and a
 22 large number of berries remained on the bushes. The mechanical parameters of the berries depend
 23 on the cultivar and weather and climatic conditions. Using the example of several red currant
 24 cultivars, changes in the F_s and F_c parameters and the duration of the biological maturation period
 25 in 2021 and 2022 are shown (Table 2).

1 The mechanical parameters of currant berries varied over two growing seasons. In 2022, the
 2 ripening period of the berries for many cultivars was quite long and began 7-10 days later than in
 3 2021. In addition, by the period of full biological maturity, the Fs and Fc indicators decreased.
 4 The duration of the optimal berry harvest period for cultivars depended on the rate of decrease in
 5 the mechanical parameters of the berries. The dependence of the harvest duration on Fs and Fc and
 6 weather conditions is shown (Table 2).

7 **Table 2.** Changes in the mechanical parameters of berries during biological maturation, depending
 8 on the cultivar and the study period.

Days	2021		2022	
	Fs, N	Fc, N	Fs, N	Fc, N
'Jonkheer Van Tets'				
1	1.47±0.12 ^a	6.04±0.80 ^a	0.65±0.25 ^a	5.57±0.4 ^{9a}
4	0.89±0.16 ^b	4.71±0.85 ^b	0.69±0.07 ^a	4.86±0.46 ^b
7	0.78±0.09 ^c	6.35±0.18 ^c	0.68±0.10 ^a	4.75±0.69 ^b
10	0.62±0.10 ^d	4.78±0.90 ^d	0.42±0.06 ^d	4.38±0.84 ^c
13	0.33±0.08 ^e	3.43±0.19 ^e		
'Englische Grosse Weisse'				
1	1.53±0.06 ^a	3.98±0.97 ^a	0.51±0.03 ^a	3.19±0.72 ^a
4	0.49±0.02 ^b	1.64±0.45 ^b	0.47±0.07 ^a	4.02±0.89 ^b
7			0.54±0.03 ^a	3.21±0.21 ^b
10			0.25±0.08 ^d	1.59±0.38 ^d
'Lozan'				
1	1.33±0.08 ^a	9.06±0.53 ^a	0.83±0.19 ^a	3.39±0.18 ^a
4	1.25±0.08 ^a	6.19±0.63 ^b	0.60±0.09 ^a	4.41±0.50 ^b
7	0.14±0.04 ^c	4.69±0.50 ^c	0.56±0.05 ^b	3.74±0.96 ^b
10			0.63±0.08 ^b	2.63±0.14 ^d
13			0.26±0.04 ^e	2.34±0.03 ^e
Red Lake				
1	0.80±0.08 ^a	8.59±0.33 ^a	0.52±0.12 ^a	3.37±0.36 ^a
4	0.69±0.14 ^b	4.57±0.46 ^b	0.46±0.05 ^a	3.24±0.23 ^a
7	0.63±0.12 ^b	6.03±0.41 ^c	0.50±0.08 ^a	3.14±0.80 ^a
10	0.36±0.04 ^d	4.09±0.28 ^d	0.26±0.02 ^c	2.35±0.41 ^b
13	0.29±0.02 ^e	3.52±0.47 ^c		
Vika				
1	1.44±0.04 ^a	6.57±0.65 ^a	0.62±0.09 ^a	4.10±0.21 ^a
4	0.55±0.07 ^b	5.02±0.46 ^b	0.60±0.07 ^a	5.02±0.58 ^b
7	0.51±0.08 ^b	5.96±0.48 ^c	0.42±0.02 ^c	4.19±0.09 ^a
10	0.25±0.02 ^d	4.02±0.43 ^d	0.32±0.08 ^d	2.87±0.46 ^d

9 Note: «-»- berries at the stage of physiological maturity; biosynthetic processes stop; berries lose their characteristic
 10 taste and appearance.


11
 12 The high relationship between the mechanical parameters of berries ($R=0.75-0.85$) makes it
 13 possible to calculate the prediction of the optimal period for berry harvesting, taking into account
 14 the cultivar. For 'Jonkheer Van Tets', according to the multiple regression equation, the effective
 15 period for using the harvester was 7 days in 2021 ($R=0.69$) and 4 days in 2022 ($R=0.71$).

16 Fc depends on the strength of the skin and for most cultivars it was more than 2 N. In the process
 17 of biological maturation, the strength of the skin decreased in all cultivars. Reducing the strength
 18 of the skin during the ripening period of berries reduces the resistance of berries to mechanical

1 damage and the quality characteristics of berries. Changes in the strength of the berry skin during
 2 maturation are associated with biochemical changes (for example, hydrolysis of protopectin,
 3 hemicellulose) (Giongo, L., 2019; Tousehik *et al.*, 2017; Spinei and Oroian, 2021). Of particular
 4 importance in mechanized harvesting is the separation of berries from the raceme without damage.
 5 This separation is called "dry berry separation". This separation allows to maintain the appearance
 6 and quality characteristics of berries at a high level during transportation (Kazakhmedov *et al.*,
 7 2017; Rivera, 2022). The strength of the attachment of berries to the raceme depends on F_s and
 8 F_c. According to the criterion of A.F. Aleynikov and V.V. Mineev (2016), most of the studied
 9 cultivars are considered suitable for mechanized harvesting ($C \geq 0.8$).

10 Relationships among study cultivars

12 Distance measures showed that 'Niva', 'Asya' and 'Jonkheer Van Tets' were similar in many studied
 13 indicators and differed from 'Rondom'. At the same time, 'Rondom', 'Rolan' and 'Vika' were
 14 different from 'Red Lake'. 'Viksne' and 'Osipovskaya' were similar in a number of indicators
 15 (Figure 3A). Cluster analysis combined the cultivars according to the studied parameters into 4
 16 clusters (Figure 3B).



17
 18 A




Figure 3. A. Distance measure (orange color shows differences, teal – how’s similarities) between cultivars; B. Hierarchical clustering to group the cultivars.

The first cluster included 'E. Grosse Weisse'. According to PCA-Biplot analysis, this cultivar was very different from all the studied cultivars (Fig. 4). The second cluster combined cultivars in terms of bush compactness (K); the third cluster consisted of cultivars with high morphological parameters of the bush (A, B, and V) and average K values; the fourth cluster combined cultivars in terms of mechanical parameters of berries (Fc, Fs, σ , and C).




Figure 4. Correlation (A) and PCA - biplot (B) analyses of characters and cultivars. Note: K - compactness of the bush, H - the height of the bush (m), B - width of the bush across the row (m), A - length of the bush diagonally along the row (m), and V - the volume of the bush (m³), Fs - separation force, Fc -crushing force, C - coefficient of the relative strength of berries, and σ - static crushing force.

1 DISCUSSION

2 In this experiment, the Joonas-2000 combine harvester (Finland) was used, its work is based on
3 the vibration effect on shoots. Therefore, the habitus of the bush is important. For the normal
4 operation of the harvester, the bushes should be either erect or slightly spreading and consist of
5 10-15 perennial fruiting shoots (Pluta *et al.*, 2008; Sava and Bodiu, 2012). For this trait, 'Viksne',
6 'Englische Grosse Weisse', 'Osipovskaya' (Table 1) did not meet these requirements and should
7 not be included in currant breeding programs for suitability for mechanized harvesting, as well as
8 for industrial cultivation. In experiments with blueberry cultivars, it has also been shown that
9 certain parameters of plant height and bush volume are necessary for the use of berry harvesting
10 equipment. The minimum yield losses were in blueberry varieties with a straight-growing,
11 compact bush shape, which also made it possible to extend the life of these plantations (Patrick,
12 and Li, 2017). The second disadvantage of the combine harvester of this model is the collection of
13 ripe and unripe berries during vibration on shoots. This result was observed in this experiment
14 with 'Osipovskaya' red currant and when harvesting sea buckthorn berries (Khabarov, 2014;
15 Zubarev., 2022). When harvesting berries in bulk, it is important to determine the effective period
16 of using the harvester, when berry losses will be minimal and the quality of the berries will meet
17 the quality standard (Yu, 2012; Brondino, 2021). The dependence of the effective harvest period
18 on climatic conditions and the cultivar has been proven in this experiment (Table 2), as well as on
19 other cultivars of red and white currants (Papstein *et al.*, 2016) and grape cultivars (Jobbágy,
20 2021). This experiment shows the use of mechanical parameters of berries as an informative
21 feature for assessing the quality of berry products (Table 2). For blueberry cultivars, the possibility
22 of using various upgraded combine harvester models is determined by the strength of the berries
23 (Rivera, 2022.). However, it is not entirely correct to recommend using only the mechanical
24 parameters of berries to predict the optimal period of using harvester, since it is also necessary to
25 take into account long-term weather data and biometric characteristics of the bush. A similar
26 conclusion was made by a comprehensive assessment of the mechanical parameters of the berry
27 skin, the morphology of the blueberry bush (Rivera, 2022) and grapes (Zouid, 2010; Brillante,
28 2015). The current study confirmed the relationship between high F_c values of red currant berries
29 and product quality. These indicators tended to decrease during the maturation process. A decrease
30 in the strength of the skin of berries and the quality of products in the process of biological ripeness
31 were noted in grapes (Rolle *et al.*, 2012). In addition, the current experiment did not reveal a
32 relationship between F_s and the strength of the berry skin, which is consistent with the data
33 obtained on grape cultivars (Giacosa *et al.*, 2013)

34

1 **CONCLUSION**

2 The suitability of red currant cultivars for mechanized harvesting should be assessed by the
3 productivity of a berry harvester. In this study, the cultivars were rejected according to indicators
4 that determine the profitability of berry production, including a high percentage of branch breakage
5 during two years of using the harvester, low regenerative ability of the bush after sanitary pruning,
6 as well as low mechanical parameters of berries. According to the architectonics of the bush,
7 mechanical parameters of berries and testing of the harvester, 'Vika', 'Asya', 'Red Lake', 'Niva',
8 'Rolan', and 'Jonkheer Van Tets' are technological cultivars suitable for mechanized harvesting and
9 promising for industrial cultivation and breeding programs.

10
11 **REFERENCES**

- 12 1. Aleynikov, A. F. and Mineyev, V. V. 2016. Measurement of mechanical properties of sea
13 buckthorn and black currant berries. *Sib. Herald of Agr. Sci.*, **4**: 105-111.
- 14 2. Asănică, A. 2019. Growing berries in containers - a new perspective for urban horticulture.
15 *Sci. Papers-Ser. B, Hort.*, **63(1)**: 97-102.
- 16 3. Brillante, L., Tomasi, D., Gaiotti, F., Giacosa, S., Torchio, F., Segade, S. R., and Rolle, L.
17 2015. Relationships between skin flavonoid content and berry physical-mechanical
18 properties in four red wine grape cultivars (*Vitis vinifera* L.). *Sci.Hort.*, **197**: 272-279.
- 19 4. Brondino, L., Borra, D. and Giuggioli, N. R. 2021. Massaglia, S. Mechanized Blueberry
20 Harvesting: Preliminary Results in the Italian Context. *Agriculture*, **11(12)**: 1197.
- 21 5. Brondino, L., Borra, D., Giuggioli, N. R., and Massaglia, S. 2021. Mechanized blueberry
22 harvesting: preliminary results in the Italian context. *Agriculture*, **11(12)**: 1197.
- 23 6. Čejka, B., Matějčiček, A., Matějčičková, J. and Paprštein, F. 2013. Red currant cultivar
24 'Rubigo'. *Vědecké Práce Ovocnářské*, **23**: 79-86.
- 25 7. Djordjević, B., Rakonjac, V., Akšić, M. F., Šavikin, K. and Vulić, T. 2014. Pomological
26 and biochemical characterization of European currant berry (*Ribes* sp.) cultivars. *Sci. Hort.*,
27 **165**: 156-162.
- 28 8. Draper, N. R. and Smith, H. 1998. Selecting the "best" regression equation, in: chap. 15 in
29 Applied Regression Analysis, 3rd Edn., John Wiley & Sons, 327-368.
- 30 9. **FAO**, 2021. Food and Agriculture Organization of the United Nations, 2021. Available
31 online: <https://www.fao.org/sustainable-development-goals/background/ru/>
- 32 10. Giacosa, S., Torchio, F., Segade, S.R., Gaiotti, F., Tomasi, D., Lovat, L., Vincenzi, S. and
33 Rolle L. 2013. Physico- mechanical evaluation of the aptitude of berries of red wine grape

- 1 varieties to resist the compression in carbonic maceration vinification. *Int. J. of food sci. &*
2 *tech.*, **48(4)**: 817-825.
- 3 11. Giongo, L., Ajelli, M., Poncetta, P., Ramos-García, M., Sambo, P., and Farneti, B. 2019.
4 Raspberry texture mechanical profiling during fruit ripening and storage. *Posth. Biol. and*
5 *Tech.*, **149**: 177-186.
- 6 12. Gurin, A.G. 2000. Prediction of the duration of mechanized harvesting of blackcurrant.
7 *Hort. and viticulture*, **3**: 13-15.
- 8 13. Jobbágy, J., Dočkalík, M., Krištof, K., and Burg, P. 2021. Mechanized Grape Harvest
9 Efficiency. *Appl. Sci.*, **11(10)**: 4621.
- 10 14. Kahramanoğlu, İ., Panfilova, O., Kesimci, T. G., Bozhüyük, A. U., Gürbüz, R. and
11 Alptekin, H., 2022. Control of Postharvest Gray Mold at Strawberry Fruits Caused by
12 *Botrytis cinerea* and Improving Fruit Storability through *Origanum onites* L. and *Ziziphora*
13 *clinopodioides* L. Volatile Essential Oils. *Agronomy*, **12(2)**: 389.
- 14 15. Kirina, I. B., Belosokhov, F. G., Titova, L. V., Suraykina, I. A. and Pulpitow, V. F. 2020.
15 Biochemical assessment of berry crops as a source of production of functional food
16 products. *IOP Conf. Ser.: Earth Environ. Sci.*, **548(8)**: 082068.
- 17 16. Korovin, K. L. 2011. Evaluation of black currant varieties and hybrids on some parameters
18 of suitability to mechanized fruit harvesting. *Fruit Grow.: sci. papers*, **23**: 198-202.
- 19 17. Masny, A., Pluta, S. and Seliga, Ł. 2018. Breeding value of selected blackcurrant (*Ribes*
20 *nigrum* L.) genotypes for early-age fruit yield and its quality. *Euphytica*, **214(6)**: 1-21.
- 21 18. Paprstein, F., Sedlak, J. and Kaplan, J. 2016. Rescue of red and white currant germplasm
22 in the Czech Republic. *Acta Hort.*, **1133**: 49-52.
- 23 19. Patrick, A., and Li, C. 2017. High throughput phenotyping of blueberry bush
24 morphological traits using unmanned Aerial systems. *Remote Sensing*, **9 (12)**: 1250.
- 25 20. Perekopskiy, A. N., Zыkov, A. V. and Egorova, K.I. 2021. Evaluation of black currant
26 varieties suitability for combine harvesting. *The Agr. Sci. J.* **7**: 35–39.
- 27 21. Pluta, S., Zurawicz, E., Krawiec, A. and Salamon, Z. 2008. Evaluation of the suitability of
28 polish blackcurrant cultivars for commercial cultivation. *J. of Fruit and Orn. Plant Res.*,
29 **16**: 153-166
- 30 22. Rakonjac, V., Djordjević, B., Fotirić-akšić, M., Vulić, T. and Djurović, D. 2015.
31 Estimation of variation and correlation analysis for yield components in black currant
32 cultivars. *Genetika*, **(3)**: 785-794.

- 1 23. Rivera, S. 2022. Identification of mechanical parameters to be used as a firmness standard
2 on quality evaluations of stored blueberry. Thesis of Doctor of Philosophy in Plant Science.
3 Manawatu, New Zealand.
- 4 24. Rivera, S., Giongo, L., Cappai, F., Kerckhoffs, H., Sofkova-Bobcheva, S., Hutchins, D.,
5 and East, A. 2022. Blueberry firmness-A review of the textural and mechanical properties
6 used in quality evaluations. *Posth. Biol. and Tech.*, **192**: 112016.
- 7 25. Rivera, S., Kerckhoffs, H., Sofkova-Bobcheva, S., Hutchins, D., and East, A. 2022.
8 Influence of harvest maturity and storage technology on mechanical properties of
9 blueberries. *Posth. Biol. and Tech.*, **191**: 111961
- 10 26. Rolle, L., Torchio, F., Ferrandino, A. and Guidoni, S. 2012. Influence of wine-grape skin
11 hardness on the kinetics of anthocyanin extraction. *Int. J. of Food Properties*, **15(2)**: 249-
12 261.
- 13 27. Sarig, Y. 2012. Mechanical harvesting of fruit - past achievements. Current status and
14 prospects. *Acta Hort.*, **965**: 163-169.
- 15 28. Sasnauskas, A., Rugienius, R., Mazeikiene, I., Bendokas, V. and Stanys, V. 2019. Small
16 fruit breeding tendencies in Lithuania: a review. *Acta Hort.*, **1265**: 225-231.
- 17 29. Sava, P. and Bodi, G. 2012. Growing technology implementation of black currant
18 varieties for berries production in District Soroca, Republic of Moldova. *Sci. Papers – Ser.*
19 *B Hort.*, **56**: 167-170.
- 20 30. Spinei, M. and Oroian, M. 2021. The potential of grape pomace varieties as a dietary source
21 of pectin substances. *Foods*, **10(4)**: 867.
- 22 31. Szmajda, G. and Nowakowski T. 2020. Analysis of the Harvesting Quality of Redcurrant
23 with a Trailed Combine. *Agr. Eng.*, **24 (1)**: 91-102.
- 24 32. Tousehik, S.H., Lee, K.T., Lee, J.S. and Kim, K.S. 2017. Functional applications of
25 lignocellulolytic enzymes in the fruit and vegetable processing industries. *J. of food sci.*,
26 **82(3)**: 585- 593.
- 27 33. Wang, Y., Chen, H. and Lin, Q. 2009. Optimization of parameters of blackcurrant
28 harvesting mechanism. *Trans. of the Chinese Soci. of Agr. Eng.*, **25(3)**: 79-83.
- 29 34. Wróblewska, W., Pawlak, J. and Paszko, D. 2019. Economic aspects in the raspberry
30 production on the example of farms from Poland, Serbia and Ukraine. *J. of Hort. Res.*,
31 **27(2)**: 71-80.
- 32 35. Yu, P., Li, C., Takeda, F., Krewer, G., Rains, G., and Hamrita, T. 2012. Quantitative
33 evaluation of a rotary blueberry mechanical harvester using a miniature instrumented
34 sphere. *Comp. and elect.in agriculturae*, **88**: 25-31.

- 1 36. Zouid, I., Siret, R., Mehinagic, E., Maury, C., Chevalier, M., and Jourjon, F. 2010.
2 Evolution of grape berries during ripening: Investigations into the links between their
3 mechanical properties and the extractability of their skin anthocyanins. *OENO One*, **44(2)**:
4 87-99.
- 5 **37.** Zubarev, Yu. A., Gunin, A. V., Panteleeva, E. I., and Vorobyeva, A. V. 2022. Sea
6 buckthorn cultivars promising for mechanized harvesting by cutting fruit-bearing
7 branches. *Proc. on applied botany, genetics and breeding*. **183(2)**:43-50. (In Russ.)