

## Synthesis of Poly- $\gamma$ -Glutamate in Solid-State Fermentation and Its Application in Biocontrol

Sareh Hashemi<sup>1</sup>, Masoud Ahmadzadeh<sup>1\*</sup>, Hossein Saremi<sup>1</sup>, Soleiman Ghasemi<sup>2</sup>, Azad Omrani<sup>3</sup>

### ABSTRACT

Poly-  $\gamma$ - glutamic acid ( $\gamma$ -PGA) is a natural polymer with diverse applications across multiple industries. However, its use in agriculture is limited due to high production costs. This study aimed to optimize the cost-effective production of  $\gamma$ -PGA through Solid-State Fermentation (SSF) using *Bacillus velezensis* UTB96, evaluate the concentration and molecular weight of  $\gamma$ -PGA suitable for agricultural applications, particularly in strawberry cultivation, and explore the impact of  $\gamma$ -PGA on extending the shelf-life of strawberry fruits during cold storage. Initially, the production of  $\gamma$ -PGA using SSF with *B. velezensis* UTB96 was investigated, along with an evaluation of the influence of physicochemical factors on the molecular weight of  $\gamma$ -PGA. Based on the results, three different molecular weights of  $\gamma$ -PGA were identified: 1156.43 kDa, 734.38 kDa, and 296.55 kDa. These were selected for greenhouse trials to assess their effectiveness in controlling gray mold on strawberry plants. The results showed that by utilizing agricultural wastes, including sesame flour, wheat straw, and banana peel in SSF methodology,  $\gamma$ -PGA could be produced at a rate of 70 g/kg of dry weight of the culture medium. Analyzing the impact of  $\gamma$ -PGA on reducing gray mold revealed that this compound could enhance the plant's defense. A significant increase in the activity of ascorbate peroxidase and phenylalanine ammonia-lyase (PAL) enzymes was observed, along with the production of polyphenolic compounds such as ellagic acid. Consequently, these mechanisms improved the plant's flexibility and tolerance to the fungus, helping to maintain the quality of the fruits during cold storage.

**Keywords:** *Botrytis cinerea*, Molecular weight,  $\gamma$ -PGA, Solid-state fermentation.

### INTRODUCTION

There is increasing interest in finding effective and sustainable alternatives to chemical

---

<sup>1</sup> Department of Plant Protection, Faculty of Agriculture, College of Agriculture and Natural Resources, University of Tehran, Alborz, Karaj, Islamic Republic of Iran.

<sup>2</sup> Research and Development Section of Nature Biotechnology Co. (Biorun), Karaj. Alborz, Islamic Republic of Iran.

<sup>3</sup> Research and Development Section of Fruit Science Co., Karaj, Alborz, Islamic Republic of Iran.

\* Corresponding author: e-mail: ahmadz@ut.ac.ir,

30 pesticides. One promising option is biological control. Various microbial Biological Control  
31 Agents (BCAs) have been developed in recent decades to tackle fungal and bacterial diseases.  
32 Many studies have focused on microorganisms such as *Pseudomonas* spp., *Bacillus* spp., and  
33 *Streptomyces* spp. (Bonaterra *et al.*, 2022). *Bacillus velezensis* is a well-known strain  
34 recognized for its beneficial effects on plant growth and its role in biocontrol as a gram-positive  
35 rhizobacterium (Fan *et al.*, 2018). Recent research has investigated the microencapsulation of  
36 this bacterium with natural polymers and nanoparticles to control diseases like the *Rhizoctonia*  
37 *solani* fungus in beans (Moradi Pour *et al.*, 2021) and pistachio gum (Moradi Pour *et al.*, 2022).  
38 Numerous studies have highlighted the potential of bacteria, particularly *Bacillus* species, in  
39 direct antibiosis, competition, and the secretion of various secondary metabolites in the  
40 rhizosphere (Fan *et al.*, 2018). A significant secondary metabolite produced by *B. velezensis* is  
41 poly-gamma-glutamic acid ( $\gamma$ -PGA). This biopolymer is biodegradable, non-toxic,  
42 environmentally safe, and hypoallergenic, making it highly sought after in various industries.  
43 Research on this biological compound has shown that  $\gamma$ -PGA can significantly enhance plant  
44 performance and morphological characteristics. It also improves Soil Microbial Biomass  
45 Carbon (SMBC) and Nitrogen (SMBN), boosts soil enzyme activity, and increases plant  
46 resistance to both biotic and abiotic stresses (Song *et al.*, 2019).

47 Given the widespread application of  $\gamma$ -PGA across various industries, its industrial  
48 production through Submerged Fermentation (SmF) using *Bacillus* bacteria is common.  
49 However, this method is expensive, which restricts its use in agriculture. In contrast, Solid-  
50 State Fermentation (SSF) presents a cost-effective alternative by utilizing inexpensive and  
51 readily accessible raw materials. Furthermore, SSF provides various economic and engineering  
52 benefits, such as lower energy consumption and simplified equipment and infrastructure  
53 requirements (Chen *et al.*, 2005).

54 Therefore, this study investigates innovative approaches for the cost-effective production of  
55  $\gamma$ -PGA through solid-state fermentation, utilizing affordable waste materials for agricultural  
56 purposes. Additionally, it examines the effects of environmental factors on the molecular  
57 weight of  $\gamma$ -PGA. Another critical aspect addressed in this article is the mechanisms by which  
58  $\gamma$ -PGA enhances the resistance of strawberry plants to necrotrophic fungi, particularly *Botrytis*  
59 *cinerea*, the most prevalent and destructive fungal pathogen affecting strawberries.  
60 Additionally, the study investigates the effect of pre-harvest application of  $\gamma$ -PGA on the shelf  
61 life of strawberry fruits during cold storage.

62

63 **MATERIALS AND METHODS**

64 **1:  $\gamma$ -PGA Production in Solid-State Fermentation (SSF) and Its Identification**

65 **Preparation of *Bacillus velezensis* UTB96 Bacteria:**

66 In this study, the *Bacillus velezensis* strain UTB96 was obtained from the microbial collection  
67 at Tehran University. The bacteria were cultured in a nutrient broth medium (NB) at 37°C and  
68 200 rpm for 16 hours to serve as the inoculum. This medium contained approximately  $3 \times 10^7$   
69 cells per mL.

70  
71 **Substrate for Solid-State Fermentation**

72 To achieve optimal  $\gamma$ -PGA production, a balanced combination of protein, sugar, and carbon  
73 sources is essential. For the economical production of  $\gamma$ -PGA, the recommended medium  
74 consists of a blend of sesame flour (as the protein source), wheat straw (as the carbon source),  
75 banana peel (providing sugar and minerals), and cow manure (as a mineral source). Various  
76 experiments were conducted with different ratios of these components to determine the most  
77 effective combination for production.

78  
79 **Extraction and Purification of  $\gamma$ -PGA**

80 The extraction and purification of  $\gamma$ -PGA were performed according to the method described  
81 by Goto and Kunioka (Goto and Kunioka, 1999).

82  
83 **Identification and Characterization of  $\gamma$ -PGA**

84 The produced  $\gamma$ -PGA was identified and characterized using SDS-PAGE (Yu *et al.*, 2016), FT-  
85 IR (Khalil *et al.*, 2018), and spectrophotometric methods (Zeng *et al.*, 2012).

86  
87 **Determining the Molecular Weight and Investigating the Effect of Environmental**  
88 **Physicochemical Factors on the Molecular Weight of  $\gamma$ -PGA**

89 The effects of the following physicochemical factors on the molecular weight of  $\gamma$ -PGA were  
90 studied by varying one factor at a time:

- 91 a) Incubation temperature (27 to 42°C)
- 92 b) Initial moisture level of the fermentation medium (50 to 75 %)
- 93 c) Initial pH level of the fermentation medium (5 to 8)
- 94 d) Fermentation time (12 to 96 hours)

95 The average molecular weights of the  $\gamma$ -PGA product obtained under these specified  
96 environmental conditions were determined using Gel Permeation Chromatography (GPC). The

97 mobile phase consisted of 50 mM phosphate buffer at pH 6.8, with a flow rate of 1.0 mL/min.  
98 Molecular weights were calculated relative to polystyrene standards.

99  
100 **2: Studying the Impact of  $\gamma$ -PGA on Strawberry Resistance to *B. cinerea* Fungus and**  
101 **Plant Biochemical Mechanisms**

102 The effect of  $\gamma$ -PGA on the resistance of strawberry plants to *B. cinerea* was evaluated under  
103 greenhouse conditions. For this research, all fermented culture media containing  $\gamma$ -PGA in three  
104 different molecular weights (high: 1156.43 kDa, medium: 734.38 kDa, and low: 296.55 kDa)  
105 were selected for experimentation. These media were dried in an oven at 70°C for 3 days and  
106 then crushed using a mill. The resulting powders were mixed with a combination of perlite and  
107 cocopeat in a 50:50 ratio to serve as the substrate for strawberry cultivation.

108 The concentrations of  $\gamma$ -PGA investigated were 10, 20, 50, and 100 mg/kg of the cultivation  
109 bed. Each treatment involved five 1.5-liter pots, with two strawberry cv. 'Camarosa' seedlings  
110 planted in each pot, which were irrigated daily with 100 cc of drinking water. The light-dark  
111 photoperiod was set to 8 hours of light and 16 hours of darkness. At the end of the flowering  
112 stage, the plants were inoculated with a *B. cinerea* fungus suspension containing  $10^6$  spores per  
113 mL. The control treatment (Sh0) was inoculated with water.

114  
115 **Treatments**

116 The treatments consisted of two control groups and  $\gamma$ -PGA groups:

117 a) Control treatment group:

- 118 ➤ Sh0: Control treatment without fermentation substrate, with or without  $\gamma$ -PGA, and  
119 without contamination by *B. cinerea* fungus.  
120 ➤ Sh-: Control treatment without fermentation substrate, with or without  $\gamma$ -PGA, and with  
121 contamination by *B. cinerea* fungus.  
122 ➤ Sh 10,20, 50, and 100 mg: In this investigation, the entire solid-state fermentation  
123 substrate was added to the pots. To assess the impact of compounds produced by bacteria  
124 other than  $\gamma$ -PGA, the fermentation substrate without the  $\gamma$ -PGA compound was added to the  
125 culture medium at concentrations of 10, 20, 50, and 100 mg/kg of soil. This group of control  
126 treatments was inoculated with the *B. cinerea* fungus.

127 b)  $\gamma$ -PGA treatment group: The treatments consist of fermentation substrate containing  $\gamma$ -PGA  
128 with molecular weights of 296.55, 734.38, and 1156.43 kDa. Each of these molecular weights  
129 was investigated at four concentrations: 10, 20, 50, and 100 mg/kg of soil. The  $\gamma$ -PGA group  
130 treatments were inoculated with the fungus *B. cinerea*.

131 **Ascorbate Peroxidase (APX) Enzyme Activity Assay**

132 Sampling of strawberry leaves for the investigation of biochemical characteristics was  
133 conducted immediately before inoculation and on days 3, 7, 10, 20, and 30 post-inoculations,  
134 coinciding with the first fruit harvest. The samples were promptly transferred to a -80 freezer  
135 for storage until examination. The activity of the APX enzyme was determined using the  
136 method described by Braga *et al.* (Braga *et al.*, 2009).

137  
138 **Phenylalanine Ammonia Lyase (PAL) Enzyme Activity Assay**

139 The activity of phenylalanine ammonia-lyase was assessed by converting L-phenylalanine into  
140 trans-cinnamic acid following the protocol outlined by Tovar *et al.* (Tovar *et al.*, 2002). PAL  
141 enzymatic activity was reported as units per gram of fresh leaf weight.

142  
143 **Assay of Ellagic Acid (EA)**

144 The concentration of ellagic acid was measured using the spectrophotometric method  
145 established by Wilson and Hagerman (Wilson and Hagerman, 1990).

146  
147 **3: Measurement of Physicochemical Characteristics of Fruits in Storage Conditions**

148 To determine the effectiveness of  $\gamma$ -PGA on the shelf life and quality of fruits in storage,  
149 physicochemical tests were conducted. These tests included assessing weight loss, firmness,  
150 Total Soluble Solids (TSS), and Titratable Acidity (TA) using established methods (Farida *et al.*, 2023). Forty fruits were harvested from each treatment and stored in a cold room at a  
151 temperature of 4°C and a humidity of 90% for 10 days. The samples were analyzed on days 0,  
152 3, 6, and 10.

154  
155 **Apparent Decay of Fruits**

156 To assess decay, 20 fruits from each treatment were randomly selected, and the average decay  
157 for each treatment was calculated using the numerical scale defined by Babalare *et al.* (2007).  
158 Fungal decay was evaluated through visual examination and microscopic observation of fungal  
159 mycelium growth. The numerical scale ranged from 5 (indicating no decay) to 1 (indicating  
160 more than 16% decay), with intermediate values representing different levels of decay: 4 (less  
161 than 5%), 3 (6-10%), and 2 (11-15%) (Babalar *et al.*, 2007).

162  
163 **4. Statistical Analysis**

164 This study utilized a completely randomized design. The data were analyzed using SAS  
165 9.1.3 statistical software (2001). Duncan's multiple range test was employed at the 5%

166 significance level to compare the means. The experiments were conducted in triplicate.

167

## 168 RESULTS AND DISCUSSION

### 169 1: $\gamma$ -PGA Production by SSF and Its Identification

#### 170 Substrate of SSF

171 Studies have shown that nitrogen and carbon sources, particularly sugars, play a crucial role in  
 172  $\gamma$ -PGA production (Sung *et al.*, 2005). Therefore, in this study, the optimal ratio between  
 173 sesame flour, which serves as a nitrogen source, and banana peel, which acts as a sugar source,  
 174 was determined (Table 1, Table 2). In all the examined samples, the amounts of wheat straw  
 175 was 150 g and manure in the substrate - 50 g and remained constant. Subsequently, based on  
 176 this ratio, the proportion of cow manure to wheat straw was calculated. To determine the  
 177 optimal ratio, various combinations were examined (Table 3). The most effective composition  
 178 of the fermentation substrate comprised (g/kg dry weight of substrate): 600g sesame flour, 200g  
 179 banana peels, 160g wheat straw, and 40g cow manure, resulting in a  $\gamma$ -PGA production rate of  
 180 70 g/kg dry weight (DW) of the substrate.

181 **Table 1.** Main composition in selected substances for solid-state fermentation.

The main composition	Minerals	Carbohydrate	Carbon source (sugar)	Protein
Substances	Cow manure	Wheat straw	Banana peel	Sesame flour
Amount	34 %	42.56 g/100 g	39.8 g/100g	45.8 g/100g

182

183 **Table 2.** The optimal ratio of sesame flour to banana peel (g).

Sample	Sesame flour	Banana peel	Kg of the	The average production of $\gamma$ -PGA per dry weight of the substrate
1	200	600		55/75 <sup>d</sup>
2	300	500		61/5 <sup>c</sup>
3	400	400		65 <sup>b</sup>
4	500	300		68/33 <sup>a</sup>
5	600	200		69/16 <sup>a</sup>
6	700	100		60/2 <sup>c</sup>

184

185

186

187

188

189

190

191

192

193

194

195

196 **Table 3.** The optimal ratio of wheat straw to manure and between sesame flour and banana peel  
 197 (g).

Sample	Sesame flour	Banana peel	Wheat straw	Manure	The average production of $\gamma$ -PGA per Kg of the dry weight of the substrate
1	600	200	180	20	62/5 <sup>c</sup>
2	600	200	170	30	62/5 <sup>c</sup>
3	600	200	160	40	70 <sup>a</sup>
4	600	200	150	50	69 <sup>b</sup>
5	600	200	140	60	52/5 <sup>e</sup>
6	600	200	100	100	33/25 <sup>g</sup>
7	600	200	160	40	67/5 <sup>b</sup>
8	400	200	380	20	54 <sup>e</sup>
9	400	200	360	40	57/5 <sup>d</sup>
10	400	200	340	60	56/75 <sup>d</sup>
11	400	200	320	80	42/5 <sup>f</sup>

198 Different letters indicate significant differences between production values of  $\gamma$ -PGA ( $P < 0.05$ ).

### 199 Identification and Characterization of $\gamma$ -PGA

200 Various analytical tools were employed to identify and characterize the  $\gamma$ -PGA produced by *B.*  
 201 *velezensis* UTB96 through SSF. As shown in Figure 1- a, the Fourier-Transform Infrared (FT-  
 202 IR) spectrum of  $\gamma$ -PGA displayed amide bands at 1648.6  $\text{cm}^{-1}$  and carbonyl group C=O at  
 203 1402.43  $\text{cm}^{-1}$ . Additionally, it exhibited C-N stretching vibrations at 1075.76  $\text{cm}^{-1}$  and an O-H  
 204 stretching band at 3447.26  $\text{cm}^{-1}$ .


205 The chemical structure of the  $\gamma$ -PGA obtained in this study was consistent with the structure  
 206 elucidated by Ho *et al.* (2006) (Khalil *et al.*, 2018). Notably, based on previous studies (Rajan  
 207 *et al.*, 2014) and the absorption bands of amide groups, the secondary structure of  $\gamma$ -PGA was  
 208 identified as an  $\alpha$ -helix motif.

209 Figure 1- b displays the UV absorption spectrum of  $\gamma$ -PGA in deionized water, covering the  
 210 range of 190 – 340 nm. According to the orbital law, peptide excitation absorption peak,  
 211 resulting from the presence of carbonyl and amide groups, typically occurs around 200 nm  
 212 (Braga *et al.*, 2009). This aligns with our study's findings that the  $\gamma$ -PGA biopolymer exhibited  
 213 maximum absorption at 200 nm (216 nm).

214 Studies have demonstrated the effectiveness of methylene blue staining as a qualitative  
 215 method for measuring  $\gamma$ -PGA production (Yu *et al.*, 2016). In this investigation, the production  
 216 of  $\gamma$ -PGA by *B. velezensis* UTB96 was successfully confirmed using methylene blue staining,  
 217 as shown in Figure 1- c.

218

219



**Figure 1.**  $\gamma$ -PGA Identification, a: FT-IR absorption peaks ( $\text{cm}^{-1}$ ), b: UV spectral analysis, c: SDS-PAGE for methylene blue stained samples containing purified  $\gamma$ -PGA.

The mechanism by which the cationic dye methylene blue is absorbed by  $\gamma$ -PGA is linked to the active sites ( $\text{COO}^-$ ) present on the surface of the polymer. The absorption of methylene blue occurs spontaneously (Ogata *et al.*, 2017). The increases of the concentration or molecular weight of  $\gamma$ -PGA is indicating on a longer peptide chain and increasing the number of active sites ( $\text{COO}^-$ ). This increase in active sites enhances dye absorption, as demonstrated by the results.

### Poly- $\gamma$ - Glutamic Acid Molecular Weight

Microbial production of  $\gamma$ -PGA can result in molecular weights ranging from 100 to over 2,000 kDa. Several factors influence the efficiency and molecular weight of  $\gamma$ -PGA (Sung *et al.*, 2005). To investigate the effect of temperature on the molecular weight of  $\gamma$ -PGA, five experiments were conducted at varying temperatures. As shown in Figure 2- a, the maximum  $\gamma$ -PGA molecular weight (1117.53 kDa) was observed at  $37^\circ\text{C}$ , with similar values around 1112.9 kDa at  $32^\circ\text{C}$ . A decrease in the molecular weight was noted at  $42^\circ\text{C}$ , although this change was not significant compared to the  $37^\circ\text{C}$ . This reduction may be attributed to rapid cell growth at  $42^\circ\text{C}$  during the initial stages, which depletes nutrients in the medium and leads to the utilization of  $\gamma$ -PGA as a nitrogen and carbon source for bacterial cells in the later stages of

255 fermentation. Additionally, temperature can influence the activity of enzymes involved in  
256 synthesizing the  $\gamma$ -PGA amino acid chain, potentially affecting the molecular weight (Ajayeoba  
257 *et al.*, 2019).


258 The effect of moisture on the molecular weight of  $\gamma$ -PGA is illustrated in Figure 2- b. There  
259 was a direct relationship between humidity and the molecular weight of  $\gamma$ -PGA; as humidity  
260 increased to 65%, the molecular weight of  $\gamma$ -PGA also rose. However, a subsequent decrease  
261 in molecular weight was observed beyond this point. Moisture plays a vital role in oxygen and  
262 mass transfer during solid-state fermentation (SSF). Reports indicate that the activity of the  
263 pgsBCA enzyme complex is ATP-dependent, with ATP levels being influenced by oxygen  
264 availability. Therefore, when media humidity is at its optimal level, both enzyme complex  
265 activity and the molecular weight of  $\gamma$ -PGA are expected to increase (Sung *et al.*, 2005).

266 In Figure 2- c, it is shown that as the pH increased from 5 to 8, the molecular weight of  $\gamma$ -  
267 PGA rose from 630 kDa to 1121.4 kDa. There was no significant difference in molecular weight  
268 between pH values from 6.5 to 8. Under acidic conditions, the carboxylic acid groups do not  
269 ionize, causing the  $\gamma$ -PGA structure to adopt an  $\alpha$ -helical conformation, which results in  
270 decreased stability and decomposition of the compound (Seo *et al.*, 2008), leading to a reduction  
271 in molecular weight. In contrast, at higher pH levels, the ionic hydration of  $\gamma$ -PGA induces a  
272 conformational shift from  $\alpha$ -helix to random coil (Seo *et al.*, 2008), thereby increasing the  
273 compound's molecular weight.

274 At 96 hours of fermentation, the molecular weight of  $\gamma$ -PGA increased significantly between  
275 12 and 36 hours. However, after 48 hours, a downward trend in molecular weight was observed.  
276 By the end of the 96-hour fermentation period, the molecular weight of  $\gamma$ -PGA showed a 46%  
277 decline compared to the 36-hour fermentation period (Figure 2- d). When  $\gamma$ -PGA consists of  
278 longer monomeric chains, its molecular weight is higher. Therefore, extending the fermentation  
279 time up to 36 hours provides the polymerase enzyme systems with more time to polymerize  
280 monomers, increasing the molecular weight of  $\gamma$ -PGA. Prolonging fermentation beyond 36  
281 hours can create adverse conditions such as nutrient scarcity, oxygen depletion, and low  
282 humidity levels, which subsequently reduce the activity of the  $\gamma$ -PGA polymerase enzyme. In  
283 response, bacteria may hydrolyze  $\gamma$ -PGA as a source of carbon and nitrogen to ensure their  
284 survival (Cao *et al.*, 2018), leading to a decrease in the molecular weight of  $\gamma$ -PGA.

285  
286  
287

288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322



**Fig 2.** Effect of different factors on  $\gamma$ -PGA molecular weight, a: temperature, b: initial moisture, c: Initial pH, d: incubation time. Different letters in each figure indicate significant differences between the states of the investigated factor ( $P < 0.05$ ).

## 2: $\gamma$ -PGA Effect on Strawberry Plant Resistance to Gray Mold Disease


### Ascorbate Peroxidase (APX) Enzyme Assay

The activity of the APX enzyme in the leaves of plants treated with  $\gamma$ -PGA showed an increasing trend until the tenth day. Notably, the increase in enzyme activity was significantly higher in the  $\gamma$ -PGA treatments compared to the infected control (Sh-). On the tenth day, the highest enzyme activity was recorded in the treatment of 20 mg 296.55 kDa, which was 50 times higher than that of the negative control (Fig. 3- a). After the tenth day, the activity of the ascorbate peroxidase enzyme began to decline. In contrast, the control treatments continued to show an increase in enzyme activity until the twentieth day (Fig. 3- a). This continued increase in the control could be attributed to the emergence of new infections or the spread of the fungal pathogen *B. cinerea* within the plant tissues.

APX is a key enzyme that converts ascorbate to dehydroascorbate, effectively removing peroxides, particularly  $H_2O_2$ , from plant cells (Navari-Izzo *et al.*, 1997). Based on the results obtained, it can be concluded that  $\gamma$ -PGA positively influences the plant's antioxidant system, facilitating the metabolism of reactive oxygen species (ROS) and preventing the penetration and spread of the necrotrophic fungus *B. cinerea* in strawberry plants.

323 **Phenylalanine Ammonia Lyase (PAL) Enzyme Assay**

324 The application of  $\gamma$ -PGA significantly increased the activity of the PAL enzyme from the third  
 325 to the seventh day following the inoculation of *B. cinerea* in the  $\gamma$ -PGA treatments compared to  
 326 the control groups. The highest increases in PAL activity were observed in the treatments with  
 327 20 and 50 mg of 296.5 kDa, as well as 50 and 20 mg of 734.38 kDa. In contrast, the uninfected  
 328 (Sh0) and negative (Sh-) control groups exhibited the lowest levels of phenylalanine ammonia-  
 329 lyase activity (Fig. 3- b). PAL plays a crucial role in the biosynthesis of polyphenolic compounds,  
 330 including flavonoids, phenylpropanoids, and lignin in plants. Research suggests that  $\gamma$ -PGA can  
 331 enhance the activity of the PAL enzyme by activating the ROS signaling pathway, leading to  
 332 increased enzyme activity (Lei *et al.*, 2015). Furthermore, studies have indicated that stimulating  
 333 the phenylpropanoid pathway to produce polyphenolic and flavonoid compounds can help  
 334 neutralize and prevent excessive ROS production caused by *B. cinerea* infection (Kumar *et al.*,  
 335 2020; Perkowski and Warpeha., 2019).



353 \*The order of the columns in the days examined is: Sh0, Sh-, Sh10, Sh20, Sh50, Sh100, 10 mg 296.5 kDa, 20 mg  
 354 296.5 kDa, 50 mg 296.5 kDa, 100 mg 296.5 kDa, 10 mg 734.3 kDa, 20 mg 734.3 kDa, 50 mg 734.3 kDa, 100 mg  
 355 734.3 kDa, 10 mg 1156.4 kDa, 20 mg 1156.4 kDa, 50 mg 1156.4 kDa and 100 mg 1156.4 kDa.

356 **Figure 3.** The effect of  $\gamma$ -PGA on the activity of enzymes involved in strawberry plant defense,  
 357 a: Ascorbate peroxidase (APX), b: Phenylalanine ammonia- lyase (PAL).

358 **Impact of  $\gamma$ -PGA on the Ellagic Acid (EA) Content in Leaves**

359 The level of ellagic acid in strawberry leaves continued to rise until the seventh day, with the  
 360 greatest increase observed in the treatment with 50 mg of 296.55 kDa (18.6 mg/100 g FW).  
 361 Subsequently, a significant decline in ellagic acid was noted in the control treatments, while in  
 362 the  $\gamma$ -PGA treatments, this decline exhibited a more gradual trend (Fig. 4).

363 Within plant cells, ellagic acid exists in both free and covalently-bound forms, such as EA  
 364 glycosides and ellagitannins. Although the level of free ellagic acid in cells is initially low, it  
 365 tends to increase during biotic and abiotic stresses. Ellagic acid has demonstrated efficiency in  
 366 absorbing free radicals (Williams *et al.*, 2014), Therefore, the rise in ellagic acid levels during  
 367 the compatible interaction between *B. cinerea* and strawberries may indicate its direct  
 368 involvement in protecting plant cells through the modulation of the redox balance. Additionally,  
 369 this compound contributes to defense against fungal pathogens by damaging ergosterol (Lei *et*  
 370 *al.*, 2015). As a result, ellagic acid leads to pathogen death by disrupting ergosterol and  
 371 compromising the integrity of the fungal membrane.

372

373

374

375

376

377

378


379

380

381

382

383



384 \*The order of the columns in the days examined is: Sh0, Sh-, Sh10, Sh20, Sh50, Sh100, 10 mg 296.5 kDa, 20 mg  
 385 296.5 kDa, 50 mg 296.5 kDa, 100 mg 296.5 kDa, 10 mg 734.3 kDa, 20 mg 734.3 kDa, 50 mg 734.3 kDa, 100 mg  
 386 734.3 kDa, 10 mg 1156.4 kDa, 20 mg 1156.4 kDa, 50 mg 1156.4 kDa and 100 mg 1156.4 kDa.

387 **Figure 4.**  $\gamma$ -PGA effect on the activity of Ellagic acid (EA) in strawberry leaves inoculated with  
 388 *B. cinere*.

389

390 **3: Measurement of Physicochemical Characteristics of Fruits in Storage Conditions**

391 As storage time increased, fruits exhibited a decrease in weight, firmness, and titratable  
 392 acidity. Notably, the control treatments showed the most significant reduction compared to the

393  $\gamma$ -PGA treatments (Table 4). The lowest reduction was observed in the treatments with 50 mg  
394 and 20 mg of 296.55 kDa, as well as 50 mg of 734.38 kDa.

395 In terms of Total Soluble Solids (TSS) analysis, a gradual increase was noted across all  
396 treatments throughout the storage period. The smallest increase was recorded in the treatment  
397 with 50 mg of 296.55 kDa, which was 27.4% lower than the Sh- treatment.

398 Physicochemical changes during fruit storage are primarily caused by respiration, water loss,  
399 the consumption and breakdown of stored compounds and metabolites during metabolic  
400 activities, and the activity of degrading enzymes (Fawole *et al.*, 2020). Research has shown that  
401  $\gamma$ -PGA operates through various mechanisms, including reducing the activity of cell wall  
402 degrading enzymes (Wang *et al.*, 2020), inhibiting abscisic acid signal transmission (Shan *et*  
403 *al.*, 2023), and enhancing the antioxidant capacity in fruits. These actions collectively  
404 contribute to maintaining cell integrity, preventing water loss, reducing decay, and delaying  
405 aging in fruits. As a result,  $\gamma$ -PGA plays a crucial role in preserving the quality of fruits during  
406 storage.

407

#### 408 **Apparent Decay of Fruits**

409 At the end of ten days, no symptoms of fungal infection were observed in the  $\gamma$ -PGA  
410 treatments, while the fruits in the control treatments exhibited moderate to severe rotting (Table  
411 4). As storage time increases, the texture of strawberries softens due to changes in the cell wall  
412 structure (Brummell *et al.*, 1999). The antioxidant activity of  $\gamma$ -PGA and its ability to maintain  
413 the integrity of the cell wall, helps prevent rapid deterioration of the fruit tissue and maintains  
414 its hardness. This, in turn, inhibits the penetration and spread of fungi into the fruit tissue.

415

416

417

418

419

420


421

422

423

424

425



**Figure 5.** Examination of the apparent decay in strawberry fruits after 10 days of storage at 4°C and 90% humidity, a: Control treatments, b: Microscopic examination of fungal mycelium grown on the fruits of control treatments, and c:  $\gamma$ -PGA treatments.

**Table 4.** The impact of different treatments on the physicochemical properties of strawberry fruits after 10 days of cold storage.

Indicator Treatment	Weight loss (%)	TA (%)	TSS (%)	Firmness (kg/m <sup>2</sup> )	Apparent decay
Sh 0	5.3 <sup>b</sup>	0.62 <sup>cdefg</sup>	8.3 <sup>a</sup>	0.64 <sup>g</sup>	3.11 <sup>b</sup>
Sh-	5.9 <sup>a</sup>	0.47 <sup>g</sup>	8.4 <sup>a</sup>	0.6 <sup>e</sup>	1.04 <sup>e</sup>
Sh 10 mg	4.5 <sup>c</sup>	0.5 <sup>fg</sup>	8 <sup>ab</sup>	0.62 <sup>g</sup>	1.95 <sup>d</sup>
Sh 20 mg	4.28 <sup>cd</sup>	0.53 <sup>efg</sup>	7.9 <sup>ab</sup>	0.73 <sup>fg</sup>	1.95 <sup>d</sup>
Sh 50 mg	4.2 <sup>cd</sup>	0.58 <sup>defg</sup>	7.6 <sup>bc</sup>	0.81 <sup>ef</sup>	2.83 <sup>c</sup>
Sh 100 mg	4.12 <sup>d</sup>	0.6 <sup>defg</sup>	7.3 <sup>cd</sup>	0.84 <sup>def</sup>	2.52 <sup>c</sup>
10 mg 296.5 kDa	2.9 <sup>hi</sup>	0.72 <sup>bcd</sup>	6.7 <sup>ef</sup>	0.95 <sup>cde</sup>	4.79 <sup>a</sup>
20 mg 296.5 kDa	1.6 <sup>kl</sup>	1.15 <sup>a</sup>	6.5 <sup>fg</sup>	1.14 <sup>ab</sup>	4.79 <sup>a</sup>
50 mg 296.5 kDa	1.5 <sup>l</sup>	0.8 <sup>b</sup>	6.1 <sup>g</sup>	0.9 <sup>de</sup>	4.79 <sup>a</sup>
100 mg 296.5 kDa	2.3 <sup>j</sup>	0.72 <sup>bcd</sup>	6.7 <sup>ef</sup>	0.98 <sup>cd</sup>	4.79 <sup>a</sup>
10 mg 734.3 kDa	2.9 <sup>hi</sup>	0.69 <sup>cde</sup>	7 <sup>def</sup>	0.88 <sup>de</sup>	4.79 <sup>a</sup>
20 mg 734.3 kDa	3.06 <sup>gh</sup>	0.74 <sup>bcd</sup>	6.5 <sup>fg</sup>	0.83 <sup>ef</sup>	4.79 <sup>a</sup>
50 mg 734.3 kDa	1.9 <sup>k</sup>	0.7 <sup>dc</sup>	6.5 <sup>fg</sup>	1.2 <sup>a</sup>	4.79 <sup>a</sup>
100 mg 734.3 kDa	3.23 <sup>fg</sup>	0.64 <sup>cdef</sup>	6.8 <sup>def</sup>	0.91 <sup>cde</sup>	4.79 <sup>a</sup>
10 mg 1156.4 kDa	2.7 <sup>i</sup>	0.77 <sup>bc</sup>	6.6 <sup>ef</sup>	1.05 <sup>bc</sup>	4.79 <sup>a</sup>
20 mg 1156.4 kDa	2.8 <sup>hi</sup>	0.68 <sup>cde</sup>	7.1 <sup>cde</sup>	0.9 <sup>de</sup>	4.79 <sup>a</sup>
50 mg 1156.4 kDa	3.6 <sup>e</sup>	0.66 <sup>cdef</sup>	6.9 <sup>def</sup>	0.92 <sup>cde</sup>	4.79 <sup>a</sup>
100 mg 1156.4 kDa	3.5 <sup>ef</sup>	0.64 <sup>cdef</sup>	7.2 <sup>cde</sup>	0.98 <sup>cd</sup>	4.79 <sup>a</sup>

\* Different letters in columns show significant differences (P<0.001).

## CONCLUSIONS

The findings of this research demonstrate that through solid-state fermentation and

449 manipulation of fermentation conditions, it is possible to produce  $\gamma$ -PGA compounds in  
450 significant quantities with diverse molecular weights. The results indicate that  $\gamma$ -PGA,  
451 particularly at low to medium molecular weights and concentrations, could enhance plant  
452 defense against necrotrophic pathogens, such as *B. cinerea*, by activating antioxidant  
453 mechanisms and boosting the generation of defense compounds. This enhancement could also  
454 prolong the storage life of strawberry fruit. The outcomes of this study present a promising  
455 avenue for researchers and experts in agriculture to further investigate the potential applications  
456 of  $\gamma$ -PGA in various fields, including biological control.

457

458 **REFERENCES**

- 459 (1) Ajayeoba, T. A., Dula, S. and Ijabadeniyi, O. A. 2019. Properties of Poly-Glutamic Acid  
460 Producing- *Bacillus* Species Isolated from Ogi Liquor and Lemon- Ogi Liquor. *Front.*  
461 *Microbiol.*, **10**: 771-778.
- 462 (2) Babalar, M. A., sghari, M., Talaei, A. and Khosroshahi, A. 2007. Effect of Pre-and  
463 Postharvest Salicylic Acid Treatment on Ethylene Production, Fungal Decay and Overall  
464 Quality of Selva Strawberry Fruit. *Food Chem.*, **105(2)**: 449-453.
- 465 (3) Bonaterra, A., Badosa, E., Daranas, N., Francés, J., Roselló, G. and Montesinos, E. 2022.  
466 Bacteria as Biological Control Agents of Plant Diseases. *Microorganisms*, **10(9)**:1759.
- 467 (4) Braga, L. F., Sousa, M. P., Ferreira, L. C., Delachieve, M. E. A., Cataneo, A. C. and Braga,  
468 J. F. 2009. Proline Level and Amylase and Ascorbate Peroxidase Activity in Germination of  
469 *Plantago ovata* Forsk (Plantaginaceae) Seeds. *JABS*, **4(6)**: 49-54.
- 470 (5) Brummell, D. A., Harpster, M. H., Civello, P. M., Palys, J. M., Bennett, A. B. and Dunsmuir,  
471 P. 1999. Modification of Expansin Protein Abundance in Tomato Fruit Alters Softening and  
472 Cell Wall Polymer Metabolism During Ripening. *Plant Cell*, **11(11)**: 2203-16.
- 473 (6) Cao, M., Feng, J., Sirisansaneeyakul, S., Song, C. and Chisti, Y. 2018. Genetic and  
474 Metabolic Engineering for Microbial Production of Poly-  $\gamma$ - Glutamic Acid. *Biotechnol. Adv.*,  
475 **36**: 1424-1433.
- 476 (7) Chen, X., Chen, S., Sun, M. and Yu, Z. 2005. High Yield of Poly- $\gamma$ -Glutamic Acid From  
477 *Bacillus subtilis* by Solid-State Fermentation Using Swine Manure as the Basis of a Solid  
478 Substrate. *Bioresource Technol.*, **96**: 1872-1879.
- 479 (8) Fan, B., Wang, C., Song, X., Ding X., Wu, L., Wu, H., Gao, X. and, Borriss R. 2018.  
480 *Bacillus velezensis* FZB42 in 2018: The Gram-Positive Model Strain for Plant Growth  
481 Promotion and Biocontrol. *Front Microbiol.*, **9**:2491.

- 482 (9) Farida, F., Hamdani, J. S., Mubarok, S., Akutsu, M., Noviyanti, K. and Nur Rahmat, B. P.  
483 2023. Variability of Strawberry Fruit Quality and Shelf Life with Different Edible Coatings.  
484 *Horticulturae*, **9(7)**:741.
- 485 (10) Fawole, O. A., Atukuri, J., Arendse, E. and Opara, U. O. 2020. Postharvest Physiological  
486 Responses of Pomegranate Fruit (cv. Wonderful) to Exogenous Putrescine Treatment and  
487 Effects on Physico-Chemical and Phytochemical Properties. *Food Science and Human*  
488 *Wellness*, **9(2)**: 146-161.
- 489 (11) Goto, A. and Kunioka, M. 1992. Biosynthesis and Hydrolysis of Poly(Gamma-Glutamic  
490 Acid) from *Bacillus subtilis* IFO3335. *Biosci. Biotechnol. Biochem.*, **156**:1031-1035.
- 491 (12) Khalil, I. R., Khechara, M. P., Kurusamy, S., Armesilla, A. L., Gupta, A., Mendrek, B.,  
492 Khalaf, T., Scandola, M., Focarete, M. L. and Kowalczyk, M. 2018. Poly- Gamma- Glutamic  
493 Acid ( $\gamma$ - PGA)- Based Encapsulation of Adenovirus to Evade Neutralizing Antibodies.  
494 *Molecules*, **23**: 2565.
- 495 (13) Kumar, V., Hatan, E., Bar, E., Davidovich-Rikanati, R., Doron-Faigenboim, A., Spitzer-  
496 Rimon, B., Elad, Y., Alkan, N. and Lewinsohn, E. 2020. Phenylalanine Increases  
497 Chrysanthemum Flower Immunity Against *Botrytis cinerea* Attack. *The plant journal*, **104(1)**:  
498 226-240.
- 499 (14) Lei, P., Xu, Z., Ding, Y., Tang, B., Zhang, Y., Li, H., Feng, X. and Xu, H. 2015. Effect of  
500 Poly ( $\gamma$ -Glutamic Acid) on the Physiological Responses and Calcium Signaling of Rape  
501 Seedlings (*Brassica napus* L.) under Cold Stress. *J. Agric. Food Chem.*, **63 (48)**:10399-10406.
- 502 (15) Moradi Pour, M., Saberi-Riseh, R., Esmailzadeh-Salestani, K., Mohammadinejad, R. and  
503 Loit, E. (2021). Evaluation of *Bacillus velezensis* for Biological Control of *Rhizoctonia solani*  
504 in Bean by Alginate/Gelatin Encapsulation Supplemented with Nanoparticles. *J Microbiol*  
505 *Biotechnol.*, **31(10)**:1373-1382.
- 506 (16) Moradi Pour, M., Saberi Riseh, R. and Skorik, YA. (2022). Sodium Alginate-Gelatin  
507 Nanoformulations for Encapsulation of *Bacillus velezensis* and Their Use for Biological  
508 Control of Pistachio Gummosis. *Materials (Basel)*, **15(6)**:2114.
- 509 (17) Navari-Izzo, F., Meneguzzo, S., Loggini, B., Vazzana, C. and Sgherri, C. L. 1997. The  
510 Role of the Glutathione System During Dehydration of *Boea hygroskopica*. *Physiol. Plant*,  
511 **99(1)**: 23-30.
- 512 (18) Ogata, F., Nagai, N. and Kawasaki, N. 2017. Adsorption Capability of Cationic Dyes  
513 (Methylene Blue and Crystal Violet) onto Poly- $\gamma$ -Glutamic Acid. *Chem. Pharm. Bull.*, **65(3)**:  
514 268-275.

- 515 (19) Perkowski, M. C. and Warpeha, K. M. 2019. Phenylalanine Roles in the Seed-to-Seedling  
516 Stage: not Just an Amino Acid. *Plant Sci.*, **289**.
- 517 (20) Rajan, Y. C., Inbaraj, B. S. and Chen, B. H. 2014. In vitro Adsorption of Aluminum by an  
518 Edible Biopolymer Poly ( $\gamma$ - Glutamic Acid). *J. Agric. Food Chem.*, **62**: 4803-4811.
- 519 (21) Seo, J. H., Kim, C. H. and Lee, S. P. 2008. Physicochemical Properties of Poly-  $\gamma$ - Glutamic  
520 Acid Produced by a Novel *Bacillus subtilis* HA Isolated from Cheonggukjan. *J. Food Sci. Nutr.*,  
521 **13(4)**: 354-361.
- 522 (22) Shan, C., Luo, Y., Yang, C. and Gao, X. 2023. The Effects of Poly- $\gamma$ -Glutamic Acid on  
523 the Postharvest Physiology and Quality of Strawberry cv. Hongyan during Cold Storage.  
524 *Foods*, **12 (15)**.
- 525 (23) Song, D. Y., Reddy, L. V., Charalampopoulos, D. and Wee Y. J. 2019. Poly-( $\gamma$ -Glutamic  
526 Acid) Production and Optimization from Agro-Industrial Bioresources as Renewable  
527 Substrates by *Bacillus* sp. FBL-2 through Response Surface Methodology. *Biomolecules*,  
528 **9**:754-762.
- 529 (24) Sung, M. H., Park, C., Kim, C. J., Poo, H., Soda, K. and Ashiuchi, M. 2005. Natural and  
530 Edible Biopolymer Poly-  $\gamma$  - Gutamic Acid: Synthesis, Production, and Applications. *Chem*  
531 *Rec.* **5**: 352-66.
- 532 (25) Tovar, M. J., Romero, M. P., Girona, J. and Motilva, M. J. 2002. L-  
533 Phenylalanineammonia-lyase and Concentration of Phenolics in Developing Olive  
534 (*Olea europaea* L. cv. Arbequina) Fruit Grown under Different Irrigation Regimes. *J. Sci. Food*.  
535 *Agric.*, **82**: 892- 898.
- 536 (26) Wang, K., Li, T., Chen, S., Li, Y. and Rashid, A. 2020. The Biochemical and Molecular  
537 Mechanisms of Softening Inhibition by Chitosan Coating in Strawberry Fruit (*Fragaria*  
538  $\times$  *ananassa*) during Cold Storage. *Sci. Hortic.*, **271(2)**:109483.
- 539 (27) Wilson, T. C. and Hagerman, A. E. 1990. Quantitative Determination of Ellagic Acid. *J.*  
540 *Agric. Food Chem.*, **38**: 1678-1683.
- 541 (28) Williams, D. J., Edwards, D., Sharon, P. and Chaliha, M. 2014. Sultambawa. Profiling  
542 Ellagic Acid Content: The Importance of Form and Ascorbic Acid Levels. *Food Res Int.*,  
543 **66**:100-106.
- 544 (29) Yu, Y., Yan, F., Chen, Y., Jin, C., Guo, J-H. and Chai, Y. 2016. Poly- Glutamic Acids  
545 Contribute to Biofilm Formation and Plant Root Colonization in Selected Environmental  
546 Isolates of *Bacillus subtilis*. *Front. Microbiol.*, **7**: 1811.
- 547 (30) Zeng, W., Chena, G., Zhang, Y., Wuc, K. and Lianga, Z. 2012. Studies on the UV

548 Spectrum of Poly ( $\gamma$ - Glutamic Acid) Based on Development of a Simple Quantitative Method.  
549 *Int. J. Biol. Macromol.*, **51**: 83-90.

550

551 تولید پلی- $\gamma$ -گلوتامات به روش فرماتتاسیون بستر جامد و کاربرد آن در کنترل زیستی

552

553 ساره هاشمی ، مسعود احمدزاده، حسین صارمی ، سلیمان قاسمی، و آزاد عمرانی

554

555

#### چکیده

556 پلی- گاما- گلوتامیک اسید ( $\gamma$ -PGA) یک پلیمر طبیعی با کاربردهای متنوع در صنایع مختلف است. با این حال، استفاده  
557 از آن در کشاورزی به دلیل هزینه‌های بالای تولید محدود است. اهداف این مطالعه: تولید بهینه و مقرون به صرفه  $\gamma$ -  
558 PGA از طریق تخمیر حالت جامد (SSF) با استفاده از باکتری *Bacillus velezensis* UTB96، تعیین غلظت و وزن  
559 مولکولی مناسب  $\gamma$ -PGA برای کاربردهای کشاورزی، به‌ویژه در کشت توت فرنگی، و در نهایت بررسی تأثیر  $\gamma$ -PGA  
560 تولید شده بر افزایش عمر مفید میوه‌های توت‌فرنگی در طول انبارمانی است. ابتدا تولید  $\gamma$ -PGA با استفاده از SSF و  
561 باکتری *Bacillus velezensis* UTB96 مورد بررسی قرار گرفت و تأثیر عوامل فیزیکی و شیمیایی بر وزن مولکولی  $\gamma$ -  
562 PGA ارزیابی شد. بر اساس نتایج، سه وزن مولکولی مختلف  $\gamma$ -PGA شناسایی شد: 1156.43، 734.38 و 296.55  
563 کیلودالتون. این وزن‌ها برای آزمایش‌های گلخانه‌ای به منظور ارزیابی اثربخشی آن‌ها در کنترل کپک خاکستری روی گیاه  
564 توت فرنگی انتخاب شدند. نتایج نشان داد که با استفاده از ضایعات کشاورزی، از جمله کنجاله کنجد، کاه گندم و پوست  
565 موز در روش SSF، می‌توان  $\gamma$ -PGA را با نرخ ۷۰ گرم در کیلوگرم وزن خشک محیط کشت تولید کرد. تحلیل تأثیر  $\gamma$ -  
566 PGA بر کاهش بیماری کپک خاکستری نشان داد که این ترکیب می‌تواند مقاومت گیاه را بهبود بخشد. افزایش قابل توجهی  
567 در فعالیت آنزیم‌های آسکوربات پراکسیداز و فنیل‌آلانین آمونیلایز (PAL) همراه با تولید ترکیبات پلی‌فنولی مانند اسید  
568 الاژیک مشاهده شد. در نتیجه، این مکانیسم‌ها انعطاف‌پذیری و تحمل گیاه را در برابر قارچ بهبود بخشیدند و منجر به حفظ  
569 کیفیت میوه‌ها در طول نگهداری در سردخانه شدند.

570