

Effect of dietary energy source and level on the performance, antibody titers and the relative expression of *IL-2* and *IL-6* gene in broilers under heat stress

Nematollah Dayani¹, Mohammad Chamani^{1*}, Parvin Shawrang², Asa Ebrahimi³, and Ali Asghar Sadeghi¹

ABSTRACT

This study aimed to determine the effects of energy levels and sources on growth performance, antibody titers, and the gene expression of pro-inflammatory cytokines in broilers exposed to heat stress. 450 one-day-old Ross chickens were assigned to six dietary treatments and five replicates in a completely randomized design. Chickens have received diets differentiated by main energy source (corn grain and soybean oil) and energy level (equal, 3 or 6% lower or higher than Ross 308 recommendation). Treatments were as follows: corn grain and equal as control (CON), corn grain, 3% lower (T1), corn grain, 6% lower (T2), corn grain and soybean oil, equal (T3), corn grain and soybean oil, 3% higher (T4), corn grain and soybean oil, 6% higher (T5). The room temperature was increased to 34 °C (6-h daily) from day 12 to 42 of age to induce heat stress. The highest corticosterone level was observed in the T1, T2, and T5 groups. The lowest antibody titers were observed in the T2 group and the highest expression levels of pro-inflammatory cytokines genes were in chickens receiving T5 diet. The highest feed conversion ratio (FCR) during the grower and finisher periods was observed in T2, and the lowest FCR was observed in T3 and T4 groups. It was recommended to feed Ross broiler chickens with a diet containing oil instead of a part of grain based on energy recommended by the strain recommendation.

Keywords: Chicken. Corticosterone. Inflammation. Interleukin. Ross strain.

INTRODUCTION

The major broiler farms exist in subtropical and tropical regions of the world (Kpomasse *et al.*, 2021). In these regions, farmers have to use various strategies to control the temperature of their houses, to reduce the negative effects of heat stress on the health and performance of broilers (Costantino *et al.*, 2018). After exposing broilers to high ambient temperatures, some toxic

¹ Department of Animal Science, Science and Research Branch, Islamic Azad University, Tehran, Islamic Republic of Iran.

² Nuclear Agriculture Research School, Nuclear Science and Technology Research Institute, Karaj, Islamic Republic of Iran.

³ Department of Biotechnology, Science and Research Branch, Islamic Azad University, Tehran, Islamic Republic of Iran.

* Corresponding author; e-mail: , poultsai@gmail.com or m.chamani@srbiau.ac.ir

mechanisms may be induced in the body, including the generation of reactive oxygen species, which finally results in oxidative stress. Oxidative stress could affect the metabolic pathways liver and small intestine health, which reduce the nutrient digestion and absorption, and the merit of substrates for metabolism (Mancinelli *et al.*, 2023). Various management techniques, such as cooling systems, have been used to reduce the negative effects of heat stress on broiler chickens (Fisinin and Kavtarashvili, 2015). The cost of cooling broiler houses is high in many regions; hence, some researchers focused on nutritional management (Daghir, 2009). The manipulation of dietary energy level and source has been considered as a useful method in broiler farms to overcome the negative effects of heat stress (Daghir, 2009; Raghebian *et al.*, 2016). Seifi *et al.* (2018) reported that feeding a high-fat diet could improve the heat tolerance in broiler chickens, and dietary inclusion of palm oil improved the growth performance and survivability of heat stressed broiler chickens (Zulkifli *et al.*, 2007).

Moreover, Kim *et al.* (2019) reported that fat supplementation had preventative effects on weight loss for hens raised under heat stress. In contrast, Rafiei-Tari *et al.* (2021) reported that feeding oils containing n-6 fatty acids had detrimental effects on the health of broilers exposed to heat stress. On the other hand, when chickens were fed with low energy diets, deviations from physiological homeostasis occurred, leading to impaired bird welfare (Cheng and Jefferson, 2008) and significant reductions in production capabilities (Jariyahatthakij *et al.*, 2018). In an interesting study, Raghebian *et al.* (2016) reported that high energy in a broiler diet could enhance heat resistance and improve performance parameters.

Today, the effect of nutrition on gene expression is very important (Goel *et al.*, 2021). The effect of energy level and source on the expression of genes related to heat resistance has been investigated (Raghebian *et al.*, 2016), but its effect on gene expression of interleukins are related to the immune response, was not evaluated. The interleukin-2 (*IL-2*) and interleukin-6 (*IL-6*) are pro-inflammatory cytokines that play an important role in the inflammatory response in the body of broiler chickens under heat stress (Goel *et al.*, 2021) and prolong inflammation responses cause tissue damage, especially in the liver and immune system tissues (Helwig and Leon, 2011; Goel *et al.*, 2021). Finding the relationship between the level and energy source of the diet with the relative expression of genes of these two cytokines helps to understand better the cause of the effects observed in the body of broiler chickens.

In the literature, the effects of energy source and level on the antibody titers and relative expression of pro-inflammatory cytokine genes (*IL-2* and *IL-6*) in chickens under heat stress have not been completely evaluated (Ndlebe *et al.*, 2023). Taleb *et al.* (2017) reported that

antibody titers decreased in broiler chickens (Cobb 500 strain) raised under hot environmental conditions receiving soybean oil. In contrast, Sadeghi *et al.* (2013) reported that including soybean oil could enhance the immune response in broiler chickens (Ross 308). It is unclear what effects the soybean oil inclusion and the dietary energy concentration have on the expression of genes related to the immune system and the antibody titer.

It was hypothesized that in the heat stress condition, including soybean oil and formulation of high-energy diet could enhance health, immune responses, and performance compared to a diet containing the main energy source from a carbohydrate or low-energy diet. In the present study, low and high levels of dietary energy were considered factors that cause metabolic stress in the body to mimic the conditions chickens face in different breeding centers.

Therefore, the present study aimed to assess the effects of energy source and level on the growth performance, liver health, immune responses, and the relative expression of IL-2 and IL-6 genes in broiler chickens exposed to heat stress.

MATERIALS AND METHODS

Chickens Management

A total of 465 one-day-old male Ross 308 broiler chickens (average weight of 40 g) were purchased from a local hatchery and allocated randomly to thirty one floor pens (200 cm × 180 cm) covered with wood shaving. Chicks were randomly assigned to 6 dietary treatments with 5 replicates and 15 chicks per each. Except ambient temperature, chicks were raised under controlled conditions, lighting program, and feed recommendations based on Ross 308 broiler guides. Chickens (n=450) were exposed to heat stress from day 12 to 42 of age in the relative humidity of 65%. During heat stress, temperatures were raised daily to 34±1 °C for 6 hours from 08:00 to 14:00 and then decreased to 24 ± 1 °C. Fifteen chicks were kept in a room at normal temperature to assess whether experimental chicks were exposed to heat stress. These chickens received corn grain and energy density based on Ross 308 recommendation (3100 and 3200 kcal/kg during grower and finisher periods, respectively). Blood samples were taken from these chickens to measure corticosterone levels as a biological marker of heat stress. All chickens had access *ad libitum* to feed and fresh water, especially throughout the heat challenge period. Chickens were vaccinated with the Newcastle disease (ND) vaccine and infectious bursal disease (IBD) vaccine. In the experiment's initial and end, the amount of feed intake and body weight were measured, and the feed conversion ratio (FCR) was calculated. Dead chicken was weighed and the weight was included in the calculations of FCR.

Experimental Design

Dietary treatments were included in: the control group (CON), chickens receiving the main energy source from corn grain and energy density based on Ross 308 recommendation; T1: chickens receiving the main energy source from corn grain and 3% lower energy density than Ross 308 recommendation; T2: chickens receiving the main energy source from corn grain and 6% lower energy density than Ross 308 recommendation; T3: chickens receiving the main energy source from corn grain and soybean oil and energy density based on Ross 308 recommendation; T4: chickens receiving the main energy from corn grain and soybean oil and 3% higher energy density than Ross 308 recommendation, and T5: chickens receiving the main energy from corn grain and soybean oil and 6% higher energy density than Ross 308 recommendation. Metabolizable energy levels of diets were balanced using starch or washed sand. Chickens were raised at three feeding periods: starter (days 1 to 10), growers (days 11 to 24), and finishers (days 25-42) periods.

Sample Collection and Measurements

On days 24 and 42 of age, blood samples (6 mL) were collected using sterile Venoject directly from the heart of two chickens in each replicate. The serum of the blood sample was separated using a centrifuge ($1500 \times g$, 15 minutes) and stored at -20°C until further analysis. Two days after sampling, antibody titers against viruses of ND and IBD were determined in all serum samples. Biochemical measurements were done on samples taken on day 42 of age.

On day 24 of age, immediately after blood sampling, chicks were sacrificed by cervical dislocation, then the spleen and liver were removed and sampled. Five spleen samples from each treatment were collected to analyze the relative expression of the *IL-2* and *IL-6* genes. Spleen tissues were transferred in a cry-protectant tube, snap-frozen in liquid nitrogen, and stored at -70°C until RT-PCR analysis.

Blood Sample Analysis

Serum corticosterone level was measured enzymatically using an enzyme-linked immunosorbent assay kit (Enzo Life Sciences, NY, USA). Serum concentrations of glucose, total protein, albumin, creatinine, and uric acid were measured using the photometric method by autoanalyzer (BS-120 model, Minbray Co., USA) and commercial kits (Pars Azmon Co., Tehran, Iran).

Serology

The titers of antibodies against Newcastle disease virus were measured by hemagglutination-inhibition test (Allan and Gough, 1974) and against Infectious Bursal Disease virus by ELISA kit, IDEXX FlockChek standard (IDEXX Corporation, Westbrook, ME, USA). The value of antibody titers was transformed to $\log_2(x)$ before statistical analysis.

Analysis of the Gene Expression of *IL-2* and *IL-6*

The relative abundances of *IL-2* and *IL-6* mRNA were determined by the RT-PCR technique described by Paraskeuas and Mountzouris (2019) and Long *et al.* (2011). The frozen spleen sample was crushed in a sterile mortar, and the powder was applied for total RNA extraction using a suitable kit (Bioneer Co., Seoul, South Korea). Then, each gene's cDNA was synthesized using a suitable kit using the reverse transcription technique (Bioneer Co., Seoul, South Korea). Quantitative PCR was performed with specific primer pairs for *IL-2* (Paraskeuas and Mountzouris, 2019) and *IL-6* (Long *et al.*, 2011) using Quanti Fast SYBER Green PCR kit (QIAGEN, Cat. No. 204052). GAPDH was chosen as a housekeeping gene. The relative gene expression of *IL-2* and *IL-6* as target genes was normalized to the GAPDH gene using method as previously described by Livak and Schmittgen (2001). Quantification for each treatment group was performed in triplicates.

Statistical Analysis

Statistical analyses were done using the General Linear Model procedure of the SAS for Windows version 9.1 (SAS Institute Inc., Cary, NC) appropriate for a completely randomized design. To evaluate the normal distribution of data, the Kolmogorov-Smirnov test was done. Duncan multiple range tests were used to compare the means. Effects between the control and experimental groups were considered significant when $P < 0.05$.

RESULTS

Effect on Serum Corticosterone Level and Biochemical Measurements

Table 1 shows the serum corticosterone levels and biochemical parameters of broilers receiving different dietary energy levels and sources. The highest corticosterone level was observed in the T1, T2, and T5 groups, and no difference was observed among other treatments with the control group. Broilers in the T5 group had the highest serum glucose level, and those in the T2 group had the lowest. Broilers receiving the T4 diet had the highest albumin, globulin, and total protein, and broilers in the T2 group had the lowest protein sections. The highest concentration

of creatinine and uric acid was observed in the T2 group, and broilers of CON, T3, and T4 had the lowest creatinine and uric acid concentrations.

Effect on Antibody Titers and the Relative Expression Levels of Interleukins

Table 2 shows the effect of dietary energy level and source on antibody titers against Newcastle disease virus and infectious bursa disease virus determined on days 24 and 42 of age. There was no difference among treatments for ND titer at day 24, but differences were observed for ND titer at day 42. At day 42 of age, the lowest ND titer was observed in the T2 group and the highest in the T3 and T4 groups. At day 24 of age, the IBD titer was the highest in the T3 and T4 groups and the lowest in the T2 group. At day 42 of age, the highest IBD titer was observed in the CON, T3, T4, and T5 groups, and the lowest titer was found in T1 and T2 groups.

Figure 1 shows the relative expression level of IL-2 and IL-6 genes in the spleen of broiler chickens under heat stress receiving different energy densities and sources. Significant differences were observed among treatments for the relative expression of IL-2 and IL-6 genes. The highest relative expression of the IL-2 gene was observed in chickens receiving T5 and then in T4 diets, and the lowest expression was observed in chickens receiving CON, T1, T2, and T3 diets. Chickens receiving the T5 diet had the highest relative gene expression of IL-6, and the lowest expression was observed in chickens receiving CON, T1, and T2 diets.

Effect on Performance Parameters

The performance parameters of broiler chickens are presented in Table 3. There were no differences among treatments for performance parameters during the starter period. Differences appeared among treatments for performance parameters at grower and finisher periods. The daily gain of broilers in the T3 and T4 groups was higher than in T1 and T2 groups. The lowest daily gain during grower and finisher periods was observed in the T2 group and the highest daily gain was observed in broilers receiving T3 diet. Feed intake of broilers during grower and finisher periods was the highest in the T2 group, and there was no difference in feed intake among other treatments. The highest FCR during the grower and finisher periods was observed in T2, and the lowest FCR was observed in the T3 and T4 groups. Broilers in the T5 group had the same FCR as T3 and T4 groups.

DISCUSSION

In the present study, serum corticosterone levels were high in broilers receiving the control and experimental diets (heat stress condition) compared to the level of corticosterone in chickens

kept in normal temperature conditions (6.78 ng/ml). Consistent with our results, previous studies have reported that acute heat stress elevates corticosterone levels in the serum of broiler chickens (Quinteiro-Filho *et al.*, 2010; Soleimani *et al.*, 2011). In contrast to our findings, broilers' exposure to heat stress has not been shown to influence serum corticosterone levels (Mack *et al.*, 2013; Xie *et al.*, 2015). Possible reasons for the discrepancies among the results of various studies might be differences in temperature and humidity set, time of blood sampling, and chicken genotypes. Increases in corticosterone levels in broilers' serum are linked to the hypothalamic-pituitary-adrenal (HPA) axis. The HPA axis controls the adaptability of broilers in response to various stressors (He *et al.*, 2018).

In the present study, broilers receiving diets with energy restriction (T1 and T2) showed higher corticosterone levels than the control group. In stressful conditions, a change in the energy density of the diet causes additional metabolic stress in the body of chicks and may increase the generation of free radicals (Raghebian *et al.*, 2017; He *et al.*, 2018). Based on reports (Emami *et al.*, 2021, Rafiei-Tari *et al.*, 2021), chickens receiving a diet with restricted energy experience higher protein turnover, and those receiving a diet with surplus energy experience higher metabolic rate, which both processes increase the heat production and expose body to intense heat stress. Chicken receiving diets containing soybean oil (T4 and T5) had no difference in corticosterone concentration compared to the control group. Also, chicken in T3 and T4, which received soybean oil instead of a part of starch from corn grain, showed lower corticosterone levels than T1 and T2, which may be related to lower heat increment. In a previous study (Sadeghi *et al.*, 2013), a shift from starch to lipid during heat stress decreased heat increment. Many researchers recommend replacing soybean oil with starch (Yaqoob, 2004; Cherian, 2015) to reduce the heat increment and the negative effects of heat stress on the animal body.

The serum glucose level of chickens in heat stress was higher than those raised in normal conditions (185 mg/ml), which might be an adaptation for survivability and tolerance. In agreement with our finding, Bogin *et al.* (1996) reported that chickens that survived under intense heat stress had higher blood glucose levels than the non-surviving chickens. The reductions in serum albumin, globulin, and total protein levels in the chicken receiving low dietary energy (T1 and T2) and high dietary energy (T5) compared to the control group can be linked to elevation of serum corticosterone levels. Corticosterone can change metabolic pathways, reduce protein synthesis (Sadeghi *et al.*, 2013), and increases the catabolism of proteins to use as fuel in broilers receiving low dietary energy (Kitaysky *et al.*, 1999). In broilers receiving high dietary energy (T5), reduced total protein in the serum may be linked to liver

inflammation. The result of a previous study (Ozbey *et al.*, 2004) is consistent with the reductions observed in our study after the heat challenge.

The marked increase in the serum uric acid of broilers receiving low dietary energy (T2) may be linked to an increase of protein turnover and, in those receiving surplus energy (T5), linked to liver inflammation and oxidative stress. Previous studies reported increases (Ozbey *et al.*, 2004), reductions (Bogin *et al.*, 1996), and no alteration (Xie *et al.*, 2014) in the serum levels of uric acid after heat stress and energy restriction or surplus. The discrepancies in responses among various studies may be related to the differences in metabolic rates and physiological states and also signify protein catabolism for energy generation in energy-restricted birds resulting from increased corticosterone levels (Vandana *et al.*, 2021).

Chicken receiving T1, T3, T4, and T5 had the same performance parameters, but the chickens in the T2 group had higher feed intake and FCR than the control group. To compensate for the energy dilution of the diet, chickens receiving the T2 diet try to feed more. As feed intake increased, the activity of eating and the digestive tract increased, resulting in increased heat production (Herd and Arthur, 2009). In the heat stress condition, heat dissipating from the body decreases, and the animal body is exposed to oxidative stress (Teeter and Belay, 1996). Chickens exposed to oxidative stress could not grow perfectly and showed a lower feed conversion ratio than CON, T3 and T4 groups. In consistent with our finding, Classen (2017) and Azizi *et al.* (2021) reported that chickens increased feed intake in response to dietary energy dilution. However, Yuan *et al.* (2008) reported that the weight gain of chickens was not altered by dietary energy level, which contrasts with our results.

Antibody titers against ND and IBD were the highest in the T3 and T4 diet formulated with soybean oil and the lowest in the T1 and T2 diet formulated with low energy density, which is inconsistent with the findings of Taleb *et al.* (2017). They reported that an increase in soybean oil level in the Cobb strain diet resulted in lower antibody titers against ND and IBD. In the current study, an increase in the level of soybean oil in the Ross broiler diet had no negative effect on the antibody titers against ND and IBD.

In broilers receiving, low-energy diets, the effect of metabolic stress caused by energy level on the high corticosterone level and low blood glucose level may play an important role in reducing the immune response and antibody production (Yang *et al.*, 2015). The factors above cause disturbances in the process of growth and maturation of T and B cells in primary and secondary lymphoid tissues, which ultimately causes numerous immune abnormalities in broiler chickens (Hirakawa *et al.*, 2020).

Pro-inflammatory cytokines such as *IL-2* and *IL-6* have been found to play an active role in the inflammatory response under stressful conditions (Helwig and Leon, 2011). In the literature, limited information exists concerning energy level's effect on the gene expression of pro-inflammatory cytokines. A striking finding in the present study was the low expression of genes involved in inflammation in broiler chickens' diets with low energy density (T1 and T2). This finding agrees with some studies (Trayhurn and Wood, 2004; Higami *et al.*, 2006), which reported low energy diet resulted in low inflammation and gene expression of pro-inflammatory cytokines in laboratory animals. In contrast, high-energy diets (T4 and T5) increased the relative gene expression of *IL-2* and *IL-6*. In T4 and T5 groups, high corticosterone levels may be influenced by the expression of pro-inflammatory cytokines as it could increase the proliferation of lymphocytes and macrophages (Hirakawa *et al.*, 2020; Goel *et al.*, 2021). In energy-dense diets, soybean oil is included, and higher expression of these genes may be related to oil inclusion. Previous studies (Mu *et al.*, 2018) revealed that dietary soybean oil significantly increased the gene expression of pro-inflammatory cytokines. The results observed for fasting glucose level and *IL-2* gene expression in the present study are consistence with the finding of Kochumon *et al.* (2020), who reported that the level of *IL-2* expression was associated positively with fasting blood glucose.

CONCLUSIONS

The results of the present study indicate that energy restriction and surplus negatively affect the immune response and performance of chickens raised under heat stress. Surplus energy negatively affects the relative expression levels of pro-inflammatory cytokines genes (*IL-2* and *IL-6*), and energy restriction results in higher protein catabolism (higher uric acid and creatinine), which reduces broiler performance and immune responses. The inclusion of soybean oil in the diet positively affected immune response and performance. It was recommended to feed Ross broiler chickens under heat stress with a diet containing oil instead of a part of grain based on the energy recommended by the strain recommendation.

References

- Aami Azghadi, M., Kermanshahi, H. and Golian, A. 2014. The effect of dietary energy and protein levels on growth performance and antibody responses of offspring of laying Japanese quails. *Iran. J. Appl. Anim. Sci.*, **4**(1): 185-190.

- Azizi, B., Sadeghi, G., Karimi, A., and Abed, F. 2011. Effects of dietary energy and protein dilution and time of feed replacement from starter to grower on broiler chickens performance. *J. Central Europ. Agri.*, **12(1)**: 44-52.
- Bogin, E., Avidar, Y., Pech-Waffenschmidt, V., Doron, Y., Israeli, B. A., and Kevkhayev, E. 1996. The relationship between heat stress, survivability and blood composition of the domestic chicken. *European J. Clinical Chem. Clinical Biochem.*, **34(6)**: 463-469.
- Cheng, H. W., and Jefferson, L. 2008. Different behavioral and physiological responses in two genetic lines of laying hens after transportation. *Poult. Sci.*, **87(5)**: 885-892.
- Cherian, G. 2015. Nutrition and metabolism in poultry: role of lipids in early diet. *J. Anim. Sci. Biotechnol.*, **6(1)**: 1-9.
- Classen, H. L. 2017. Diet energy and feed intake in chickens. *Anim. Feed Sci. Technol.*, **233**: 13-21.
- Costantino, A., Fabrizio, E., Ghiggini, A., and Bariani, M. 2018. Climate control in broiler houses: A thermal model for the calculation of the energy use and indoor environmental conditions. *Energy and Buildings*, **169**: 110-126.
- Daghir, N. J. 2009. Nutritional strategies to reduce heat stress in broilers and broiler breeders. *Lohmann Inform.*, **44(1)**: 6-15.
- Emami, N. K., Greene, E. S., Kogut, M. H., and Dridi, S. 2021. Heat stress and feed restriction distinctly affect performance, carcass and meat yield, intestinal integrity, and inflammatory (chemo) cytokines in broiler chickens. *Front. Physiol.*, **12**:707757.
- Fisinin, V. I., and Kavtarashvili, A. S. 2015. Heat stress in poultry. II. Methods and techniques for prevention and alleviation. *Agri. Biol.*, **50(4)**: 431-443.
- Goel, A., Ncho, C. M., and Choi, Y. H. 2021. Regulation of gene expression in chickens by heat stress. *J. Anim. Sci. Biotechnol.*, **12(1)**: 1-13.
- He, S.P., Arowolo, M.A., Medrano, R.F., Li, S., Yu, Q.F., Chen, J.Y. and He, J.H. 2018. Impact of heat stress and nutritional interventions on poultry production. *World's Poult. Sci. J.*, **74(4)**: 647-664.
- Helwig, B.G., and Leon, L.R. 2011. Tissue and circulating expression of IL-1 family members following heat stroke. *Physiol. Genom.*, **43(19)**: 1096-1104.
- Herd, R.M., and Arthur, P.F. 2009. Physiological basis for residual feed intake. *J. Anim. Sci.*, **87(suppl_14)**: E64-E71.
- Higami, Y., Barger, J. L., Page, G. P., Allison, D. B., Smith, S. R., Prolla, T. A., and Weindruch, R. 2006. Energy restriction lowers the expression of genes linked to

inflammation, the cytoskeleton, the extracellular matrix, and angiogenesis in mouse adipose tissue. *J. Nut.*, **136(2)**: 343-352.

Hirakawa, R., Nurjanah, S., Furukawa, K., Murai, A., Kikusato, M., Nochi, T., and Toyomizu, M. 2020. Heat stress causes immune abnormalities via massive damage to effect proliferation and differentiation of lymphocytes in broiler chickens. *Front. Vet. Sci.*, **7**: 46.

Jariyahatthakij, P., Chomtee, B., Poeikhampha, T., Loongyai, W., and Bunchasak, C. 2018. Effects of adding methionine in low-protein diet and subsequently fed low-energy diet on productive performance, blood chemical profile, and lipid metabolism-related gene expression of broiler chickens. *Poult. Sci.*, **97(6)**: 2021-2033.

Kim, J.H., Lee, H.K., Yang, T.S., Kang, H.K., and Kil, D.Y. 2019. Effect of different sources and inclusion levels of dietary fat on productive performance and egg quality in laying hens raised under hot environmental conditions. *Asian-Austral. J. Anim. Sci.*, **32(9)**:1407-1412.

Kitaysky, A.S., Piatt, J. F., Wingfield, J. C., and Romano, M. 1999. The adrenocortical stress-response of black-legged kittiwake chicks in relation to dietary restrictions. *J. Comp. Physiol. B*, **169**: 303-310.

Kochumon, S., Al Madhoun, A., Al-Rashed, F., Thomas, R., Sindhu, S., Al-Ozairi, E. and Ahmad, R. (2020). Elevated adipose tissue associated *IL-2* expression in obesity correlates with metabolic inflammation and insulin resistance. *Scientific Reports*, **10(1)**: 16364.

Kpomasse, C. C., Oke, O. E., Houndonougbo, F. M., and Tona, K. 2021. Broiler production challenges in the tropics: A review. *Vet. Med. Sci.*, **7(3)**: 831-842.

Long, G. L., Hao, W. X., Bao, L. F., Li, J. H., Zhang, Y., and Li, G. H. 2019. Effects of dietary inclusion levels of palm oil on growth performance, antioxidative status and serum cytokines of broiler chickens. *J. Anim. Physiol. Anim. Nut.*, **103(4)**: 1116-1124.

Mack, L. A., Felver-Gant, J. N., Dennis, R. L., and Cheng, H. W. 2013. Genetic variations alter production and behavioral responses following heat stress in 2 strains of laying hens. *Poult. Sci.*, **92(2)**: 285-294.

Mancinelli, A. C., Baldi, G., Soglia, F., Mattioli, S., Sirri, F., Petracci, M., and Zampiga, M. 2023. Impact of chronic heat stress on behavior, oxidative status and meat quality traits of fast-growing broiler chickens. *Front. Physiol.*, **14**: 1-8.

Mu, H., Shen, H., Liu, J., Xie, F., Zhang, W., and Mai, K. 2018. High level of dietary soybean oil depresses the growth and anti-oxidative capacity and induces inflammatory response in large yellow croaker *Larimichthys crocea*. *Fish Shellfish Immunol.*, **77**: 465-473.

- Ndlebe, L., Tyler, N. C., and Ciacchiariello, M. 2023. Effect of varying levels of dietary energy and protein on broiler performance: a review. *World's Poult. Sci. J.*, **79(3)**: 449-465.
- Nikraves-Masouleh, T., Seidavi, A. R., Kawka, M. and Dadashbeiki, M. 2018. The effect of dietary energy and protein levels on body weight, size, and microflora of ostrich chicks. *Trop. Anim. Health Product.*, **50(3)**: 635-641.
- Özbey, O., Yıldız, N., Aysöndü, M. H., and Özmen, Ö. 2004. The effects of high temperature on blood serum parameters and the egg productivity characteristics of Japanese quails (*Coturnix coturnix japonica*). *Inter. J. Poult. Sci.*, **3 (7)**: 485-489.
- Paraskeuas, V.V., and Mountzouris, K.C. 2019. Modulation of broiler gut microbiota and gene expression of Toll-like receptors and tight junction proteins by diet type and inclusion of phytogenics. *Poult. Sci.*, **98(5)**: 2220-2230.
- Rafiei-Tari, A., Sadeghi, A. A., and Mousavi, S. N. 2021. Inclusion of vegetable oils in diets of broiler chicken raised in hot weather and effects on antioxidant capacity, lipid components in the blood and immune responses. *Acta Scient.. Anim. Sci.*, **43**: 1-6.
- Ragheblian, M., Sadeghi, A. A., and Aminafshar, M. 2017. Impact of dietary energy density on the liver health of broilers exposed to heat stress and their performance during finisher period. *J. Livestock Sci.*, **8**: 122-130.
- Ragheblian, M., Sadeghi, A. A., and Aminafshar, M. 2016. Energy sources and levels influenced on performance parameters, thyroid hormones, and HSP70 gene expression of broiler chickens under heat stress. *Trop. Anim. Health Product.*, **48**: 1697-1702.
- Sadeghi, A. A., Mirmohseni, M., Shawrang, P., and Aminafshar, M. 2013. The effect of soy oil addition to the diet of broiler chicks on the immune response. *Turk. J. Vet. Anim. Sci.*, **37(3)**: 264-270.
- Seifi, K., Rezaei, M., Yansari, A. T., Riazi, G. H., Zamiri, M. J., and Heidari, R. 2018. Saturated fatty acids may ameliorate environmental heat stress in broiler birds by affecting mitochondrial energetics and related genes. *J. Thermal Biol.*, **78**: 1-9.
- Soleimani, A. F., Zulkifli, I., Omar, A. R., and Raha, A. R. 2011. Physiological responses of 3 chicken breeds to acute heat stress. *Poult. Sci.*, **90(7)**: 1435-1440.
- Taleb, Z., Sadeghi, A.A., Shawrang, P., Chamani, M. and Aminafshar, M. 2017. Effect of energy levels and sources on the blood attributes and immune response in broiler chickens exposed to heat stress. *J. Livestock Sci.*, **8**: 123-144.
- Trayhurn, P., and Wood, I. S. 2004. Adipokines: inflammation and the pleiotropic role of white adipose tissue. *Brit. J. Nut.*, **92(3)**: 347-355.

- Teeter, R. G., and Belay, T. 1996. Broiler management during acute heat stress. *Anim. Feed S. Technol.*, **58(1-2)**: 127-142.
- Quinteiro-Filho, W. M., Ribeiro, A., Ferraz-de-Paula, V., Pinheiro, M. L., Sakai, M., Sá, L. R. M. D. and Palermo-Neto, J. 2010. Heat stress impairs performance parameters, induces intestinal injury, and decreases macrophage activity in broiler chickens. *Poult. Sci.*, **89(9)**: 1905-1914.
- Vandana, G. D., Sejian, V., Lees, A. M., Pragna, P., Silpa, M. V., and Maloney, S. K. 2021. Heat stress and poultry production: impact and amelioration. *Inter. J. Biometeorol.*, **65**: 163-179.
- Xie, J., Tang, L., Lu, L., Zhang, L., Lin, X., Liu, H. C. and Luo, X. 2015. Effects of acute and chronic heat stress on plasma metabolites, hormones and oxidant status in restrictedly fed broiler breeders. *Poult. Sci.*, **94(7)**: 1635-1644.
- Yang, J., Liu, L., Sheikahmadi, A., Wang, Y., Li, C., Jiao, H. and Song, Z. 2015. Effects of corticosterone and dietary energy on immune function of broiler chickens. *PLoS One*, **10(3)**: e0119750.
- Yaqoob, P. 2004. Fatty acids and the immune system: from basic science to clinical applications. *Proceed. Nut. Society*, **63(1)**: 89-105.
- Yuan, L., Lin, H., Jiang, K. J., Jiao, H. C., and Song, Z. G. 2008. Corticosterone administration and high-energy feed results in enhanced fat accumulation and insulin resistance in broiler chickens. *Brit. Poult. Sci.*, **49(4)**: 487-495.
- Zulkifli, I., Siegel, H. S., Mashaly, M. M., Dunnington, E. A., and Siegel, P. B. 1995. Inhibition of adrenal steroidogenesis, neonatal feed restriction, and pituitary-adrenal axis response to subsequent fasting in chickens. *General Comparat. Endocrinol.*, **97(1)**: 49-56.
- Zulkifli, I., Htin, N. N., Alimon, A. R., Loh, T. C., and Hair-Bejo, M. 2007. Dietary selection of fat by heat-stressed broiler chickens. *Asian-Austral. J. Anim. Sci.*, **20(2)**: 245-251.

Table 1: Effect of dietary energy level and source on serum corticosterone level and biochemical parameters in broiler chickens under heat stress.

Item	CON	T1	T2	T3	T4	T5	SEM	P value
Corticosterone (ng/ml)	20.61 ^b	29.84 ^a	30.37 ^a	21.26 ^b	22.54 ^b	27.33 ^a	1.03	0.01
Glucose (mg/dl)	200.09 ^{bc}	195.12 ^{cd}	189.45 ^d	204.72 ^{bc}	210.58 ^{ab}	219.21 ^a	3.50	0.01
Albumin (mg/dl)	1.13 ^b	1.13 ^b	1.02 ^c	1.15 ^{ab}	1.23 ^a	1.07 ^{bc}	0.027	0.02
Globulin (mg/dl)	1.35 ^b	1.27 ^c	1.17 ^d	1.33 ^b	1.51 ^a	1.17 ^d	0.021	0.03
Total protein (mg/dl)	2.48 ^b	2.40 ^b	2.19 ^c	2.49 ^b	2.73 ^a	2.24 ^c	0.039	0.01
Creatinine (mg/dl)	2.54 ^c	2.73 ^{bc}	3.74 ^a	2.53 ^c	2.51 ^c	3.01 ^b	0.103	0.02
Uric acid (mg/dl)	4.22 ^c	4.58 ^b	4.94 ^a	4.20 ^c	4.24 ^c	4.83 ^a	0.023	0.01

^{a, b, c, d} Means within a row with different superscripts are significantly different ($P < 0.05$).

*CON: control, energy based on Ross standard diet with main energy from corn; T1: chickens receiving 3% lesser energy than Ross standard diet with energy from corn; T2: chickens received 6% lesser energy than Ross standard diet with main energy from corn; T3: chicken receiving Ross standard diet with main energy from corn grain and soybean oil, T4: chicken receiving 3% upper energy than Ross standard diet with main energy from corn and soybean oil and T5: chicken receiving 6% upper energy than Ross standard diet with main energy from corn and soybean oil.


Table 2: Effect of dietary energy level and source on antibody titers against viruses of Newcastle disease (ND) and infectious bursal disease (IBD) at days 24 and 42 of age.

Item*	CON	T1	T2	T3	T4	T5	SEM	P value
Day 24 of age								
ND (log 2)	4.67	4.35	4.00	3.65	3.34	4.00	0.45	0.369
IBD (log 2)	419.31 ^{ab}	414.25 ^{ab}	402.56 ^b	434.97 ^a	432.08 ^a	416.98 ^{ab}	9.20	0.016
Day 42 of age								
ND (log 2)	5.35 ^{abc}	5.09 ^{bc}	4.67 ^c	7.00 ^a	7.15 ^a	6.65 ^{ab}	0.55	0.021
IBD (log 2)	3177.30 ^a	3007.60 ^b	3017.09 ^b	3106.02 ^a	3187.72 ^a	3125.34 ^a	50.6	0.050


* ND: Newcastle disease virus; IBD: Infectious Bursal Disease virus

^{a, b, c} Means within a row with different superscripts are significantly different ($P < 0.05$).

*CON: control, energy based on Ross standard diet with main energy from corn; T1: chickens receiving 3% lesser energy than Ross standard diet with energy from corn; T2: chickens received 6% lesser energy than Ross standard diet with main energy from corn; T3: chicken receiving Ross standard diet with main energy from corn grain and soybean oil, T4: chicken receiving 3% upper energy than Ross standard diet with main energy from corn and soybean oil and T5: chicken receiving 6% upper energy than Ross standard diet with main energy from corn and soybean oil.



A



B

Figure 1. The relative expression level of *IL-2* (A) and *IL-6* (B) genes in heat-stressed broilers receiving different energy level and source

*CON: control, energy based on Ross standard diet with main energy from corn; T1: chickens receiving 3% lesser energy than Ross standard diet with energy from corn; T2: chickens received 6% lesser energy than Ross standard diet with main energy from corn; T3: chicken receiving Ross standard diet with main energy from corn grain and soybean oil, T4: chicken receiving 3% upper energy than Ross standard diet with main energy from corn and soybean oil and T5: chicken receiving 6% upper energy than Ross standard diet with main energy from corn and soybean oil.

Table 3: Effect of dietary energy level and source on performance parameters of broiler chickens under heat stress.

Item	CON	T1	T2	T3	T4	T5	SEM	P value
Starter phase								
Gain (g/d)	17.21	17.42	17.15	17.02	17.00	17.06	0.209	0.70
Feed intake (g/d)	24.10	24.30	24.20	24.00	23.90	23.90	0.161	0.29
FCR	1.40	1.41	1.41	1.41	1.40	1.40	0.023	0.98
Grower phase								
Gain (g/d)	51.05 ^{ab}	48.22 ^{bc}	46.41 ^c	52.05 ^a	53.12 ^a	50.72 ^{ab}	0.681	0.01
Feed intake (g/d)	87.95 ^{ab}	89.67 ^{ab}	90.45 ^a	85.47 ^b	85.32 ^b	86.32 ^{ab}	1.301	0.01
FCR	1.72 ^{bc}	1.86 ^{ab}	1.95 ^a	1.64 ^{bc}	1.60 ^c	1.70 ^{bc}	0.044	0.01
Finisher								
Gain (g/d)	80.41 ^{bc}	77.52 ^{cd}	73.67 ^d	87.37 ^a	85.75 ^{ab}	83.87 ^{ab}	0.985	0.01
Feed intake (g/d)	160.21 ^b	163.63 ^b	172.12 ^a	161.71 ^b	160.80 ^b	160.3 ^b	1.53	0.03
FCR	1.99 ^{bc}	2.11 ^b	2.33 ^a	1.85 ^d	1.87 ^{cd}	1.91 ^{cd}	0.033	0.01

^{a, b, c} Means within a row with different superscripts are significantly different ($P < 0.05$).

*CON: control, energy based on Ross standard diet with main energy from corn; T1: chickens receiving 3% lesser energy than Ross standard diet with energy from corn; T2: chickens received 6% lesser energy than Ross standard diet with main energy from corn; T3: chicken receiving Ross standard diet with main energy from corn grain and soybean oil, T4: chicken receiving 3% upper energy than Ross standard diet with main energy from corn and soybean oil and T5: chicken receiving 6% upper energy than Ross standard diet with main energy from corn and soybean oil.

اثر منبع و سطح انرژی جیره بر عملکرد، تیترا آنتی بادی و بیان نسبی ژن های استرلوکین 2 و 6 در جوجه های گوشتی تحت تنش گرمایی

نعمت الله دینانی، محمد چمنی، پروین شورنگ، آسا ابراهیمی، و علی اصغر صادقی

چکیده

این مطالعه با هدف تعیین تأثیر سطوح و نوع منبع انرژی بر عملکرد رشد، تیترا آنتی بادی و بیان ژن های سیتوکین های پیش التهابی در جوجه های گوشتی در معرض تنش گرمایی انجام شد. 450 قطعه جوجه راس یک روزه در قالب طرح کاملاً تصادفی در شش جیره آزمایشی و پنج تکرار قرار گرفتند. جوجه ها جیره های متمایز شده بر اساس منبع اصلی انرژی (دانه ذرت و روغن سویا) و سطح انرژی (برابر، 3 یا 6 درصد کمتر یا بالاتر از توصیه Ross 308) دریافت کرده اند. تیمارها به شرح زیر بود: دانه ذرت و برابر با شاهد (CON)، دانه ذرت، 3 درصد کمتر (T1)، دانه ذرت، 6 درصد کمتر (T2)، دانه ذرت و روغن سویا، برابر (T3)، دانه ذرت و روغن سویا، 3 درصد بیشتر (T4)، دانه ذرت و روغن سویا، 6 درصد بیشتر (T5). دمای سالن از روز 12 تا 42 درجه پرورش به 34 درجه سلسیوس (6 ساعت در روز) افزایش یافت تا تنش گرمایی ایجاد شود. بالاترین سطح کورتیکوسترون در گروه های T1، T2 و T5 مشاهده شد. کمترین تیترا آنتی بادی در گروه T2 و بالاترین سطح بیان ژن های سیتوکین های پیش التهابی در جوجه های دریافت کننده جیره T5 مشاهده شد. بیشترین ضریب تبدیل خوراک (FCR) در طول دوره رشد و پایانی در T2 و کمترین FCR در گروه

515 T3 و T4 مشاهده شد. تغذیه جوجه های گوشتی راس با جیره غذایی حاوی روغن به جای بخشی از غلات بر اساس
516 انرژی پیشنهاد شده در کاتالوگ سویه توصیه می شود.
517