

Research Note: Gill Histopathological Lesions of Common Carp in Exposed to Chlorpyrifos and Salinity

Mona Hasankhany^{1*}, Tina Hasankhani², Seyed Ali Akbar Hedayati¹, and Mohammad Mazandarani¹

ABSTRACT

The effect of poison to salinity were studied on gill tissue of common carp (*C. carpio*). Based on this, 250 common carp fry with an average weight of 21 ± 2 g were distributed in four treatments, each with three repetitions, including salinities of zero, 4, 8, and 12 ppt. After 7 days, one group was placed for 4 days in exposure to poison with an acute concentration of 150 ppm chlorpyrifos with the formulation of 40.8% EC, and the second group was placed for 7 d in the sub-acute concentration of 15 ppm chlorpyrifos. Histopathology of Gill tissue showed that the poison and salinity had such injuries as epithelial hypertrophy, lamellar aneurism, secondary connecting adjacent blades, distal hyperplasia, epithelial lifting, leukocyte infelt, and hyperplasia. Gill histopathological result showed some lesions at high concentrations as epithelial hypertrophy, distal hyperplasia and lamellar fusion, however at low concentrations lamellar aneurism, epithelial lifting and leukocyte infelt were recorded. Therefore, these pathological indicators can be used as biomarkers.

Keywords: Agriculture poison, Histological lesions, Toxicology.

INTRODUCTION

Chlorpyrifos, Dursban, Imidacloprid and Confidor insecticides are widely used in Iran (Shafiei *et al.*, 2023). Chlorpyrifos belongs to a group of organophosphorus pesticides, which exhibit a wide spectrum of biological activity (Perry *et al.*, 2020). Chlorpyrifos is the common name for the chemical 0,0-diethyl 0-(3,5,6-trichloro-2-pyridinyl)-phosphorothioate (Figure 1). Application of Chlorpyrifos for agricultural purposes results in its dispersion into various environmental components such as air, soil, rivers and lakes, disrupting ecosystem functioning (Mackay *et al.*, 2014). When pesticides enter water bodies, they can exert direct toxic effects on fish, resulting in impaired reproduction, loss of balance, impaired growth, disruptions of

physiological, convulsions and mortality (Esbaugh *et al.*, 2018). Gills are in contact with the external environment, they are directly affected by many stress factors and toxic substances in aquatic areas (Bury *et al.*, 2014). Increasing evidence indicates that fish gills are excellent biomarkers for analyzing the impact of contaminants in aquatic ecosystems due to their high permeability and large area of water exposure. Gill surfaces make up 50% of the total surface in fish. When gills are affected by pollutants in water, normal physiological activities such as respiration, excretion, and ion transport are inevitably affected, impacting fish survival and even causing death. Additionally, pesticides can bio-accumulate within the tissues of fish. As they move up the food chain, predatory fish such as larger species

¹ Faculty of Fisheries and Environmental Sciences, Gorgan University of Natural Resources and Agricultural Sciences, Gorgan, Islamic Republic of Iran.

² Department of Immunology and Microbiology, Mycology Research Center, Faculty of Veterinary Medicine, University of Tehran, Tehran, Islamic Republic of Iran.

*Corresponding's author, e-mail: mna.hasankhany@gmail.com




Figure 1. Biochemical structure of chlorpyrifos (Cocker *et al.*, 2002).

can accumulate significant levels of these chemicals, posing a threat to both their own health and that of species (Rohani, 2023).

Common carp (*C. carpio*) is commercially an important species and cultured throughout the world. It belongs to Cyprinidae family and distributed in all watersheds. *C. carpio* are important components of freshwater ecosystems and represent one of the major sources of nutrients for people in Asia, especially China. *C. carpio* is relatively insensitive and can survive and accumulate contaminants at heavily polluted sites, which is why it is used as a bio-indicator in environmental toxicology (Sanoesi *et al.*, 2020). Therefore, our aim was to detect the remarkable histopathological alterations in gills of *C. carpio* exposed to chlorpyrifos and salinity.

MATERIALS AND METHODS

Materials

This research was conducted in Shahid Fazli Aquaculture Hall, Faculty of Fisheries, Gorgan University of Agricultural Sciences and Natural Resources. Two hundred and fifty common carps, ranging 21 ± 2 g in weight, were transferred from the rearing place of the private center to Veniro. Common carp were divided into four treatments, conducting three phases of trials with each phase consisting of 84 fish in three replicates: It is important to note that the fish did not receive any poison through-out the first phase. The only variable factor was

different concentrations of salinity i.e. 0, 4, 8 and 12 ppt, for 7 days. Salinity treatments include 0, 4, 8 and 12 ppt, for treatments 1, 2, 3, and 4 respectively, were exposed to 15 ppm chlorpyrifos for 7 days. For the next phase of the experiment; salinity 0, 4, 8 and 12 ppt, were placed for 4 days in acute expose of 150 ppm chlorpyrifos. Fish were subjected. Fish were subjected to a one-week period of acclimatization prior to the start of the experiment. Fish were fed with commercial pellets twice daily at 3% body weight (Hasankhani *et al.*, 2023). Feeding was discontinued 24 hours prior to the test, and the water was changed every 24 hours from the prepared stocks. Physicochemical conditions were monitored daily. The water temperature was $21 \pm 1^\circ\text{C}$ and pH was kept at 7.6 ± 1 . No mortality was observed during this period.

Salty Caspian Sea Water

The sea water salinity from the shores of Port Turkmen was checked with a salinometer (Atago Refractometer, Japan) (Ataimehr *et al.*, 2009) to match the salinity intended for the experiment (Moustakas *et al.*, 2004). Then, juvenile fish were transferred to tanks individually and were exposed to 15 and 150 ppm of chlorpyrifos at varying salt concentration of 0, 4, 8 and 12 ppt.

Effective physicochemical parameters of water including pH, dissolved oxygen, and temperature were recorded daily (Huyben *et al.*, 2018).

Determination of Acute and Sub-Acute LC50 Toxicity Concentrations

In this study, the statically acute (150 ppm) and sub-acute (15 ppm) toxicity (Hasankhany *et al.*, 2020) of chlorpyrifos on *C. carpio* was determined according to standard methods OECD in 4 and 7 days (Hedayati *et al.*, 2013; Gao *et al.*, 2020).

Histopathological Examination

The second gill arches from opercula cavities were harvested. Gills from each fish were fixed in 10% neutral buffered formalin. The tissues were then dehydrated in graded series of ethanol, embedded in paraffin and sectioned at 5 μm , and were stained with (H&E) stains (Hedayati *et al.*, 2013). The slides were observed under a light microscope at 40X magnification and were photographed with Nikon, TS100 digital camera attached to the microscope.

Statistical Analysis

The test results were calculated in Excel and LC50 with PROBIT software. The calculated LC50 was found to be 150 ppm for 96 h. The following method was used to describe the severity of the pathological change: (-) No alteration, (+) Mild alteration, (++) Moderate alteration, and (+++) Severe alteration. These symptoms indicate the severity of complications (Riba *et al.*, 2005; Roy *et al.*, 2013).

RESULTS AND DISCUSSION

Fish Behavior

Fish applied with 150 ppm chlorpyrifos did not show any change in food starting from day 1.

Moreover, their aggression and remaining motionless on the bottom were also observed. The progressive darkening in the body

coloration with slight mucus secretion was recorded as a clinical sign of toxicity (Mazandarani *et al.*, 2015). The calculated LC50 was found to be 150 ppm for 96 hours. Mortality was observed during this period. A decline in the rate of swimming after 4 days was studied in *C. carpio*, when exposed to chlorpyrifos, as the concentration increased from 0.1 to 2.2 mg L^{-1} (Xing *et al.*, 2015) and the 96 hours LC (50) of chromium salt, potassium dichromate was determined to be 41.75 mg L^{-1} (Mishra and Mohanty, 2008). The chlorpyrifos (5 ppm) exposed group shows loss of balance, swimming pattern, food search behavior and convulsions were noticed on after 10 days exposure (Stalin *et al.*, 2019). On exposure to chlorpyrifos, the Japanese medaka showed abnormality in swimming (Khalil *et al.*, 2013).

Histopathological Findings

Histopathologic studies showed a series of changes in gill tissue, including primary lamellar edema, hyperplasia, severe secondary lamellar fusion, and clubbing (Schlenk and Benson, 2001) (Figure 2). Moreover, the imposed stress was responded in a tissue-specific manner and histological lesions became more severe with increasing concentration (Dogan *et al.*, 2022).

Figure 2 illustrates an increase in the lamellar fusion originating from the mucus cells located at the base of the lamellae, leading to the merging between secondary lamellae (Kakade *et al.*, 2020). Another damage that occurred was hyperplasia. The interlamellar space, which is the aqueduct and the mucus or mucus production space, can be blocked due to hyperplasia of epithelial cells originating from the primary filaments. Gill is usually the first target tissue for waterborne contaminants, and prolonged exposure results in the absorption of these pollutants through the gills, producing visible damage effects in this tissue.

Through increasing the duration of exposure and an increasing dose of the poisoning agent, epithelium thoroughly




Figure 2. Light microscope image gill of common carp in the exposure to the salinity and chlorpyrifos. [Nikon Eclipse TS100 (40X)]: (a): Up and down (arrow), Epithelial hypertrophy, (b): Up and down (arrow), Epithelial lifting, (c): Left to right (arrow), Distal hyperplasia, (d): Left to right (arrow), Lamellar aneurism, (e): Up and down (arrow), Lamellar fusion.

separates and necrosis of gill tissues is performed. The pronounced degenerative changes observed in gill indicate the vulnerability of the tissue, possibly due to its role as the first contact and entry point for the pesticide. Consequently, chlorpyrifos exerted its toxic effects by altering normal behavior,

causing neurotoxicity and disturbing osmoregulation.

Complications from Gill Histological Studies

Histopathological results indicated that gill was the primary target tissue affected by chlorpyrifos and salinity. No histopathological changes were observed in the gill of the control fish. Epithelial hypertrophy, epithelial lifting, lamellar fusion and distal hyperplasia were observed in the 4 and 8 ppt salinity and chlorpyrifos groups. Exposure to 8, 12 ppt and 15 ppm salinity and chlorpyrifos resulted in lamellar fusion, distal hyperplasia, epithelial hypertrophy, lamellar aneurism and epithelial lifting (Table 1). Also, epithelial

exhibited lamellar aneurism, rupture of the lamellar epithelium (Pal *et al.*, 2012), necrosis, epithelial lifting, epithelial hypertrophy, lamellar fusion, (Stalin *et al.*, 2019), hyperplasia (Samanta *et al.*, 2015) and excessive secretion of mucus. Alteration in pattern histopathology of gill was studied in *Channa punctatus*, after acute exposure to hexavalent chromium (Mishra and Mohanty, 2008). Katuli *et al.* (2014) reported that the impact of diazinon and sodium dodecyl sulfate leads to severe necrosis lamellar in gill tissues of *Rutilus rutilus* and *Scophthalmus maximus*. Similarly, diazinon exposure in gills of *Scatophagus argus* exhibited, epithelial lifting, hyperplasia and lamellar fusion (Ghasemzadeh *et al.*, 2015). These

Table 1. Common complications of carp gill tissue exposure to salinity and chlorpyrifos poison.^a

Treatments (T)	0 ppt				4 ppt				8 ppt				12 ppt			
	0 ppm				15 ppm				150 ppm							
	T1	T2	T3	T4	T1	T2	T3	T4	T1	T2	T3	T4	T1	T2	T3	T4
Lamellar fusion	(-)	(-)	(+)	(+)	(+)	(+)	(+)	(+)	(+)	(+)	(+)	(+)	(+)	(+)	(++)	(+++)
Distal hyperplasia	(-)	(+)	(+)	(+)	(+)	(+)	(++)	(++)	(+)	(+)	(++)	(++)	(+)	(+)	(++)	(+++)
Lamellar aneurism	(-)	(-)	(-)	(-)	(-)	(-)	(+)	(+)	(-)	(+)	(+)	(+)	(+)	(+)	(+)	(++)
Epithelial hypertrophy	(-)	(-)	(+)	(+)	(-)	(+)	(++)	(++)	(+)	(+)	(++)	(++)	(+)	(+)	(+++)	(+++)
Leukocyte infiltr	(-)	(-)	(-)	(-)	(-)	(-)	(-)	(-)	(-)	(-)	(-)	(-)	(-)	(-)	(-)	(+)
Epithelial lifting	(-)	(-)	(+)	(+)	(-)	(-)	(+)	(+)	(-)	(-)	(+)	(+)	(-)	(-)	(+)	(++)

^a No complications (-), 1 to 3 complications (+), 3 to 5 complications (++), 5 to 11 complications (+++). (Riba *et al.*, 2005; Roy *et al.*, 2013).

hypertrophy, lamellar aneurism, epithelial lifting, lamellar fusion, leukocyte infiltr and distal hyperplasia were observed in the 12 ppt, 150 ppm salinity and chlorpyrifos groups. Exposure to 8 ppt, 150 ppm salinity and chlorpyrifos resulted in lamellar fusion, distal hyperplasia, epithelial hypertrophy, lamellar aneurism and epithelial lifting and lamellar fusion, distal hyperplasia, lamellar aneurism and epithelial hypertrophy were observed in the 4 ppt, 150 ppm salinity and chlorpyrifos groups (Table 1). The most common gill changes at all doses of chlorpyrifos and salinity were lamellar fusion, distal hyperplasia and epithelial hypertrophy.

Histopathological alterations in gill of common carp, *C. carpio*, intoxicated with sub-lethal concentrations of chlorpyrifos

changes lead to reduced oxygen consumption in fish and, ultimately, their death.

CONCLUSIONS

In the present study, gill histological changes have been related to poison and salinity concentrations. It can be concluded that gill alterations as a result of salinity and Chlorpyrifos exposition of fish may serve as a sensitive biomarker for the toxicity of sublethal concentrations of Chlorpyrifos as well as other pollutants. Exposure to low concentrations altered biological parameters in fish, but long-term exposure to high concentrations caused death.



ETHICAL APPROVAL

The experimental procedure was performed according to the Guide for the Care and Use of Laboratory Animals and was approved by the research committee of Gorgan University of Agricultural Sciences and Natural Resources (Ethical Approval Code: 9721313104).

ACKNOWLEDGEMENTS

The authors thank Gorgan University of Natural Resources and Agricultural Sciences for supporting this work.

REFERENCES

1. Ataimehr, B., MajaziAmiri, B., Mirvaghefi, A., Nezami, Sh. and Riazi, Gh. 200). The Effect of Different Salinities on Solute Concentration, Osmotic Pressure, Body Tissue Water, Gill Chloride Cells and Percent Mortality of Whitefish (*Rutilus frisii kutum Kamensky 1901*). *Sci. J. Iran. Fish.*, **19(2)**: 115-130.
2. Bury, N. R., Schnell, S. and Hogstrand, C. 2014 Gill Cell Culture Systems as Models for Aquatic Environmental Monitoring. *J. Exp. Biol.*, **(5)**: 639-650.
3. Esbaugh, A. J., Khursigara, A. and Johansen, J. 2018. Toxicity in Aquatic Environments: the Cocktail Effect. *Environ. Sci. Pollut. Res*, 203-234.
4. Cocker, J., Mason, H. J., Garfitt, S. J. and Jones, K. 2002, Biological Monitoring of Exposure to Organophosphate Pesticides. *Toxicol. Lett.*, **134(1-3)**: 97-103.
5. Dogan, D., Nur, G., and Ahmet Deveci, H. 2022. Tissue-Specific Toxicity of Clothianidin on Rainbow Trout (*Oncorhynchus mykiss*). *Drug Chem. Toxicol.*, **45(4)**: 1851-1861.
6. Ghasemzadeh, J., Sinaei, M. and Bolouki, M. 2015. Biochemical and Histological Changes in Fish, Spotted Scat (*Scatophagus argus*) Exposed to Diazinon. *Bull. Environ. Contam. Toxicol.*, **94(2)**: 164-170.
7. Gao, X. Q., Fei, F. and Huo, H. H. 2020. Effect of Acute Exposure to Nitrite on Physiological Parameters, Oxidative Stress, and Apoptosis in *Takifugu rubripes*. *Ecotoxicol. Environ. Saf.*, **188**: 109878.
8. Hasankhani, T., Nikaein, D., Khosravi, A., Rahmati-holasoo, H. and Hasankhany, M. 2023. The Effect of *Echinacea Purpurea* L. (Eastern Purple Coneflower) Essential Oil on Hematological Parameters and Gut Microbial Population of Zebrafish (*Danio Rerio*) With Aflatoxicosis. *Iran. J. Vet. Med.*, **17(2)**: 173-182.
9. Hasankhany, M., Hedayati, A., Mazandarani, M., Bagheri, T. and Jafar, A. 2020. Interaction between Different Levels of Salinity and Exposure to Chlorpyrifos on Some Serum and Mucus Biochemical Indices of Common Carp (*Cyprinus carpio*). *Iran. Sci. Fish. J.*, **29(3)**: 37-44.
10. Hedayati, A., JahanBakhshi, A. and Qadri, A. 2013. *Aquatic Toxicology*. Gorgan University of Agricultural Sciences and Natural Resources Publications, 212 PP. [in Persian]
11. Huyben, D., Bevan, D., Stevenson, R., Zhou, H. and Moccia, R. 2018. Evaluation of Membrane Filtration and UV Irradiation to Control Bacterial Loads in Recirculation Aquaculture Systems. *Aquac. Int.*, **26(6)**: 1531-1540.
12. Kakade, A., Salama, E. S., Pengya, F., Liu, P. and Li, X. 2020. Long-Term Exposure of High Concentration Heavy Metals Induced Toxicity, Fatality, and Gut Microbial Dysbiosis in Common Carp, *Cyprinus carpio*. *Environ. Pollut.*, **266(3)**: 115293.
13. Katuli, K. K., Amiri, B. M., Massarsky, A. and Yelghi, S. 2014. Impact of a Shortterm Diazinon Exposure on the Osmoregulation Potentiality of Caspian Roach (*Rutilus rutilus*) Fingerlings. *Chemosphere*, **(108)**: 396-404
14. Khalil, F., Kang, I. J., Undap, S., Tasmin, R., Qiu, X. and Shimasaki, Y. 2013. Alterations in Social Behavior of Japanese Medaka (*Oryzias latipes*) in Response to Sublethal Chlorpyrifos Exposure. *Chemosphere*, **92**: 125-130
15. Mackay, D., Giesy, J. P. and Solomon, K. R. 2014. Fate in the Environment and Long-Range Atmospheric Transport of the Organophosphorus Insecticide, Chlorpyrifos and Its Oxon. *Rev. Environ. Contam. Toxicol.*, **231**: 35-76.
16. Mazandarani, M., Sudagar, M. and Namroodi, S. 2015. Histopathological Effects of Acute Copper Sulphate Exposure

- on Kidney, Liver and Gill of Common Carp, *Cyprinus carpio*, Fingerlings. *J. Aqua. Ecol.*, **5(1)**: 9-16. (in Persian with English Abstract)
17. Mishra, A. K. and Mohanty, B. 2008. Acute Toxicity Impacts of Hexavalent Chromium on Behavior and Histopathology of Gill, Kidney and Liver of the 422 Freshwater Fish, *Channa punctatus* (Bloch). *Environ. Toxicol. Pharmacol.*, **26**: 136-141.
 18. Moustakas, C. T., Watanabe, W. O. and Xopeland, K. A. 2004. Combined Effects of Photoperiod and Salinity on Growth, Survival and Osmoregulatory Ability of Larval Southern Flounder. *Aquaculture*, **229**: 159-179.
 19. Perry, J., Cotton, J., Rahman, M. A. and Brumby, S. 2020. Organophosphate Exposure and the Chronic Effects on Farmers: A Narrative Review. *Rural Remote Health.*, **20**: 4508.
 20. Riba, I., Blasco, J., Jiménez-Tenorio, N. and DelValls, T. Á. 2005. Heavy Metal Bioavailability and Effects: I. Bioaccumulation Caused by Mining Activities in the Gulf of Cádiz (SW, Spain). *Chemosphere*, **58(5)**: 659-669.
 21. Pal, S., Kokushi, E., Koyama, J., Uno, S. and Ghosh A. R. 2012. Histopathological Alterations in Gill, Liver and Kidney of Common Carp Exposed to Chlorpyrifos. *J. Environ. Sci. Health B.*, **47(3)**: 180-195.
 22. Rohani, M. F. 2023. Pesticides Toxicity in Fish: Histopathological and Hemato-Biochemical Aspects – A Review. *Emerg. Contam.*, **9(3)**: 100234.
 23. Roy, D., Ghosh, D. and Mandal, D. K. 2013. Cadmium Induced Histopathology in the Olfactory Epithelium of a Snakehead Fish, *Channa punctatus* (Bloch). *Int. J. Aquat. Biol.*, **1(5)**: 221-227.
 24. Samanta, P., Bandyopadhyay, N., Pal, S. and Mukherjee A. K. and Ghosh, A. R. 2015. Histopathological and Ultramicroscopical Changes in Gill, Liver and Kidney of *Anabas testudineus* (Bloch) after Chronic Intoxication of Almix (Metsulfuron Methyl 10.1%+Chlorimuronethyl10.1%) Herbicide. *Ecotoxicol. Environ. Saf.*, **122**: 360-367.
 25. Schlenk, D. and Benson, W. H. 2001. Target Organ Toxicity in Marine and Freshwater Teleosts: Systems. 1st Edition, CRC Press.
 26. Shafiei, E., Ghadamyari, M. and Mosallanejad, H. 2023. Levels of Imidacloprid and Chlorpyrifos Resistance and Associated Mechanisms in Greenhouse Populations of *Trialeurodes vaporariorum* (Hemiptera: Aleyrodidae) from Iran. *Toxin. Rev.*, **4(42)**: 255-265.
 27. Stalin, A., Suganthi, P., Mathivani, S., Paray, B. A., Al-Sadoon, M. K., Gokula, V. and Musthafa, M. S. 2019. Impact of Chlorpyrifos on Behavior and Histopathological Indices in Different Tissues of Freshwater Fish *Channa punctatus* (Bloch). *Environ. Sci. Pollut. Res. Int.*, **26(17)**: 17623-17631.
 28. Sanoesi, E., Asmara, S. D., Haromain, A. F., Nurcholis, A., Mafruch and Wijanarko, E. 2020. Hematological Analysis of Common Carp (*Cyprinus carpio*) Using Hematology Analyzer Tools and Manual at Fish Seed Center, Pasuruan, East Java. *IOP Conf. Ser.: Earth Environ. Sci.*, **493(1)**: 012011.
 29. Xing, H., Liu, T., Zhang, Z., Wang, X. and Xu, S. 2015. Fish and Shellfish Immunology Acute and Subchronic Toxic Effects of Atrazine and Chlorpyrifos on Common Carp (*Cyprinus carpio* L.): Immunotoxicity Assessments. *Fish Shellfish Immunol.*, **45(2)**: 327-33.



یادداشت پژوهشی: تغییرات هیستوپاتولوژیک آبشش ماهی کپور معمولی در مواجهه با کلرپیریفوس و شوری

مونا حسن خانی، تینا حسن خانی، سید علی اکبر هدایتی، و محمد مازندرانی

چکیده

در این مطالعه، اثر سم بر شوری بر بافت آبشش ماهی کپور معمولی (C. *carpio*) بررسی شد. بر این اساس، ۲۵۰ قطعه بچه ماهی کپور معمولی با وزن متوسط 21 ± 2 گرم در چهار تیمار هر کدام با سه تکرار شامل: شوری صفر، ۴، ۸ و ۱۲ ppt به مدت ۷ روز توزیع شدند. سپس یک گروه به مدت ۴ روز در معرض سم با غلظت حاد ۱۵۰ ppm کلرپیریفوس با فرمولاسیون ۴۰.۸٪ EC و گروه دوم به مدت ۷ روز در معرض غلظت تحت حاد ۱۵ ppm کلرپیریفوس قرار گرفتند. هیستوپاتولوژی بافت آبشش نشان داد که سم و شوری آسیب‌هایی مانند هیپرتروفی اپیتلیال، آنوریسم لایه‌ای، اتصال ثانویه تیغه‌های مجاور، هیپرپلازی انتهایی، بلند شدن اپیتلیال، نفوذ لکوسیتی و هیپرپلازی ایجاد کرده‌اند. نتایج هیستوپاتولوژی آبشش در غلظت‌های بالا به عنوان هیپرتروفی اپی تلیال، هیپرپلازی دیستال و جوش خوردن لایه‌ای برخی ضایعات را نشان داد، اما در غلظت‌های پایین آنوریسم لایه‌ای، جدا شدن اپیتلیال و نفوذ لکوسیت‌ها ثبت شد. بنابراین، این شاخص‌های پاتولوژیک می‌توانند به عنوان نشانگرهای زیستی مورد استفاده قرار گیرند.