

## Evaluating Raw and Heat-Treated Hempseed (*Cannabis sativa* L.) with Enzyme Supplementation for Broiler Chicken

Aref Mahmoodtabar<sup>1</sup>, Farid Shariatmadari<sup>1\*</sup>, Hamed Ahmadi<sup>1</sup>, and Mohammad Amir Karimi Torshizi<sup>1</sup>

### ABSTARCT

A total of 480 seven-day-old male Arian broiler chickens were divided into five treatment groups with six replicates each. The treatments were offered to the birds for three weeks (days 7 to 28) and included a control group, 10% Raw Hempseed (*Cannabis sativa* L) supplementation (RH), 10% RH with Enzyme addition (RHE), 10% Heat-treated Hempseed (HH) in the diet, and 10% HH with Enzyme supplementation (HHE). A completely randomized design with a 2×2 factorial arrangement (raw vs. heat-treated hempseed, with vs. without enzyme supplementation), plus a control group, was used. While dietary treatments (hemp supplementation) significantly increased body weight and feed intake, the heat processing decreased weight gain. Hemp supplementation significantly lowered Coliform and increased *Lactobacillus* content in the ileum, while processing increased *Lactobacillus* and enzyme addition decreased *E. coli* ( $P < 0.05$ ). Digestibility parameters were positively affected by enzyme addition ( $P < 0.05$ ), but protein digestibility was reduced by heating. There were no significant interaction effects (enzyme supplementation and heat treatment), except for the total Aerobes count of intestinal micro flora ( $P < 0.05$ ). In conclusion, hempseed addition in the diet of broiler chickens during 7-28 days of age improved broiler performance and enzyme supplementation improved microbiology and, more profoundly, digestibility parameters.

**Keywords:** Arian broiler, Coliform, *E. coli*, Hemp supplementation.

### INTRODUCTION

Hempseed (*Cannabis sativa* L) along with its by-products, such as hempseed oil and meal, show potential as feed ingredient for livestock. It contains about 25% crude protein, 33-35% oil, 34% carbohydrates (mostly as fiber), and 18.3 MJ kg<sup>-1</sup> (4308 kcal kg<sup>-1</sup>) of metabolizable energy, and is rich in essential minerals and vitamins. Its primary protein, edestin (a highly-digestible, hexameric legumin protein), is particularly noted for its high essential amino acid content (Gakhar *et al.*, 2012; Wang *et al.*, 2008).

Historically, hemp cultivation was restricted in many countries until the early

2000s. Although industrial hempseed (with less than 0.3% tetrahydrocannabinol) is now approved for human consumption, its use in animal feed was once deemed "unsafe". Recent research, however, has begun to explore its potential benefits as hempseed has become more widely legalized (Shariatmadari, 2023).

Early studies on hempseed's impact on broiler performance revealed varied outcomes. For instance, Khan *et al.* (2010) reported that including 10% hempseed significantly improved body weight and feed efficiency. Mahmoudi *et al.* (2015) noted that while 2.5% hempseed had no effect, 7.5% was optimal for weight gain. Skřivan *et al.* (2020) found no effect with 4%

<sup>1</sup> Department of Poultry Science, Faculty of Agriculture, Tarbiat Modares University, Tehran, Islamic Republic of Iran.

\* Corresponding author; e-mail: shariatf@modares.ac.ir



hempseed, but noted improved tibia bone strength. Parr *et al.* (2020) observed that a 20% hemp heart led to increased weight gain and better feed efficiency compared to a soybean meal-based diet. These studies support the safety of up to 10% hempseed inclusion, despite the presence of some anti-nutritional factors such as trypsin inhibitors, fiber, condensed tannins, phytic acid, and saponins (Russo and Reggian, 2013).

To address these anti nutritional factors, heating and exogenous carbohydrase enzyme supplementations have been proposed as strategies to improve hempseed's effectiveness. Konca *et al.* (2019) demonstrated enhanced performance and egg quality in layer chickens fed 15% heat-treated hempseed as compared to similar amount of row hemp seed. As there was no report on the effect of exogenous enzyme supplementation, an enzyme cocktail containing two main commonly supplemented carbohydrase (glucanase and xylanase) was hypothesized to enhance chicken performances (Munyaka *et al.*, 2016; Mathlouthi, *et al.*, 2002). The combined effects of heat-treated hempseed and enzyme supplementation were also considered (Amerah *et al.*, 2011).

Thus, the aim of this study was to evaluate the combined effects of heat-treated hempseed and an exogenous enzyme cocktail on broiler performance, ileum nutrient digestibility, and microbiota composition at a 10% dietary inclusion

**Table 1.** Chemical composition of the raw hempseed used for the formulation of the diets (as-is basis).

Item	Content
Dry matter (%)	94.62
Gross energy (kcal kg <sup>-1</sup> )	5925
Crude protein (%)	24.7
Ether extract (%)	30.5
Ash (%)	5.37
Crude fiber, %	29.6
Neutral detergent fiber (%)	32.4
Acid detergent fiber (%)	22.1
Total lysine (%)	1.02
Total methionine (%)	0.43
Total threonine (%)	0.62

level. As young chicks have a less developed digestive tract, they are unable to produce enzyme in sufficient quantities by themselves, and may not tolerate high fibrous diet. According to Wang *et al.* (2021), chicks benefit more from enzyme addition at a younger age. Therefore, this experiment was designed to assess performance criteria up to 28 days.

## MATERIALS AND METHODS

The experimental protocols were approved (IR.MODARES.REC.1400.032) by the Biomedical Research Ethics Committee of Tarbiat Modares University.

### Diets, Birds and Housing

Samples of hemp seed (Table 1) and experimental diets (Table 2) were chemically analyzed in duplicate according to standard methods of the Association of Official Analytical Chemists (AOAC, 2000) for dry matter (at 105°C overnight), ash (oven at 600 overnight), crude protein (N×6.25 - Kjeldhal), crude fat (Soxhlet extraction), gross energy by bomb calorimetric (Gallenkamp Autobomb, UK) and crude fiber. Neutral Detergent Fiber (NDF) and Acid Detergent Fiber (ADF) were measured according to the procedures of Van Soest *et al.* (1991) and Robertson and Van Soest (1981), respectively. The method used for Amino Acid (AA) profiling was based on the standard protocol of the Pico-Tag method from Waters Corporation (Deng, *et al.*, 2004). High-performance liquid chromatography (Waters, Model: 2695E, USA) was used to determine samples following hydrolysis by hydrochloric acid (6N) and derivatization by orthophaldialdehyde. Metabolisable energy was estimated according to van der Klis and Fledderus (2007). Other nutrient compositions are calculated based on NRC (1994) data of feedstuffs nutrient tables.

A total of 480 one-day-old male Arian broiler chickens were randomly divided into five treatment groups, with each group housed in six replicate pens containing 16 chickens each. The treatment groups were:

**Control (No hempseed),**

- 10% Raw Hempseed (RH) in the diet,
- 10% RH with Enzyme supplementation (RHE),
- 10% Heat-treated Hempseed (HH) in the diet,
- 10% HH with Enzyme supplementation (HHE).

The birds were given a starter diet from days 0 to 6, then, they were weighed randomly divided and switched to the

experimental diets from days 7 to 28. Hempseed underwent heat treatment at 120°C for 60 minutes, following the method as described by Konca *et al.* (2019). Two NSPase enzymes, Econase XT (endo-1,4- $\beta$ -xylanase, with a minimum activity of 4,000,000 BXU g<sup>-1</sup>) and Econase GT 200 (endo-1,3(4)- $\beta$ -glucanase, with a minimum activity of 200,000 BU g<sup>-1</sup>), were obtained from AB Vista, in the Netherlands. The enzyme treatments consisted of a mixture of 4 g ton<sup>-1</sup> of Econase XT and 100 g ton<sup>-1</sup> of Econase GT 200.

All birds had free access to feed and water throughout the experiment. The diet's composition and nutrient content are outlined in Table 1 and were formulated based on the Arian breeding guide (Corporation Support of Animal Affairs,

**Table 2.** Composition and nutrient contents of experimental diets (as-fed basis) offered days 8-28.

Ingredients (kg ton <sup>-1</sup> feed)	Control diet	Raw hempseed diet	Heat-treated hempseed diet
Corn	558.65	536	536
Soybean meal (43%)	374	320	320
Hemp	0	100	100
Vegetable oil	27	0	0
Dicalcium phosphate	17.3	16	16
Limestone	9.8	10.5	10.5
Salt	1.8	1.75	1.75
NaHco3	2.75	2.85	2.85
DL-Methionine	2.55	2.8	2.8
L-Lysine HCl	0.9	1.45	1.45
L-Threonine	0.25	0.6	0.6
Vitamin/Mineral premix <sup>a</sup>	5	5	5
Filler	0	3.05	3.05
Calculated nutrient composition			
ME (kcal kg <sup>-1</sup> )	2950	2950	2950
Crude protein (%)	20.52	20.59	20.59
Ca (%)	0.87	0.87	0.87
Available p (%)	0.44	0.44	0.44
Na (%)	0.16	0.16	0.16
Lysine (%)	1.18	1.18	1.18
Methionine+Cystine (%)	0.9	0.9	0.9
Threonine (%)	0.8	0.8	0.8
Analyzed nutrient composition <sup>b</sup>			
Moisture (%)	7.93	7.18	6.70
Crude protein (%)	20.4	20.18	20.26
Ash (%)	6.26	6.08	6.05
Ether extract	7.06	7.11	7.40

<sup>a</sup> Each kg of vitamin and mineral premix contained: Vitamin A 4000000 IU, vitamin E 26000 IU, vitamin D3 1800000 IU, vitamin K 1200 mg, vitamin B1 1000 mg, vitamin B2 2600 mg, Niacin 5400 mg, Pantothenic Acid 7500 mg, vitamin B6 1280 mg, Folic acid 760 mg, Biotin 72 mg, vitamin B12 6.8 mg, choline chlorohide 320000 mg and antioxidant 1000 mg, Fe, 8000 mg, Mn, 48000 mg, Cu, 6400 mg, I, 500 mg, Zn, 44000 mg, Se, 120 mg. <sup>b</sup> Analyzed according to the AOAC (1995).



2008). Arian is a descendant of the Hybro Normal breed, originally developed by the Dutch company, and now widely bred in Iran.

The initial temperature in the house was set to  $33\pm 1^{\circ}\text{C}$  for the first week and gradually lowered to  $24\pm 1^{\circ}\text{C}$  in the subsequent weeks. The humidity level was consistently maintained at 60% throughout the study. The lighting schedule began with 23 hours of light and 1 hour of darkness from days 1 to 3, increasing by 2 hours of light each day until a final schedule of 16 hours of light and 8 hours of darkness was established. This continued for the remainder of the experiment. Light intensity ranged from 3–4 lux during the first week and increased to 5–7 lux thereafter. Each pen, measuring  $1\times 2$  m, had a stocking density of less than  $25\text{ kg m}^{-2}$  and was fitted with a nipple drinker and feeder to ensure continuous access to food and water for the birds. Regular cleaning was performed in the broiler room to maintain hygiene standards throughout the experiment.

### Measurements and Sampling

During the experimental period from days 7 to 28, we measured performance metrics including Feed Intake (FI), Body Weight (BW), Body Weight Gain (BWG), and Feed Conversion Ratio (FCR). We tracked daily mortality to adjust FI, live weight, and FCR calculations. At the conclusion of the study, we randomly selected 12 birds from each treatment group, with two birds chosen from each replicate pen. After a 2-hour fasting period with water access, the birds were killed by cervical dislocation and exsanguination. The carcasses were plucked, and ileum (from Meckel's diverticulum to 5 cm before the ileocecal colonic junction) of each bird was cut open and ileum content

were taken (to evaluate digestibility, and intestinal bacterial populations).

To determine the Colony Forming Units (CFU), the drop count method was used in saline solution (Miles and Misra, 1938). Each sample of ileal contents were homogenized, and then 1g of each sample was collected and transferred into 9 mL sterile saline solution to prepare serial dilutions. Plate count agar (Merck, Darmstadt, Germany), MacConkey agar (Himedia laboratories, Mumbai, India) and MRS agar (Merck, Darmstadt, Germany) were used for enumeration of total aerobes, *Escherichia coli* and lactic acid bacteria, respectively, following 24-hour aerobic incubation at  $37^{\circ}\text{C}$  (Jabbar *et al.*, 2024).

For digestibility trial, the diet was top-dressed with 3 g Marker (Titanium dioxide)  $\text{kg}^{-1}$  in last 4 days of experiment. Frozen ileal contents were thawed and dried at  $60^{\circ}\text{C}$  using a hot-air oven. Similar methods as for dietary component analysis (above) were applied for these samples. Apparent ileal digestibility for nutrients and energy was calculated following the methods described by Del Alamo *et al.* (2008) and Latifi *et al.* (2023).

### Statistical Analysis

A completely randomized design with a  $2\times 2$  factorial arrangement (raw vs. heat-treated hempseed and with vs. without enzyme supplementation), plus a control group, was used. Data were analyzed using a two-way ANOVA through the GLM procedure in SAS (SAS, 1990) for the factorial part. Additionally, a one-way ANOVA was performed to compare the control group with all other treated diets. A significance level of  $P < 0.05$  was applied, and significant differences were identified using Tukey's test.

**Table 3.** Effects of dietary treatments on growth performance (days 7-28).<sup>a</sup>

Treatment	Body weight (g)	Body weight gain (g)	Feed intake (g)	Feed conversion ratio	Viability
Control	1003 <sup>b</sup>	852.1 <sup>b</sup>	1233 <sup>b</sup>	1.447	91.6
RH	1044 <sup>a</sup>	896.7 <sup>a</sup>	1278 <sup>a</sup>	1.425	95.0
RHE	1041 <sup>a</sup>	892.2 <sup>a</sup>	1280 <sup>a</sup>	1.435	96.6
HH	1027 <sup>a</sup>	879.5 <sup>a</sup>	1270 <sup>a</sup>	1.444	96.6
HHE	1023 <sup>ab</sup>	874.7 <sup>a</sup>	1259 <sup>ab</sup>	1.439	95.0
SEM	7.00	7.01	9.25	0.01	2.08
P values	0.002	0.001	0.009	NS	NS
Process					
RH	1042 <sup>a</sup>	894.4 <sup>a</sup>	1279	1.44	95.8
HH	1025 <sup>b</sup>	877.2 <sup>b</sup>	1264	1.43	95.8
SEM	5.34	5.28	6.30	0.007	1.53
P-value	0.034	0.031	NS	NS	NS
Enzyme					
E0	1035	888.1	1274	1.43	95.8
E1	1032	883.5	1269	1.44	95.8
SEM	5.34	5.28	6.30	0.007	1.53
P-value	NS	NS	NS	NS	NS
Process×Enzyme					
SEM	7.55	7.47	8.19	0.01	2.17
P-value	NS	NS	NS	NS	NS

<sup>a</sup> Control (No hempseed), RH= 10% Raw Hempseed in the diet, RHE= 10% RH with Enzyme supplementation, HH= 10% Heat-treated Hempseed in the diet, HHE= 10% HH with Enzyme supplementation, E0= Without Enzyme, E1= With Enzyme, NS= Not Significant.

<sup>a-b</sup>: Different letters in the same column indicate significantly different means according to Tukey's HSD test at the 0.05 significance level.

## RESULTS AND DISCUSSION

The composition of hempseed analyzed in this study (Table 1) was consistent with previous reports by Callaway (2004) and House *et al.* (2010). While hempseed and its by-products have been utilized for medicinal purposes for centuries (della Rocca and Di Salvo, 2020), there remains considerable lack of understanding regarding their nutritional value and impact on poultry performance.

The effects of 10% RHS and HHS with and without multi-enzyme supplementation (G and X) on broiler chicken performance from 7 to 28 days of age were evaluated. The initial average live weight of day-old broiler chickens was 41.0±1.7 g, increasing to approximately 153±4.4 g by 7 days of age. The performance metrics of chickens during the experiment are presented in Table 3. Chickens fed a diet supplemented with

raw hempseed had a significantly higher ( $P < 0.05$ ) body weight at 28 days of age compared to those fed the control diet. No additional benefit of the enzyme addition or heat treatment was observed in this group. The pattern for feed intake was similar to body weight gain, with no significant effect of treatment (heating and enzyme addition) on feed efficiency ratio. Mortality was only observed in the control group, and hempseed supplementation did not affect livability.

The existing research on the effects of hempseed supplementation is relatively sparse, which complicates detailed comparative assessments. The initial scientific investigation into hempseed's impact on poultry (layer chicken) was conducted by Silversides and Lefrançois (2005). The earliest study specifically examining the influence of hempseed on broiler performance was conducted by Khan *et al.* (2010). According to a review by Shariatmadari (2023), there are only a



limited number of studies that directly explore the effects of hempseed on broiler performance.

While hempseed and its by-products have been utilized for medicinal purposes for centuries (della Rocca and Di Salvo, 2020), there remains considerable lack of understanding regarding their nutritional value and impact on poultry performance. The literature shows varying results regarding the impact of hempseed on feed intake. Mahmoudi *et al.* (2015) and Bahari *et al.* (2014) reported no significant change in feed intake with hempseed supplementation. In contrast, Skrivan *et al.* (2020) observed an increase in feed intake among broilers consuming hempseed. However, Khan *et al.* (2010) found that hempseed-fed broilers had reduced feed intake. Some believe that hempseed's Tetrahydrocannabinol (THC) content may stimulate appetite and feed intake, impacting eating behavior and body weight regulation (Mahmoudi *et al.* 2015). However, at high inclusion levels (20%), elevated THC levels can have adverse effects on appetite and body weight (Vispute *et al.* 2019).

High hempseed inclusion may depress feed intake due to its high crude ash (8.8%) and cellulose content (House *et al.*, 2010), which can be particularly problematic for younger birds. Vispute *et al.* (2019) reported reduced feed intake and body weight gain in early life stages, likely due to less developed gut mucosa and digestive enzymes. Konca *et al.* (2019) attributed lower feed intake to the characteristic flavor of raw hempseed, while heating enhanced flavor and increased feed intake.

Regarding enzyme supplementation, Doskoviv *et al.* (2013) found no impact on feed intake, while Francesch *et al.* (2009) suggested enzymes might decrease feed intake by increasing energy availability. Alternatively, enzymes could increase feed intake by reducing digestive content viscosity, enhancing nutrient digestibility (Lázaro *et al.*, 2004; Wiśniewska *et al.*, 2023).

The observed improvement in broiler performance with hempseed indicates its nutritive value. Hempseed is recognized for its excellent protein quality and amino acid profile (Callaway, 2004), along with beneficial fatty acids, vitamins, and minerals, contributing to better performance. However, Konca *et al.* (2014) suggested that excessive amino acids from hempseed might imbalance amino acid ratios, reducing bioavailability. Roasting and enzyme supplementation mitigated some negative effects of hempseed inclusion.

All birds, except those in the control group (8% mortality), remained healthy throughout the experiment. Potential health benefits of hempseed may be due to orexigenic, anti-inflammatory, antipyretic, and antiparasitic effects of tetrahydrocannabinol (Callaway 2004; Mechoulam and Hanus, 2001). Cannabis sativa is reported to alleviate stress, improve immunity, and exhibit antimicrobial and antiviral properties (Novak *et al.*, 2001; Sakakibara *et al.*, 1991).

### Microflora of the Ileum

Dietary treatments, while reducing coliform content, increased *Lactobacillus* content of the ileum ( $P < 0.05$ ) due to dietary hempseed inclusion (Table 4). Total aerobes were not influenced by raw hempseed inclusion but were reduced with enzyme addition and heat-treated hempseed diets ( $P < 0.05$ ). Heat-treated hempseed significantly increased *Lactobacillus* while enzyme inclusion reduced coliform counts ( $P < 0.05$ ). There was significant ( $P < 0.05$ ) enzyme and heating interaction effect on total aerobes counts.

The poultry industry faces challenges from pathogenic diseases, impacting mortality and production. Microbial content in the digestive tract plays a crucial role in gut health (Markovi *et al.*, 2009). Industrial hempseed contains essential oils and cannabinoids that inhibit microbial growth (Nissen *et al.*, 2010). However, Stastnik *et*

**Table 4.** Effects of dietary treatments on micro-flora composition of ileum (Log 10 CFU g<sup>-1</sup>) at day 28 of broilers.<sup>a</sup>

Treatment	<i>E. coli</i>	Total Aerobes	<i>Lactobacillus</i> spp.
Control	10.06 <sup>a</sup>	9.49 <sup>a</sup>	8.62 <sup>c</sup>
RH	9.23 <sup>b</sup>	9.68 <sup>a</sup>	8.87 <sup>b</sup>
RHE	8.75 <sup>c</sup>	9.17 <sup>b</sup>	8.97 <sup>b</sup>
HH	9.41 <sup>b</sup>	8.93 <sup>b</sup>	9.32 <sup>a</sup>
HHE	8.77 <sup>c</sup>	9.57 <sup>a</sup>	9.37 <sup>a</sup>
SEM	0.086	0.099	0.070
P-value	< 0.001	< 0.001	< 0.001
Process			
RH	8.99	9.42	8.92 <sup>b</sup>
HH	9.09	9.25	9.35 <sup>a</sup>
SEM	0.060	0.072	0.052
P-value	NS	NS	< 0.001
Enzyme			
E0	9.32 <sup>a</sup>	9.31	9.10
E1	8.76 <sup>b</sup>	9.37	9.17
SEM	0.060	0.072	0.052
P-value	< 0.001	NS	NS
Process×Enzyme			
SEM	0.085	0.102	0.074
P-value	NS	< 0.001	NS

<sup>a</sup> Control (No hempseed), RH= 10% Raw Hempseed in the diet, RHE= 10% RH with Enzyme supplementation, HH= 10% Heat-treated Hempseed in the diet, HHE= 10% HH with Enzyme supplementation, E0= Without Enzyme, E1= With Enzyme, NS= Not Significant.

<sup>a-c</sup>: Different letters in the same column indicate significantly different means according to Tukey's HSD test at the 0.05 significance level.

*al.* (2016) found that higher cannabidiol levels did not affect microbiological parameters in the ileum. Conversely, Vispute *et al.* (2019) reported decreased coliform counts and increased *Lactobacillus* counts in the caecum and jejunum with hempseed supplementation. Enzyme supplementation in our study reduced coliforms and heating increased *Lactobacillus* counts. Bedford and Cowieson (2012) noted that exogenous enzymes can influence nutrient partitioning and bacterial populations, though effectiveness varies based on several factors such as the strain, age, health status/disease challenge of the animals, presence of antibiotics, quality of ingredients fed, along with the type (and levels) of enzyme employed.

### Ileal Digestibility

No general increase in digestibility parameters was observed with RHS inclusion (Table 5). Digestibility was largely unaffected by treatment groups, except for enzyme supplementation. Heat treatment lowers anti-nutritional compounds, increases protein availability, and enhances enzyme susceptibility (Marsman *et al.*, 1995). Overheating can damage heat-sensitive amino acids and reduce the bioavailability of some minerals and vitamins (Hurrel, 1990). Although heating did not affect digestibility, the heated group showed increased weight gain, likely due to higher feed intake. Previous studies suggest that heating may not significantly alter nutrient fractions (Rocha *et al.*, 2014). Newkirk *et al.* (2003) noted that non-heat-treated canola meals might contain higher levels of digestible amino acids. It is possible that heating's

**Table 5.** Effects of dietary treatments on ileal digestibility.

Treatment	DM (%)	Organic matter (%)	Fat (%)	NDF (%)	ADF (%)	Crude protein (%)	Gross energy (kcal kg <sup>-1</sup> )
Control	60.78 <sup>b</sup>	61.41 <sup>b</sup>	69.91 <sup>b</sup>	27.31 <sup>c</sup>	15.24 <sup>bc</sup>	67.66 <sup>c</sup>	62.59 <sup>b</sup>
RH	60.80 <sup>b</sup>	61.30 <sup>b</sup>	71.59 <sup>b</sup>	29.39 <sup>bc</sup>	14.28 <sup>c</sup>	65.62 <sup>c</sup>	60.50 <sup>b</sup>
RHE	62.85 <sup>a</sup>	69.25 <sup>a</sup>	79.64 <sup>a</sup>	34.05 <sup>a</sup>	19.32 <sup>a</sup>	78.26 <sup>a</sup>	70.16 <sup>a</sup>
HH	60.75 <sup>b</sup>	63.27 <sup>b</sup>	72.99 <sup>b</sup>	27.90 <sup>c</sup>	13.92 <sup>c</sup>	64.97 <sup>c</sup>	61.32 <sup>b</sup>
HHE	63.16 <sup>a</sup>	70.33 <sup>a</sup>	78.64 <sup>a</sup>	31.06 <sup>b</sup>	17.86 <sup>b</sup>	72.61 <sup>b</sup>	71.20 <sup>a</sup>
SEM	0.284	0.921	0.945	0.947	1.070	1.301	1.506
P-value	< 0.001	< 0.001	< 0.001	0.003	0.019	< 0.001	< 0.001
Process							
RH	0.946	65.27	75.61	31.71	16.80	71.94 <sup>a</sup>	65.33
HH	61.95	66.79	75.81	29.48	15.89	68.79 <sup>b</sup>	66.26
SEM	0.225	0.728	0.734	0.742	0.788	0.967	1.19
P-value	NS <sup>4</sup>	NS	NS	NS	NS	0.049	NS
Enzyme							
E0	60.77 <sup>b</sup>	62.28 <sup>b</sup>	72.28 <sup>b</sup>	28.64 <sup>b</sup>	14.10 <sup>b</sup>	65.29 <sup>b</sup>	60.91 <sup>b</sup>
E1	63.00 <sup>a</sup>	69.79 <sup>a</sup>	79.13 <sup>a</sup>	32.55 <sup>a</sup>	18.59 <sup>a</sup>	75.43 <sup>a</sup>	70.68 <sup>a</sup>
SEM	0.225	0.728	0.734	0.742	0.788	0.967	1.19
P-value	0.001	< 0.001	0.002	0.005	0.003	< 0.001	0.004
Process×Enzyme							
SEM	0.318	1.03	1.04	1.05	1.11	1.36	1.68
P-value	NS	NS	NS	NS	NS	NS	NS

<sup>a</sup> Control (No hempseed), RH= 10% Raw Hempseed in the diet, RHE= 10% RH with Enzyme supplementation, HH= 10% Heat-treated Hempseed in the diet, HHE= 10% HH with Enzyme supplementation, ADF= Acid Detergent Fiber, NDF= Neutral Detergent Fiber, E0= Without Enzyme, E1= With Enzyme, NS= Not Significant.

<sup>a-c</sup>: Different digestibility values are minimal and not significantly different enough according to the test at the 0.05 significance level. The differences in digestibility processes are needed for optimal hempseed digestibility.

Digestibility of nutrients is affected by gut micro-flora and exogenous enzyme supplementation (Bedford and Cowison, 2012). According to Lazaro *et al.* (2004), enzyme supplementation mainly enhances performance by improving nutrient digestibility (Lazaro *et al.*, 2004). Evidently the enzyme supplementation (to raw and heated hemp) had improved all digestibility parameters. Yet this was not reflected in growth and feed efficiency as may arguable expected. It has to be noted that the digestibility trial was in the last 4 days of the experiment, while growth performances criteria was over a 3 weeks period. A positive correlation might be observed if the trial was conducted over the longer period. It may also be that the extent of digestibility

was not enough to be reflected in the performance parameters.

Age plays a crucial role in digestibility issues (Wang *et al.*, 2021). Lu *et al.* (2013) reported lower nutrient digestibility values for younger broiler chickens. Young birds have a less developed digestive tract, cannot produce enough enzymes on their own, and may not tolerate high fiber diet (Olukosi *et al.* (2007). According to Jozefiak *et al.* (2004), during the starter phase, undigested fiber limits the accessibility of digestive enzymes to feed substrates. Exogenous enzyme supplementation overcomes these short-comings, reduces the requirement for the enzyme and makes more nutrients and energy available for chicks' growth. However, the beneficial effect of exogenous supplementation diminishes as chickens get older (Olukosi *et al.*, 2007). Wang *et al.* (2021) reported that chicks benefit more

from enzyme addition at a younger age and the contribution of enzymes to nutrient retention decreases with age in chickens.

### CONCLUSIONS

Raw hempseed can be promising and beneficial in broiler feeding, improving performance and feed intake. However, heat-treated hempseed and adding enzymes did not offer additional benefits beyond those provided by raw hempseed alone. Exogenous enzyme supplementation did improve all digestibility parameters, while heat treatment of hempseed reduced protein digestibility. Further research is needed to evaluate the effects of higher hempseed inclusion levels and varying types and doses of enzyme supplementation at older ages on broiler chicken performance. Additionally, efforts could also focus on optimizing heating programs to reduce anti-nutritional factors and improve the nutritional digestibility of hempseed.

### ACKNOWLEDGEMENTS

We would like to thank AB Vista for supplying the enzymes used in this study.

### REFERENCES

1. Amerah, A. M., Gilbert, C., Simmins, P. H. and Ravindran, V. 2011. Influence of Feed Processing on the Efficacy of Exogenous Enzymes in Broiler Diets. *World Poult. Sci. J.*, **67(1)**: 29-46.
2. AOAC. 2000. *Official Method of Analysis*. 15<sup>th</sup> Edition, Assoc. Off. Anal. Chem., Washington, DC, USA.
3. Bahari, M., Bahari, M., Khorshidi, K. J., Abdollahpour, R. and Gharahveysi, S. 2014. Effect of Using Medicinal Cannabis Seed (*Nigella sativa*) in Diets on Performance and Parameters of Broiler Chickens ROSS 308. *Adv. Environ. Biol.*, **8(13)**: 931-935.
4. Bedford, M. and Cowieson, A. J. 2012. Exogenous Enzymes and Their Effects on Intestinal Microbiology. *Anim. Feed Sci. Tech.*, **173(1-2)**: 76-85.
5. Callaway, J. C. 2004. Hempseed as a Nutritional Resource: An Overview. *Euphytica*, **140(1)**: 65-72.
6. Corporation Support of Animal Affairs. 2008. *Arian Broiler Chicken Management Guide Breeding*. Tehran, Iran. (in Persian)
7. Del Alamo, A. G., M. W. A. Verstegen, L. A. Den Hartog, P. Perez De Ayala, and M. J. Villamide. 2008. Effect of Wheat Cultivar and Enzyme Addition to Broiler Chicken Diets on Nutrient Digestibility, Performance, and Apparent Metabolizable Energy Content. *Poult. Sci.* **87**: 759-767.
8. della Rocca, G., and Di Salvo, A. 2020. Hemp in Veterinary Medicine: From Feed to Drug. *Front. Vet. Sci.*, **7**: 1-11.
9. Deng, S. G., Peng, Z. Y., Chen, F., Yang, P. and Wu, T. 2004. Amino Acid Composition and Anti-Anaemia Action of Hydrolyzed Offal Protein from *Harengula zunasi* Bleeker. *Food Chem.*, **87(1)**: 97-102.
10. Dosković, V., Bogosavljević-Bosković, S., Pavlovski, Z., Milošević, B., Škrbić, Z., Rakonjac, S. and Petričević, V. 2013. Enzymes in Broiler Diets with Special Reference to Protease. *World Poult. Sci. J.*, **69(2)**: 343-360.
11. Francesch, M., Liu, K., Dalibard, P. and Geraert, P. A. 2009. Combining NSP-Enzymes and Phytase: The Formulation Challenge in Broiler Nutrition. *Aust. Poult. Sci. Symp.*, **20**: 36-39.
12. Gakhar, N., Goldberg, E., Jing, E., Gibson, M. and House, J. D. 2012. Effect of Feeding Hempseed and Hempseed Oil on Laying Hen Performance and Egg Yolk Fatty Acid Content: Evidence of Their Safety and Efficacy for Laying Hen Diets." *Poult. Sci.*, **91**: 701-711.
13. House, J. D., Neufeld, J. and Leeson, G. 2010. Evaluating the Quality of Protein from Hemp Seed (*Cannabis sativa* L.) Products Through the use of the Protein Digestibility-Corrected Amino Acid Score Method. *J. Agric. Food Chem.*, **58(22)**: 11801-11807.
14. Hurrell, R. F. 1990. Influence of the Maillard Reaction on the Nutritional Value of Foods. In: "The Maillard Reaction in Food Processing, Human Nutrition and Physiology". Birkhauser Verlag, Basel, Switzerland, PP. 245-259.



15. Jabbar, M., Baboo, I., Majeed, H., Farooq, Z. and Palangi, V., 2024. Characterization and Antibacterial Application of Peppermint Essential Oil Nanoemulsions in Broiler. *Poult. Sci.*, **103(12)**: 1-15.
16. Józefiak, D., Rutkowski, A. and Martin, S. A. 2004. Carbohydrate Fermentation in the Avian Ceca: A Review. *Anim. Feed Sci. Technol.*, **113(1-4)**: 1-15.
17. Khan, R. U., Durrani, F. R., Chand, N. and Anwar, H. 2010. Influence of Feed Supplementation with *Cannabis sativa* on Quality of Broilers Carcass.” *Pak. Vet. J.*, **30(1)**: 34-38.
18. Konca, Y., Cimen, B., Yalcin, H., Kaliber, M. and BuyukkilicBeyzi, S. 2019. Effects of Heat Treated Hempseed Supplementation on Performance, Egg Quality, Sensory Evaluation and Antioxidant Activity of Laying Hens. *Br. Poult. Sci.* **60(1)**:39-46.
19. Konca, Y., Cimen, B., Yalcin, H., Kaliber, M. and BuyukkilicBeyzi, S. 2014. Effect of Hempseed (*Cannabis Sativa L.*) on Performance, Egg Traits and Blood Biochemical Parameters and Antioxidant Activity in Laying Japanese Quail (*Coturnix coturnix Japonica*). *Br. Poult. Sci.*, **55(6)**: 785-794.
20. Latifi, M., Moravej, H., Ghaziani, F. and Kim, W. K. 2023. Determination of Prediction Equations for Apparent Metabolizable Energy Corrected for Nitrogen of Corn Gluten Meal and Canola Meal in Broilers. *Poult. Sci.*, **102(5)**: 102587.
21. Lazaro, R., Latorre, M. A., Medel, P., Gracia, M. and Mateos, G. G. 2004. Feeding Regimen and Enzyme Supplementation to Rye-Based Diets for Broilers. *Poult. Sci.*, **83(2)**: 152-160.
22. Lu, H., Adedokun, S. A., Preynat, A., Legrand-Defretin, V., Geraert, P. A., Adeola, O. and Ajuwon, K. M. 2013. Impact of Exogenous Carbohydrases and Phytase on Growth Performance and Nutrient Digestibility in Broilers. *Can. J. Anim. Sci.*, **93(2)**: 243-249.
23. Mahmoudi, M., Farhoomand, P. and Nourmohammadi, R. 2015. Effects of Different Levels of Hemp Seed (*Cannabis sativa L.*) and Dextran Oligosaccharide on Growth Performance and Antibody Titer Response of Broiler Chickens. *Ital. J. Anim. Sci.*, **14(1)**: 1-6.
24. Marković, R., Šefer, D., Krstić, M. and Petrujkić, B. 2009. Effect of Different Growth Promoters on Broiler Performance and Gut Morphology. *Arch. Med. Vet.*, **41(2)**: 163-169.
25. Marsman, G. J. P., Gruppen, H., van der Poel, A. F. B., Resink, J. W., Verstegen, M. W. A. and Voragen, A. G. J. 1995. The Effect of Shear Forces and Addition of a Mixture of a Protease and a Hemicellulase on Chemical, Physical and Physiological Parameters during Extrusion of Soybean Meal. *Anim. Feed Sci. Technol.*, **56**: 21-35.
26. Mathlouthi, N., Mallet, S., Saulnier, L., Quemener, B. and Larbier, M. 2002. Effects of Xylanase and Beta-Glucanase Addition on Performance, Nutrient Digestibility, and Physico-Chemical Conditions in the Small Intestine Contents and Caecal Microflora of Broiler Chickens Fed a Wheat and Barley-Based Diet. *Anim. Res.*, **51(05)**: 395-406.
27. Mechoulam, R. and Hanus, L. 2001. The Cannabinoids: An Overview. Therapeutic Implications in Vomiting and Nausea after Cancer Chemotherapy, in Appetite Promotion, in Multiple Sclerosis and in Neuroprotection. *Pain Res. Manag.*, **6(2)**: 67-73.
28. Miles, A. and Misra, S. 1938 The Estimation of the Bactericidal Power of the Blood. *J. Hyg.*, **38**: 732-748.
29. Munyaka, P. M., Nandha, N. K., Kiarie, E., Nyachoti, C. M. and Khafipour, E. 2016. Impact of Combined  $\beta$ -Glucanase and Xylanase Enzymes on Growth Performance, Nutrients Utilization and Gut Microbiota in Broiler Chickens Fed Corn or Wheat-Based Diets. *Poult. Sci.*, **95(3)**: 528-540.
30. Newkirk, R. W., Classen, H. L., Scott, T. A. and Edney, M. J. 2003. The Digestibility and Content of Amino Acids in Toasted and Non-Toasteds Canola Meals. *Can. J. Anim. Sci.*, **83(1)**: 131-139.
31. Nissen, A., Zatta, I., Stefanini, S., Grandi, B., Sgorbati, B., Biavati, A. and Monti, A. 2010. Characterization and Antimicrobial Activity of Essential Oils of Industrial Hemp Varieties (*Cannabis sativa L.*). *Fitoterapia*, **81(5)**: 413-419.
32. Novak, J., Zitterl-Eglseer, K., Deans, S. G. and Franz, C. M. 2001. Essential Oils of Different Cultivars of *Cannabis sativa L.*

- and Their Antimicrobial Activity. *Flavour Fragr. J.*, **16(4)**: 259-262.
33. NRC (National Research Council, & Subcommittee on Poultry Nutrition). 1994. *Nutrient Requirements of Poultry: 1994*. National Academies Press.
  34. Olukosi, O. A., Cowieson, A. J. and Adeola, O. 2007. Age-Related Influence of a Cocktail of Xylanase, Amylase, and Protease or Phytase Individually or in Combination in Broilers. *Poult. Sci.*, **86(1)**: 77-86.
  35. Parr, B., Hook, C. Brannon., T., Chae, H. and Snider, M. 2020. Effects of Using Hemp Hearts as a Feed Additive in Broiler Chicks. *J. Agric. Hemp Res.*, **2(1)**: 1-11.
  36. Robertson, J. B. and van Soest, P. J. 1981. The Detergent System of Analysis and Its Applications to Human Foods. In: *"The Analysis of Dietary Fiber in Food"*, (Eds.): James, W. P. and Theander, O. Marcel Dekker, New York, PP. 123-158.
  37. Rocha, C., Durau, J. F., Barrilli, L. N. E., Dahlke, F., Maiorka, P. and Maiorka, A. 2014. The Effect of Raw and Roasted Soybeans on Intestinal Health, Diet Digestibility, and Pancreas Weight of Broilers. *J. Appl. Poult. Res.*, **23(1)**: 71-79.
  38. Russo, R. and Reggian, R. 2013. Variability in Antinutritional Compounds in Hempseed Meal of Italian and French Varieties. *Plant*, **1(2)**: 25-29.
  39. Sakakibara, I., Katsuhara, T., Ikeya, Y., Hayashi, K., and Mitsunashi, H. 1991. Cannabisin A, an Arylnaphthalene Lignanamide from Fruits of *Cannabis sativa*. *Phytochemistry*, **30(9)**: 3013-3016.
  40. SAS Institute. 1990. *SAS/STAT® User's Guide, Release 6.03 Edition*. SAS institute Inc., Cary, NC, USA.
  41. Shariatmadari, F. 2023: Emergence of Hemp as Feed for Poultry, *World Poult. Sci. J.*, **79(4)**: 769-782.
  42. Silversides, F. G., and Lefrancois, M. R. 2005. The Effect of Feeding Hemp Seed Meal to Laying Hens. *Br. Poult. Sci.*, **46(2)**: 231-235.
  43. Skřivan, M., Englmaierová., M., Taubner, T. and Skřivanová, E. 2020. Effects of Dietary Hempseed and Flaxseed on Growth Performance, Meat Fatty Acid Compositions, Liver Tocopherol Concentration and Bone Strength of Cockerels. *Animals*, **10**: 458.
  44. Stastník, O., Karasek, F., Stenclová, H., Trojan, V., Vyhnanek, T. and Pavlata, L. 2016. The Effect of Hempseed Cakes on Broiler Chickens Performance Parameters. *MendelNet*, **22**: 157-164.
  45. van der Klis, J. D. and Fledderus, J. 2007. Evaluation of Raw Materials for Poultry: What's up?. In *Proc. 16th Eur. Symp. Poult. Nutr. (ESPN)*, Strasbourg, France, PP. 123-130.
  46. Van Soest, P. J., Robertson, J. B. and Lewis, B. A. 1991. Symposium: Carbohydrate Methodology, Metabolism, and Nutritional Implications in Dairy Cattle. *J. Dairy Sci.*, **74**: 3583- 3597.
  47. Vispute, M. M., Sharma, D., Mandal, A. B., Rokade, J. J., Tagyi, P. K. and Yadav, A. S. 2019. "Effect of Dietary Supplementation of Hemp (*Cannabis sativa*) and Dill Seed (*Anethum graveolens*) on Performance, Serum Biochemicals and Gut Health of Broiler Chickens. *J. Anim. Physiol. Anim. Nutr.*, **103**: 525-533.
  48. Wang, J., Patterson, R. and Kim, W. K. 2021. Effects of Phytase and Multicarbhydrase on Growth Performance, Bone Mineralization, and Nutrient Digestibility in Broilers Fed a Nutritionally Reduced Diet. *J. Appl. Poult. Res.*, **30(2)**: 100146.
  49. Wang, X. S., Tang, C. H., Yang, X. O. and Gao, W. R. 2008. Characterization, Amino Acid Composition and in vitro Digestibility of Hemp (*Cannabis sativa* L.) Proteins. *Food Chem.*, **107**: 11-18.
  50. Wiśniewska, Z., Kołodziejcki, P., Pruszyńska-Oszmałek, E., Konieczka, P., Kinsner, M., Górka, P., Flaga, J., Kowalik, K., Hejdysz, M., Kubiś, M., Jarosz, Ł. S., Ciszewski, A. and Kaczmarek, S. 2023. Combination of Emulsifier and Xylanase in Triticale-Based Broiler Chickens Diets. *Arch. Anim. Nutr.*, **77(3)**: 187-204.



## ارزیابی دانه شاهدانه خام و حرارت دیده (*Cannabis sativa* L.) با مکمل آنزیمی برای مرغ گوشتی

عارف محمودتبار، فرید شریعتمداری، حامد احمدی، و محمد امیر کریمی  
ترشیزی

### چکیده

در مجموع ۴۸۰ قطعه جوجه گوشتی نر هفت روزه سویه آرین به پنج گروه تیماری با شش تکرار در هر گروه تقسیم شدند. تیمارها به مدت سه هفته (روزهای ۷ تا ۲۸) به پرندگان ارائه شد و شامل یک گروه کنترل، ۱۰٪ مکمل دانه شاهدانه خام (*Cannabis sativa* L.) (RH)، ۱۰٪ RH با افزودن آنزیم (RHE)، ۱۰٪ دانه شاهدانه فرآوری شده با حرارت (HH) در جیره و ۱۰٪ HH با افزودن آنزیم (HHE) بود. یک طرح کاملاً تصادفی با آرایش فاکتوریل ۲×۲ (دانه شاهدانه خام در مقابل فرآوری شده با حرارت، با در مقابل بدون افزودن آنزیم) به علاوه یک گروه کنترل استفاده شد. در حالی که تیمارهای غذایی (مکمل شاهدانه) به طور قابل توجهی وزن بدن و مصرف خوراک را افزایش دادند، فرآوری حرارتی افزایش وزن را کاهش داد. مکمل شاهدانه به طور قابل توجهی میزان کلی فرم را کاهش و میزان لاکتوباسیلوس را در ایلئوم افزایش داد، در حالی که فرآوری باعث افزایش لاکتوباسیلوس و افزودن آنزیم باعث کاهش اشریشیا کلی ( $P < 0.05$ ) شد. پارامترهای قابلیت هضم به طور مثبت تحت تأثیر افزودن آنزیم ( $P < 0.05$ ) قرار گرفتند، اما قابلیت هضم پروتئین با حرارت دادن کاهش یافت. هیچ اثر متقابل معنی داری (مکمل آنزیم× عملیات حرارتی) به جز تعداد کل باکتری‌های هوازی میکروفلور روده ( $P < 0.05$ ) وجود نداشت. در نتیجه، افزودن دانه شاهدانه به جیره جوجه‌های گوشتی در طول 7 تا 28 روزگی عملکرد جوجه‌های گوشتی را بهبود بخشید و مکمل آنزیمی، میکروبیولوژی و به طور عمیق‌تر، پارامترهای قابلیت هضم را بهبود بخشید.