

## Antimicrobial and Antioxidant Effects of Emulsions and Nanoemulsions of *S. officinalis*, *P. anisum*, *D. moldavica*, and *S. aromaticum* Against Foodborne Bacteria

Mojtaba Raeisi<sup>1\*</sup>, Bahador Haji Mohammadi<sup>1</sup>, and Negin Mehdinejad<sup>2</sup>

### ABSTRACT

Due to their antimicrobial and antioxidant properties, essential oils are used as natural preservatives. The purpose of this study was to investigate the chemical composition, antioxidant properties, and antimicrobial activity of emulsion and nano-emulsion forms of *Salvia officinalis*, *Pimpinella anisum*, *Dracocephalum moldavica*, and *Syzygium aromaticum* essential oils. The agar well-diffusion assay results obtained from the experiment suggested that nano-emulsion of *Dracocephalum moldavica* essential oil had the maximum antimicrobial activity against the pathogenic microorganisms drawn in the experiment. The inhibition zone diameters of the nanoemulsion of this essential oil against *Shigella dysenteriae*, *Salmonella Typhimurium*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Listeria monocytogenes*, *Escherichia coli*, and *Bacillus cereus* were 11.0, 11.8, 13.0, 13.1, 13.1, 13.6, and 14.1 mm, respectively. In contrast, the inhibition zone diameters of this essential oil against *S. dysenteriae*, *S. Typhimurium*, *P. aeruginosa*, *S. aureus*, *L. monocytogenes*, *E. coli*, and *B. cereus* were 9.6, 10.3, 10.8, 11.8, 11.3, 11.1, and 11.2 mm, respectively. The major components of *Dracocephalum moldavica* essential oil included geraniol (27.24%), geranial (10.75%), alpha-copaene (8.16%), alpha-pinene (7.37%), carvacrol (7.41%), limonene (6.86%), and nerol (6.45%). The nanoemulsion form possessed a significantly greater antioxidant potential compared to their emulsion form. Also, the nanoemulsions exhibited significantly lower IC<sub>50</sub> compared to the emulsions. The nanoemulsion form of *D. moldavica* had the lowest IC<sub>50</sub> and EC<sub>50</sub> values of 22.1 and 4.51 µg mL<sup>-1</sup>, respectively.

**Keywords:** Antioxidant activity, *Dracocephalum moldavica*, Essential oil.

### INTRODUCTION

Statistics published by WHO shows that, every year a large number of people across the world die from alimentary diseases. Also, huge amounts of antibiotics have given rise to resistant microbes. However, food spoilage threatens consumer health and incurs a big economic loss from countries (Burt, 2004). Some of the methods to tackle pathogenic as well as spoilage microorganisms include introducing chemical preservatives. In the past few decades, the growing awareness of consumers about harmful effects of chemical

preservatives, especially the carcinogenic properties, has therefore increased the demand of foods containing natural preservatives (Skandamis *et al.*, 2001). Medicinal plants have been used since ancient times in medicine, in the preparation of aromatic cosmetics, in the control of spoilage and pathogenic microorganisms in food, and in enhancing the flavor of foods. Plant secondary metabolites such as essential oils, aromatic compounds and volatile compounds have extensive applications in traditional medicine, flavoring and food preservation. These substances have shown

<sup>1</sup> Food, Drug and Natural Products Health Research Center, Golestan University of Medical Sciences, Gorgan, Islamic Republic of Iran.

<sup>2</sup> Faculty of Veterinary Medicine, Science and Research Branch, Islamic Azad University, Tehran, Islamic Republic of Iran.

\* Corresponding author; e-mail: drmraeisi@goums.ac.ir



antibacterial, antifungal, antiviral and antiparasitic effects. The biological effects of medicinal plants and their essential oils or extracts are largely due to their chemical compounds, particularly to the phenolic compounds they contain (Kelen and Tepe, 2008; Shahbazi *et al.*, 2016). The chemical compositions of essential oils in various plants may vary due to genetic and environmental factors such as geographical conditions, climatic and seasonal changes, and growth stages of the plant (Ruiz-Navajas *et al.*, 2012).

*Salvia officinalis*, *Pimpinella anisum*, *Eugenia caryophyllata*, and *Moldavian dragonhead* contain active substances that can effectively inhibit the growth of pathogenic microorganisms and reduce food oxidation. These effects have been the subject of investigations in various studies. In one study, to investigate the antibacterial activity of an essential oil mixture from some medicinal plants, including *Malva Sylvestris* and *Salvia officinalis*, against bacteria responsible for common oral infections, it was found that these oils exhibited significant growth inhibitory activity against both Gram-positive and Gram-negative bacteria and represent a natural substitute to chemical mouthwashes such as chlorhexidine (Eghbal *et al.*, 2021). The extract of *Pimpinella anisum*, beyond exhibiting antimicrobial effects, has demonstrated a considerable degree of antioxidant activity in oil and emulsion systems by means of free radical-scavenging activity, which positively reflected the oxidative stability of oil during storage, showing promise for application in food preservation (Singh *et al.*, 2008). *Eugenia caryophyllata* is another herbal plant extensively studied for its antimicrobial and antioxidant properties. The study on the bioactive properties and composition of clove (*Syzygium aromaticum*) essential oil reported the presence of a plethora of phenolic and terpenoid compounds that possess potent antimicrobial activity against many bacteria, fungi, and yeasts. In addition, the essential oil from this herb is able to scavenge free radicals and serve as a natural antioxidant

(Kennouche *et al.*, 2015). In another study, the chemical composition, antioxidant activity, and antimicrobial properties regarding the essential oil of Moldavian dragonhead (*Dracocephalum moldavica* L.) were evaluated. The results indicated that the main components of Moldavian dragonhead essential oil are terpenoids and phenolic compounds, thus giving it a very high potency against free radical scavenging, resulting in its importance as a natural antioxidant. Antimicrobial assays also proved that both essential oil and hydrolate of Moldavian dragonhead were successful against various microorganisms, mainly pathogenic bacteria (Acimovic *et al.*, 2022).

Though essential oils possess their own biological properties, in their application within food products, they face certain challenges. For instance, high concentrations of these oils may lead to undesirable organoleptic properties in foods due to the inherent antimicrobial properties of these oils. Additionally, their hydrophobic character, low solubility in water, poor chemical stability, and volatile nature add tremendous challenges in the technology applicable for essential oils in food products (Shavisi *et al.*, 2017). To solve these problems, encapsulation of essential oils into nanoparticles is considered a way to improve their applicability, stability, and efficacy. Of all systems of nanoparticles, nanoemulsions are found to be most advantageous in food, health, and cosmetic applications because of easy preparation and favorable functional properties. Nanoemulsions, whose particles have diameters in the range of 10–200 nm, are one of the most successful carrier systems for lipophilic compounds, including drugs, flavorings, antioxidants, and antimicrobial agents (Rao and McClements, 2011; McClements and Rao, 2011). A nanoemulsion is an oil phase dispersed in an aqueous phase, with fine droplets surrounded by a thin interfacial layer of surfactant and/or amphiphilic molecules, providing stability to the system (Borrin *et al.* 2016). Therefore, decreasing the size of the oil phase in the structure of the nanoemulsion, while

increasing its surface area, increases the efficiency with which it improves interaction between active compounds and biological membranes for their transport (Perugini Biasi-Garbin *et al.* 2015). It was reported that the antimicrobial property of nanoemulsions is directly related to their formation methods (Shavisi *et al.*, 2017). There are many methods for nano-emulsion production, including many high-energy and low-energy approaches. One example of a high-energy method is ultrasonication, which produces quickly and effectively nanoemulsions with small particle sizes and uniform distribution (McClements, 2012).

Iran is vastly rich in special resources, especially in plant cover and plant diversity. In terms of biodiversity, Iran ranks among the top eight countries in the world, with 8423 plant species, two-thirds of which are European plant species. Out of more than 2300 plant species in Iran, 1730 have specific medicinal properties, and all of these are unique to, and endemic, in this country where, naturally, they grow in about 8.84 million hectares of dense, semi-dense and sparse rangelands. Given that, this study aimed to prepare essential oils and nanoemulsions of the plants *Salvia officinalis*, *Pimpinella anisum*, *Eugenia caryophyllata* and Moldavian dragonhead grown in various regions of Iran, and to evaluate and compare their activities with respect to antibacterial and antioxidant activity.

## MATERIALS AND METHODS

### Plant Materials

The plants used in this research, namely, *Salvia officinalis*, *Pimpinella anisum*, *Syzygium aromaticum*, and *Dracocephalum moldavica*, were freshly harvested in the harvesting season and authenticated in the Botany Department of Gorgan University of Agricultural Sciences and Natural Resources. Then, the plants were dried under shade in a

well-ventilated place, avoiding direct sunlight.

### Essential Oil Extraction

Powdered plant samples were used for the extraction of essential oils using a Clevenger-type apparatus (Avijeh, Iran) according to the standard method. The powdered plant samples were put into distilled water, and the extraction began as soon as the water boiled; extraction continued for a total of 180 minutes. The oil was subsequently dried over sodium sulfate and purified by passing through filters of 0.22  $\mu\text{m}$  pore size. Finally, the purified oils were kept in dark vials at a temperature of 4°C to maintain their integrity until analysis (Purkait *et al.*, 2018).

### GC/MS Analysis

A GC/MS instrument (Agilent 7890B, USA) with a nonpolar capillary column - Agilent Hp5Ms (30 m length, 0.25 mm inner diameter, and 0.25  $\mu\text{m}$  film thickness) was used. The GC injector was kept at 80°C for 5 minutes, then, its interface was kept at 270°C. The injection volume into the device was 1  $\mu\text{L}$ . The carrier gas was also Helium with a flow rate of 1  $\text{mL min}^{-1}$  (Chamorro *et al.*, 2012). The analysis was conducted at Golestan University of Medical Sciences.

### Preparation and Characterization of the Emulsion and Nanoemulsion of Essential Oils

It has now become 0.5% w/v of essential oil in sterile distilled water, and with TWEEN 80 (0.2 g w/w EO) used as an emulsifier. This emulsion was stirred continuously for 10 minutes to form a clear, stable, and uniform emulsion. This has been prepared according to the method explained by Ghosh *et al.* (2013) with slight modifications. Thereafter, the emulsion was given 3 minutes treatment using Ultra Turrax homogenizer (OPTIMA,



XL100K, Clausthal, Germany) at 3,000 r min<sup>-1</sup> and subsequently sonic emulsifier at 50°C; frequency: 50 kHz; pulse: 45 s; rest: 15 s for a total of 6 minutes (Probe diameter: 15 mm). The size of the particles was analyzed using a Dynamic Light Scattering (DLS) device (Nanophox Sympatec GmbH, Clausthal, Germany), which measures the size distribution of particles suspended under conditions of Brownian motion characterized by a low charge density and negligible hydrodynamic interactions. The analysis was conducted at Mashhad University of Medical Sciences.

### Total Phenolic Contents

The Folin-Ciocalteu reagent assay was employed to measure the total phenolic content in the extracted essential oils. To begin, a solution of each essential oil dissolved in methanol at a concentration of 2 mg mL<sup>-1</sup> was combined with 2.25 mL of distilled water and 250 µL of Folin-Ciocalteu reagent. The mixture was vortexed thoroughly and left to react for five minutes. Then, two mL of a 7.5% Na<sub>2</sub>CO<sub>3</sub> solution was added. After allowing the mixture to sit at room temperature for 120 minutes, its absorbance was measured at 760 nm using a spectrophotometer (Model: LKB Novaspec II; Pharmacia, Cambridge, England). The results were reported as mg of gallic acid equivalent per gram of essential oil, based on a standard curve generated from known concentrations of gallic acid (Alizadeh *et al.*, 2013).

### Antibacterial Effects of the Essential Oils

Various essential oils were evaluated for their antibacterial activity against seven foodborne pathogenic bacteria, including four Gram-negative and three Gram-positive strains. The bacterial strains used in this study are presented in Table 1. The antibacterial activity was assessed using agar well-diffusion and microdilution methods. All bacterial strains were obtained from the culture collection of the Food and Drug Deputy at Golestan University of Medical Sciences, Gorgan, Iran.

### Agar Well-Diffusion Assay

To begin, 100 mL of nutrient agar (Merck, Darmstadt, Germany) was seeded with 1 mL of actively growing 18 hours bacterial broth cultures (1.5×10<sup>6</sup> CFU mL<sup>-1</sup>). The combined mixture, which was cooled to 45°C, was mixed well for 2 minutes, transferred to sterile plates, and allowed to solidify. Four wells were then aseptically prepared in the agar with a sterile cork-borer, and 10 µL of the essential oils were placed in each well. Dimethyl Sulfoxide (DMSO) served as the negative control. The plates were incubated for 72 hours at 37°C, then, the inhibition zone around the wells was measured in mm using a manual caliper (Vernier, Mitutoyo, Japan) (Mirtaghi *et al.*, 2016).

**Table 1.** Bacterial strains used for evaluation of the antibacterial activity of the essential oils.

Bacterial species	Strain code	Gram reaction
<i>Escherichia coli</i>	PTCC 1399	Gram-negative
<i>Salmonella typhimurium</i>	ATCC 13311	Gram-negative
<i>Pseudomonas aeruginosa</i>	PTCC 1616	Gram-negative
<i>Shigella dysenteriae</i>	PTCC 1188	Gram-negative
<i>Staphylococcus aureus</i>	PTCC 1112	Gram-positive
<i>Listeria monocytogenes</i>	PTCC 1298	Gram-positive
<i>Bacillus cereus</i>	PTCC 1154	Gram-positive

### Determining the Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC)

The assay was conducted as per Turgis *et al.* (2012), with some changes. Prior to the assay, the essential oils were dissolved in 10% DMSO so that the maximum concentration could reach 10,000  $\mu\text{g mL}^{-1}$ , then, serially diluted twofold, from 10 to 10,000  $\mu\text{g mL}^{-1}$ . A total of 125  $\mu\text{L}$  of each of the above essential oils solutions was introduced into wells 1-11 of a 96-well microplate (Sarstedt, Montreal, QC, Canada). A total volume of 140  $\mu\text{L}$  was achieved from each well by adding 15  $\mu\text{L}$  of mueller hinton broth (Merck, Germany) containing  $10^6$  CFU  $\text{mL}^{-1}$  of the actively growing cultures of the target microorganisms into all wells. Three rows of the microplate were used for each bacterium. The negative controls (two rows) contained 15  $\mu\text{L}$  of sterile physiological saline solution (0.85% NaCl) instead of the bacteria, while the positive controls contain 125  $\mu\text{L}$  of growth media and 15  $\mu\text{L}$  of the bacterial cultures. The samples were incubated for 24 hours at 37°C, after which absorbance was determined using a Biotech ELX8000 microplate reader (Biotek Instruments Inc., Winooski, VT, USA) at 595 nm. The Minimum Inhibitory Concentration (MIC) was defined as the lowest concentration of the antimicrobial agent that completely inhibited visible bacterial growth. The Minimum Bactericidal Concentration (MBC) is the lowest concentration required to kill at least 99.9% of the initial number of bacteria, determined by subculturing 10  $\mu\text{L}$  from each well showing no visible growth onto nutrient agar plates, and incubating for 72 hours at 37°C.

### Antioxidant Activity of the Essential Oils

The antioxidant activity of essential oils was evaluated using two methods, including

the 2,2-Diphenyl-1-Picrylhydrazyl (DPPH) assay and reducing power assay.

#### DPPH Assay

The free radical scavenging activity of the emulsion and nanoemulsion forms essential oils was evaluated using the DPPH assay (Sigma-Aldrich, Steinheim, Germany), following the protocol established by Erkan *et al.* (2008) with minor changes. Initially, 50  $\mu\text{L}$  of each essential oil at various concentrations (10-10,000  $\mu\text{g mL}^{-1}$ ) and the reference antioxidant Butylated Hydroxytoluene (BHT) were mixed with 2 mL of a 0.2 mM methanolic DPPH solution. After shaking well, they were kept at room temperature in the dark place for 60 minutes. Subsequently, the absorbance of the mixture was measured at 517 nm using a spectrophotometer (model: LKB Novaspec II, Pharmacia, Cambridge, England). The blank sample contained methanol solvent with DPPH. The radical scavenging activity was calculated using the following formula:

$$\% I = [A_{\text{blank}} - A_{\text{sample}} / A_{\text{blank}}] \times 100$$

Where, I represents the percentage of free radical DPPH scavenging,  $A_{\text{blank}}$  is the Absorbance of the control, and  $A_{\text{sample}}$  is the Absorbance of the sample, all measured in nm.

The IC<sub>50</sub> value was derived from the scavenging activity curve plotted against the concentrations of the essential oils, indicating the total antioxidant activity required to achieve a 50% reduction in the initial DPPH radical concentration.

#### Reducing Power Assay

The reducing power of the essential oils was evaluated using the method established by Oyaizu (1986). In this procedure, the essential oils were prepared by different concentrations of essential oils (10- 1000  $\mu\text{g mL}^{-1}$ ) and 2.5 mL 0.2M phosphate buffer pH= 6.6 were added with potassium ferricyanide ( $\text{K}_3\text{Fe}[\text{CN}]_6$ ) were added as 1%



potassium ferricyanide ( $K_3Fe[CN]_6$ ). The mixtures were incubated for 20 minutes at 50°C. The samples were added with 2.5 mL of the 10% trichloroacetic acid after incubation, which was then centrifuged for 10 minutes at 1,036 rpm. The upper layer was subsequently transferred to 2.5 mL Distilled water and combined with 2.5 mL ferric chloride (1%). The absorbance at 700 nm was measured by a double-test UV-Vis spectrophotometer (Model: LKB Novaspec II) (Pharmacia, Cambridge, England), the blank contained all reagents except the essential oils. Increased absorbance is shown to have more reducing power Butylated hydroxytoluene (BHT) as a positive control. Additionally, this contained the absorbance of each essential oil fraction which provided an EC50 (absorbance value of 0.5), against the standard antioxidant (BHT).

### Statistical Analysis

Data analysis was performed in SPSS version 16 (SPSS Inc., Chicago, IL, USA), and all the assays were performed in triplicate. Tukey's test was used to compare the differences between the mean values obtained from the experiments at the significance level of  $P < 0.05$ .

## RESULTS AND DISCUSSION

### Identification of Essential Oil Components

Using mass spectrum similarity and the mass library of the Gas Chromatography-Mass Spectrometry (GC-MS) device, 34, 9, 24, and 13 compounds were identified in the essential oils of *Salvia officinalis* (Table 2), *Pimpinella anisum* (Table 3), *Dracocephalum moldavica* (Table 4), and *Syzygium aromaticum* (Table 5), respectively. According to the results in Table 2, the main compounds of *Salvia officinalis* essential oil include Beta Thujene (13.74%), 1,8-Cineole (12.13%), Alpha

Thujene (9.52%), Alpha Fenchyl Acetate (6.12%), Camphor Bicyclo Heptan (7.36%), Caryophyllene (4.52%), Viridifloral (3.15%),  $\alpha$ -Pinene (3.11%) and  $\beta$ -Pinene (2.15%). These compounds together made up 61.8% of the identified compounds.

*Salvia officinalis* is one of the most important medicinal and aromatic plants and has antioxidant, antimicrobial, spasmolytic, astringent, antihydrotic and sensory properties. The essential oil of the plant, which is mainly formed in the very young leaves, is partly responsible for these activities. This essential oil is mainly composed of mono-terpenes such as 1,8-Cineole,  $\alpha$ - and  $\beta$ -Thujene, and Camphor, and it is produced and stored in epidermal glands (Grausgruber-Gröger *et al.*, 2012). Other researchers in Morocco studied the chemical properties of *Salvia officinalis* and found that 36 compounds were present in the chemical structure of this plant. Among them, 1,8-Cineole, camphor, borneol,  $\alpha$ -pinene,  $\beta$ -pinene,  $\alpha$ -humulene,  $\alpha$ -caryophyllene,  $\beta$ -caryophyllene, and viridiflorol were the dominant compounds (Delamare *et al.*, 2007). In another study, Couladis *et al.* (2002) examined the chemical properties of *Salvia officinalis* medicinal plant samples in Serbia and Montenegro, and reported that  $\alpha$ -thujone,  $\beta$ -thujone, 1,8-cineol, camphor, borneol, and bornyl acetate were the dominant compounds in the leaves of this medicinal plant, while in the flowers, the percentages of camphor and thujone were much lower. Different parts of medicinal plants vary in their chemical composition, but the diversity and quantity of compounds are much higher in the reproductive organs of plants than in other parts. A comparison of the chemical analysis results of *Salvia officinalis* leaf essential oil showed many similarities between these researchers' analyses and the results of the present study, confirming the aforementioned points.

**Table 2.** Identified Compounds in *Salvia officinalis* essential oil.

Number	Compound	RT <sup>a</sup> (Min)	% Of total essential oil composition
1	Cis salvene	3.12	0.48
2	Tricyclene	4.17	0.53
3	$\alpha$ -Thujene	4.28	0.17
4	$\alpha$ -Pinene	4.72	3.11
5	$\beta$ -Pinene	5.18	2.15
6	$\beta$ -Myrcene	5.42	1.18
7	L Phellanderene	5.53	1.84
8	$\alpha$ -Terpinene	6.12	0.70
9	Benzene 1 Methyl 3 (1-Methylether)	6.28	0.56
10	1,8 Cineole	6.41	12.13
11	Gamma Terpinene	6.72	2.17
12	$\alpha$ -Terpinolene	7.12	0.10
13	$\beta$ -Thujene	7.33	13.74
14	$\alpha$ -Thujene	7.84	9.52
15	Camphor Bicyclo Heptan	8.17	7.36
16	Isoborneol	8.34	1.14
17	Borneol L	8.89	1.52
18	$\alpha$ -Terpineol	9.11	0.30
19	Trans Carveol	9.29	0.56
20	1,3,6 Octatriene, 3, 7 Dimethyl	10.18	2.17
21	Cis 3 Hexenyl Isovalerate	10.74	2.52
22	$\alpha$ -Fenchyl Acetate	12.49	6.12
23	Sabiny Acetate	13.09	1.50
24	Myrtenyl Acetate	14.17	0.35
25	Eugenol	15.82	2.17
26	$\alpha$ -Copaene	16.13	1.15
27	Caryophyllene	17.21	4.52
28	Aromadendrene	17.38	1.15
29	$\alpha$ -Caryophyllene	19.12	1.19
30	Spathulenol	21.17	0.10
31	Caryophyllene Oxide	22.33	1.78
32	viridifloral	22.84	3.15
33	Camphene	23.14	2.72
34	$\beta$ -Clovane	25.54	0.92
Total			90.77

<sup>a</sup> RT: Retention Times.**Table 3.** Identified compounds in *Pimpinella anisum* essential oil.

Number	Compound	RT (Min)	% Of total essential oil composition
1	D limonene	11.72	2.86
2	Meta anisaldehyde	11.96	3.58
3	Stragole	12.06	1.96
4	Trans anethole	12.15	60.17
5	Trans ocimenone	12.26	5.85
6	Murolene	13.50	3.31
7	Curcumene	14.11	2.37
8	$\gamma$ -himachalene	14.53	14.19
9	Beta biabolene	14.92	2.75
Total			97.04

**Table 4.** Identified compounds in *Dracocephalum moldavica* essential oil.

Number	Compound	RT (Min)	% Of total essential oil composition
1	$\alpha$ - Thujene	9.20	1.18
2	$\alpha$ -Pinene	9.27	7.37
3	Camphene	9.36	2.58
4	$\beta$ -Pinnene	9.54	1.17
5	$\alpha$ -Phellandrene	9.72	0.59
6	Limonene	9.81	6.86
7	Linalool oxide	9.93	0.94
8	Linalool	10.52	1.15
9	Terpinen 4 ol	11.02	3.14
10	$\alpha$ -Terpineol	11.15	1.12
11	Nerol	11.43	6.54
12	Neral	11.55	1.90
13	Geraniol	11.62	27.24
14	Geranial	11.71	10.75
15	Thymol	12.12	3.15
16	Carvacrol	12.31	7.41
17	Geranyl acetate	12.54	1.25
18	$\alpha$ -Copaene	12.67	8.16
19	$\beta$ -Bourbonene	12.75	0.78
20	$\beta$ -Caryophyllene	13.14	1.25
21	$\gamma$ -Muurolene	13.36	0.50
22	$\gamma$ -Cadinene	13.95	0.88
23	Caryophyllene oxide	14.30	0.33
24	Viridifloral	14.78	0.56
Total			96.8

**Table 5.** Identified compounds in *Syzygium aromaticum* essential oil.

Number	Compound	RT (Min)	% Of total essential oil composition
1	Carvacrol	1270	1.18
2	Alpha Cubebin	1313	2.14
3	Eugenol	1326	28.13
4	M Eugenol	1348	3.72
5	Trans Caryophyllene	1382	0.70
6	Caryophyllene	1403	22.17
7	Trans Isoeugenol	1417	0.38
8	Isoeugenol	1426	3.17
9	Alpha Humulene	1441	1.15
10	Eugenol acetate	1486	17.75
11	Caryophyllene oxide	1519	9.51
12	Hexadecanoic acid	1718	0.80
13	Eicosane	1829	0.56
Total			91.36

In another part of this study, using gas chromatography-mass spectrometry, nine compounds were identified in the essential oil of *Pimpinella anisum* (Table 3). Trans-anethole (60.17%) with a retention time of 12.15 minutes was the most abundant

compound identified in this essential oil. Additionally,  $\gamma$ -himachalene (14.19%) and Trans-ocimene (5.85%) were ranked next. The identified compounds in *Pimpinella anisum* essential oil have been reported in various forms in previous studies. Orav *et al.*

(2008) reported that the main compound in all essential oil samples of *Pimpinella anisum* collected from different European countries was Trans-anethole, ranging from 76.9 to 93.7%, and other major compounds included  $\gamma$ -himachalene, trans-pseudoisoeugenyl 2-methylbutyrate, p-anisaldehyde, and methylchavicol. Furthermore, Ullah *et al.* (2014) reported Trans-anethole at 82.1% and  $\gamma$ -himachalene at 7% as the main compounds in the essential oil of this plant. Abdel-Reheem and Oraby (2015) also identified the main constituents of *Pimpinella anisum* essential oil as trans-anethole (82.1%), cis-anethole (5.8%), methylchavicol (2.5%), linalool (2.3%),  $\alpha$ -terpineol (1.5%), and methyl eugenol (1.3%). It could be said that there are differences, not great, among major constituents of the essential oil of this plant, but in general, any difference in the composition may arise from variation in climate among different eco-regions. Also, the parts of a plant may widely differ in their chemical composition at a particular time, and it is essential to pick up the part when it has the highest concentration of the active ingredient. Other important considerations for the level of active compounds during harvesting are the appropriate harvest time depending on the plant's genetics, type of cultivation, and times of cultivation, location and soil used, irrigation level, light exposure, altitude, and many others (Hendawy *et al.*, 2018, Ali-Shtayeh *et al.*, 2018).

Also, identification of the active compounds in the essential oil of *Dracocephalum moldavica* (Table 4) showed that 74.23% consisted of Geraniol (27.24%), Geranial (10.75%), Alpha Copaene (8.16%), Alpha Pinene (7.37%), Carvacrol (7.41%), Limonene (6.86%), and Nerol (6.45%). Yousefzadeh *et al.* (2013) and Kakasy *et al.* (2006) reported that Geranyl acetate, Geranial, and Geraniol constituted the major components of the essential oil in *Dracocephalum moldavica*. According to the findings of Holm *et al.* (1988), the main components of the essential oil in *Dracocephalum moldavica* were oxygenated cyclic monoterpenes, including Geraniol,

Geranial, Geranyl acetate, and Nerol. Sonboli *et al.* (2008) reported that Nerol, Geranial, Geranyl acetate, and Geraniol, with 32.1%, 21.6%, 19.9%, and 17.6%, respectively, were the major components of *Dracocephalum moldavica* essential oil.

*Syzygium aromaticum* was another essential oil studied in this research. Based on the results in Table 5, the compounds identified in this essential oil with their respective Retention Times (RT) were Eugenol (28.13%) with RT of 13.26 minutes, Caryophyllene (22.17%) with RT of 14.03 minutes, Eugenol acetate (17.75%) with RT of 14.86 minutes, and Caryophyllene oxide (9.51%) with RT of 15.19 minutes.

Such was the state of affairs with respect to the highest identified compounds found in the essential oil of *Syzygium aromaticum*. A lot of researchers have worked on *Syzygium aromaticum* essential oil components. For instance, Myint *et al.* (1996) reported that Eugenol, Caryophyllene, and Eugenol acetate were the main components in the ethanolic extract of *Malaysian Eugenia caryophyllata*. Another study conducted by Nassar *et al.* (2007) revealed that Eugenol, Eugenol acetate, and Caryophyllene were the major constituents among 16 volatile compounds isolated from hexane extract of *Syzygium aromaticum*. Fichi *et al.* (2007) reported similar results for chemical analysis of *Syzygium aromaticum* essential oil, where Eugenol (59.3%) and Caryophyllene (24.9%) were the major compounds.

Particle Size of Emulsion and Nanoemulsion of Essential Oils

The diameter of emulsion and nano-emulsion particles of essential oils is shown in Figure 1. Based on the results, the size of the emulsion and nano-emulsion particles was 874-973 and 50.4-64.9 nm, respectively.


#### Antimicrobial Activity of Emulsion and Nanoemulsion Forms Essential Oils

In order to compare the antimicrobial activity of the emulsion and nano-emulsion of plant essential oils studied in this research,



the well diffusion and micro-dilution methods were used. The results of the antibacterial activity assessment of these essential oils are shown in Tables 6 and 7, respectively.

liberation of ions by a nano-emulsion may react with thiol groups of proteins located on bacterial cell surfaces. Following this, these proteins become inactivated; lessening membrane permeability, which eventually



**Figure 1.** The particle diameter of emulsion and nano-emulsion of different essential oils.

The results from the well diffusion method (Table 6) and the micro-dilution method (Table 7) indicate that both emulsion and nano-emulsion forms exhibited significant antimicrobial effects against all tested microorganisms. Moreover, in all the studied essential oils, the antimicrobial properties significantly increased ( $P < 0.05$ ) when the particle size of the essential oil emulsion was reduced and converted into the nano-emulsion form. Antibacterial activity is most likely to be improved upon finely converting various essential oils into nano-sized particles. This is likely since, because of their low solubility in water, they cannot interact with the cell membrane easily. Nano-emulsions through their smaller size can easily approach the surfaces of the cell membranes (Moghimi *et al.*, 2016).

Generally, nanoemulsions could cause damage or kill bacteria by different mechanisms. Essential oils' mode of action is cited to involve the disruption and destabilization of the phospholipid bilayer structure, leading to the impairment of the cell membrane, interaction with membrane enzymes and proteins as proton carriers, and pH reduction across the membrane (Burt, 2004; Nazzaro *et al.*, 2013). For instance, the

results in death for the bacteria. Electrostatic interaction is possible between the positively charged nanoparticle and the negatively charged cell membrane and can result in disruption of the membrane through nanoparticle binding. The accumulation of nanoparticles in both the cytoplasm and outer membrane can also interfere with bacterial growth and their survival. Nanoemulsions formed through high-pressure homogenization are even more potent than pure essential oils because of their size reduction (Topuz *et al.*, 2016). Moghimi *et al.* (2016) also reported that the nano-emulsion of *Thymus daenensis* essential oil was ten times more antibacterial than the pure essential oil. Results depend on many variables, including type and physical properties of essential oil, method of preparation, and type of the tested microorganism. As cited by Shahabi *et al.* (2017), the nano-emulsion of essential oil extracted from *Zataria multiflora* had an enhanced antimicrobial potency toward *L. monocytogenes* compared to that of *S. Typhimurium* when compared to the same essential oil. The antimicrobial activity of the nano-emulsion, however, depends on the

**Table 6.** Inhibitory zone diameter (mm) of emulsion and nano-emulsion forms of essential oils against bacterial pathogens using the well diffusion method.<sup>a</sup>

Bacteria	Forms of essential oils	<i>Salvia officinalis</i>	<i>Pimpinella anisum</i>	<i>Dracocephalum moldavica</i>	<i>Syzygium aromaticum</i>
<i>S. Typhimurium</i>	Emulsion	9.28±0.12	10.11±0.05	10.34±0.2	9.82±0.17
	Nanoemulsion	10.13±0.05 <sup>*Cc</sup>	10.94±0.22 <sup>*Bd</sup>	11.82±0.12 <sup>*d</sup>	10.78±0.2 <sup>*Bc</sup>
<i>S. aureus</i>	Emulsion	10.12±0.07	11.61±0.11	11.84±0.22	10.74±0.31
	Nanoemulsion	11.34±0.15 <sup>*Dd</sup>	12.86±0.15 <sup>*Bb</sup>	13.13±0.10 <sup>*Ac</sup>	11.89±0.23 <sup>*Cb</sup>
<i>S. dysenteriae</i>	Emulsion	8.62±0.17	9.24±0.09	9.66±0.13	8.76±0.22
	Nanoemulsion	9.92±0.31 <sup>*Cf</sup>	10.86±0.27 <sup>*Ad</sup>	11.03±0.09 <sup>*Ae</sup>	10.3±0.17 <sup>*Bd</sup>
<i>L. monocytogenes</i>	Emulsion	10.12±0.04	10.92±0.17	11.34±0.12	10.73±0.25
	Nanoemulsion	11.78±0.26 <sup>*Cc</sup>	12.36±0.2 <sup>*Bc</sup>	13.13±0.13 <sup>*Ac</sup>	12.32±0.27 <sup>*Bb</sup>
<i>P.aeruginosa</i>	Emulsion	10.26±0.15	10.92±0.27	10.84±0.12	10.13±0.09
	Nanoemulsion	11.82±0.26 <sup>*Cc</sup>	12.38±0.15 <sup>*Bc</sup>	13.02±0.12 <sup>*Ac</sup>	12.9±0.18 <sup>*Aa</sup>
<i>B. cereus</i>	Emulsion	10.63±0.26	11.36±0.02	11.24±0.10	10.82±0.16
	Nanoemulsion	12.14±0.1 <sup>*Cb</sup>	13.76±0.28 <sup>*Ba</sup>	14.10±0.13 <sup>*Aa</sup>	12.33±0.20 <sup>*Cb</sup>
<i>E. coli</i>	Emulsion	10.54±0.22	10.8±0.34	11.17±0.10	10.36±0.26
	Nanoemulsion	13.12±0.15 <sup>*Ba</sup>	13.81±0.23 <sup>*Aa</sup>	13.62±0.27 <sup>*Ab</sup>	12.14±0.19 <sup>*Cb</sup>

<sup>a</sup> The asterisk (\*) in each column for each bacterium separately indicates a significant difference between the inhibitory effect of the emulsion and nano-emulsion forms of the studied essential oils ( $P < 0.05$ ). Similar uppercase letters in each row and similar lowercase letters in each column indicate no significant difference between the mean antibacterial effects of the nanoemulsion forms of the studied essential oils ( $P < 0.05$ ).

**Table 7.** Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) ( $\mu\text{g mL}^{-1}$ ) of emulsion and nanoemulsion forms of essential oils against pathogenic bacteria.

Bacteria	Forms of essential oils	<i>Salvia officinalis</i>		<i>Pimpinella anisum</i>		<i>Dracocephalum moldavica</i>		<i>Syzygium aromaticum</i>	
		MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
<i>S. typhimurium</i>	Emulsion	10000	10000	2500	5000	2500	5000	5000	10000
	Nanoemulsion	5000	5000	1250	2500	1250	2500	2500	5000
<i>S. aureus</i>	Emulsion	2500	5000	1250	2500	1250	2500	2500	5000
	Nanoemulsion	1250	2500	625	1250	312.5	625	625	1250
<i>S. dysenteriae</i>	Emulsion	5000	10000	2500	5000	2500	5000	2500	5000
	Nanoemulsion	2500	2500	625	1250	625	625	625	1250
<i>L. monocytogenes</i>	Emulsion	2500	5000	1250	2500	1250	2500	2500	5000
	Nanoemulsion	1250	1250	625	1250	312.5	625	625	1250
<i>P.aeruginosa</i>	Emulsion	2500	5000	2500	5000	1250	2500	2500	5000
	Nanoemulsion	1250	2500	2500	5000	1250	2500	1250	2500
<i>B. cereus</i>	Emulsion	1250	2500	1250	2500	625	1250	2500	5000
	Nanoemulsion	625	1250	312.5	625	312.5	625	1250	2500
<i>E. coli</i>	Emulsion	5000	10000	2500	5000	2500	5000	5000	5000
	Nanoemulsion	1250	2500	625	1250	625	1250	1250	2500

type of bacteria, concentration of the nano-emulsion, and the exposure period to bacteria. In 2016, Moghimi *et al.* (2016) evaluated and validated the antibacterial activity of thyme essential oil in its pure form and in nano-emulsion form against *E. coli* by stating that, for the nano-emulsion, the

antibacterial activity of the essential oil increased as it gained faster access to the bacterial cells.

The nano-emulsion of *Dracocephalum Moldavica* indicated the highest action upon the pathogenic microorganisms, followed by the nano-emulsion of *Pimpinella anisum*. The



inhibition diameters for the nano-emulsion against *S. dysenteriae*, *S. Typhimurium*, *P. aeruginosa*, *S. aureus*, *L. monocytogenes*, *E. coli*, and *B. cereus* were 11.03, 11.82, 13.02, 13.13, 13.13, 13.62, and 14.1 mm, respectively (Table 6). The findings indicate that this essential oil had the highest inhibitory effect against *B. cereus* and the least effect against *S. dysenteriae* and *S. Typhimurium* ( $P < 0.05$ ). The nano-emulsion of *Pimpinella anisum* also showed a significantly lower inhibitory effect against both *S. Typhimurium* and *S. dysenteriae* ( $P < 0.05$ ).

Microdilution method was also used to compare the antimicrobial activity of emulsified and nano-emulsified essential oils studied in this work. Results in Table 7 indicate that the antimicrobial activity of all essential oils was enhanced as a result of the decreasing Minimum Inhibitory Concentration (MIC) of the nano-emulsion compared to the emulsified form against all studied microorganisms. In fact, in many cases, the MIC of nanoemulsions was half of those of the emulsified essential oils. Moreover, as shown in Table 6, the highest antibacterial effects were observed of the nano-emulsion forms of *Dracocephalum Moldavica* and *Pimpinella anisum* against gram-positive bacteria.

In this study, antibacterial effects were evaluated by liquid dilution and well diffusion methods. Tables 6 and 7 show that among the aforementioned essential oils, *Dracocephalum moldavica* showed a relatively stronger antibacterial activity on the Gram positive than on the Gram-negative organisms. Antimicrobial activity of essential oils from plants is dependent on their chemical constituents. Analysis of essential oils originating from different plants using gas chromatography revealed that these oils contained many compounds such as monoterpenes, sesquiterpenes and other oxygenated compounds such as alcohols, aldehydes, esters, ethers, ketones and phenols (Nazemisalman et al., 2024). The main constituents of essential oil of *Dracocephalum moldavica* included

terpenoids; citral accounted for a substantial part of them, as investigated by Maham et al. (2013). Citral is well known for its prominent antimicrobial activity against Gram-positive and Gram-negative bacteria, the results of this research agree with those analyzed (Wójtowicz et al. 2017). El-Baky and El-Baroty (2008) evaluated the antibacterial activity of *Dracocephalum moldavica* essential oil through bio-autography and reported that, among its components, geraniol, neral, geranyl acetate, geranial, nerol, neryl acetate, and methyl nerolate exhibited antibacterial activity. They also showed that the essential oil of *Dracocephalum moldavica* possessed a significant antibacterial activity against *S. aureus*, *Micrococcus luteus* and *Serratia marcescens*. The present study further showed that 74.23% of the essential oil of *Dracocephalum moldavica* was composed of geraniol. Geraniol is an aliphatic monoterpene structure mainly related to the functional alcohol group in its organic composition. The possible mechanism of the antimicrobial action of geraniol, via its lipophilicity, would be its ability to attach to cell membrane lipids of the microorganism. Then, it would increase membrane permeability while binding to sites that are essential in cells, and eventually disrupts their structures (Lira et al., 2020).

#### Antioxidant Activity of the Essential Oils

The measured phenol content of the studied essential oils and their antioxidant activity are shown in Table 8. No significant difference was observed in the total phenol content between the emulsion and nano-emulsion forms of each essential oil ( $P < 0.05$ ). The results indicated that the essential oils of *Dracocephalum moldavica* and *Pimpinella anisum* had the highest, while the essential oil of *Salvia officinalis* had the lowest total phenol content.

**Table 8.** Comparison of the total phenolic content and antioxidant activity of various essential oils in two forms: emulsion and nano-emulsion.<sup>a</sup>

Parameter	Forms of essential oils	<i>Salvia officinalis</i>	<i>Pimpinella anisum</i>	<i>Dracocephalum moldavica</i>	<i>Syzygium aromaticum</i>	BHT
DPPH (IC50)	Emulsion	43.78±0.09 <sup>b</sup>	36.23±0.11 <sup>c</sup>	29.76±0.18 <sup>d</sup>	56.25±0.13 <sup>a</sup>	28.92±0.16 <sup>d</sup>
	Nanoemulsion	31.82±0.18 <sup>*b</sup>	29.56±0.21 <sup>*c</sup>	22.17±0.1 <sup>*c</sup>	47.16±0.12 <sup>*a</sup>	28.22±0.16 <sup>d</sup>
Reducing power (EC50)	Emulsion	6.27±0.17 <sup>a</sup>	5.36±0.13 <sup>b</sup>	5.82±0.10 <sup>c</sup>	6.11±0.08 <sup>a</sup>	1.31±0.12 <sup>d</sup>
	Nanoemulsion	5.13±0.14 <sup>*a</sup>	4.62±0.28 <sup>*b</sup>	4.51±0.20 <sup>*b</sup>	5.15±0.10 <sup>*a</sup>	1.31±0.12 <sup>c</sup>
Total phenol content (mg of gallic acid per g of essential oil)	Emulsion	259.14±0.52 <sup>d</sup>	317.12±0.38 <sup>b</sup>	392.51±0.41 <sup>a</sup>	289.53±0.12 <sup>c</sup>	
	Nanoemulsion	260.21±0.36 <sup>d</sup>	319.28±0.45 <sup>b</sup>	397.82±0.28 <sup>a</sup>	292.39±0.40 <sup>c</sup>	

<sup>a</sup> The asterisk (\*) in each column of the table, for each parameter separately, indicates a significant difference between the emulsion and nano-emulsion forms ( $P < 0.05$ ). Similar lowercase letters in each row indicate no significant difference ( $P < 0.05$ ).

In this study, the antioxidant effect of the essential oil emulsions and nanoemulsions was also evaluated using two methods: DPPH and reducing power. The results from both methods are presented in Table 8. As observed, although the antioxidant activity of both emulsions and nanoemulsions is clearly lower than the standard group (BHT) in both methods, the overall antioxidant activity of the nanoemulsions is higher than that of the emulsions, and this difference is statistically significant ( $P < 0.05$ ). The study also demonstrated that the IC50 of the essential oil nanoemulsions was significantly lower than that of the emulsions. According to the results, the lowest IC50 and EC50 values were found in the emulsion form of *Dracocephalum moldavica*, with values of 22.17 and 4.51  $\mu\text{g mL}^{-1}$ , respectively.

This is because of the phenolic content of the essential oils that are responsible for high-radical scavenging activity. Generally, increasing the concentration of phenolic compounds directly enhances the activity of different essential oils for inhibiting free radicals. It is inferred that at high concentrations of phenolic compounds, there is an increased number of hydroxyl groups present in the reaction environments, thus, increasing the probability of hydrogen donation to free radicals and leading to an increased scavenging capability of the extract. Many studies have shown the relationship between the electron-donating ability of bioactive compounds and their

scavenging activity against free radicals. Results indicate that the highest phenolic essential oils exhibit the strongest antioxidant (Baliyan *et al.*, 2022). In fact, the radical scavenging power of different essential oils depends to a large extent on the number and position of hydroxyl groups as well as the molecular weight of phenolic compounds. The more hydroxyl groups a phenolic compound has, the more readily accessible it becomes when dealing with other compounds in reactions and, therefore, is considered to be of lower molecular weights. Reducing properties in essential oils are also related to the presence of electron-donating compounds. In other words, as the amount of phenolic compounds in the extract increases, its reducing power also increases (Kumar and Goel, 2019; Liu and Yao, 2007).

## CONCLUSIONS

The findings of this study highlighted the potent antimicrobial activity of *Dracocephalum moldavica* essential oil against the major Gram-positive and Gram-negative foodborne pathogens. This effectiveness is attributed to its rich composition of bioactive compounds such as geraniol, geranial, alpha-copaene, and alpha-pinene. Considering the demonstrated antimicrobial and antioxidant properties of the nano-emulsion form of this essential oil, it holds great potential as a natural



preservative and antioxidant in the food industry. Its application could effectively control the proliferation of key foodborne pathogens, while enhancing the shelf life of food products. Furthermore, the high concentration of geraniol makes this essential oil valuable for use in other industries, including pharmaceuticals, cosmetics, and chemical manufacturing.

## REFERENCES

1. Abdel-Reheem, M. A., and Oraby, M. M. 2015. Anti-Microbial, Cytotoxicity, and Necrotic Ripostes of *Pimpinella anisum* Essential Oil. *Ann. Agric. Sci.*, **60(2)**: 335-340.
2. Aćimović, M., Šovljanski, O., Šeregelj, V., Pezo, L., Zheljazkov, V. D., Ljujić, J., Tomić, A., Četković, G., Čanadanović-Brunet, J., Miljković, A. and Vujisić, L. 2022. Chemical Composition, Antioxidant, and Antimicrobial Activity of *Dracocephalum moldavica* L. Essential Oil and Hydrolate. *Plants*, **11(7)**: 941.
3. Ali-Shtayeh, M. S., Jamous, R. M., Abu-Zaitoun, S. Y., Akkawi, R. J., Kalbouneh, S. R., Dudai, N., and Bernstein, N. 2018. Secondary Treated Effluent Irrigation Did Not Impact Chemical Composition, and Enzyme Inhibition Activities of Essential Oils from *Origanum syriacum* var. *Syriacum*. *Ind. Crops Prod.*, **111**: 775-786.
4. Alizadeh, A., Alizadeh, O., Amari, G., and Zare, M. 2013. Essential Oil Composition, Total Phenolic Content, Antioxidant Activity and Antifungal Properties of Iranian *Thymus daenensis* subsp. *daenensis* Celak. as in Influenced by Ontogenetical Variation. *J. Essent. Oil Bear. Plants*, **16(1)**: 59-70.
5. Baliyan, S., Mukherjee, R., Priyadarshini, A., Vibhuti, A., Gupta, A., Pandey, R. P., and Chang, C. M. 2022. Determination of Antioxidants by DPPH Radical Scavenging Activity and Quantitative Phytochemical Analysis of *Ficus religiosa*. *Molecules*, **27(4)**: 1326.
6. Borrin, T. R., Georges, E. L., Moraes, I. C. and Pinho, S. C. 2016. Curcumin-Loaded Nanoemulsions Produced by the Emulsion Inversion Point (EIP) Method: An Evaluation of Process Parameters and Physico-Chemical Stability. *J. Food Eng.*, **169**: 1-9.
7. Burt, S. 2004. Essential Oils: Their Antibacterial Properties and Potential Applications in Foods— A Review. *Int. J. Food Microbiol.*, **94(3)**: 223-253.
8. Chamorro, E. R., Zambón, S. N., Morales, W. G., Sequeira, A. F. and Velasco, G. A. 2012. Study of the Chemical Composition of Essential Oils by Gas Chromatography. *Gas Chromatography in Plant Science, Wine Technology, Toxicology and Some Specific Applications*, **1**: 307-324.
9. Couladis, M., Tzakou, O., Mimica-Dukić, N., Jančić, R. and Stojanović, D. 2002. Essential Oil of *Salvia officinalis* L. from Serbia and Montenegro. *Flavour Fragr. J.*, **17(2)**: 119-126.
10. Delamare, A. P. L., Moschen-Pistorello, I. T., Artico, L., Atti-Serafini, L. and Echeverrigaray, S. 2007. Antibacterial Activity of the Essential Oils of *Salvia officinalis* L. and *Salvia triloba* L. Cultivated in South Brazil. *Food Chem.*, **100(2)**: 603-608.
11. Eghbal, H., Mohammadi, A., Mohammad Nejad Khiavi, N., Ahmadi Sabegh, M. and Jahani, N. 2021. Comparison of the Antibacterial Properties of Essential Oils of *Malva sylvestris* and *Salvia officinalis* on Common Bacteria of Oral Infection with Chlorhexidine Mouthwash. *J. Mashhad Dent. Sch.*, **45(3)**: 217-229.
12. El-Baky, H. A., and El-Baroty, G. S. 2008. Chemical and Biological Evaluation of the Essential Oil of Egyptian Moldavian Balm. *Int. J. Essent. Oil Ther.*, **2**: 78-81.
13. Erkan, N., Ayranci, G., and Ayranci, E. 2008. Antioxidant Activities of Rosemary (*Rosmarinus officinalis* L.) Extract, Blackseed (*Nigella sativa* L.) Essential Oil, Carnosic Acid, Rosmarinic Acid and Sesamol. *Food Chem.*, **110(1)**: 76-82.
14. Fichi, G., Flamini, G., Giovanelli, F., Otranto, D. and Perrucci, S. 2007. Efficacy of an Essential Oil of *Eugenia caryophyllata* against *Psoroptes cuniculi*. *Exp. Parasitol.*, **115(2)**: 168-172.
15. Ghosh, V., Mukherjee, A. and Chandrasekaran, N. 2013. Formulation and Characterization of Plant Essential Oil Based Nanoemulsion: Evaluation of Its Larvicidal Activity against *Aedes aegypti*. *Asian J. Chem.*, **25(Suppl. Issue)**: S321.

16. Grausgruber-Gröger, S., Schmiderer, C., Steinborn, R. and Novak, J. 2012. Seasonal Influence on Gene Expression of Monoterpene Synthases in *Salvia officinalis* (Lamiaceae). *J. Plant Physiol.*, **169(4)**: 353-359.
17. Hakkak, M., Hozni S., Morovati, H. and Akhlaghi, T. 2018. Research Diagnosis in Postgraduate Students of Guilan University of Medical Sciences. *Res. Med. Educ.*, **10(2)**: 46-57
18. Hendawy, S. F., El Gendy, A. G., Omer, E. A., Pistelli, L. and Pistelli, L. 2018. Growth, Yield and Chemical Composition of Essential Oil of *Mentha piperita* var. Multimentha Grown under Different Agro-Ecological Locations in Egypt. *J. Essent. Oil Bear. Plants*, **21(1)**: 23-39.
19. Holm, Y., Hiltunen, R. and Nykanen, I. 1988. Capillary Gas Chromatographic-Mass Spectrometric Determination of the Flavour Composition of Dragonhead (*Dracocephalum moldavica* L.). *Flavour Fragr. J.*, **3(3)**: 109-112.
20. Kakasy, A. Z., Lemberkovics, E., Simandi, B., Lelik, L., Hethelyi, E., Antal, I. and Szöke, É. 2006. Comparative Study of Traditional Essential Oil and Supercritical Fluid Extracts of Moldavian Dragonhead (*Dracocephalum moldavica* L.). *Flavour Fragr. J.*, **21(4)**, 598-603.
21. Kelen, M. and Tepe, B. 2008. Chemical Composition, Antioxidant and Antimicrobial Properties of the Essential Oils of Three *Salvia* Species from Turkish Flora. *Bioresour. Technol.*, **99(10)**: 4096-4104.
22. Kennouche, A., Benkaci-Ali, F., Scholl, G. and Epe, G. 2015. Chemical Composition and Antimicrobial Activity of the Essential Oil of *Eugenia caryophyllata* Cloves Extracted by Conventional and Microwave Techniques. *J. Biol. Active Prod. Nat.*, **5(1)**: 1-11.
23. Kumar, N. and Goel, N. 2019. Phenolic Acids: Natural Versatile Molecules with Promising Therapeutic Applications. *Biotech. Rep.*, **24**: e00370.
24. Lira, M. H. P. D., Andrade Júnior, F. P. D., Moraes, G. F. Q., Macena, G. D. S., Pereira, F. D. O. and Lima, I. O. 2020. Antimicrobial Activity of Geraniol: An Integrative Review. *J. Essent. Oil Res.*, **32(3)**: 187-197.
25. Liu, Q., and Yao, H. 2007. Antioxidant Activities of Barley Seeds Extracts. *Food Chem.*, **102(3)**: 732-737.
26. Maham, M., Akbari, H. and Delazar, A. 2013. Chemical Composition and Antinociceptive Effect of the Essential Oil of *Dracocephalum moldavica* L. *Pharm. Sci.*, **18(4)**:187-192.
27. McClements, D. J. 2012. Nanoemulsions versus Microemulsions: Terminology, Differences, and Similarities. *Soft Matter*, **8(6)**: 1719-1729.
28. McClements, D. J. and Rao, J. 2011. Food-Grade Nanoemulsions: Formulation, Fabrication, Properties, Performance, Biological Fate, and Potential Toxicity. *Crit. Rev. Food Sci. Nutr.*, **51(4)**: 285-330.
29. Mirtaghi, S. M., Nejad, P. T., Masoumeh Mazandarani, M., Livani, F. and Bagheri, H. 2016. Evaluation of Antibacterial Activity of *Urtica dioica* L. Leaf Ethanolic Extract Using Agar Well Diffusion and Disc Diffusion Methods. *Med. Lab. J.*, **10(5)**: 15-21.
30. Moghimi, R., Ghaderi, L., Rafati, H., Aliahmadi, A., and McClements, D. J. 2016. Superior Antibacterial Activity of Nanoemulsion of *Thymus daenensis* Essential Oil against *E. coli*. *Food Chem.*, **194**: 410-415.
31. Myint, S., Daud, W. R. W., Mohamad, A. B. and Kadhum, A. A. H. 1996. Gas Chromatographic Analysis of Eugenol in Alcohol Clove Extract of Cloves. *J. Chromatogr. B.*, **679**: 193-195.
32. Nassar, M. I., Gaara, A. H., El-Ghorab, A. H., Farrag, A., Shen, H., Huq, E. and Mabry, T. J. 2007. Chemical Constituents of Clove (*Syzygium aromaticum*, Fam. Myrtaceae) and Their Antioxidant Activity. *Rev. Latinoam. Quím.*, **35(3)**: 47.
33. Nazemismalman, B., Taheri, S. S., Heidari, F., Yazdinezhad, A., Hagh, F., Shabouei Jam, M. and Basir Shabestari, S. 2024. Comparison of *Dracocephalum Moldavica* Essential Oil with Chlorhexidine on Cariogenic Bacteria. *J. Dent.*, **25(3)**: 223-228.
34. Nazzaro, F., Fratianni, F., De Martino, L., Coppola, R. and De Feo, V. 2013. Effect of Essential Oils on Pathogenic Bacteria. *Pharmaceuticals*, **6(12)**: 1451-1474.
35. Orav, A., Raal, A. and Arak, E. 2008. Essential Oil Composition of *Pimpinella*



- anisum* L. Fruits from Various European Countries. *Nat. Prod. Res.*, **22(3)**: 227-232.
36. Oyaizu, M. 1986. Antioxidative Activities of Browning Reaction Prepared from Glucosamine. *Jap. J. Nutr. Diet.*, **44**, 307-315.
37. Perugini Biasi-Garbin, R., Saori Otaguiri, E., Morey, A. T., Fernandes da Silva, M., Belotto Morguette, A. E., Armando Contreras Lancheros, C., Kian, D., Perugini, M. R. E., Nakazato, G., Durán, N., Nakamura, C. V., Yamauchi, L. M. and Yamada-Ogatta, S. F. 2015. Effect of Eugenol against *Streptococcus agalactiae* and Synergistic Interaction with Biologically Produced Silver Nanoparticles. *Evid. Based Complement. Alternat. Med.*, **2015(1)**: 861497.
38. Purkait, S., Bhattacharya, A., Bag, A. and Chattopadhyay, R. R. 2018. Evaluation of Antibacterial and Antioxidant Activities of Essential Oils of *Fve* Spices. *J. Food Qual. Hazards Control*, **5(2)**: 61–71.
39. Rao, J. and McClements, D. J. 2011. Food-Grade Microemulsions, Nanoemulsions and Emulsions: Fabrication from Sucrose Monopalmitate and Lemon Oil. *Food Hydrocoll.*, **25(6)**: 1413-1423.
40. Ruiz-Navajas, Y., Viuda-Martos, M., Sendra, E., Perez-Alvarez, J. A. and Fernández-López, J. 2012. Chemical Characterization and Antibacterial Activity of *Thymus moroderi* and *Thymus piperella* Essential Oils, Two *Thymus* Endemic Species from Southeast of Spain. *Food Control*, **27(2)**: 294-299.
41. Shahabi, N., Tajik, H., Moradi, M., Forough, M. and Ezati, P. 2017. Physical, Antimicrobial and Antibiofilm Properties of *Zataria multiflora* Boiss Essential Oil Nanoemulsion. *Int. J. Food Sci. Technol.*, **52(7)**: 1645-1652.
42. Shahbazi, Y., Shavisi, N., Modarresi, M. and Karami, N. 2016. Chemical Composition, Antibacterial and Antioxidant Activities of Essential Oils from the Aerial Parts of *Ferulago angulata* (Schlecht.) Boiss and *Ferulago bernardii* Tomk. and *M. pimen* from Different Parts of Iran. *J. Essent. Oil Bear. Plants*, **19(7)**: 1627-1638.
43. Shavisi, N., Khanjari, A., Basti, A. A., Misaghi, A. and Shahbazi, Y. 2017. Effect of PLA Films Containing Propolis Ethanolic Extract, Cellulose Nanoparticle and *Ziziphora clinopodioides* Essential Oil on Chemical, Microbial and Sensory Properties of Minced Beef. *Meat Sci.*, **124**: 95-104.
44. Singh, G., Kapoor, I. P. S., de Heluani, C. S. and Catalan, C. A. N. 2008. Chemical Composition and Antioxidant Potential of Essential Oil and Oleoresins from Anise Seeds (*Pimpinella anisum* L.). *Int. J. Essent. Oil Ther.*, **2**: 122-130
45. Skandamis, P., Koutsoumanis, K., Fasseas, K. and Nychas, G. J. E. 2001. Inhibition of Oregano Essential Oil and EDTA on *Escherichia coli* O157: H7. *Ital. J. Food Sci.*, **13(1)**: 65-75.
46. Sonboli, A., Mojarrad, M., Gholipour, A., Ebrahimi, S. N. and Arman, M. 2008. Biological Activity and Composition of the Essential Oil of *Dracocephalum moldavica* L. Grown in Iran. *Nat. Prod. Commun.*, **3(9)**: 1934578X0800300930.
47. Topuz, O. K., Özvural, E. B., Zhao, Q., Huang, Q., Chikindas, M. and Gölükçü, M. 2016. Physical and Antimicrobial Properties of Anise Oil Loaded Nanoemulsions on the Survival of Foodborne Pathogens. *Food Chem.*, **203**: 117-123.
48. Turgis, M., Vu, K. D., Dupont, C. and Lacroix, M. 2012. Combined Antimicrobial Effect of Essential Oils and Bacteriocins against Foodborne Pathogens and Food Spoilage Bacteria. *Food Res. Int.*, **48(2)**, 696-702.
49. Ullah, H., Mahmood, A., and Honermeier, B. 2014. Essential Oil and Composition of Anise (*Pimpinella anisum* L.) with Varying Seed Rates and Row Spacing. *Pak. J. Bot.*, **46(5)**: 1859-1864.
50. Wójtowicz, A., Oniszczyk, A., Oniszczyk, T., Kocira, S., Wojtunik, K., Mitrus, M., Kocira, A., Widelski, J. and Skalicka-Woźniak, K. 2017. Application of Moldavian Dragonhead (*Dracocephalum moldavica* L.) Leaves Addition as a Functional Component of Nutritionally Valuable Corn Snacks. *J. Food Sci. Technol.*, **54**: 3218-3229.
51. Yousefzadeh, S., Modarres-Sanavy, S. A. M., Sefidkon, F., Asgarzadeh, A., Ghalavand, A., and Sadat-Asilan, K. 2013. Effects of Azocompost and Urea on the Herbage Yield and Contents and Compositions of Essential Oils from Two Genotypes of Dragonhead (*Dracocephalum moldavica* L.) in Two Regions of Iran. *Food Chem.*, **138(2-3)**: 1407-1413.

اثرات ضد میکروبی و آنتی اکسیدانی امولسیون و نانوامولسیون گیاهان مریم گلی، بادیان رومی، بادرشبی و میخک بر علیه باکتری های با منشأ مواد غذایی

مجتبی رئیسی، بهادر حاجی محمدی، و نگین مهدی نژاد

### چکیده

اسانس های گیاهی به دلیل اثرات ضد میکروبی و آنتی اکسیدانی به عنوان نگهدارنده های طبیعی مورد استفاده قرار می گیرند. هدف از انجام این مطالعه، بررسی ترکیبات شیمیایی، اثرات آنتی اکسیدانی، و ضد میکروبی امولسیون و نانوامولسیون گیاهان مریم گلی، بادیان رومی، بادرشبی و میخک می باشد. نتایج آزمون انتشار از چاهک مشخص کرد که فرم نانوامولسیون اسانس بادرشبی بیشترین اثر ضد میکروبی را بر علیه باکتری های مورد بررسی داشته است. قطر هاله عدم رشد فرم نانوامولسیون اسانس بادرشبی بر علیه شیگلا دیسانتری، سالمونلا تایفی موریوم، سودوموناس آنروژینوزا، استافیلوکوکوس اورئوس، لیستریا مونوسیژنوزا، اشرشیا کلی و باسیلوس سرئوس به ترتیب 11.03، 11.82، 13.02، 13.13، 13.13، 13.62 و 14.10 بوده است. در مقابل، قطر هاله عدم رشد فرم امولسیون این اسانس بر علیه شیگلا دیسانتری، سالمونلا تایفی موریوم، سودوموناس آنروژینوزا، استافیلوکوکوس اورئوس، لیستریا مونوسیژنوزا، اشرشیا کلی و باسیلوس سرئوس به ترتیب 9.66، 10.34، 10.84، 11.84، 11.34، 11.17 و 11.24 بود. ترکیبات غالب در اسانس بادرشبی شامل ژرانیول (27.24%)، ژرانیال (10.75%)، آلفاکوپائن (8.16%)، آلفاپنین (7.37%)، کارواکرول (7.41%)، لیمونن (6.86%) و نرول (6.45%) بود. فرم نانوامولسیون اسانس های گیاهی پتانسیل اثرات آنتی اکسیدانی بالاتری نسبت به فرم امولسیون آن ها نشان دادند. این مطالعه همچنین نشان داد که فرم نانوامولسیون اسانس های گیاهی نیمه حداکثر غلظت بازدارندگی (Ic50) پایین تری نسبت به فرم امولسیون آن ها داشتند. بر اساس نتایج نشان داده شد فرم نانوامولسیون اسانس بادرشبی کمترین نیمه حداکثر غلظت بازدارندگی (Ic50) و غلظت موثر 50 درصد بازدارنده رشد (Ec50) با اعداد به ترتیب 22.17 میکروگرم در میلی لیتر و 4.51 میکروگرم در میلی لیتر داشت.