

## Influence of Pre-Imaginal Stages of *Bemisia tabaci* on Development, Life Table Parameters, and Predation Rate of *Euseius scutalis* (Acari: Phytoseiidae)

Mohammad Gravandian<sup>1</sup>, Parviz Shishehbor<sup>1\*</sup>, Mehdi Esfandiari<sup>1</sup>, and Elham Riahi<sup>2</sup>

### ABSTRACT

The predatory mite, *Euseius scutalis* (Athias-Henriot), was studied in terms of its development, survival, and life table parameters on two preimaginal stages (egg and first instar nymph) of its prey, *Bemisia tabaci* (Gennadius) (Hem.: Aleyrodidae). The first instar nymphs of *B. tabaci* were found to be the preferred food for *E. scutalis*, resulting in decreased developmental time from egg to adult, as well as a shorter pre-oviposition period and a higher rate of oviposition. The intrinsic rate of increase ( $r$ ) was found to be  $0.1503 \text{ d}^{-1}$  on the first instar nymphs and  $0.0843 \text{ d}^{-1}$  on eggs of the prey. On average, females' *E. scutalis* consumed 16.30 eggs and 29.40 nymphs from their emergence to death. When first instar nymphs of *B. tabaci* were provided, *E. scutalis* showed a higher net predation rate ( $C_0$ ) and finite predation rate ( $\omega$ ) compared to feeding on eggs. On average, it consumed 3.52 eggs or 2.76 first instar nymphs of *B. tabaci* to produce a single egg of *E. scutalis*. In terms of sex ratio of the progeny, predatory females that fed on the first instar nymphs produced more females.

**Keywords:** Cotton whitefly, Growth rate, Phytoseiidae, Predation rate, Sex ratio.

### INTRODUCTION

The agricultural pest known as the cotton whitefly, *Bemisia tabaci* (Gennadius) (Hem.: Aleyrodidae) is one of the major global threats to different crops. *B. tabaci* damages greenhouse vegetables and ornamental plants by sucking sap from the plants, which can reduce production (De Barro *et al.*, 2011). This is primarily due to their direct damage to host plants through leaf piercing, sap sucking, and honeydew secretion (Jones, 2003). It has a high reproductive capacity and is capable of causing significant damage through the secretion of honeydew, which promotes the rapid growth of molds (Gangwar and

Gangwar, 2018). The use of insecticides to eliminate this pest raises production costs and contributes to the development of pest resistance.

The crucial role of predatory mites of the family Phytoseiidae in biological control is widely recognized (McMurtry *et al.*, 2013). These efficient predators play a significant role in managing pest mites and other small arthropods in crops grown in both greenhouses and open fields worldwide. With over 2,700 species reported on all continents, except Antarctica, the mites from this family make up 60% of the global biological control market (Momen *et al.*, 2020). They are considered a vital solution for sustainable plant protection and for promoting integrated pest management

<sup>1</sup> Department of Plant Protection, Faculty of Agriculture, Shahid Chamran University of Ahvaz, Ahvaz, Islamic Republic of Iran.

<sup>2</sup> Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, P. O. Box: 14115-336, Tehran, Islamic Republic of Iran.

\*Corresponding author; e-mail: pshishehbor@scu.ac.ir



against spider mites, thrips, and whiteflies (Yazdanpanah and Fathipour, 2022).

*Euseius scutalis* (Athias-Henriot) is a common phytoseiids in Middle East and North Africa (Bounfour and McMurtry, 1987). This predatory mite is one of the most dominant predator on economic crops in Iran (Kamali et al., 2001), and it is one of the pollen-feeding generalist species of the Phytoseiid mite (McMurtry et al., 2013). It is a significant predatory mite in biological control due to its ability to effectively manage pest populations. Although this predatory mite has the highest reproductive capacity when feeding on pollen, it is also capable of consuming a variety of small insects and mites, such as eriophyid, tenuipalpid and spider mites, eggs, and immature stages of whiteflies, thrips and scale insects (Abou-Elella et al., 2013; Stathakis et al., 2021). *E. scutalis* exhibits optimal life cycle parameters and fecundity when feeding on *Tetranychus turkestanii* (Ugarov and Nikolskii), followed by date palm pollen, making it a versatile and effective biocontrol agent (Shishehbor et al., 2022). Previous studies have shown a close association between this species and *B. tabaci* (Meyerdirk and Coudriet, 1986). The use of *E. scutalis* in integrated pest management programs can reduce reliance on chemical pesticides, promoting a safer and more sustainable agricultural ecosystem.

While some researchers have provided information on the life table of *E. scutalis* when fed on *B. tabaci* compared to other prey species or pollen grains (Nomikou et al., 2001, Nawar 2017), there is a lack of data regarding the life table of this predatory mite in relation to the different growth stages of *B. tabaci*. Therefore, this study aimed to assess the influence of two different developmental stages (eggs and first instar nymphs) of the *B. tabaci* on the biological characteristics and predation rate of *E. scutalis*, which plays a vital role in biological control.

## MATERIALS AND METHODS

### Mite Rearing

The population of *E. scutalis* was obtained from a laboratory-grown culture at the Faculty of Agriculture, Shahid Chamran University of Ahvaz, Iran. The initial population, identified by Dr. Farid Faraji from Mitox (Amsterdam, The Netherlands), was collected in February 2021 from marshmallow plants (*Althea officinalis* L.) infested with *T. turkestanii* on the university campus. For laboratory rearing, we used plastic green sheets measuring 3 cm in diameter, placed on a water-saturated sponge in a Petri dish measuring 6 cm in diameter and filled with water. The edges of the sheet were surrounded by strips of tissue paper. Cotton threads were placed on the plastic sheets to provide egg-laying sites and shelter for the predators (Walzer and Schausberger, 1999). As a food source, we provided date palm pollen (*Phoenix dactylifera* L.) on the plastic sheets. The mites were fed this pollen grain for approximately one month before starting the experiments. Date palm pollen was chosen because it has been recommended as one of the most suitable alternative diet sources for the easy and cost-effective rearing of *E. scutalis* (Shishehbor et al., 2022). The rearing units were held in a growth chamber at  $25\pm 1^{\circ}\text{C}$ ,  $60\pm 10\%$  RH, and a photoperiod of 16:8 (L:D).

### Whitefly Rearing

A number of mature cotton whiteflies were collected using an aspirator from infested plants in the greenhouse at Shahid Chamran University of Ahvaz, Faculty of Agriculture. The whiteflies were then reared on cucumber plants inside cages (60×60×120 cm). Identification of this species was carried out using the taxonomic keys of Martin (1987). The cucumber plants (Var. Negin) used for colony maintenance

were reared at  $25\pm 1^\circ\text{C}$ ,  $60\%\pm 10\%$  RH, and a photoperiod of 16L:8D. Every week, a new cucumber plant with at least eight fully developed leaves was added to each cage.

### Experimental Setup

Leaves of a cucumber plant infested with *B. tabaci* were carefully selected and cut into smaller pieces. Each piece was then examined under a stereomicroscope to confirm the presence of only 10 fresh *B. tabaci* eggs. For the first instar nymphs, the same procedure was repeated, but the leaves were cut after the eggs had developed to the first instar nymph stage (10 first instar stable nymphs). The test units were prepared the same as the units used for mite rearing, except using a piece of cucumber leaf (3 cm in diameter) containing whitefly eggs or first instar nymphs pad instead of using plastic sheet.

#### Effect of *B. tabaci* Immature Stages on Predatory Mite *E. scutalis*

About 50 females of *E. scutalis* were separated from the rearing unit using a fine brush and transferred to a new experimental unit. After 24 hours, the eggs that were less than 24 hour-old were placed individually in petri dishes containing one of the two pre-adult stages of *B. tabaci*. The petri dishes were checked daily to record survival and developmental durations until adulthood. Once they became adults, the females and males were paired and transferred to a new petri dish. Each day, we checked the experimental units to confirm oviposition and note the consumption. The leaf in each arena was replaced daily with a new piece of fresh food. The number of prey consumed by each pair of predatory mite was recorded daily. Additionally, the number of eggs laid by each female was recorded daily until all the females died. The survival of both females and males was monitored daily until all individuals had died. Each experiment

was repeated 60 times. The tests were conducted in a controlled-temperature cabinet at  $25\pm 1^\circ\text{C}$ ,  $60\%\pm 10\%$  RH, and a photoperiod of 16:8 (L:D). *E. scutalis* eggs collected daily and were transferred to new petri dishes containing a piece of cucumber leaf containing whitefly eggs or first instar nymphs. This procedure was repeated on all days when oviposition occurred. After they matured, the gender of the predators was determined to estimate the daily sex ratio.

### Data Analysis

The computer programs TWSEX-MSChart and CONSUME-MS Chart were utilized for analyzing the life history and the daily consumption of *E. scutalis* on the two stages of *B. tabaci*, respectively (Chi and Liu, 1985; Chi, 1988; Chi, 2023a, b). The age-stage-specific survival rate ( $s_{xj}$ ), age-stage-specific fecundity ( $f_{xj}$ ), age-specific survival rate ( $l_x$ ), age-specific fecundity ( $m_x$ ), age-stage life expectancies ( $e_{xj}$ ), and age-stage reproductive value ( $v_{xj}$ ) along with population growth parameters were calculated using TWSEX-MS Chart program (Chi, 2023a). Based on Chi and Yang (2003), net predation rate ( $C_0$ ), transformation rate ( $Q_p$ ), stable predation rate ( $\psi$ ), and finite predation rate ( $\omega$ ), were computed using CONSUME-MS Chart program (Chi and Yang, 2003; Chi, 2023b). Means and standard errors of the all parameters were calculated by bootstrap method with 100000 resamplings. Paired bootstrap test using the TWSEX-MS Chart program (Chi, 2023a) was used for mean comparisons.

## RESULTS

### Immature Developmental Time and Survival

Individuals of *E. scutalis* successfully developed from larvae to adults when fed on eggs and first instar nymphs of *B. tabaci*



(Table 1). The time spent by *E. scutalis* in each developmental stage was affected by the immature stage of *B. tabaci*. Development from egg to adult was significantly faster for both females and males when feeding on the first instar nymphs of *B. tabaci* compared to eggs. While the adult longevity and total lifespan of female *E. scutalis* fed on the first instar nymphs of *B. tabaci* were not different from those reared on eggs, these parameters for males were significantly longer on the first instar nymphs of *B. tabaci* than on its eggs (Table 1). Additionally, there were no significant differences between the sexes in terms of immature development on either food item. When *B. tabaci* eggs were the food source, males exhibited shorter adult longevity and total lifespan compared to the females (Table 1).

The overlapping observed in Figure 1A can be explained by the fact that individuals at different stages had varying development rates. *Euseius scutalis* exhibited high survivorship during the immature development stage. Females that were fed eggs and the first instar nymphs of *B. tabaci* survived for 39 and 32 days, respectively. The males lived for 24 days when fed on the eggs of *B. tabaci*, while they survived for 31 days when fed on the first instar nymphs.

### Fecundity and Life Table Parameters


The consumption of the first instar nymphs of *B. tabaci* resulted in a shorter Average Pre-Oviposition Period (APOP) of *E. scutalis* compared to its eggs (Table 1). Similarly, significant differences in the Total

**Table 1.** Comparative duration (Mean±SE) of developmental time, adult longevity, total life span, APOP, TPOP, and oviposition days (days), as well as, fecundity and sex ratio of females and males of *Euseius scutalis* fed on egg and first-instar nymph of *Bemisia tabaci*.<sup>a</sup>

Parameter	Prey stage	
	Egg	Nymph (L1)
Female egg	2.17 ±0.10 a A	2.07 ±0.05 a A
Male egg	2.27 ±0.19 a A	2.26±0.10 a A
Female larva	1.50 ±0.09 a A	1.13 ±0.06 b A
Male larva	1.64 ±0.15 a A	1.21±0.12 b A
Female protonymph	1.90 ±0.12 a A	1.57 ±0.09 b A
Male protonymph	2.00 ±0.13 a A	1.53 ±0.12 b A
Female deutonymph	2.17 ±0.14 a A	2.10 ±0.13 a A
Male deutonymph	2.36 ±0.24 a A	1.68 ±0.17 b A
Female developmental time	7.73 ±0.21 a A	6.87 ±0.15 b A
Male developmental time	8.27 ±0.24 a A	6.68 ±0.19 b A
Female adult longevity	12.80 ±1.29 a A	14.93 ±0.85 a A
Male adult longevity	9.45 ±1.27 b B	14.74 ±1.27 a A
Female total life span	20.53 ±1.40 a A	21.80 ±0.86 a A
Male total life span	17.73 ±1.28 b B	21.42 ±1.25 a A
APOP*	3.92 ±0.35 a	2.67 ±0.15 b
TPOP*	11.73 ±0.45 a	7.87 ±2.00 b
Oviposition days	5.42 ±0.83 b	8.93 ±0.54 a
Fecundity (Eggs/Female)	6.50 ± 1.09 b	15.67 ± 1.47 a
Sex ratio (Nf/N)*	0.67 ± 0.07 a	0.56 ± 0.07 a

<sup>a</sup> Means (±SE) within rows followed by lowercase different letters are significantly different (100,000 resamplings, paired bootstrap test,  $P < 0.05$ ). Mean values followed by different capital letters shows the significant difference between female and male for each parameter ( $P < 0.05$ ), based on paired bootstrap test with 100,000 samples.

\* APOP: Adult Pre-Oviposition Period, TPOP: Total Pre-Oviposition Period. Nf/N: Number of females to Number of individuals.



**Figure 1.** The age-stage survival rate ( $s_{xj}$ ) (A) and the age-specific survivorship ( $l_x$ ), age-stage specific fecundity of female ( $f_{xj}$ ) (eggs) and age-specific fecundity ( $m_x$ ) (B) of *Euseius scutalis* fed on egg and first-instar nymph of *Bemisia tabaci*.

Pre-Oviposition Period (TPOP) were observed between the different immature stages of *B. tabaci*. When the predator was fed the first instar nymphs of *B. tabaci*, the adult females laid eggs more rapidly ( $2.67 \pm 0.15$  days) compared to when eggs of this prey were offered ( $3.92 \pm 0.35$  days). *Euseius scutalis* that fed *B. tabaci* eggs had a shorter oviposition duration compared to those that used *B. tabaci* first instar nymphs.

Figure 1B summarizes the age-specific survivorship ( $l_x$ ), the age-specific fecundity rate ( $m_x$ ), and female age-specific fecundity ( $f_x$ ) for *E. scutalis* reared on different life stages of *B. tabaci*. The maximum values of  $f_x$  for *E. scutalis* on the first instar nymphs


and eggs of *B. tabaci* were 2.54 and 2 eggs/individual/day, respectively, which occurred on the 13<sup>th</sup> and 33<sup>rd</sup> days, respectively.

All the population parameters of *E. scutalis* were significantly influenced by the prey stage, except for the Gross Reproductive Rate (GRR) (Table 2). When *B. tabaci* first instar nymphs were offered as food,  $R_0$ ,  $r$ , and  $\lambda$  of the predator mite were higher compared to when whitefly eggs were offered. Similarly, the mean generation time of *E. scutalis* on the former stage of prey was shorter than on the latter stage (Table 2).

**Table 2.** Population parameters of *Euseius scutalis* fed on egg and first-instar nymph of *Bemisia tabaci*.<sup>a</sup>


Parameter	Egg	Nymph (L1)
$GRR$ (Eggs/Individuas)	17.77 $\pm$ 3.27 a	11.12 $\pm$ 1.54 a
$R_0$ (Eggs/Individuas)	4.33 $\pm$ 0.85 b	8.70 $\pm$ 1.34 a
$r$ ( $d^{-1}$ )	0.084 $\pm$ 0.01 b	0.150 $\pm$ 0.01 a
$\lambda$ ( $d^{-1}$ )	1.088 $\pm$ 0.01 b	1.162 $\pm$ 0.01 a
$T$ (Day)	17.38 $\pm$ 1.11 a	14.39 $\pm$ 0.18 b

<sup>a</sup> Means ( $\pm$ SE) within rows followed by different letters are significantly different (100,000 resamplings, paired bootstrap test,  $P < 0.05$ ).  $GRR$ : Gross Reproductive Rate,  $R_0$ : Net reproductive rate,  $r$ : Intrinsic rate of increase,  $\lambda$ : Finite rate of increase,  $T$ : Mean generation time.

**Figure 2.** The age-stage specific life expectancy ( $e_{xj}$ ) (A) and the age-stage reproductive value ( $v_{xj}$ ) (B) of *Euseius scutalis* fed on egg and first-instar nymph of *Bemisia tabaci*.

The age-stage life expectancies ( $e_{xj}$ ) of *E. scutalis*, which represent the expected lifespan of individuals at age  $x$  and stage  $j$  after age  $x$ , are illustrated in Figure 2A for two different life stages of *B. tabaci*. As age increased, the  $e_{xj}$  values gradually decreased. Notably,  $e_{xj}$  was higher for nymphs compared to eggs (Figure 2-A), indicating that *E. scutalis* feeding on nymphs of *B. tabaci* lived longer than those

that fed on eggs. The age-stage reproductive value ( $v_{xj}$ ), which indicates the contribution of an individual at age  $x$  and stage  $j$  to the future population, increased significantly when adults emerged and the peak of  $v_{xj}$  for the predator occurred at ages 28 and 12 days on the egg and nymph stages of *B. tabaci*, respectively (Figure 2-B).



**Figure 3.** Offspring sex ratio of females of *Euseius scutalis* fed on egg and first-instar nymph of *Bemisia tabaci*. White and black bars indicate the percentages of male and female offspring, respectively.

### Sex Ratio

During the first days of oviposition, sex ratio was males-based (Figure 3). The percentage of female to male progeny per female parent reared on *B. tabaci* eggs during the initial three 24-hour periods were 00:00, 40:60 and 44:55 (on the first day, all collected eggs died before adult emergence). For *E. scutalis* females reared on *B. tabaci* first instar nymphs, similar measurements were 0:100, 61:38, and 36:63.

### Consumption Rate

The larval stage of *E. scutalis* did not prey on the eggs and the first instar nymphs of *B. tabaci*. The amount of prey consumed by *E. scutalis* immatures when *B. tabaci* eggs were provided did not differ from that

obtained when *B. tabaci* first instar nymphs were offered (Table 3). The predation rate of *E. scutalis* adults and their total life span were significantly affected by the prey stage. The consumption rate of both female and male *E. scutalis* was higher when they were fed on the first instar nymphs of *B. tabaci* rather than its eggs. When the first instar nymphs of *B. tabaci* were provided, *E. scutalis* showed a higher net predation rate ( $C_0$ ) and finite predation rate ( $\omega$ ) compared to when its eggs were provided. On average, it takes either 3.52 eggs or 2.76 first instar nymphs of *B. tabaci* to produce a single egg of *E. scutalis*, respectively (Table 3).

The age-specific predation rate ( $k_x$ ) refers to the average number of *B. tabaci* consumed by *E. scutalis* at age  $x$  (Figure 4). The highest value of this parameter was 2.08 and 1.62 on the first instar nymphs and eggs of *B. tabaci*, respectively, which was



observed at age 10. The age-specific net predation rate ( $q_x$ ) can be determined by taking into account the survivorship (Figure 4). Its highest value was estimated to be 1.88 and 1.40 on the aforementioned prey stages.

## DISCUSSION

In our study, we found that the different immature stages of *B. tabaci* as prey had a significant impact on the duration of *E. scutalis*' pre-adult stages and its total lifespan. We observed that motile prey stage (nymph L1) was more conducive to the growth of *E. scutalis*. Regardless of previous studies showing that the type of prey influences the fecundity and reproductive parameters of *E. scutalis* (Momen and Hussein, 2011), our research has revealed that the developmental stage of the prey can also impact these parameters. Female *E. scutalis* that fed on the first instar nymphs of *B. tabaci* had a shorter pre-oviposition period, longer oviposition period, and produced more eggs than those reared on eggs of *B. tabaci*. This suggests that prey on the first instar nymphs are more favorable to this predator than prey on eggs.

Prey stage preferences vary considerably among generalist and specialist phytoseiid predatory mites, with specialists typically preferring the egg stage over other stages of their prey (Blackwood *et al.*, 2001). According to the literature, some other predators, including *Macrolophus caliginosus* (Wagner) (Hem.: Miridae) and *Serangium parcesetosum* Sicard (Col.: Coccinellidae) also preferred the nymph stage of *B. tabaci* for optimal reproductive success (Bonato *et al.*, 2006; Firas and Sengonca, 2007). However, our findings differ from those reported by Meyerdirk and Coudriet (1986), who found that eggs, followed by the first instars of *B. tabaci*, were the most suitable whitefly host stage for *E. scutalis*, while the second instar


nymph was the least suitable stage. The predation of *E. scutalis* on the eggs and first instar larvae of *B. tabaci* was not significantly different (Nomikou *et al.*, 2004), which contrasts with our results. This discrepancy can be explained by the fact that Nomikou *et al.* (2004) reported the predation rate over only one day, while we examined the predation rate throughout the lifespan. Additionally, variations in predator and prey strains, as well as the previous food source of the predatory mite, can also account for this difference. Furthermore, it appears that insect nymphs have a higher nutritional content compared to insect eggs. Shah *et al.* (2022) indicated that insect nymphs can contain a protein content ranging from 40-73% of their body weight, depending on the species. Additionally, higher concentrations of fatty acids are found in insect nymphs compared to their eggs. However, information on the nutritional value of insect eggs is not as extensively covered in the literature. It has been noted that the hardness of the egg's chorion and the length of time required for handling them may pose difficulties for the mites when feeding on them (Meyerdirk and Coudriet, 1986; Carrillo and Pena, 2012; Ganjisaffar and Perring, 2015). Furthermore, it is evident that there is a difference in size and biomass between the eggs and nymphs of *B. tabaci*. The nymphs were found to be larger than the eggs. However, the potential benefits of this size difference are somewhat diminished by the fact that the nymphs are active stages, requiring the predator to exert energy and time in capturing them. Nevertheless, due to the smaller size of the leaf discs compared to normal leaves and the high density of prey, the predator females did not have to exert much energy in searching for active prey (Bruce-Oliver and Hoy, 1990).

Similar to previous studies on *E. scutalis* (Bounfour and McMurtry, 1987;

**Table 3.** Mean ( $\pm$ SE) predation rates of *Euseius scutalis* fed on egg and first-instar nymph of *Bemisia tabaci*.

Parameter	Egg	Nymph (L1)
Protonymph female	0.27 $\pm$ 0.07 a A	0.23 $\pm$ 0.05 a A
Protonymph male	0.36 $\pm$ 0.08 a A	0.26 $\pm$ 0.08 a A
Deutonymph female	1.33 $\pm$ 0.10 a A	1.47 $\pm$ 0.07 a A
Deutonymph male	1.18 $\pm$ 0.12 a A	1.11 $\pm$ 0.15 a A
Pre-adult female	1.60 $\pm$ 0.19 a A	1.70 $\pm$ 0.18 a A
Pre-adult male	1.55 $\pm$ 0.25 a A	1.37 $\pm$ 0.22 a B
Adult female	16.30 $\pm$ 5.48 b A	29.40 $\pm$ 5.48 a A
Adult male	11.33 $\pm$ 3.32 b B	17.67 $\pm$ 4.36 a B
Total longevity female	17.90 $\pm$ 1.55 b A	32.10 $\pm$ 1.65 a A
Total longevity male	12.87 $\pm$ 1.55 b B	19.04 $\pm$ 1.29 a B
$C_0$	15.26 $\pm$ 1.30 b	23.99 $\pm$ 1.62 a
$Q_p$	3.52 $\pm$ 0.54 a	2.76 $\pm$ 0.34 a
$\psi$	0.58 $\pm$ 0.03 a	0.63 $\pm$ 0.03 a
$\omega$	0.63 $\pm$ 0.03 b	0.73 $\pm$ 0.04 a

<sup>a</sup> Mean values followed by different lowercase letters within the same row are significantly different ( $P < 0.05$ ), based on paired bootstrap test with 100,000 samples. Mean values followed by different capital letters shows the significant difference between female and male for each parameter ( $P < 0.05$ ), based on paired bootstrap test with 100,000 samples.  $C_0$ : Net predation rate;  $\omega$ : Finite predation rate;  $Q_p$ : Transformation rate;  $\psi$ : Stable predation rate.

**Figure 4.** Age-specific predation rate ( $k_x$ ), age-specific net predation rate ( $q_x$ ) and cumulative predation rate ( $C_x$ ) of predator mite *Euseius scutalis* fed on egg and first-instar nymph of *Bemisia tabaci*.



Bazazzadeh *et al.*, 2025), the pre-adult development for male was incomparable to that of females in our study. This developmental similarity can be attributed to some biological and genetic factors. Research indicates that the sex-lethal gene (*Ppsxl*) plays a role in sex determination and reproductive processes, but its influence on sexual differentiation in *Phytoseiulus persimilis* Athias-Henriot is not as pronounced as in insect (Li *et al.*, 2023). This suggests that both sexes may share similar developmental pathways during their immature stages. While the similarities in immature development are evident, it is important to consider that environmental factors and specific ecological roles may also influence the developmental trajectories of these mites, potentially leading to variations not solely explained by genetic mechanisms.

Based on the results of the current study and regardless of the tested immature stages, most of the eggs give rise to male offspring at the beginning of the oviposition period and to female offspring throughout the rest of the oviposition period. A similar trend has also been reported for a Morocco (Marrakech) strain of *E. scutalis* (Bounfour and McMurtry, 1987) and for other phytoseiid such as *Typhlodromus caudiglans* Schuster (Putman, 1962), *Phytoseiulus persimilis*, *Amblyseius bibens* Blommers (Schulten *et al.*, 1978), *Galendromus helveolus* (Chant) (Caceres and Childers, 1991), and *Kampimodromus aberrans* (Boroufas *et al.*, 2007). Sabelis (1985) stated that increased male offspring production at the beginning of the oviposition period could result in the early insemination of females that would afterwards start to search for suitable food.

The predatory mite *E. scutalis* showed a higher *r*-value when preying on the first instar nymphs of *B. tabaci* compared to its predation on *B. tabaci* eggs. The *r*-value estimated in this study was lower than the *r*-value of *E. scutalis* on *B. tabaci* (0.191 d<sup>-1</sup>) (Fouly *et al.*, 2013). By contrast to our study, when *Amblyseius orientalis* Ehara

was fed on *B. tabaci*, the *r* was negative, indicating that this prey is not a sufficient food source for sustaining *A. orientalis* populations (Zhang *et al.*, 2015). In contrast, when *Typhlodromus negevi* Swirski and Amitai was fed on *B. tabaci* eggs, the intrinsic rate of increase (0.271 d<sup>-1</sup>) was higher than our study (Momen *et al.*, 2009), indicating that this prey is more suitable for *T. negevi* than *E. scutalis*.

## CONCLUSIONS

The pre-immature stage of the prey appears to be a crucial factor in developing an integrated pest management program for *B. tabaci* using *E. scutalis*. Our results demonstrate that *E. scutalis* has the potential to control *B. tabaci* due to its ability to feed on both eggs and the first instar nymphs of the prey. Predators that can consume multiple life stages of their prey are more effective at controlling pest populations. Furthermore, *E. scutalis* showed a preference for feeding on nymphs rather than eggs. Field evaluations should be conducted to further assess its biological control potential, particularly with strategies aimed at enhancing its population establishment and long-term control efficiency, such as providing supplementary food.

## ACKNOWLEDGEMENTS

Financial support (Grant no. SCU.AP1403.400) provided by the research deputy of Shahid Chamran University of Ahvaz, Ahvaz, Iran is gratefully acknowledged.

## REFERENCES

1. Abou-Elella, G. M., Saber, S. A. and El-Sawi, S. A. 2013. Biological Aspects and Life Tables of the Predacious Mites, *Typhlodromips swirskii* (Athias-Henriot) and *Euseius scutalis* (Athias-Henriot)

- Feeding on Two Scale Insect Species and Plant Pollen. *Arch. Phytopathol. Plant Protec.*, **46(14)**: 1717–1725.
2. Bazazzadeh, F., Shishehbor, P., Hemmati, S. A. and Riahi, E. 2025. Artificial Diets for Rearing *Euseius scutalis* (Acari: Phytoseiidae). *Exp. Appl. Acarol.*, **95**: 7.
  3. Blackwood, J. S., Schausberger, P. and Croft, B. A. 2001. Prey-Stage Preference in Generalist and Specialist Phytoseiid Mites (Acari: Phytoseiidae) When Offered *Tetranychus urticae* (Acari: Tetranychidae) Eggs and Larvae. *Environ. Entomol.*, **30(6)**: 1103–1111.
  4. Bonato, O., Couton, L. and Fargues, J. 2006. Feeding Preference of *Macrolophus caliginosus* (Heteroptera: Miridae) on *Bemisia tabaci* and *Trialeurodes vaporariorum* (Homoptera: Aleyrodidae). *J. Econ. Entomol.*, **99(4)**: 1143-1151.
  5. Bounfour, M. and McMurtry, J. A. 1987. Biology and Ecology of *Euseius scutalis* (Athias-Henriot) (Acarina: Phytoseiidae). *Hilgardia*, **55(5)**: 1- 23.
  6. Boroufas, G. D., Pappas, M. L. and Koveos, D. S. 2007. Development, Survival and Reproduction of *Kampimodromus aberrans* (Acari: Phytoseiidae) at Different Constant Temperatures. *Environ. Entomol.*, **36(4)**: 657- 665.
  7. Bruce-Oliver, S. J. and Hoy, M. A. 1990. Effect of Prey Stage on Life Table Attributes of a Genetically Manipulated Strain of *Metaseiulus occidentalis* (Acari: Phytoseiidae). *Exp. Appl. Acarol.*, **9**: 201–217.
  8. Caceres, S. and Childers, C. C. 1991. Biology and Life Tables of *Galendromus helveolus* (Acari: Phytoseiidae) on Florida Citrus. *Environ. Entomol.*, **20**: 224- 229.
  9. Carrillo D. and Pena J. E. 2012. Prey-Stage Preferences and Functional and Numerical Responses of *Amblyseius largoensis* (Acari: Phytoseiidae) to *Raoiella indica* (Acari: Tenuipalpidae). *Exp. Appl. Acarol.*, **57**: 361-372.
  10. Chi, H. 1988. Life-Table Analysis Incorporating Both Sexes and Variable Development Rates Among Individuals. *Environ. Entomol.*, **17**: 26-34.
  11. Chi, H. 2023a. *TWOSEX-MSChart: A Computer Program for the Age Stage, Two-Sex Life Table Analysis*. 2019.01.09 Edition. <http://140.120.197.173/Ecology/prod02.htm> . Accessed June 2024
  12. Chi, H. 2023b. *CONSUME-MSChart: Computer Program for Consumption Rate Analysis Based on the Age Stage, Two-Sex Life Table*. 2019.02.01 ed. <http://140.120.197.173/Ecology/prod02.htm> . Accessed June 2024
  13. Chi, H. and Liu, H. 1985. Two New Methods for the Study of Insect Population Ecology. *Bull. Inst. Zool. Acad. Sin.*, **24**: 225-240.
  14. Chi, H. and Yang, T. 2003. Two-Sex Life Table and Predation Rate of *Propylaea japonica* Thunberg (Coleoptera: Coccinellidae) Fed on *Myzus persicae* (Sulzer). *Environ. Entomol.*, **32**: 327-333.
  15. De Barro, P. J., Liu, S. S., Boykin, L. M. and Dinsdale, A. B. 2011. *Bemisia tabaci*: a Statement of Species Status. *Annu. Rev. Entomol.*, **56**: 1–19.
  16. Firas, A. and Sengonca, C. 2007. Prey Preference and Evaluation of the Predator *Serangium parcesetosum* Sicard (Col., Coccinellidae) as a Bio-Agent of *Bemisia tabaci* (Genn.) (Hom., Aleyrodidae) under Greenhouse Conditions. Sixth Agricultural Scientific Jordanian Conference. p. 18.
  17. Fouly, A. H., Nassar, O. A. and Osman, M. A. 2013. Biology and Life Table of *Euseius scutalis* (A.-H.) Reared on Different Kinds of Food. *J. Entomol.*, **10**: 199-206.
  18. Ganjisaffar F. and Perring T.M. 2015. Prey Stage Preference and Functional Response of the Predatory Mite *Galendromus flumenis* to *Oligonychus pratensis*. *Biol. Control.*, **82**: 40-45. doi:10.1016/j.biocontrol.2014.12.004
  19. Gangwar, R. K. and Gangwar, Ch. 2018. Lifecycle, Distribution, Nature of Damage and Economic Importance of Whitefly, *Bemisia tabaci* (Gennadius). *Acta Sci. Agric.*, **2**: 36–39.
  20. Jones, D. R. 2003. Plant Viruses Transmitted by Whiteflies. *Eur. J. Plant Pathol.*, **109**: 195–219.
  21. Kamali, K., Ostovan, H. and Atamehr, A. 2001. A Catalogue of Mites and Ticks (Acari) of Iran. Islamic Azad University Scientific Publication Center, 192 pp. <https://doi.org/10.13140/2.1.4825.8244>
  22. Martin, J. H. 1987. An Identification Guide to Common Whitefly Pest Species of the World (Homoptera,



- Aleyrodidae). *Trop. Pest Manag.*, **33**: 298–322.
23. Li, M., Wang, Z., Wang, E., Zhang, B., Lv, J. and Xu, X. 2023. Expression and Function Analyses of Sexlethal Gene in *Phytoseiulus persimilis*. *Syst. Appl. Acarol.*, **28(8)**: 1356–1367.
  24. McMurtry, J. A., De Moraes, G. J. and Sourassou, N. F. 2013. Revision of the Lifestyles of Phytoseiid Mites (Acari: Phytoseiidae) and Implications for Biological Control strategies. *Syst. Appl. Acarol.*, **18 (4)**: 297- 320.
  25. Meyerdirk, D. E. and Coudriet, D. L. 1986. Evaluation of Two Biotypes of *Euseius scutalis* (Acarina: Phytoseiidae) as Predators of *Bemisia tabaci* (Homoptera: Aleyrodidae). *J. Econ. Entomol.*, **79(3)**: 659-663.
  26. Momen, F. M. and Hussein, H. 2011. Influence of Prey Stage on Survival, Development and Life Table of the Predacious Mite, *Neoseiulus barkeri* (Hughes) (Acari: Phytoseiidae). *Acta Phytopathol. Entomol. Hung.*, **46 (2)**: 319–328.
  27. Momen, F., Fahim, S. and Barghout, M. 2020. Mass Production of Predatory Mites and Their Efficacy for Controlling Pests. In: El-Wakeil, N., Saleh, M. and Abu-Hashim, M. (eds). *Cottage Industry of Biocontrol Agents and Their Applications*. Springer Nature, Cham, pp 157–202.
  28. Momen, F., Abdel-Khalek, A. and El-Sawi, S. 2009. Life Tables of the Predatory Mite *Typhlodromus negevi* Feeding on Prey Insect Species and Pollen Diet (Acari: Phytoseiidae). *Acta Phytopathol. Entomol. Hung.*, **44 (2)**: 353-361.
  29. Nawar, M.A. 2017. Biology and Thermal Requirements of *Euseius scutalis* (Athias-Henriot) Fed on Three Pest Prey Types and Pollen. *Acarines*, **11**: 21-27.
  30. Nomikou, M., Janssen, A., Schraag, R. and Sabelis, M. W. 2001. Phytoseiid Predators as Potential Biological Control Agents for *Bemisia tabaci*. *Exp. Appl. Acarol.*, **25**: 271 - 291.
  31. Nomikou, M., Janssen, A., Schraag, R. and Sabelis, M. W. 2004. Vulnerability of *Bemisia tabaci* Immatures to Phytoseiid Predators: Consequences for Oviposition and Influence of Alternative Food. *Entomol Exp Appl.*, **110**: 95–102.
  32. Putman, W. L. 1962. Life-History and Behavior of the Predacious Mite *Typhlodromus (T.) caudiglans* (Acarina: Phytoseiidae) in Ontario, with Notes on the Prey of Related Species. *Can. Entomol.*, **94**: 163- 177.
  33. Sabelis, M.W. 1985. *Sex Allocation*. In: “*World Crop Pests, Spider Mites, Their Biology, Natural Enemies and Control*”, (Eds.): Helle W. and Sabelis M. W. Volume 1B, Elsevier, Amsterdam, The Netherlands, PP. 83- 94.
  34. Schulten, G. G. M., Vanarendonk, R. C. M., Russell, V. M. and Roorda, F. A. 1978. Copulation, Egg Production and Sex Ratio in *Phytoseiulus persimilis* and *Amblyseius bibens* (Acari: Phytoseiidae). *Entomol. Exp. Appl. Acarol.*, **24**: 145- 153.
  35. Shah, A. A., Totakul, P., Matra, M., Cherdthong, A., Hanboonsong, Y. and Wanapat, M. 2022. Nutritional Composition of Various Insects and Potential Uses as Alternative Protein Sources in Animal Diets. *Anim Biosci.*, **35(2)**: 317-331.
  36. Shishehbor, P., Rahmani Piyani, A. and Riahi, E. 2022. Effect of Different Pollen Diets in Comparison to a Natural Prey, *Tetranychus turkestanii* (Acari: Tetranychidae), on Development, Survival, and Reproduction of *Euseius scutalis* (Acari: Phytoseiidae). *Syst. Appl. Acarol.*, **23(10)**: 2111- 2122.
  37. Stathakis, T. I., Kapaxidi, E. V., Papadoulis, G. Th. and Papanikolaou, N. E. 2021. Predation by *Euseius scutalis* (Acari: Phytoseiidae) on *Tetranychus urticae* and *Eutetranychus orientalis* (Acari: Tetranychidae): Effect of Prey Density and Developmental Stage. *Syst. Appl. Acarol.*, **26(10)**: 1940-1951.
  38. Walzer, A. and Schausberger, P. 1999. Cannibalism and Interspecific Predation in the Phytoseiid Mites *Phytoseiulus persimilis* and *Neoseiulus californicus*: Predation Rates and Effects on Reproduction and Juvenile Development. *BioControl*, **43**: 457–468.
  39. Yazdanpanah, S. and Fathipour, Y. 2022. Predators of Mite Pests. In: “*Insect Predators in Pest Management*”, (Ed.): Omkar. CRC Press, Taylor and Francis, USA, PP. 245-283.
  40. Zhang, X., LV., J., Hu, Y., Wang, B., Chen, X., Xu, X. and Wang, E. 2015. Prey

Preference and Life Table of *Amblyseius orientalis* on *Bemisia tabaci* and *Tetranychus cinabarinus*. *PLoS One*, **10(10)**: e0138820.

## تأثیر دو مرحله رشدی *Bemisia tabaci* بر رشد، پارامترهای جدول زندگی، و نرخ شکارگری کنه شکارگر، (*Euseius scutalis* (Acari: Phytoseiidae)

محمدگراوندیان، پرویز شیشه‌بر، مهدی اسنفدیاری، و الهام ریاحی

### چکیده

رشد، بقا و پارامترهای جدول زندگی کنه شکارگر، *Euseius scutalis* - (Athias-Henriot)، با تغذیه از دو مرحله پیش از بلوغ (تخم و پوره سن اول) (*Bemisia tabaci* (Gennadius) مورد مطالعه قرار گرفت. پوره‌های سن اول *B. tabaci* به عنوان غذای ارجح برای *E. scutalis* شناخته شد زیرا منجر به کاهش دوره رشد از تخم تا بلوغ و همچنین دوره پیش از تخم‌گذاری و افزایش نرخ تخم‌گذاری شد. نرخ ذاتی افزایش جمعیت ( $r$ ) با تغذیه از پوره‌های سن اول 0/1503 بر روز و با تغذیه از تخم 0/0843 بر روز بود. به طور متوسط ماده‌های *E. scutalis، 16/30 تخم و 29/40 پوره را در کل دوره زندگی (از زمان ظهور تا مرگ) تغذیه کردند. هنگامی که پوره‌های سن اول *B. tabaci* ارائه شد، *E. scutalis* نرخ خالص شکارگری ( $C_0$ ) و نرخ متناهی شکارگری ( $\omega$ ) بالاتری را در مقایسه با تغذیه از تخم‌ها نشان داد. به طور متوسط کنه شکارگر 3/52 عدد تخم یا 2/76 عدد پوره اول *B. tabaci* را برای تولید یک تخم *E. scutalis* مصرف کرد. از نظر نسبت جنسی نتاج، ماده‌هایی که از پوره‌های سن اول تغذیه کردند، ماده‌های بیشتری تولید کردند.*