

## Herbicidal and Insecticidal Activity of Secondary Metabolites from Endophytic and Soil Fungi

S. A. M. Abdelgaleil<sup>1\*</sup>, Y. Shiono<sup>2</sup>, N. E. M. Taktak<sup>3</sup>, and M. M. G. Saad<sup>1</sup>

### ABSTRACT

Fungi are a good source for lead molecules in drug discovery and development. However, many compounds derived from fungi were not evaluated for their bioactivity against economic, agricultural, and public health pests. Twelve fungal secondary metabolites (1-12) were evaluated for herbicidal activity against *Silybum marianum* and insecticidal activity against *Culex pipiens* larvae. Among the tested metabolites, brefeldin A (6) and 6-epoxy-4-hydroxy-3-methoxy-5-methyl-cyclohex-2-en-1-one (11) showed potent herbicidal activity against *S. marianum* with complete inhibition of seed germination at 500 mg L<sup>-1</sup>. Compound 6 revealed an exceptional herbicidal activity as it caused complete inhibition of root growth and strong reduction in shoot growth (I = 74.5%) and germination (10.0%) at 25 mg L<sup>-1</sup>. In addition, dehydroaustin (9), phomoxanthone A (4) and deacetylphomoxanthone A (5) displayed a potent toxicity against fourth larval instar of *C. pipiens* with LC<sub>50</sub> values of 3.27, 57.03 and 63.50 mg L<sup>-1</sup>, respectively. Based on the results of this study, compounds 4-6, 9 and 11 should be developed as natural pesticides.

**Keywords:** Bioactivity, *Culex pipiens*, Fungal metabolites, Natural products, *Silybum marianum*,

### INTRODUCTION

Fungi are a rich source of diverse natural compounds that have the potential for the discovery of new drugs and agrochemicals (Porrás-Alfaro *et al.*, 2011). It has been estimated that only 5% (around 70,000) of fungi total populations (1.5 million) have been explored (Sharma *et al.*, 2016). This confirms fungi as possible alternatives to plants for producing valuable natural products. Although the soil fungi are well-known as a source for important antibiotics and drugs (Wasser, 2002; Sethi *et al.*, 2013), much effort has been devoted in the last two

decades to study the secondary metabolites from marine and endophytic fungi as new sources for drugs and agrochemicals (Butler *et al.*, 2014; Segaran *et al.*, 2019).

Interest in microbial metabolites as pesticides arises to overcome the problems associated with the continuous use of synthetic pesticides, such as adverse effects of pesticide residues on human, non-target organisms and environment, and emergence of pest resistance (Aktar *et al.*, 2009). In fact, the microbial metabolites have several advantages to be good candidates in pest control, such as structure diversity, wide-spectrum of bioactivity, new modes of

<sup>1</sup> Department of Pesticide Chemistry and Technology, Faculty of Agriculture, El-Shatby, Alexandria University, Alexandria 21545, Egypt.

<sup>2</sup> Department of Food, Life, and Environmental Science, Faculty of Agriculture, Yamagata University, Tsuruoka, Yamagata, 997-8555, Japan.

<sup>3</sup> Department of Tropical Health, High Institute of Public Health, Alexandria University, Alexandria, Egypt.

\* Corresponding author; e-mail: samirabdelgaleil@gmail.com



action, and biodegradability (Gerwick and Sparks, 2014). Several pesticides of microbial origin that have been introduced into the field of pest management include avermectin, bialaphos, milbemycin, kasugamycin, blastidicin S, polyoxin, mildiomyacin, validamycin, and tetranactin (Tanaka and Omura, 1993). Therefore, it is highly important to investigate new fungal metabolites and examine their bioactivities against economic pests.

*Silybum marianum* (L.) Gaertn. (Asteraceae) is a dangerous weed in many parts of Middle East, Africa, Australia, and North and South America (Holm et al., 1997). In Egypt, it is considered as an invasive weed that spread through wheat fields in both new reclaimed lands and desert areas. *Culex pipiens* L. (Diptera: Culicidae) is the most widespread mosquito species in Egypt (Zahran et al., 2017; Abdelgaleil et al., 2017). This insect is able to transmit several pathogens, such as West Nile virus, *Bancroftian filariasis* and Rift Valley Fever virus (Meegan et al., 1980).

In our continuous efforts to search for new natural products with possible application in pest management, in this research, twelve fungal secondary metabolites (1-12) were examined for their effect on the germination and growth of roots and shoots of *S. marianum*. In addition, we aimed to test toxicity of these compounds against the fourth larval stage of *C. pipiens* to explore their potential for the control of these economic pests.

## MATERIALS AND METHODS

### Test Organisms

Field biotype seeds of Holy thistle, *Silybum marianum* (L.) Gaertn. (Asteraceae), were collected from Faculty of Agriculture Farm, Alexandria (31° 12' 56.30" N, 29° 57' 18.97" E). The plant seeds were identified by Prof. Fath Allah Zaitoon of the Plant Pathology Department, Faculty of Agriculture, Alexandria University.

Voucher specimen (SM-1909) was deposited in the Department of Pesticide Chemistry and Technology. Uniform and undamaged seeds were used for the germination and seedling growth tests. Seeds were examined for their germination before experiments. The seed germination was 80% after 12 days of sowing. *Culex pipiens* L. (Diptera: Culicidae) was reared in an insectary at 27±2°C and 75±5% Relative Humidity (RH) at the Department of Applied Entomology and Zoology, Alexandria University, Egypt as described by Zahran et al. (2017).

### Fungal Strains and Isolation of Secondary Metabolites

Strains of different endophytic and soil fungi were isolated from different sources (Table 1), identified by BEX Co. Ltd., Japan, using a DNA analysis of the 18S rDNA regions, and were deposited in the laboratory of natural products at the Faculty of Agriculture of Yamagata University. After fermentation on unpolished rice media (unpolished rice 1000 g+ 35 ml of 3.5% NaCl at 25°C for four weeks), the fungal growth media were extracted with MeOH. These extracts were concentrated and partitioned into *n*-hexane and EtOAc. The EtOAc extracts were subjected to silica gel column chromatography with mixtures of *n*-hexane-EtOAc (100:0-0:100, v/v), and mixtures of EtOAc-MeOH (50:50-0:100, v/v). The rustling fractions were further chromatographed on ODS by eluting with H<sub>2</sub>O and an increasing ratio of MeOH (100:0 to 0:100) or silica gel by eluting with a mixture of CHCl<sub>3</sub>-MeOH (90:10, v/v). For HPLC analysis, a reversed phase HPLC (Semipreparative HPLC with Shimadzu pump and UV LC-10A detector (set at 210 nm) on a Mitysil ODS column (150×6.0 mm id) at a flow rate of 2.0 mL min<sup>-1</sup>, solvent systems, MeOH:H<sub>2</sub>O, 60:40-80:20, v/v ) and PTLC (EtOAc – *n*-hexane, 70:30-30:70, v/v) were used. Purity of the isolated compounds was higher than 95%.

**Table 1.** Names, chemical class and source of fungal secondary metabolites.

Name	Chemical group	Fungus (Fungal origin)	Reference
Nodulisporone B (1)	Phenylisobenzofuranones	<i>Nodulisporium</i> sp. SH-1 ( <i>Xylaria polymorpha</i> )	Hayasaka <i>et al.</i> , 2011
19-( $\alpha$ -D-glucopyranosyloxy)isopimar-7,15-dien-3 $\beta$ -ol (2)	Isopimarane diterpenoids	<i>Paraconiothyrium</i> sp. MY-42 (unidentified plant)	Shiono <i>et al.</i> , 2011
Pencolide (3)	Maleimides	<i>Penicillium sclerotiorum</i> (Soil)	Lucas <i>et al.</i> , 2007
Phomaxanthone A (4)	Dimeric xanthenes	<i>Phomopsis</i> sp. ( <i>Rhizophora mucronata</i> )	Elsaesser <i>et al.</i> , 2005
Deacetylphomaxanthone A (5)	Dimeric xanthenes	<i>Phomopsis longicolla</i> ( <i>Sonneratia caseolaris</i> )	Ronsberg <i>et al.</i> , 2013
Brefeldin A (6)	Macrolides	<i>Penicillium brefeldianum</i> ( <i>Pinellia ternata</i> )	Hutchinson <i>et al.</i> , 1983
Anthracobic acid A (7)	Polyketides	<i>Anthracobia</i> sp. (unidentified plant)	Shiono. 2006
Fasciculol C (8)	Lanostane triterpenoids	<i>Neamatoloma fasciculare</i> (Mushroom)	Ikeda <i>et al.</i> , 1977; Kim <i>et al.</i> , 2013
Dehydroaustin (9)	Meroterpenoids	<i>Penicillium brasilianum</i> (Soil)	Schürmann <i>et al.</i> , 2010
Pyrrocidine A (10)	Alkaloids	<i>Acremonium zeae</i> ( <i>Zea mays</i> )	Wicklow and Poling, 2009
6-Eopxy-4-hydroxy-3-methoxy-5-methyl-cyclohex-2-en-1-one (11)	Cyclohexenones	Xylariaceous endophytic fungus (YUA 026) (unidentified plant)	Shiono <i>et al.</i> , 2005
Secalonic acid A (12)	Dimeric xanthenes	<i>Claviceps purpurea</i> ( <i>Secale cereale</i> )	Masters and Bräse, 2002

### Germination and Seedling Growth Inhibition Bioassay

The effect of fungal secondary metabolites (1-12) on seed germination, and root and shoot growth of *S. marianum* was evaluated using a bioassay method explained by Abdelgaleil *et al.* (2009). The tested compounds were dissolved in DMSO and diluted with distilled water containing 0.02% of Triton-X 100 to give final concentration of 500 mg L<sup>-1</sup>. Three replicates with 10 seeds in each one were prepared for each compound. The compound solutions (6 mL) were transferred to each Petri dish (9 cm) lined with filter paper (Whatman No. 2). Petri dishes were then placed in polyethylene bags, which were expanded and closed to prevent moisture loss. In the control treatment, a solution of 0.5% of DMSO and 0.02% of Triton-X 100

in distilled water was added. All treatments were kept at 20±2°C and a 12-hour photoperiod for 12 days. Then, the number of germinated seeds and length of root and shoot were taken. Growth Inhibition (I %) of root and shoot lengths was calculated from this equation:

$$I(\%) = [1 - T/C] \times 100$$

Where, T and C are the root or shoot lengths (cm) in treatment and control. Moreover, two compounds (6 and 11) were further evaluated on germination and seedling growth at a series of concentrations of 25, 50, 100, 250 and 500 mg L<sup>-1</sup>.

### Larvicidal Bioassay

The toxicity of fungal secondary metabolites (1-12) was carried out against *C. pipiens* larvae following a recommended method of the World Health Organization (WHO, 1996) with slight modifications. The



tested compounds were dissolved in dimethyl sulfoxide (DMSO) and diluted with distilled water containing 0.1% of Tween-20 to give three final concentrations (100, 250 and 500 mg L<sup>-1</sup>). Water in the control treatments was mixed with DMSO (0.5%) and Tween-20 (0.1%). Twenty *C. pipiens* larvae were separately put into 200-mL plastic cups containing 100 mL of the compound solutions. Each concentration was replicated three times. All treatments were kept under the same insect rearing conditions for 24 h. Then, the number of dead larvae was counted and mortality (%) was calculated. Three compounds (4, 5 and 9) were further tested at concentrations of 10, 25, 50, 70 and 100 mg L<sup>-1</sup>. Mortality data of these compounds were subjected to probit analysis to calculate LC<sub>50</sub> values for these compounds (Finney, 1971).


## Statistical Analysis

Significant differences among mean values of germination percentages, root and shoot lengths were determined ( $P=0.05$ ) by using a one-way Analysis Of Variance (ANOVA) followed by Tukey's HSD test. The LC<sub>50</sub> values of the compounds were estimated by probit analysis using the SPSS 21.0 software (Statistical Package of Social Sciences Inc., USA).

## RESULTS

### Isolation, Structure Elucidation of Isolated Secondary Metabolites

Twelve secondary metabolites (1-12) were isolated from different fungal strains (Table 1) using different chromatographic



**Figure 1.** Chemical structure of fungal metabolites (1-12).

techniques, including silica gel and ODS columns, PTLC and HPLC. Among the isolated metabolites, nine compounds (1, 2, 4-7 and 10-12) were isolated from endophytic fungi and three compounds (3, 8, and 9) were isolated from soil fungi. The chemical structure (Figure 1) of the isolated compounds was elucidated based on their spectroscopic data of Ultraviolet (UV), Infrared (IR), High Resolution Mass Spectroscopy (HRMS) and Nuclear Magnetic Resonance (NMR). The isolated compounds belonged to ten different chemical groups of natural products (Table 1).

#### Effect of Fungal Metabolites on Germination and Seedling Growth of *S. marianum*

Effect of fungal secondary metabolites (1-4 and 6-12) on germination, root and shoot growth of *S. marianum* at concentration of 500 mg L<sup>-1</sup> is shown in Table 2. Among the 11 tested metabolites, compounds 3, 4, 6, 8, 9, 10 and 11 caused reduction in seed

germination after 12 days of treatment. Compounds 6 and 11 induced complete inhibition of seed germination at this concentration. In addition, seven compounds (2, 3, 4, 6, 8, 11 and 12) revealed significant inhibition in root growth compared with the control treatment. Similarly, compounds 6 and 11 showed complete inhibition of germination and compounds 3, 4, and 12 strongly inhibited root growth with 87.5, 80.4, and 74.8% inhibition, respectively. Conversely, compounds 1 and 10 significantly increased the root growth. Similarly, compounds 3, 4, 8 and 12, significantly reduced shoot growth. The results revealed that the tested compounds caused higher inhibitory effects on plant establishment by reducing root and shoot growth than on seed germination. Also, some of the tested compounds had greater inhibitory effects on root growth than on shoot growth.

The previously presented results showed that compounds 6 and 11 caused complete inhibition of seed germination at 500 mg L<sup>-1</sup>. Therefore, these two compounds were further evaluated at lower concentrations

**Table 2.** Effect secondary metabolites isolated from endophytic fungi on germination and seedling growth of *Silybum marianum* after 12 days of sowing at 500 mg L<sup>-1</sup>.<sup>a</sup>

Compound	Germination (%±SE)	Root		Shoot	
		Length (cm) (Mean±SE)	I <sup>b</sup> (%)	Length (cm) (Mean±SE)	I (%)
Control	86.7±3.34ab	11.39±0.22bc	0.0	2.16±0.06abc	0.0
1	86.7±3.34ab	14.32±0.14a	-28.2	2.33±0.13ab	-7.87
2	86.7±3.34ab	5.47±0.23d	52.0	1.86±0.24bc	13.9
3	76.7±3.34b	1.42±0.06g	87.5	1.33±0.17d	38.4
4	80.0±5.78b	2.23±0.25f	80.4	1.30±0.10d	39.8
6	0.0±0.0c	0.0±0.0h	100.0	0.0±0.0e	100.0
7	96.71±3.34a	11.93±0.11b	-4.74	2.33±0.25ab	-7.87
8	83.3±3.34ab	5.05±0.42d	55.7	1.35±0.09d	37.5
9	80.0±5.78b	11.12±0.17c	2.37	2.53±0.06a	-17.1
10	83.3±6.67ab	13.82±0.17a	-21.3	1.80±0.26c	16.7
11	0.0±0.0c	0.0±0.0h	100.0	0.0±0.0e	100.0
12	90.0±5.78ab	2.87±0.11e	74.8	1.31±0.16d	39.4
df <sup>c</sup>	11	11		11	
F <sup>d</sup>	63.74	801.70		29.30	
P <sup>e</sup>	< 0.00	< 0.00		< 0.00	

<sup>a</sup> Values are means±SE of three replicates with 10 seeds in each one. <sup>(a-g)</sup> Values followed by different letters are significantly different at 0.05 probability level. <sup>b</sup> I= Inhibition; <sup>c</sup> df= Degrees of freedom; <sup>d</sup> F= F-statistic, <sup>e</sup> P= P-value.

**Table 3.** Effect of Brefeldin A and 6-eopxy-4-hydroxy-3-methoxy-5-methyl-cyclohex-2-en-1-one on germination and seedling growth of *Silybum marianum* after 12 days of sowing at different concentrations.<sup>a</sup>

Compound	Conc. (mg L <sup>-1</sup> )	Germination (%±SE)	Root		Shoot	
			Length (cm) (Mean±SE)	I <sup>b</sup> (%)	Length (cm) (Mean±SE)	I (%)
Control	0.0	75.0±2.89a	6.33±0.49a	0.0	1.96±0.21a	0.0
Brefeldin A (6)	25	10.0±5.78d	0.0±0.0d	100.0	0.50±0.28c	74.5
	50	0.0±0.0d	0.0±0.0d	100.0	0.0±0.0d	100.0
	100	0.0±0.0d	0.0±0.0d	100.0	0.0±0.0d	100.0
	250	0.0±0.0d	0.0±0.0d	100.0	0.0±0.0d	100.0
	500	0.0±0.0d	0.0±0.0d	100.0	0.0±0.0d	100.0
6-Eopxy-4-hydroxy-3-methoxy-5-methyl-cyclohex-2-en-1-one (11)	25	60.0±5.78b	6.27±0.40a	0.95	1.63±0.03a	16.8
	50	56.7±3.33b	4.50±0.17b	28.9	1.87±0.20a	4.59
	100	43.3±3.33c	0.83±0.27c	86.9	1.20±0.17b	38.8
	250	6.7±3.33d	0.02±0.02d	99.7	0.17±0.17cd	91.3
	500	0.0±0.0d	0.0±0.0d	100.0	0.0±0.0d	100.0
df <sup>c</sup>		10	10		10	
F <sup>d</sup>		88.34	139.35		32.99	
P <sup>e</sup>		< 0.00	< 0.00		< 0.00	

<sup>a</sup> Values are means±SE of three replicates with 10 seeds in each one. (<sup>a-d</sup>) Values followed by different letters are significantly different at 0.05 probability level. <sup>b</sup> I= Inhibition; <sup>c</sup> df= Degrees of freedom; <sup>d</sup> F= F-statistic, <sup>e</sup> P= P-value

(25, 50, 100, 250 and 500 mg L<sup>-1</sup>). Compound 6 caused complete inhibition of root growth at all tested concentrations (Table 3). Likewise, it induced complete inhibition of germination and shoot growth at all tested concentrations, except 25 mg L<sup>-1</sup>. This compound showed strong inhibition of germination (10.0%) and shoot growth (74.5%) at 25 mg L<sup>-1</sup>. In addition, compound 11 induced significant reduction in seed germination at all of the tested concentrations. Also, it caused strong inhibition of root and shoot growth at 100 and 250 mg L<sup>-1</sup>.

#### Larvicidal Activity of Fungal Metabolites against *C. pipiens*

Toxicity of twelve fungal metabolites against the fourth larval instar of *C. pipiens* at concentrations of 100, 250 and 500 mg L<sup>-1</sup> is presented in Table 4. Four compounds (4, 5, 8 and 9) caused complete mortality (100%) of larvae at 500 mg L<sup>-1</sup>, while three

compounds (4, 5 and 9) caused complete mortality of larvae at 250 mg L<sup>-1</sup>. At 100 mg L<sup>-1</sup>, compound 9 was the only compound that caused complete mortality of larvae. Also, compounds 4 and 5 induced strong larval toxicity at this concentration with 95.0 and 90.0 % mortality, respectively. Comparative toxicity results (Table 5) indicated that compound 9 revealed the highest larval toxicity (LC<sub>50</sub>= 3.27 mg L<sup>-1</sup>), followed by compound 4 (LC<sub>50</sub>= 57.03 mg L<sup>-1</sup>) and compound 5 (LC<sub>50</sub>= 63.50 mg L<sup>-1</sup>).

#### DISCUSSION

New pest management strategies are based on decreasing the use of synthetic pesticides and increasing the use of natural alternatives. Therefore, nowadays, introducing new natural products with low risk on human and environment, and new modes of action is highly needed in the field pest management (Schrader *et al.*, 2010). In this regard, microorganisms are a rich source

**Table 4.** Mortality percentages of *Culex pipiens* fourth instar larvae after 24 hours of treatment with secondary metabolites isolated from endophytic fungi at different concentrations.

Compound	Mortality (%) after 24 hours		
	100 (mg L <sup>-1</sup> )	250 (mg L <sup>-1</sup> )	500 (mg L <sup>-1</sup> )
Control	0.0	0.0	0.0
Nodulisporone B (1)	0.0	0.0	0.0
19-( $\alpha$ -D-glucopyranosyloxy) isopimar-7,15-dien-3 $\beta$ -ol (2)	0.0	- <sup>a</sup>	-
Pencolide (3)	0.0	0.0	0.0
Phomaxanthone A (4)	95.0	100.0	100.0
Deacetylphomaxanthone A (5)	90.0	100.0	100.0
Brefeldin A (6)	0.0	0.0	46.7
Anthracobic acid A (7)	0.0	0.0	10.0
Fasciculol C (8)	0.0	6.25	100.0
Dehydroaustin (9)	100	100.0	100.0
Pyrrocidine A (10)	0.0	0.0	15.0
6-Eopxy-4-hydroxy-3-methoxy-5-methyl-cyclohex-2-en-1-one (11)	0.0	0.0	38.9
Secalonic acid A (12)	0.0	0.0	0.0

<sup>a</sup> Not tested.

**Table 5.** Comparative toxicity of fungal metabolites against *Culex pipiens* fourth instar larvae after 24 hours of exposure.

Compound	LC <sub>50</sub> <sup>a</sup> (mg L <sup>-1</sup> )	95% confidence limits (mg L <sup>-1</sup> )		Slope $\pm$ SE	Intercept $\pm$ SE	$(\chi^2)^b$	P <sup>c</sup>
		Lower	Upper				
Phomaxanthone A (4)	57.03	52.62	61.39	6.75 $\pm$ 0.81	-11.85 $\pm$ 1.47	0.001	0.978
Deacetylphomaxanthone A (5)	63.50	58.75	68.41	6.50 $\pm$ 0.72	-11.73 $\pm$ 1.31	0.006	0.941
Dehydroaustin (9)	3.27	0.38	6.53	1.11 $\pm$ 0.30	-0.57 $\pm$ 0.40	0.181	0.671

<sup>a</sup> The concentration causing 50% mortality; <sup>b</sup> Chi square value, <sup>c</sup> Probability value.

for natural compounds with successful use in pest management (Singh and Yadav, 2020; Saad *et al.*, 2021). Several hundred species of fungi are known to infect and kill pests. The most important groups are the Hyphomycetes within genera like *Beauveria*, *Metarhizium*, *Verticillium*, *Paecilomyces* and *Hirsutella* (Keller, 1998). Moreover, fungi are also known for producing toxic compounds against pests, mainly soil fungi of genera *Aspergillus* and *Penicillium*. Secondary metabolites with insecticidal and antifeedant properties were also found in endophytic and phytopathogenic fungi (Berestetskiy and Hu, 2021).

In this study, the herbicidal and insecticidal activities of twelve secondary metabolites isolated from fungi were

investigated to explore their possible use in pest control programs. The results showed that most of the tested compounds reduced seed germination and seedling growth of *S. marianum* at concentration of 500 mg L<sup>-1</sup>. Among the eleven tested metabolites, compounds 6 and 11 revealed excellent herbicidal activity with complete inhibition of germination and seedling growth at this concentration. Moreover, compound 6 showed strong herbicidal activity at 25 mg L<sup>-1</sup>. Based on the obtained results, this compound is among the most effective natural herbicides reported so far. In our previous study, compounds 6 and 11 were also the most potent inhibitors of seed germination, and growth of root and shoot of *Echinochloa crus-galli* at concentration of 2 mM (Saad *et al.*, 2021). Also, compound 6



was found to be more effective than compound 11 against *E. crus-galli*. The herbicidal activity of compound 11 was further supported by the studies of Shiono and Murayama (2005) who stated that this compound inhibited the root growth of lettuce by 46% at concentration of 50 mg L<sup>-1</sup>. In addition, compound 6 has been described to inhibit wheat germination and the root growth of *Allium cepa* (Betina, 1992)

Comparing the inhibitory effects of tested compounds on the seed germination with seedling growth of *S. marianum* indicated that the tested compounds were more effective against seedling growth than germination. These findings are in agreement with previous studies on many natural compounds in which the seedling growth were more sensitive than germination (Leather and Einhellig, 1984; Abdelgaleil and Hashinaga, 2007; Saad et al., 2012). Also, the results of current study revealed that the tested compounds were more potent inhibitors to root than to shoot, with few exceptions. The higher sensitivity of roots could be due to the roots being the first to expose and absorb tested compounds from growing media (Turk et al., 2002). Similar observations were reported on the inhibitory effects of other natural compounds on seedling growth (Wang et al., 2007; Gouda et al., 2016).

The results of larvicidal bioassays against *C. pipiens* demonstrated that among the tested metabolites, compounds 9, 4 and 5 showed a promising larvicidal activity. In particular, compound 9 displayed the highest toxicity with LC<sub>50</sub> = 3.27 mg L<sup>-1</sup>. Based on our results, this compound is among the most toxic natural compounds reported so far against *C. pipiens* or other mosquito species. This finding is supported by previous study of Geris et al. (2008) who found that compound 9 was highly toxic against third instar larvae of *Aedes aegypti* with LC<sub>50</sub> value of 2.9 mg L<sup>-1</sup>. Compound 9 was more toxic to *C. pipiens* larvae than thymol (LC<sub>50</sub> = 37.95 mg L<sup>-1</sup>), carvacrol (LC<sub>50</sub> = 44.38 mg L<sup>-1</sup>), cinnamaldehyde (LC<sub>50</sub> = 58.97 mg L<sup>-1</sup>), eugenol (LC<sub>50</sub> = 86.22 mg L<sup>-1</sup>) and

cuminaldehyde (LC<sub>50</sub> = 38.94 mg L<sup>-1</sup>) (Radwan et al., 2008; Zahran and Abdelgaleil, 2011). Similarly, compound 9 showed higher toxicity than curcumin (LC<sub>50</sub> = 19.07 mg L<sup>-1</sup>) isolated from *Curcuma longa* (Sagnou et al., 2012).<sup>[44]</sup> Moreover, this compound showed similar toxicity to caulerpin (LC<sub>50</sub> = 1.99 mg L<sup>-1</sup>) and caulerpinic acid (LC<sub>50</sub> = 4.89 mg L<sup>-1</sup>) isolated from *Caulerpa racemosa* (Alarif et al., 2010).

The results of the current study indicate that the tested compounds showed high selectivity against the examined pests. For example, compounds 6 and 11 showed promising herbicidal activity against *S. marianum* while had no insecticidal activity. Likewise, compounds 5 and 9 possessed potent insecticidal activity against *C. pipiens*, but were not active against the tested weed. This selectivity is crucial for the development of these compounds as new bio-pesticides.

In conclusion, this study described, for the first time, the herbicidal and insecticidal activities of twelve fungal metabolites against the economic agricultural pests, *S. marianum* and *C. pipiens*. Among the tested metabolites, brefeldin A (6) and 6-epoxy-4-hydroxy-3-methoxy-5-methyl-cyclohex-2-en-1-one (11) revealed promising herbicidal activity against *S. marianum*, in particular, brefeldin A (6) which could be developed as a new natural herbicide. In addition, dehydroaustin (9) exhibited remarkable larvicidal activity against *C. pipiens* indicating its possible use in integrated management programs of mosquitoes.

## ACKNOWLEDGEMENTS

The authors are grateful to Eng. Hagar Youssef for her technical assistance.

## REFERENCES

1. Abdelgaleil, S. A. M., Abdel-Razeek, N. and Soliman, S. A. 2009. Herbicidal Activity of Three Sesquiterpene Lactones on Wild Oat (*Avena fatua*) and Their



- Possible Mode of Action. *Weed Sci.*, **57**: 6-9.
2. Abdelgaleil, S. A. M. and Hashinaga, F. 2007. Allelopathic Potential of Two Sesquiterpene Lactones from *Magnolia grandiflora* L. *Biochem. Syst. Ecol.*, **35**: 737-742.
3. Abdelgaleil, S. A. M., Zoghroban, A. A. M., El-Bakry, A. M. and Kassem, S. M. I. 2019. Insecticidal and Antifungal Activities of Crude Extracts and Isolated Compounds from Rhizomes of *Curcuma longa* (Zingiberaceae). *J. Agric. Sci. Technol.*, **21**: 1049-1061.
4. Aktar, W., Sengupta, D. and Chowdhury, A. 2009. Impact of Pesticides Use in Agriculture: Their Benefits and Hazards. *Interdiscip. Toxicol.*, **2**: 1-12.
5. Alarif, W. M., Abou-Elnaga, Z. S., Ayyad, S. E. N. and Al-lihaibi, S. S. 2010. Insecticidal Metabolites from the Green Alga *Caulerpa racemosa*. *Clean: Soil Air Water*, **38**: 548-557.
6. Betina, V. 1992. Biological Effects of the Antibiotic Brefeldin A (Decumbin, Cyanein, Ascotoxin, Synergisidin): A Retrospective. *Folia Microbiol.*, **37**: 3-11.
7. Berestetskiy, A. and Hu, Q. 2021. The Chemical Ecology Approach to Reveal Fungal Metabolites for Arthropod Pest Management. *Microorganisms*, **9**: 1379.
8. Butler, M. S., Robertson, A. A. and Cooper, M. A. 2014. Natural Product and Natural Product Derived Drugs in Clinical Trials. *Nat. Prod. Rep.*, **31**: 1612-1661.
9. Elsaesser, B., Krohn, K., Floerke, U., Root, N., Aust, H. J., Draeger, S., Schulz, B., Antus, S. and Kurtán, T. 2005. X-ray Structure Determination Absolute Configuration and Biological Activity of Phomoxanthone. *Eur. J. Org. Chem.*, **21**: 4563-4570.
10. Finney, D. J. 1971. Probit Analysis, 3rd Ed. Cambridge University Press, London, p. 318.
11. Geris, R., Rodrigues-Fo, E., da Silva, H. H. G. and da Silva, I. G. I. G. 2008. Larvicidal Effects of Fungal Meroterpenoids in the Control of *Aedes aegypti* L., the Main Vector of Dengue and Yellow Fever. *Chem. Biodivers.*, **5**: 341-345.
12. Gerwick, B. C. and Sparks, T. C. 2014. Natural Products for Pest Control: An Analysis of Their Role, Value and Future. *Pest Manag. Sci.*, **70**: 1169-1185.
13. Gouda, N. A. A., Saad, M. M. G. and Abdelgaleil, S. A. M. 2016. Pre and Post Herbicidal Activity of Monoterpenes against Barnyard Grass (*Echinochloa crus-galli*). *Weed Sci.*, **64**: 191-200.
14. Hayasaka, S., Koseki, T., Murayama, T., Kwon, E. and Shiono, Y. 2011. Phenylisobenzofuranones from *Fungicolous nodulisporium* sp. SH-1. *Z. Naturforsch.*, **66b**: 961-964.
15. Holm, L. G., Doll, J., Holm, E., Pancho, J. and Herberger, J. 1997. *World Weeds*. Natural Histories and Distribution. Wiley, New York.
16. Hutchinson, C. R., Shu-Wee, L., McInnes, A. G. and Walter, J. A. 1983. Comparative Biochemistry of Fatty Acid and Macrolide Antibiotic (Brefeldin A) Formation in *Penicillium brefeldianum*. *Tetrahedron*, **39**: 3507-3513.
17. Ikeda, M., Niwa, G.-I., Tohyama, K., Sassa, T. and Miura, Y. 1977. Structures of Fasciculol C and Its Depsipeptides, New Biologically Active Substances from *Neamatoloma fasciculare*. *Agric. Biol. Chem.*, **41**: 1803-1805.
18. Keller, S. 1998. Use of Fungi for Pest Control in Sustainable Agriculture. *Phytoprotection*, **79**: 56-60.
19. Kim, K. H., Moon, E., Choi, S. U., Kim, S. Y. and Lee, K. R. 2013. Lanostane Triterpenoids from the Mushroom *Neamatoloma fasciculare*. *J. Nat. Prod.*, **76**: 845-851.
20. Leather, G. R. and Einhellig, A. 1985. Mechanisms of Allelopathic Action in Bioassay. In: "The Chemistry of Allelopathy". American Chemical Society, Washington, DC, PP. 197-205.
21. Lucas, E. M. F., de Castro, M. C. M. and Takahashi, J. A. 2007. Antimicrobial



- Properties of Sclerotiorin, Isocromophilone VI and Pencolide, Metabolites from Brazillian Cerrado Isolate of *Penicillium sclerotiorum* van beyma. *Braz. J. Microbiol.*, **38**: 785-789.
22. Masters, K. S. and Bräse, S. 2002. Xanthones from Fungi, Lichens, and Bacteria: The NaturalP and Their Synthesis. *Chem. Rev.*, **112**: 3717–3776.
23. Meegan, J. M., Khalil, G. M., Hoogstraal, H. and Adham, F. K. 1980. Experimental Transmission and Field Isolation Sstudies Implicating *Culex pipiens* as a Vector of Rift Valley Fever Virus in Egypt. *Am. J. Trop. Med. Hyg.*, **29**: 1405–1410.
24. Porras-Alfaro, A. and Bayman, P. 2011. Hidden Fungi, Emergent Properties: Endophytes and Microbiomes. *Annu. Rev. Phytopathol.*, **49**: 291-315.
25. Radwan, M. A., El-Zemity, S. R., Mohamed, S. A. and Sherby, S. M. 2008. Larvicidal Activity of Some Essential Oils, Monoterpenoids and Their Corresponding N-methyl Carbamate Derivatives against *Culex pipiens* (Diptera: Culicidae). *Int. J. Trop. Insect Sci.*, **28**: 61–68.
26. Ronsberg, D., Debbab, A., Mandi, A., Vasylyeva, V., Bohler, P., Stork, B., Engelke, L., Hamacher, A., Sawadogo, R., Diederich, M., Wray, V., Lin, W., Kassack, M. U., Janiak, C., Scheu, S., Wesselborg, S., Kurtan, T., Aly, A. H. and Proksch, P. 2013. Pro-Apoptotic and Immunostimulatory Tetrahydroxanthone Dimers from the Endophytic Fungus *Phomopsis longicolla*. *J. Org. Chem.* **78**: 12409–12425.
27. Saad, M. M. G., Abdelgaleil, S. A. M. and Shiono, Y. 2021. Antibacterial and Herbicidal Properties of Secondary Metabolites from Fungi. *Nat. Prod. Res.* **35**: 5446–5451.
28. Saad, M. M. G., Abdelgaleil, S. A. M. and Suganuma, T. 2012. Herbicidal Potential of Pseudoguaninolid Sesquiterpenes on Wild Oat, *Avena fatua* L. *Biochem. Syst. Ecol.*, **44**: 333-337.
29. Sagnou, M., Mitsopoulou, K. P., Koliopoulos, G., Pelecanou, M., Couladouros, E. A. and Michaelakis, A. 2012. Evaluation of Naturally Occurring Curcuminoids and Related Compounds against Mosquito Larvae. *Acta Trop.*, **123**: 190–195.
30. Schrader, K. K., Andolfi, A., Cantrell, C. L., Cimmino, A., Duke, S. O., Osbrink, W., Wedge, D. E. and Evidente, A. 2010. A Survey of Phytotoxic Microbial and Plant Metabolites as Potential Natural Products for Pest Management. *Chem. Biodivers.*, **7**: 2261-2280.
31. Schürmann, B. T. M., Sallum, W. S. T. and Takahashi, J. A. 2010. Austin, Dehydroaustin and Other Metabolites from *Penicillium brasilianum*. *Quim Nova.*, **33**: 1044–1046.
32. Segaran, G. and Sathivelu, M. 2019. Fungal Endophytes: A Potent Biocontrol Agent and a Bioactive Metabolite Reservoir. *Biocatal. Agric. Biotechnol.*, **21**: 101284.
33. Sethi, S., Kumar, R. and Gupta, S. 2013. Antibiotic Production by Microbes Isolated from Soil. *Int. J. Pharm. Sci. Res.*, **4**: 2967.
34. Sharma, D., Pramanik, A. and Agrawal, P. K. 2016. Evaluation of Bioactive Secondary Metabolites from Endophytic Fungus *Pestalotiopsis neglecta* BAB-5510 Isolated from Leaves of *Cupressus torulosa* D. Don. *3 Biotech*, **6**: 210.
35. Shiono, Y., Kikuchi, M., Koseki, T., Murayama, T., Kwon, E., Aburai, N., Kimura, K.-I. 2011. Isopimarane Diterpene Glycosides, Isolated from Endophytic Fungus *Paraconiothyrium* sp. MY-42. *Phytochemistry*, **72**: 1400-1405.
36. Shiono, Y. and Murayama, T. 2005. New Eremophilane-type Sesquiterpenoids, Eremoxylarins A and B from *Xylariaceae* endophytic Fungus YUA-026. *Z. Naturforsch.*, **60b**: 885–890.
37. Shiono, Y., Murayama, T., Takahashi, K., Okada, K., Katohda, S. and Ikeda, M. 2005. Three Oxygenated Cyclohexanone Derivatives Produced by an Endophytic

- Fungus. *Biosci. Biotechnol. Biochem.*, **69**: 287–292.
38. Shiono, Y. 2006. Anthracobic Acids A and B, Two Polyketides, Produced by an Endophytic Fungus *Anthracobia* sp. *Chem. Biodivers.*, **3**: 217–223.
  39. Silvie, P. J., Martin, P., Huchard, M., Keip, P., Gutierrez, A. and Sarter, S. 2021. Prototyping a Knowledge-Based System to Identify Botanical Extracts for Plant Health in Sub-Saharan Africa. *Plants*, **10**: 896.
  40. Singh, J. and Yadav, A. N. 2020. *Natural Bioactive Products in Sustainable Agriculture*. Springer, Singapore.
  41. Tanaka, Y. and Omura, S. 1993. Agroactive Compounds of Microbial Origin. *Annu. Rev. Microbiol.*, **47**: 57–87.
  42. Turk, M. A., Abdel-Rahman and Tawaha, M. 2002. Inhibitory Effects of Aqueous Extracts of Black Mustard on Germination and Growth of Lentil. *Pak. J. Agron.* **1**: 28–30.
  43. Wang, W., Zhu, T., Tao, H., Lu, Z., Fang, Y., Gu, Q. and Zhu, W. 2007. Two New Cytotoxic Quinone Type Compounds from the Halotolerant Fungus *Aspergillus varicolor*. *J. Antibiot.*, **60**: 603–607.
  44. Wasser, S. P. 2002. Medicinal Mushrooms as a Source of Antitumor and Immunomodulating Polysaccharides. *Appl. Microbiol. Biotechnol.*, **60**: 258–274.
  45. WHO1996. Report of the WHO Informal Consultation on the Evaluation on the Testing of Insecticides CTD/WHO PES/IC/96.1, p 69.
  46. Wicklow, D. T. and Poling, S. M. 2009. Antimicrobial Activity of Pyrrocidines from *Acremonium zeae* against Endophytes and Pathogens of Maize. *Phytopathol.*, **99**: 109–115.
  47. Zahran, H. A. and Abdelgaleil, S. A. M. 2011. Insecticidal and Developmental Inhibitory Properties of Monoterpenes on *Culex pipiens* L. (Diptera: Culicidae) *J. Asia-Pac. Entomol.*, **14**: 46–51.
  48. Zahran, H. E. D., Abou-Taleb, H. K. and Abdelgaleil, S. A. M. 2017. Adulticidal, Larvicidal and Biochemical Properties of Essential Oils against *Culex pipiens* L. *J. Asia-Pac. Entomol.*, **20**: 133–139.

## فعالیت علف کش و حشره کش متابولیت های ثانویه قارچ های اندوفیت و خاک

س.ا.م. عبدالجلیل، ی. شیونو، ن.ا.م طقطک، و م.م. گ. سعد

### چکیده

قارچ ها منبع خوبی برای مولکول های مهم (lead molecules) در کشف و توسعه دارو هستند. با این حال، بسیاری از ترکیبات مشتق شده از قارچ ها از نظر زیست فعالی در برابر آفات اقتصادی، کشاورزی و بهداشت عمومی مورد ارزیابی قرار نگرفته اند. در این بررسی، ۱۲ متابولیت ثانویه قارچی (۱-۱۲) از نظر فعالیت علف کشی در برابر *Silybum marianum* و فعالیت حشره کشی علیه لارو *Culex pipiens* مورد ارزیابی قرار گرفت. در میان متابولیت های آزمایش شده، برفلدین A (۶) و ۶-۵-methoxy-3-hydroxy-4-epoxy-methyl-cyclohex-2-en-1-one (۱۱) با غلظت ۵۰۰ میلی گرم در لیتر فعالیت علف کشی قوی علیه *S. marianum* با مهار کامل (complete inhibition) در جوانه زنی بذر نشان داد. ترکیب ۶ یک فعالیت علف کش استثنایی را نشان داد زیرا در ۲۵ میلی گرم در لیتر باعث مهار کامل رشد ریشه و کاهش شدید رشد ساقه



( $I = 74.5$ ) و جوانه زنی (۱۰۰٪) شد. علاوه بر این، دهیدروآستین (۹)، فوماکساتون A (۴) و داستیلفوماکساتون A (۵) سمیت قوی در برابر سن چهارم لارو *C. pipiens* با مقادیر  $LC_{50}$  برابر ۳/۲۷، ۶۳/۵۰ و ۵۷/۰۳ میلی گرم در لیتر نشان دادند. بر اساس نتایج این پژوهش، ترکیبات ۴-۶، ۹ و ۱۱ باید به عنوان آفت کش طبیعی تولید شوند.