

## Antifungal Potential of *Lactiplantibacillus plantarum* AM2 against Banana Pathogen *Fusarium oxysporum* f. sp. *cubense* Tropical Race 4

Wan Anati Nabilah Wan Tajudin Shah<sup>1</sup>, Nur Baiti Abd Murad<sup>1</sup>, Jia Xin Ong<sup>1</sup>, Shin Huey Ang<sup>1</sup>, Nur Laili<sup>2</sup>, and Noor Baity Saidi<sup>1,3\*</sup>

### ABSTRACT

*Fusarium* wilt of bananas is a serious disease affecting banana plantations worldwide. In an effort to sustainably manage the disease, biological control is considered a promising alternative to agrochemicals that can cause detrimental effects on humans and the ecosystem. In this study, we investigated the biological control potential of the present collection of beneficial bacteria, including *Lactiplantibacillus plantarum* AM2, *Streptomyces morookaensis* NRRL B-12429, *Bacillus velezensis* B4158, *B. atrophaeus* B363B, and *B. amyloliquefaciens* B942, through a dual culture assay and a greenhouse experiment. The studied causal agent of *Fusarium* wilt in banana was *Fusarium oxysporum* f. sp. *cubense* Tropical Race 4 (*Foc* TR4). The inhibition range *in vitro* was between 31.0 to 42.1%, and the highest growth inhibition of *Foc* TR4 was observed for *L. plantarum* AM2. Infected banana plantlets that received the treatment with *L. plantarum* AM2 also showed a significant reduction in disease severity index, as low as 24% compared to the treatment with other beneficial bacteria. This study showed that *L. plantarum* AM2 has a good antagonistic effect on *Foc* TR4 mycelial growth and the most potential to control *Fusarium* wilt disease in bananas.

**Keywords:** Biological control, *Fusarium* wilt, *in Planta*, Lactic acid bacteria, Suppression.

### INTRODUCTION

*Musa* spp. (banana) serves as an important cash crop for income generation and employment creation, especially in banana-producing countries (Caro, 2020). However, the recent discovery of *Fusarium* wilt disease in Latin America in 2021, together with continuing COVID-19 pandemic constraints and rising production costs had deflated the world banana export (FAO, 2022). The *Fusarium* Wilt of Bananas (FWB) is caused by a soil-borne pathogenic fungus *Fusarium*

*oxysporum* f. sp. *cubense* (*Foc*). The fungus invades the vascular system of the host, blocking the transportation of nutrients and water throughout the plant. This leads to wilting symptoms and browning of the xylem (Ordóñez, 2018). *Foc* belongs to a species complex and consists of four races (1-4), each with different pathogenicity towards different banana cultivars. Race 4 mostly infects Cavendish, with Tropical Race 4 (TR4) being the most pathogenic (Pérez-Vicente *et al.*, 2014; Ordóñez, 2018). *Foc* TR4 easily spreads through planting material, soil, and other substrates originating from infected farms.

<sup>1</sup> Department of Cell and Molecular Biology, Faculty of Biotechnology and Biomolecular Sciences, University of Putra Malaysia, Serdang, Selangor, Malaysia.

<sup>1</sup> Research Center for Applied Microbiology, Research Organization for Life Sciences, National Research and Innovation Agency (BRIN), Jl. Raya Jakarta-Bogor Km. 46, Cibinong, Bogor, West Java 16911, Indonesia.

<sup>1</sup> Laboratory of Sustainable Agronomy and Crop Protection, Institute of Plantation Studies, University of Putra Malaysia, Serdang, Selangor, Malaysia.

\*Corresponding author; email: norbaity@upm.edu.my



Eradication of the pathogen is difficult once it is established, due to its ability to survive in several alternative hosts and produces chlamydospores that persist in the soil for a long time (Ploetz, 2015).

Due to the significant impact on the global economy, the management of FWB has been a focus of the scientific community worldwide. The management approaches include manipulation of cultural practices, chemical control, breeding for resistant cultivars, and biological control. The latter has been gaining interest recently due to increasing awareness of sustainable management of plant disease with less impact on the environment (Scortichini, 2022). The growth of the organic market, in conjunction with a reduction of pesticides, further drives the demand for more effective biocontrol agents and promotes the expansion of the biocontrol industry (Lahlali *et al.*, 2022). The Biological Control Agents (BCA) of FWB are dominated by endophytes such as *Trichoderma* spp. and *Bacillus* spp. (Bubici *et al.*, 2019; Sánchez-Espinosa *et al.*, 2020). Regardless, it does not put a hold on the quest to find new BCA candidates with different modes of action or unique secondary metabolites with better biocontrol efficacy as well as plant-growth-promoting effects. However, the majority of studies involving BCA against *Foc* TR4 were only conducted *in vitro*, with only a small number at the greenhouse level and very few that reached the field trial stage. Interestingly, based on data mining from the literature, Bubici *et al.* (2019) reported that biocontrol for FWB under field conditions exhibits similar disease control efficacy as observed in pot experimental conditions.

Lactic Acid Bacteria (LAB) are an intriguing group of microorganisms frequently present in plant-associated microbiomes (Jaffar *et al.*, 2023). Moreover, it has been shown that LAB can generate compounds that are effective against a broad range of phytopathogens, including *F. oxysporum* (Raman *et al.*, 2022). Compared to other common groups of BCAs, LAB possesses the upper hand, since its application in food crop production presents no health risks to humans.

Hence, it was given the Generally Recognized as Safe (GRAS) status by the US Food and Drug Administration (USFDA). *Lactiplantibacillus plantarum* is a type of LAB belonging to the novel *Lactiplantibacillus* genus (Zheng *et al.*, 2020). On top of the GRAS status, *L. plantarum* was also given the Qualified Presumption of Safety (QPS) status from the European Food Safety Authority (EFSA) (EFSA BIOHAZ Panel, 2023). Interestingly, *L. plantarum* carries more genes in its large genome compared to other LAB species, indicating its strong adaptability to different environments and high versatility (Seddik *et al.*, 2017). Despite its huge potential, reports on the involvement of *L. plantarum* as BCA for plant pathogens have been scarce. Riolo *et al.* (2023) were the first to discover the potential of the fermentates of LABs from drupes of olive oil, including *L. plantarum*, as bio-fungicide against several plant pathogenic fungi and oomycetes. However, the study could not find an obvious correlation between the metabolic profile of the tested LABs and their antifungal efficacy. Not long after, Kavková *et al.* (2023) reported a notable inhibition of mycelial growth and conidial germination by *L. plantarum* and *L. pentosus* against *Fusarium* spp. from legumes by their cell-free supernatants.

In this study, we explored the potential of beneficial bacteria from a public culture collection and locally isolated *L. plantarum* from tempoyak, a fermented food made from durian flesh, to inhibit the growth of *Foc* TR4 *in vitro* and suppress the FWB in the greenhouse. The study aimed to include more biological control candidates with proven efficacy *in planta* to the current biocontrol resources for FWB.

## MATERIALS AND METHODS

### Experimental Site

The study was carried out in the Plant Molecular Biology Laboratory and greenhouse of the Department of Cell and Molecular Biology, Faculty of

Biotechnology and Biomolecular Sciences,  
University Putra, Malaysia, from January to  
October 2023.

### Bacterial and Fungal Cultures

*Foc* TR4 isolate 9888 was obtained from the Dept. of Biology, Faculty of Science, UPM. *Bacillus velezensis* B4158, *B. atrophaeus* B363B, *B. amyloliquefaciens* B942, and *Streptomyces morookaensis* B12429 were obtained from NRRL Culture Collection, Illinois, USA. The *Bacillus* sp. was grown on Luria Bertani (LB) agar at 30°C. *S. morookaensis* B12429 was cultured on Starch-Casein agar at 30°C. *L. plantarum* AM2 was previously isolated by a postgraduate student in the Plant Molecular Biology lab from a fermented durian (local fruit) flesh, known as 'tempoyak' among the locals, and deposited to the Microbial Culture Collection Unit (UNiCC), Institute of Bioscience, UPM (UNiCC Accession Number: UPMC 1489). *L. plantarum* AM2 was cultured on Man de Rosa (MRS) agar media at 37°C.

### In Vitro Antifungal Assay

The *in vitro* antifungal activity of the bacterial candidates against *Foc* TR4 was assessed in a dual culture plate assay where *Foc* TR4 was co-cultured on Potato Dextrose Agar (PDA) with a single bacterial strain. Then, a mycelial plug was cut from the edge of a 5-day-old *Foc* TR4 culture using a 5-mm-diameter sterile cork borer. The plug was positioned in the middle of a 25mL solidified PDA in a 100-mm petri dish. On opposing ends of the plug, 10 µL of bacterial suspension was streaked in a straight line, maintaining a distance of 2.5 cm from the plug, ensuring that the streaks of bacteria ran parallel to each other. The inhibition rate (%) was measured as follows:

(Colony diameter of the untreated group-  
Colony diameter of the treated group)/Colony  
diameter of untreated group×100.

### In Planta Bioassay

In this study, three-month-old *Musa acuminata* var. Cavendish plantlets with 4 to 5 true leaves in a polybag (13 cm height×10 cm diameter) were used for the bioassay with *Foc* TR4. The plantlets were purchased from Apex Phytocultures, Bangi, Malaysia, and maintained in a shaded greenhouse with daily watering and supplemented with inorganic fertilizer twice during the bioassay period. The bioassay was conducted according to Zhu *et al.* (2021b) in five replications for each treatment. A piece of agar with actively growing *Foc* TR4 mycelia was cut from a 4-day-old PDA plate and inoculated into 100 mL Potato Dextrose Broth (PDB) in a flask. The flask was cultured for 3 days on a shaker at 180 rpm at room temperature. After 3 days, the suspension culture was filtered using a sterile filter paper to remove the mycelia. The filtered suspension was then diluted to  $1 \times 10^6$  conidia mL<sup>-1</sup> using sterile double distilled water. Meanwhile, individual bacterial culture was prepared by inoculating each bacterium in their respective broth media of 150 mL in volume as described above. The inoculated media were incubated in a shaking incubator according to the specific temperature for each bacterium, as described above, at 180 rpm for 48 h. A ten-fold serial dilution was performed using 1X phosphate buffer saline to obtain  $1 \times 10^8$  cfu mL<sup>-1</sup> spore suspension. A total of 30 mL bacterial culture was used to drench the roots of the banana plantlets. After 14 days of treatment, 30 mL of *Foc* TR4 spore suspension was applied to the banana plantlets by pouring on the potting medium. The plantlets were allowed to grow for 8 weeks, and external disease symptoms were recorded by observing the yellowing and wilting symptoms. At the end of week 8, the plantlets were up-rooted and cut longitudinally at the rhizome to record the internal symptoms. The Leaf Symptom (LSI), Rhizome Discoloration Index (RDI), and Disease Severity Index (DSI) were



calculated based on Dita *et al.* (2021). The disease suppressiveness of the bacterial treatment was calculated as follows:

Disease suppressiveness = (Control disease index – Treatment disease index) / Control disease index × 100.

### Statistical Analysis

The data for *in vitro* antifungal assay were analyzed using Analysis Of Variance (ANOVA), and differences between treatments were determined using Tukey's HSD test at a 5% error. The data for *in planta* bioassay were analysed using one-way ANOVA on ranks (Kruskal-Wallis's test).

## RESULTS AND DISCUSSION

### *In Vitro* Assay of Bacterial Isolates against *Foc* TR4

Five bacterial strains, including *B. velezensis* B4158, *B. atrophaeus* B363B, *B. amyloliquefaciens* B942, *S. morookaensis* B12429, and *L. plantarum* AM2, were evaluated for their antifungal efficacy against *Foc* TR4 by a dual culture assay. All the bacterial isolates tested showed the ability to inhibit the growth of *Foc* TR4 (Figure 1). The greatest reduction in mycelial growth was elicited by *L. plantarum* AM2 with the inhibition percentage 42.13%, whereas *S. morookaensis* B12429 showed the least inhibition (31%) (Table 1). On the other hand, all *Bacillus* spp. tested moderately inhibited the mycelial growth of *Foc* TR4.


*L. plantarum* stands out as among the most prevalent and adaptable species within the LAB family. It is well known for its role in inhibiting fungal growth and removing mycotoxins, especially in food applications (Vanitha *et al.*, 2023; Li *et al.*, 2023). Many *L. plantarum* strains can suppress the growth of fungi by disintegrating the cell structure under the action of its metabolites. It also

has the ability to detoxify and degrade mycotoxins (Bergsma *et al.*, 2022; Wei *et al.*, 2020; Zhu *et al.*, 2021a). As a hemi-biotrophic pathogen, *Foc* TR4 utilizes an array of virulent factors to infect the host plants, including phytotoxic secondary metabolites Fusaric Acid (FSA) that could induce cell death in bananas, preparing the plants for xylem invasion (Li *et al.*, 2013). Recently, the antifungal activity of *L. plantarum* was reported against the plant pathogen *F. oxysporum* (Kavková *et al.*, 2023; Riolo *et al.*, 2023). However, the underlying mechanisms of the antifungal activity were not clearly described. It is tempting to speculate that *L. plantarum* might be suppressing the growth of *Foc* TR4 mycelia via the metabolites secreted into the medium that could distort hyphal structures, as observed by Deepthi *et al.* (2016).

On the other hand, despite the potent antifungal effects of *Bacillus* spp. against *F. oxysporum*, as previously reported (Fan *et al.*, 2021; Saravanan *et al.*, 2022), the *in vitro* effects of *B. velezensis* B4158, *B. atrophaeus* B363B, and *B. amyloliquefaciens* B942 in this study were only moderate and without much variation. Fan *et al.* (2021) reported an *in vitro* inhibition rate of 79.6% with endophytic *B. amyloliquefaciens* YN0904 against *Foc* TR4, while Saravanan *et al.* (2022) observed up to 63% inhibition by *B. velezensis* YEBBR6. Conversely, *B. atrophaeus* was reported to inhibit *F. oxysporum* f. sp. *cucumerinum* with high efficacy, but there has been no report of *B. atrophaeus* against *Foc* TR4 to date. The biocontrol capacity of *Bacillus* spp. is mainly exhibited through direct inhibitory activity on the growth of pathogens, induction of systemic resistance in host plants, and niche competition with the plant pathogens (Fira *et al.*, 2018). For instance, the application of *B. amyloliquefaciens* mixed with a compost mixture was shown to induce systemic resistance of the banana plants through the upregulation of hydrolytic enzyme activities such as chitinase and  $\beta$ -1,3-glucanase (Zhang *et al.*, 2014). Additionally, *B.*

*amyloliquefaciens* and *B. velezensis* can interact with *Foc* TR4 and establish their

produce volatile compounds for the biocontrol of plant diseases (Asari *et al.*,



**Figure 1.** Dual culture assay of antagonistic bacteria against *F. oxysporum* f. sp. *cubense* (TR4) on potato dextrose agar (PDA). Control (A), culture treated with *B. velezensis* B4158 (B), *B. atrophaeus* B363B (C), *B. amyloliquefaciens* B942 (D), *S. morookaensis* B12429 (E), and *L. plantarum* AM2 (F).

**Table 1.** Inhibition percentage of *Fusarium oxysporum* f. sp. *cubense* (TR4) by antagonistic bacteria.<sup>a</sup>

Antagonistic Bacteria	Inhibition Percentage (%)
Positive Control	0.00 ± 0.00 <sup>a</sup>
<i>B. velezensis</i> B4158	36.56 ± 0.46 <sup>c</sup>
<i>B. atrophaeus</i> B363B	38.55 ± 0.31 <sup>d</sup>
<i>B. amyloliquefaciens</i> B942	36.22 ± 0.46 <sup>c</sup>
<i>S. morookaensis</i> B12429	31.00 ± 0.46 <sup>b</sup>
<i>L. plantarum</i> AM2	42.13 ± 0.93 <sup>c</sup>

<sup>a</sup> The values are the mean percentage of five replications of the inhibition diameter zone. Different letter indicates significant differences between treatments of antagonistic bacteria and control at P < 0.05 according to Turkey's HSD test.

colonies in banana plants effectively to exert their inhibitory effects (He *et al.*, 2021). Based on our findings from the *in vitro* dual culture assay, the observed inhibition could be attributed to the diffusible compounds released by *Bacillus* spp., albeit less effective. In this case, *B. amyloliquefaciens* and *B. atrophaeus* strains were known to

2016; Rajaofera *et al.*, 2019). The ability of the *Bacillus* spp. used in this study to produce volatile compounds and their efficacy against *Foc* TR4 would be interesting subjects of future studies.

Finally, our *Streptomyces* sp. candidate, *S. morookaensis* B12429 only managed to exert a mild antagonistic activity on the






growth of *Foc* TR4 *in vitro*. In contrast, another *S. morookaensis*, strain Sm4-1986, was in the limelight recently due to its ability to promote banana growth and suppress FWB (Zhu *et al.*, 2021b). This could be attributed to the different metabolites and genetic profile of each strain that leads to different levels of antagonistic activity.


### Suppressive Effects of Bacterial Strains on FWB in Planta

Despite the variation of inhibition percentage *in vitro*, all the bacterial strains were used in the greenhouse studies, since the response of BCA during *in vitro* and *in*

*planta* assay did not always correlate (Parikh *et al.*, 2018; Besset-Manzoni *et al.*, 2019). Interestingly, *L. plantarum* AM- inoculated plantlets showed delayed wilting symptoms compared to the other treatments. The disease severity for plants co-inoculated with *L. plantarum* AM2, *B. amyloliquefaciens* B942 and *S. morookaensis* B12429 was significantly reduced (Figure 2). The lowest LSI score was recorded in *L. plantarum* AM2 at 25%, while the lowest RDI score was in *B. amyloliquefaciens* B942 at 31.76% (Figure 3). Conversely, *B. atrophaeus* B363B displayed the highest disease severity, with both LSI and RDI scores at 95 and 94%, respectively.



**Figure 2.** External and internal symptoms of different treatments on week 8. Negative control (A1- External) and (A2-Internal), *B. atrophaeus* B363B (B1- External) and (B2-Internal), *B. velezensis* B4158 (C1- External) and (C2-Internal), *B. amyloliquefaciens* B942 (D1- External) and (D2- Internal), *L. plantarum* AM2 (E1- External) and (E2- Internal), *S. morookaensis* B12429 (F1- External) and (F2- Internal), Positive control (G1- External) and (G2- Internal).



**Figure 3.** Leaf Symptoms Index (LSI) and Rhizome Discolouration Index (RDI) of different treatments. Negative Control - no *Foc* TR4, no biological control (C), *B. Atrophaeus* B363B (BA), *B. Velezensis* B4158 (BV), *B. Amyloliquefaciens* B942 (BAM), *L. Plantarum* AM2 (LP), *S. Morookaensis* B12429 (SM), Positive control-Inoculated with *Foc* TR4, no biological control (TR4). Different letters show different significance values at  $P > 0.05$  using one-way ANOVA on ranks (Kruskal-Wallis's test).

**Table 2.** Effects of treatment with antagonistic bacteria on the FWB development *in planta* and their biocontrol efficacy.

Antagonistic bacteria	Disease severity index (%)	Disease suppressiveness (%)
<i>B. atrophaeus</i> B363B	100	0
<i>B. velezensis</i> B4158	68	32
<i>B. amyloliquefaciens</i> B942	32	68
<i>L. plantarum</i> AM2	24	76
<i>S. morookaensis</i> B12429	32	68
Positive control (TR4)	100	-

Banana plants inoculated with *L. plantarum* AM2 showed the lowest DSI at 24% (Table 2), which aligned with the findings from the *in vitro* assay. These findings further support the potential of *L. plantarum* AM2 as a bio-control agent against FWB. Following closely were *B. amyloliquefaciens* B942 and *S. morookaensis* B12429, with DSI at 32%. Previous research has highlighted the effectiveness of different strains of *B. velezensis* and *B. amyloliquefaciens* as bio-control agents *in planta* (Fan *et al.*, 2021; Fu *et al.*, 2017; Xiang *et al.*, 2023). However, in this study, only the *B. amyloliquefaciens* B942 strain demonstrated a significant effect against FWB. Conversely, plantlets

inoculated with *B. atrophaeus* B363B and *B. velezensis* B4158 exhibited severe disease symptoms, both externally and internally, despite the strong inhibition observed *in vitro*. On the other hand, *S. morookaensis* strain B12429 only demonstrated moderate antagonistic effects against *Foc* TR4 in the greenhouse experiment. To the best of our knowledge, this study is the first to report on the biocontrol potential of *S. morookaensis* strain B12429 and *B. amyloliquefaciens* B942 strain against *Foc* TR4 *in planta*.

Nevertheless, further testing through field trials is needed to fully assess the potential of these bacteria as biocontrol agents, since it offers a more reliable and realistic assessment. The interactions and



competition of bio-control agents with the soil microbiome, as well as the adaptation of potential strains to the abiotic conditions, limits their efficiency against pathogens (Purkayastha *et al.*, 2018). These underscore the necessity to validate *in vivo* experiments conducted in greenhouses with field trials.

## CONCLUSIONS

This study explores the antagonistic potential of selected microbes as biocontrol agents against Fusarium Wilt of Bananas (FWB) in the susceptible banana cultivar, Cavendish. The results indicate that *L. plantarum* AM2, *B. amyloliquefaciens* B942 and *S. morookaensis* B12429 inhibited the mycelial growth of *Foc* TR4 *in vitro* and decreased the severity of FWB *in planta*. Inoculation with *L. plantarum* AM2 resulted in the lowest DSI at 24%. The microbes tested in this study showed potential in suppressing FWB and were thus considered promising candidates for biological control. However, trials in open fields are necessary to further assess the efficacy of these antagonistic microbes in a more realistic environment.

## ACKNOWLEDGEMENTS

We would like to thank Mr. Malik Abd Razak for his work in isolating *L. plantarum* AM2 used in this study. This work was supported by the Ministry of Higher Education Malaysia (FRGS/1/2019/STG03/UPM/02/7).

## REFERENCES

- Asari, S., Matzén, S., Petersen, M. A., Bejai, S. and Meijer, J. 2016. Multiple Effects of *Bacillus amyloliquefaciens* Volatile Compounds: Plant Growth Promotion and Growth Inhibition of Phytopathogens. *FEMS Microbiol. Ecol.*, **92**(6): 1-11.
- Bergsma, S., Euverink, G. J. W., Charalampogiannis, N., Poullos, E., Janssens, T. K. and Achinas, S. 2022. Biotechnological and Medical Aspects of Lactic Acid Bacteria Used for Plant Protection: A Comprehensive Review. *BioTech.*, **11**(3): 1-16.
- Bessey-Manzoni, Y., Joly, P., Brutel, A., Gerin, F., Soudiere, O., Langin, T. and Prigent-Combaret, C. 2019. Does *in Vitro* Selection of Biocontrol Agents Guarantee Success *in Planta*? A Study Case of Wheat Protection against Fusarium Seedling Blight by Soil Bacteria. *PLoS One*, **14**(12): 1-18.
- Bubici, G., Kaushal, M., Prigigallo, M. I., Gómez-Lama Cabanás, C. and Mercado-Blanco, J. 2019. Biological Control Agents against Fusarium Wilt of Banana. *Front. Microbiol.*, **10**: 1-33.
- Caro, L. P. 2020. Wages and Working Conditions in the Banana Sector: The Case of Costa Rica, Ethiopia, India, Indonesia, and Viet Nam: Background Note. International Labour Organization (ILO), Geneva, 28 PP.
- Deepthi, B. V., Poornachandra Rao, K., Chennappa, G., Naik, M. K., Chandrashekara, K. T. and Sreenivasa, M. Y. 2016. Antifungal Attributes of *Lactobacillus plantarum* MYS6 against Fumonisin-Producing *Fusarium proliferatum* Associated with Poultry Feeds. *PLoS One*, **11**(6): 1-22.
- He, P., Li, S., Xu, S., Fan, H., Wang, Y., Zhou, W., Fu, G., Han, G., Wang, Y. Y. and Zheng, S. J. 2021. Monitoring Tritrophic Biocontrol Interactions between *Bacillus* spp., *Fusarium oxysporum* f. sp. *cubense*, Tropical Race 4, and Banana Plants *in Vivo* Based on Fluorescent Transformation System. *Front. Microbiol.*, **13**(12): 1-13.
- Purkayastha, G. D., Mangar, P., Saha, A. and Saha, D. 2018. Evaluation of the Biocontrol Efficacy of a *Serratia marcescens* Strain Indigenous to Tea Rhizosphere for the Management of Root Rot Disease in Tea. *PLoS One*, **13**(2): 1-27.
- Dita, M. A., Teixeira, L., Li, C., Zheng, S., O'Neill, W., Daniels, J., Pérez-Vicente, L., Carreel, F., Roussel, V., Carlier, J., Abadie, C., Carpentier, S. C., Iyyakutty, R., Kissel, E., van Wesemael, J., Chase, R., Tomekpe, K. and Roux, N. 2021. *Practical Guidelines for Early Screening and Field Evaluation of Banana against Fusarium Wilt, Pseudocercospora Leaf Spots and Drought*. Bioversity International, Montpellier, France, 83 PP.



10. EFSA BIOHAZ Panel. 2023. *Updated List of QPS-Recommended Microorganisms for Safety Risk Assessments Carried Out by EFSA*. The European Food Safety Authority (EFSA). Retrieved December 10, 2023 from <https://zenodo.org/records/8124409>
11. Fan, H., Li, S., Zeng, L., He, P., Xu, S., Bai, T., Huang, Y., Guo, Z. and Zheng, S. -J. 2021. Biological Control of *Fusarium oxysporum* f. sp. *cubense* Tropical Race 4 Using Natively Isolated *Bacillus* spp. YN0904 and YN1419. *J. Fungi*, **7(10)**: 1-17.
12. Fira, D., Dimkić, I., Berić, T., Lozo, J. and Stanković, S. 2018. Biological Control of Plant Pathogens by *Bacillus* Species. *J. Biotech.*, **285**: 44-55.
13. FAO. 2022. *Banana Market Review-Preliminary Results 2021*. Food and Agriculture Organization of the United Nations, Rome.
14. Fu, L., Penton, C. R., Ruan, Y., Shen, Z., Xue, C., Li, R., and Shen, Q. 2017. Inducing the Rhizosphere Microbiome by Biofertilizer Application to Suppress Banana Fusarium Wilt Disease. *Soil Biol. Biochem.*, **104**: 39-48.
15. Jaffar, N. S., Jawan, R. and Chong, K. P. 2023. The Potential of Lactic Acid Bacteria in Mediating the Control of Plant Diseases and Plant Growth Stimulation in Crop Production- A Mini Review. *Front. Plant Sci.*, **13**: 1-13.
16. Kavková, M., Bazalová, O., Cihlár, J., Bohatá, A., Lencová, J. and Konvalina, P. 2023. Characterization of Wild Strains of Lactic Acid Bacteria Isolated from Legumes and Their Biocontrol Potential against *Fusarium* spp. *Agronomy*, **13(12)**: 1-19.
17. Lahlali, R., Ezrari, S., Radouane, N., Kenfaoui, J., Esmael, Q., El Hamss, H., Belabess, Z. and Barka, E. A. 2022. Biological Control of Plant Pathogens: A Global Perspective. *Microorganisms*, **10(3)**: 1-33.
18. Li, C., Zuo, C., Deng, G., Kuang, R., Yang, Q., Hu, C., Sheng, O., Zhang, S., Ma, L., Wei, Y., Yang, J., Liu, S., Biswas, M.K., Viljoen, A. and Yi, G. 2013. Contamination of Bananas with Beauvericin and Fusaric Acid Produced by *Fusarium oxysporum* f. sp. *cubense*. *PLoS One*, **8(7)**: 1-11.
19. Li, Q., Zeng, X., Fu, H., Wang, X., Guo, X. and Wang M. 2023. *Lactiplantibacillus plantarum*: A Comprehensive Review of Its Antifungal and Anti-mycotoxic Effects. *Trends Food Sci. Tech.*, **136**: 224-238.
20. Ordóñez, N. 2018. A Global Genetic Diversity Analysis of *Fusarium oxysporum* f. sp. *cubense*: the Panama Disease Pathogen of Banana. Doctoral Dissertation, Wageningen University and Research.
21. Parikh, L., Eskelson, M. J. and Adesemoye, A. O. 2018. Relationship of *in Vitro* and *in Planta* Screening: Improving the Selection Process for Biological Control Agents against Fusarium Root Rot in Row Crops. *Arch. Phytopathol. Plant Protect.*, **51(3-4)**: 156-169.
22. Pérez-Vicente, L., Dita, M. A. and Martínez-de la Parte, E. 2014. Technical Manual Prevention and Diagnostic of Fusarium Wilt (Panama Disease) of Banana Caused by *Fusarium oxysporum* f. sp. *cubense* Tropical Race 4 (TR4). Food and Agriculture Organization of The United Nations.
23. Ploetz, R. C. 2015. Fusarium Wilt of Banana. *Phytopathol.*, **105(12)**: 1512-1521.
24. Rajaofera, M. J. N., Wang, Y., Dahar, G. Y., Jin, P., Fan, L., Xu, L., Liu, W. and Miao, W. 2019. Volatile Organic Compounds of *Bacillus atrophaeus* HAB-5 Inhibit the Growth of *Colletotrichum gloeosporioides*. *Pest. Biochem. Physiol.*, **156**: 170-176.
25. Raman, J., Kim, J. S., Choi, K. R., Eun, H., Yang, D., Ko, Y. J. and Kim, S. J. 2022. Application of Lactic Acid Bacteria (LAB) in Sustainable Agriculture: Advantages and Limitations. *Intl. J. Mol. Sci.*, **23(14)**: 1-22.
26. Riolo, M., Luz, C., Santilli, E., Meca, G. and Cacciola, S. O. 2023. Antifungal Activity of Selected Lactic Acid Bacteria from Olive Drupes. *Food Biosci.*, **52**: 1-11.
27. Sánchez-Espinosa, A.C., Villarruel-Ordaz, J. L. and Maldonado-Bonilla, L. D. 2020. The Cause and Potential Solution to the Fusarium Wilt Disease in Banana Plants. *Terra Latinoam*, **38(2)**: 435-442.
28. Saravanan, R., Nakkeeran, S., Saranya, N., Kavino, M., Ragapriya, V., Varanavasiappan, S., Raveendran, M., Krishnamoorthy, A. S., Malathy, V. G. and Haripriya, S. 2022. Biohardening of Banana cv. Karpooravalli (ABB; Pisang Awak) with *Bacillus velezensis* YEBBR6 Promotes Plant Growth and Reprograms the Innate Immune Response against *Fusarium oxysporum* f. sp. *cubense*. *Front. Sustain. Food Syst.*, **6**: 1-18.
29. Scortichini, M. 2022. Sustainable Management of Diseases in Horticulture:



- Conventional and New Options. *Horticulturae*, **8(6)**: 1-31.
30. Seddik, H. A., Bendali, F., Gancel, F., Fliss, I., Spano, G. and Drider, D. 2017. *Lactobacillus plantarum* and Its Probiotic and Food Potentialities. *Probiotics Antimicrob. Proteins*, **9(2)**: 111-122.
  31. Vanitha, P. R., Somashekaraiah, R., Divyashree, S., Pan, I. and Sreenivasa, M. Y. 2023. Antifungal Activity of Probiotic Strain *Lactiplantibacillus plantarum* MYSN7 against *Trichophyton tonsurans*. *Front. Microbiol.*, **14**: 1-16.
  32. Wei, C., Yu, L., Qiao, N., Wang, S., Tian, F., Zhao, J., Zhang, H., Zhai, Q. and Chen, W. 2020. The Characteristics of Patulin Detoxification by *Lactobacillus plantarum* 13M5. *Food Chem. Toxicol.*, **146**: 1-12.
  33. Xiang, D., Yang, X., Liu, B., Chu, Y., Liu, S. and Li, C. 2023. Bio-Priming of Banana Tissue Culture Plantlets with Endophytic *Bacillus velezensis* EB1 to Improve Fusarium Wilt Resistance. *Front. Microbiol.*, **14**: 1146331.
  34. Zhang, N., He, X., Zhang, J., Raza, W., Yang, X. -M., Ruan, Y. -Z., Shen, Q. -R. and Huang, Q. -W. 2014. Suppression of Fusarium Wilt of Banana with Application of Bio-Organic Fertilizers. *Pedosphere*, **24** (5): 613-624.
  35. Zheng, J., Wittouck, S., Salvetti, E., Franz, C. M. A. P., Harris, H. M. B., Mattarelli, P., O'Toole, P. W., Pot, B., Vandamme, P., Walter, J., Watanabe, K., Wuyts, S., Felis, G. E., Gänzle, M. G. and Lebeer, S. 2020. A Taxonomic Note on the Genus *Lactobacillus*: Description of 23 Novel Genera, Emended Description of the Genus *Lactobacillus* Beijerinck 1901, and Union of *Lactobacillaceae* and *Leuconostocaceae*. *Intl. J. Syst. Evol. Microbiol.*, **70(4)**: 2782-2858.
  36. Zhu, Y., Xu, Y. and Yang, Q. 2021a. Antifungal Properties and AFB1 Detoxification Activity of a New Strain of *Lactobacillus plantarum*. *J. Hazard Mater.*, **414(7)**: 1-9.
  37. Zhu, Z., Tian, Z. and Li, J. 2021b. A *Streptomyces morookaensis* Strain Promotes Plant Growth and Suppresses Fusarium Wilt of Banana. *Trop. Plant Pathol.*, **46**: 175-185.

**پتانسیل ضد قارچی *Lactiplantibacillus plantarum* AM2 علیه پاتوژن موز *Fusarium oxysporum* f. sp. cubense Tropical Race 4**

وان آتاتی نبیله وان تاج الدین شاه، نور بیٹی عبد مراد، جیاژین اونگ، شین هیوپی  
آنگ، نور لیلی، و نور بیت سعیدی

**چکیده**

پژمردگی فوزاریوم موز یک بیماری جدی است که مزارع موز را در سراسر جهان تحت تاثیر قرار می دهد. در تلاش برای مدیریت پایدار این بیماری، کنترل بیولوژیکی، جایگزینی امیدوارکننده برای مواد شیمیایی کشاورزی که می تواند اثرات مضری بر انسان و اکوسیستم داشته باشد در نظر گرفته می شود. در این پژوهش، ما پتانسیل کنترل-بیولوژیکی مجموعه فعلی باکتری های مفید، از جمله *Lactiplantibacillus plantarum* AM2، *Bacillus velezensis* B4158، *Streptomyces morookaensis* NRRL B-12429، *B. amyloliquus* a و *atrophaeus* B363B را از طریق یک روش

کشت دوگانه و یک آزمایش گلخانه ای بررسی کردیم. عامل بررسی شده، پژمردگی فوزاریوم در موز. sp. Tropical *Fusarium oxysporum* f (Foc TR4) cubense بود. محدوده مهار (inhibition range) در شرایط آزمایشگاهی بین ۳۱.۰٪ تا ۴۲.۱٪ بود و بیشترین مهار رشد Foc Race 4 TR4 برای *L. plantarum* AM2 مشاهده شد. گیاهچه‌های موز آلوده که تیمار با *L. plantarum* AM2 را دریافت کردند نیز در مقایسه با تیمار سایر باکتری‌های مفید، تا ۲۴ درصد کاهش قابل توجهی را در شاخص شدت بیماری بروز دادند. این پژوهش نشان داد که *L. plantarum* AM2 اثر آنتاگونیستی خوبی بر رشد میسلیم Foc TR4 دارد و بیشترین پتانسیل را برای کنترل بیماری پژمردگی فوزاریوم در موز دارد.