

## Evaluating *Bacillus* spp. As Biocontrol Agents against *Meloidogyne incognita* Infesting Tomato

Bommenahalli Sathish Kavya<sup>1\*</sup>, Tiptur Roopla-Nayak Kavitha<sup>2</sup>, Arati<sup>3</sup>, and Nagavath Lohith Kumar<sup>3</sup>

### ABSTRACT

Tomato is a global host for root-knot nematode (*Meloidogyne incognita*), soliciting ponderous damage. Using bio-control agents to control plant parasitic nematodes is a well-established, green approach in advance of synthetic nematicides. The role of *Bacillus* spp. in inciting physiological and biochemical alterations in nematode infestation is discussed in the present study. The susceptible (PKM-1) and resistant (Hisar Lalit) tomato cultivars treated with *Bacillus pumilus* augmented the shoot length, root length and biomass of plants compared to the standard check, *Pseudomonas fluorescens*, followed by *B. megaterium*. Accordingly, all the bio-control agent-treated susceptible plants showed reduced galling and exhibited a root gall index of 3 (moderately resistant) and reduced nematode population in soil and roots. Contrarily, all the resistant plants showed highly resistant reactions. *B. pumilus* showed the topmost expression of all the biochemical enzymes like Peroxidase (PO), Polyphenol Oxidase (PPO), Catalase (CAT), Phenylalanine Ammonia-Lyase (PAL), and total phenols. Conclusively, *B. pumilus* was found to be the most potential in reducing nematode infestation by embellishing the plant growth and enhancing defence-related enzymes in tomatoes.

**Keywords:** Biochemical enzymes, Plant parasitic nematodes, *Pseudomonas fluorescens*, Root-knot nematode.

### INTRODUCTION

Plant Parasitic Nematodes (PPNs) are one of the egregious biotic stresses that are a well-recognized threat to crop production and food security. PPNs are omnipresent, enabling them to attack all different kinds of crop plants, field crops, and vegetable and flower crops that cause annual economic losses of USD 173 billion globally (Elling *et al.*, 2013). They are also responsible for multiple pathogen infection that predisposes the susceptible crops to other pathogens like fungi, bacteria and thus indirectly contributing to

the reduced yield and crop productivity (Back *et al.*, 2002).

There exist more than 4,000 PPNs attacking crops. Of these, the most important and prominent one is the root-knot nematode (*Meloidogyne* spp.), which voraciously feeds all crops, especially vegetables (Koenning *et al.*, 1999). Global vegetable production is under threat due to the damage from four different species of *Meloidogyne* viz., *M. arenaria*, *M. javanica*, *M. incognita*, and *M. hapla*. However, *M. incognita* is one of the important pests of solanaceous crops majorly in tomatoes.

<sup>1</sup> Department of Plant Pathology, University of Agricultural Sciences, GKVK, Bengaluru-65, Karnataka, India.

<sup>2</sup> AICRP (Nematodes) Laboratory, Department of Plant Pathology, University of Agricultural Sciences, GKVK, Bengaluru-65, Karnataka, India.

<sup>3</sup> Department of Agricultural Microbiology, University of Agricultural Sciences, GKVK, Bengaluru-65, Karnataka, India.

\* Corresponding author; e-mail: kavyavishu2210@gmail.com



Tomato is the second most paramount remunerable exigent solanaceous vegetable grown worldwide after potato. Infestation of *M. incognita* in tomatoes impedes production and lowers yield, making them vulnerable to other wilt and/or rot causing pathogens (Ogwulumba et al., 2011). The habitual monitoring tactic employed by growers is the use of synthetic nematicides, which create economic and environmental constraints. Under these circumstances, substitutive strategies are gaining preponderance. The current study explains one such alternative strategy to combat root-knot nematode infesting tomatoes.

Bio-control or biological control is an economically and environmentally well-fit tactic against plant pathogens. Precisely, bio-control of nematodes can be defined as the management of nematode populations and the damage caused by them through the action of antagonists, either directly or indirectly, by manipulating the environment favourable for nematodes (Poveda et al., 2020) that includes many of fungi and bacterial species. The organism that shows antagonism against pathogens are called bio-control agents (BCA), where these organisms interact with pathogens through various mechanisms like competition, antibiosis and/or through inducing the plant resistance against pathogens.

Plant resistance is one of the defensive strategies used by most of the hosts against pest and pathogen attacks. Once the plant encounters the attack, the Pathogen-Associated Molecular Patterns (PAMPs) will be recognized by the Pattern Recognition Receptors (PRRs) on the cell surface. Thereby it triggers a complex signalling network leading to defence responses (PAMP-triggered immunity) (Ramirez-Prado et al., 2018; Hou et al., 2019). When it fails, these bio-control agents help the plant to trigger its innate immunity by producing Microbial-Associated Molecular Patterns (MAMPs) that act as elicitors for triggering the

complex defence responses by the host against pathogens (Poveda et al., 2020). It includes the production of reactive oxygen species, hypersensitive response, production of defensive enzymes (peroxidase, polyphenol oxidase, phenylalanine ammonia-lyase, catalase, etc.), phyto-hormones (salicylic acid, jasmonic acid and/or ethylene), and many signalling cascades.

In the current study, we investigated the nematocidal potential of a bacterial bio-agent, *Bacillus* spp. against *M. incognita* in tomato and the level of defensive enzymes across different *Bacillus* spp. treated plants.

## MATERIALS AND METHODS

The experiments were laid out in a glasshouse at AICRP (Nematodes), Zonal Agricultural Research Station, GKVK, Bengaluru. Precedently, the species of *Meloidogyne* was validated as *M. incognita* by using the “Perineal pattern technique” (Chitwood, 1949).

### Collection of Bacterial Culture

Three distinct species of *Bacillus* viz., *B. subtilis* IIHR Bs-2, *B. megaterium* Bm-IIHR, *B. pumilus* IIHR Bp-2 cultures were obtained from ICAR-Indian Institute of Horticultural Research, Bengaluru, whereas *Pseudomonas fluorescence* (standard check) was obtained from Pathogenomics Laboratory, Department of Plant Pathology, University of Agricultural Sciences, GKVK, Bengaluru.

### Collection of Tomato Seeds

Tomato seeds of nematode susceptible variety, PKM-1, were collected from IIHR, Bengaluru, and tomato seeds of nematode resistant variety, Hisar Lalit (NRT 8), were collected from Chaudhary Charan

Singh Agricultural University, Hisar, Haryana. Tomato plants were raised in portrays and transplanted into pots (2 kg) filled with sterilized soil after 21 days.

### Preparation of Bacterial and Nematode Suspension

The bacterial suspensions of respective bio-agents were prepared by growing a bacteria in 250 mL nutrient broth and incubating for 24 h. The turbid broth was centrifuged at 5,000 rpm for 5 minutes and cells were re-suspended with phosphate buffer (pH= 7). The concentration was maintained at the rate of  $2.5 \times 10^7$  CFU mL<sup>-1</sup> and the bacterial suspension was inoculated through the seedling dip method to 25-day-old seedlings before translation for 3 hours. Here, *Pseudomonas fluorescens* was used as a standard check.

Nematode inoculum was obtained from infected tomato roots. We extracted the nematode by following Combined Cobb's sieving and Baermann's funnel technique (Ayoub, 1977): 50 mL of nematode suspension was prepared at the rate of 20 juveniles per mL of water. After a week of transplanting, the nematode suspension of 50 mL per plant with 1,000 juveniles was inoculated by making 4 -5 holes around the seedling.

Nematicide-treated (carbofuran@ 1 g kg<sup>-1</sup> pot) plants were considered as a positive control. Complete Random Design (CRD) was maintained in a glass house (temperature: 20 °C) with three replications of seven treatments. The experiment was conducted twice.

### Studies on Physiological Alterations in Tomato

The efficacy of *Bacillus* spp. on the incidence and establishment of root-knot nematode in tomatoes was studied through various parameters of plant growth and nematode infection under pot experiments.

After 45 days of inoculation, observations viz., shoot growth (cm), root growth (g) and root-knot index (1-5 scale), nematode population in roots and soil were taken (Narasimhamurthy *et al.*, 2017).

### Status of Defence-Related Biochemicals in Tomato

The biochemical levels in tomato-treated and untreated plants were assessed. One gram of fresh root was collected from the susceptible and resistant plants after 28 days of nematode inoculation from each treatment for biochemical analysis. Fresh roots were washed gently in running tap water and homogenized in 1 mL of 0.1 M phosphate buffer (pH 7.0) at 4°C in a pre-chilled pestle and mortar. The homogenate was centrifuged at 20,000 rpm at 4°C for 15 minutes and the supernatant served as an enzyme source for further analysis (Anita *et al.*, 2004).

Then, tomato plants were tested for biochemical levels of several defence-related enzymes like Peroxidase (PO), Polyphenol Oxidase (PPO), Phenylalanine Ammonia-Lyase (PAL), Catalase (CAT) and total phenols, using spectrophotometer by following standard procedures (Prabu *et al.*, 2019).

### Estimation of Peroxidase (PO) Activity

Peroxidase activity was assayed spectrophotometrically (Chander, 1990). The reaction mixture consisted of 2.5 mL of 0.05 M potassium phosphate buffer, 0.2 mL of O-Phenyl Diamine (OPD), 0.2 mL of 0.3 per cent H<sub>2</sub>O<sub>2</sub> and 0.03 mL of enzyme extract. The reaction mixture was incubated at room temperature (28±1°C). The change in absorbance was recorded at 60 seconds intervals for 5 min. The enzyme preparation without H<sub>2</sub>O<sub>2</sub> served as blank. The enzyme activity was expressed as change in the absorbance at 450 nm min<sup>-1</sup> g<sup>-1</sup> on fresh weight basis.



### Estimation of Polyphenol Oxidase (PPO)

PPO activity was determined as per the procedure given by Selvaraj and Kumar (1995). The reaction mixture consisted of 2.9 mL of 0.05M potassium phosphate buffer (pH 6.8), 0.1 mL of 1.25 per cent pyrogallol and enzyme extract of 0.5 mL. The increase in absorbance was measured at 450 nm up to 5 minutes for 1 minute interval. Polyphenol oxidase activity was expressed as absorbance  $\text{min}^{-1} \text{g}^{-1}$  FW (Fresh Weight).

### Estimation of Phenylalanine Ammonia Lyase (PAL)

The PAL assay was conducted as per the method described by Whetten and Senderoff (1992): 0.4 mL of enzyme extract was incubated with 0.5 mL of 0.1 M borate buffer (pH 8.8) and 0.5 mL of 12 mM L-phenylalanine in the same buffer for 30 minutes at 30°C. The reaction was arrested by adding 0.5 mL of 1M TCA and incubated at 37°C for 5 minutes. The blank was prepared, which contained 0.4 mL of crude enzyme extract and 2.7 mL of 0.1 M borate buffer (pH 8.8), and absorbance was measured at 290 nm in spectrophotometer. Standard curve was drawn with graded amounts of cinnamic acid dissolved in acetone. The enzyme activity was expressed as  $\mu\text{M}$  of trans-cinnamic acid  $\text{min}^{-1} \text{g}^{-1}$  FW.

### Estimation of Catalase

The catalase activity was estimated as per the procedure given by Masia (1998). The reaction mixture contained 2.6 mL of 0.067M sodium phosphate buffer (pH 7.0), 0.3 mL of 3%  $\text{H}_2\text{O}_2$  and 0.1 mL of enzyme extract. The decrease in absorbance was measured at 240 nm up to 5 min for 1 min time interval. Catalase activity was expressed as  $\mu\text{g}$   $\text{H}_2\text{O}_2 \text{g}^{-1}$  FW.

### Estimation of Total Phenol

Total phenol content was estimated by spectrophotometric method using Folin Ciocalteu Reagent (FCR) at an absorbance of 700 nm by following procedure (Singleton and Rossi, 1965). Five grams of sample was homogenized with 20 mL of methanol (80%) in a pestle and mortar 2-3 times and volume was made to 50 mL. Then, 0.5 mL of the extract was taken in test tubes, 0.2 mL of Folin-Ciocalteu's Phenol Reagent was added followed by 3.3 mL of distilled water and mixed well. After 2 minutes, 1 mL of sodium carbonate solution was added and mixed. This was allowed to stand at room temperature for 30 minutes and blue colour was read in a spectrophotometer at 700 nm. Standard curve for phenols was prepared using Gallic Acid (GA) as standard. The content of the total soluble phenol was calculated according to a standard curve obtained from a Folin Ciocalteu reagent with a phenol solution ( $\text{C}_6\text{H}_5\text{OH}$ ) and expressed as mg gallic acid equivalent  $100 \text{g}^{-1}$  FW.

### Data Analysis

The data so generated were analyzed using WASP – Web Agri Stat Software Package 2.0 developed by ICAR. Completely Randomized Design (CRD) was used in the present study. The difference between the treatment means was compared with the critical difference values to know a significant difference.

## RESULTS

### Physiological Transfiguration Incited by *Bacillus* spp. in Tomato

*Bacillus* spp. had shown its potential to embellish the shoot and root growth in relevance with the control treated with only

nematode (Table 1). The effectuation of *Bacillus pumilus* was premier among different *Bacillus* spp. and a standard check, *Pseudomonas fluorescence*.

### Effects on Tomato Growth

In the case of susceptible variety, despite the fact that carbofuran-treated plants showed overhead expression of shoot growth, *Bacillus pumilus*-treated plants showed the highest shoot length (57.2 cm) and shoot weight (Fresh weight: 179.1 g and Dry weight: 81.7 g) among different bio-agents followed by *B. megaterium* (shoot length: 55.3 cm; fresh weight: 175.9 g and dry weight: 77.6 g). Conversely, in the case of resistant variety, regardless of the bio-agents, the nematode-challenged plants exhibited the highest shoot growth, which was comparable with healthy plants, but the carbofuran-treated plants showed the least shoot growth.

A similar trend was noticed with respect to root growth as that of shoot growth in both root-knot nematode susceptible and resistant tomato varieties, respectively. In the case of susceptible variety, the root growth of the bacterial bio-agents treated-*Meloidogyne incognita*-infested plants showed higher values than only nematode-challenged plants, among which *B. pumilus* showed the maximal root length (30.7 cm) and root weight (Fresh weight: 7.6 g and Dry weight: 4.8 g). Despite the healthy resistant variety exhibiting the greatest root growth, the bioagent treatment did not result in a statistically significant effect.

### Effects on Nematode Infestation

The root-knot index was calculated by considering the number of galls produced per root system (Table 2). Except for the fact that the resistant variety showed highly resistant to resistant reactions with a root-knot index of 1-2, the bio-agents-treated plants revealed a lesser number of galls

(23.3-28.3) in susceptible variety and a root-knot index of 3, imposing a moderately resistant reaction. Of these, *B. pumilus* eliminated at most the production of galls on the roots (Figures 1-a and -b). Further, number of egg masses per root system (Table 2) and nematode population was evaluated both in soil and roots. The nematode population was significantly reduced in the *Bacillus* spp. treated plants compared to the control, showing the significance of bio-control treatment (Table 3).

### Biochemical Levels after *Bacillus* spp. Treatment in Tomato

In this study, bio-chemicals like Peroxidase (PO), Polyphenol Oxidase (PPO), Catalase (CAT), Phenylalanine Ammonia-Lyase (PAL) and total phenols were analyzed in both susceptible and resistant varieties. The inoculation of bio-agents to *M. incognita* infested plants showed increased activity of all tested enzymes (peroxidase, polyphenol oxidase, phenylalanine lyase, catalase and total phenols) over control in both susceptible and resistant varieties (Figures 2 and 3). Among bio-agents, *B. pumilus* showed the topmost expression of all the enzymes. However, the resistant variety showed higher enzymatic activity than the susceptible variety, except for the catalase enzyme.

### Studies on Peroxidase (PO) Activity

The activity of peroxidase was assayed spectrophotometrically at 450 nm as per procedure given by Chander (1990). The increased activity of peroxidase was recorded in *Meloidogyne incognita* inoculated samples of susceptible variety along with the resistant tomato variety (0.487 and 1.461 abs min<sup>-1</sup> g<sup>-1</sup>, respectively) compared to the respective healthy plants (0.187 and 0.561 abs min<sup>-1</sup> g<sup>-1</sup>), but

**Table 1.** Physiological alterations incited by *Bacillus* spp. in *M. incognita*-infested tomato plants.

Treatments	Susceptible variety				Resistant variety				Resistant variety			
	Shoot		Shoot		Shoot		Shoot		Shoot		Shoot	
	length (cm)	Fresh	length (cm)	Fresh	length (cm)	Fresh	length (cm)	Fresh	length (cm)	Fresh	length (cm)	Fresh
<i>M. incognita</i> +Carbofuran	60.75 (±0.08) <sup>b</sup>	185.25 (±0.69) <sup>b</sup>	59.71 (±0.35) <sup>c</sup>	163.24 (±0.28) <sup>f</sup>	33.82 (±0.36) <sup>b</sup>	71.86 (±0.37) <sup>e</sup>	33.82 (±0.36) <sup>b</sup>	8.90 (±0.22) <sup>b</sup>	33.68 (±0.28) <sup>e</sup>	9.03 (±0.27) <sup>e</sup>	33.68 (±0.28) <sup>e</sup>	9.03 (±0.27) <sup>e</sup>
<i>M. incognita</i> + <i>B. pumilus</i>	57.27 (±0.51) <sup>c</sup>	179.18 (±0.33) <sup>c</sup>	67.65 (±0.43) <sup>bc</sup>	208.09 (±0.34) <sup>b</sup>	30.73 (±0.24) <sup>c</sup>	97.54 (±0.28) <sup>ab</sup>	30.73 (±0.24) <sup>c</sup>	7.67 (±0.27) <sup>c</sup>	39.30 (±0.41) <sup>b</sup>	11.74 (±0.20) <sup>b</sup>	39.30 (±0.41) <sup>b</sup>	11.74 (±0.20) <sup>b</sup>
<i>M. incognita</i> + <i>B. megaterium</i>	55.34 (±0.29) <sup>d</sup>	175.90 (±1.36) <sup>d</sup>	67.38 (±0.33) <sup>cd</sup>	206.78 (±0.31) <sup>c</sup>	29.30 (±0.33) <sup>d</sup>	96.37 (±0.65) <sup>b</sup>	29.30 (±0.33) <sup>d</sup>	7.29 (±0.22) <sup>cd</sup>	37.86 (±0.2) <sup>c</sup>	11.28 (±0.35) <sup>bc</sup>	37.86 (±0.2) <sup>c</sup>	11.28 (±0.35) <sup>bc</sup>
<i>M. incognita</i> + <i>B. subtilis</i>	53.63 (±0.69) <sup>e</sup>	171.01 (±0.89) <sup>f</sup>	66.54 (±0.3) <sup>d</sup>	203.19 (±0.30) <sup>e</sup>	27.39 (±0.38) <sup>e</sup>	92.63 (±0.53) <sup>d</sup>	27.39 (±0.38) <sup>e</sup>	6.30 (±0.24) <sup>ef</sup>	36.51 (±0.72) <sup>d</sup>	10.34 (±0.18) <sup>d</sup>	36.51 (±0.72) <sup>d</sup>	10.34 (±0.18) <sup>d</sup>
<i>M. incognita</i> + <i>P. fluorescens</i>	54.08 (±0.37) <sup>de</sup>	173.30 (±9.74) <sup>e</sup>	66.87 (±0.18) <sup>cd</sup>	204.51 (±0.31) <sup>d</sup>	27.82 (±0.37) <sup>e</sup>	94.34 (±0.70) <sup>c</sup>	27.82 (±0.37) <sup>e</sup>	6.81 (±0.26) <sup>de</sup>	36.03 (±0.19) <sup>d</sup>	10.93 (±0.32) <sup>cd</sup>	36.03 (±0.19) <sup>d</sup>	10.93 (±0.32) <sup>cd</sup>
<i>M. incognita</i> only	43.21 (±0.46) <sup>f</sup>	144.52 (±0.59) <sup>g</sup>	68.61 (±0.32) <sup>b</sup>	210.03 (±0.46) <sup>a</sup>	21.57 (±0.61) <sup>f</sup>	97.71 (±0.2) <sup>ab</sup>	21.57 (±0.61) <sup>f</sup>	5.75 (±0.02) <sup>f</sup>	40.04 (±0.22) <sup>ab</sup>	11.92 (±0.19) <sup>b</sup>	40.04 (±0.22) <sup>ab</sup>	11.92 (±0.19) <sup>b</sup>
Healthy control	66.48 (±0.33) <sup>a</sup>	200.70 (±0.52) <sup>a</sup>	69.84 (±0.22) <sup>a</sup>	210.34 (±0.59) <sup>a</sup>	37.36 (±0.32) <sup>a</sup>	98.70 (±0.32) <sup>a</sup>	37.36 (±0.32) <sup>a</sup>	10.50 (±0.31) <sup>a</sup>	40.96 (±0.199) <sup>a</sup>	12.83 (±0.14) <sup>a</sup>	40.96 (±0.199) <sup>a</sup>	12.83 (±0.14) <sup>a</sup>
SEM±	0.55	0.69	0.32	0.44	0.46	0.69	0.46	0.17	0.42	0.18	0.42	0.18
CD @ 1%	1.81	2.02	1.39	1.62	1.65	2.03	1.65	1.01	1.59	1.05	1.59	1.05

**Table 2.** Effect of *Bacillus* spp. on egg masses and number of galls.

Treatments	Susceptible variety				Resistant variety			
	No. of galls per root system		Root-knot Index		No. of galls per root system		Root-knot index	
	Per cent reduction over control	No. of egg masses per root system	Per cent reduction over control	No. of egg masses per root system	Per cent reduction over control	No. of egg masses per root system	Per cent reduction over control	No. of egg masses per root system
<i>M. incognita</i> +Carbofuran	14.00 (±0.57) <sup>c</sup>	20.67 <sup>c</sup>	78.16	1.67 (±0.33) <sup>ab</sup>	2	1	50	1
<i>M. incognita</i> + <i>B. pumilus</i>	23.33 (±0.33) <sup>d</sup>	24.33 <sup>d</sup>	74.30	1 (±0.57) <sup>b</sup>	1	1	50	1
<i>M. incognita</i> + <i>B. megaterium</i>	25.33 (±0.33) <sup>c</sup>	27.33 <sup>c</sup>	71.13	1.33 (±0.33) <sup>ab</sup>	2	1.33	33.50	1.33
<i>M. incognita</i> + <i>B. subtilis</i>	28.33 (±0.33) <sup>b</sup>	31.33 <sup>b</sup>	66.90	1 (±0) <sup>b</sup>	1	0.67	66.50	0.67
<i>M. incognita</i> + <i>P. fluorescens</i>	27.33 (±0.33) <sup>b</sup>	30.00 <sup>b</sup>	68.31	1 (±0) <sup>b</sup>	1	0.67	66.50	0.67
<i>M. incognita</i> only	103.00 (±0.57) <sup>a</sup>	94.66 <sup>a</sup>	-	2 (±0) <sup>a</sup>	2	2	-	2
Healthy control	0.00 (±0) <sup>f</sup>	0.00 <sup>f</sup>	-	0.00 (±0) <sup>c</sup>	1	0	-	0
SEM±	0.47	0.66	-	0.23	0.23	NS	-	NS
CD @ 1%	1.67	1.98	-	1.18	1.18	NS	-	NS

enzymatic activity was higher in resistant variety (Figure 2, A).

The bacterial bio-agents treated-*Meloidogyne incognita* infested plants showed the highest enzymatic activity (Susceptible: 0.697-0.793 and Resistant: 2.537-2.056  $\text{abs min}^{-1} \text{g}^{-1}$ ) compared to all other treatments in the susceptible as well as resistant varieties. Among them, *Bacillus pumilus* treated plants (Susceptible: 0.793 and Resistant: 2.537  $\text{abs min}^{-1} \text{g}^{-1}$ ) showed significantly supreme activity of peroxidase, followed by *Pseudomonas fluorescens* (Susceptible: 0.733 and Resistant: 2.482  $\text{abs min}^{-1} \text{g}^{-1}$ ) and *B. subtilis* (susceptible: 0.730 and resistant: 2.125  $\text{abs min}^{-1} \text{g}^{-1}$ ).

### Studies on Polyphenol Oxidase Estimation (PPO) Activity

The polyphenol oxidase activity was assayed spectrophotometrically at 450 nm by following the procedure given by Selvaraj and Kumar (1995). The activity of polyphenol oxidase was increased upon bacterial bio-agents treatment to *Meloidogyne incognita* inoculated susceptible and resistant tomato varieties (0.076 to 0.094 and 0.171 to 0.216  $\text{abs min}^{-1} \text{g}^{-1}$ , respectively) compared to the respective *M. incognita* infested plants (0.060 and 0.108  $\text{abs min}^{-1} \text{g}^{-1}$ ), however, enzymatic activity was higher in resistant variety compared to the susceptible variety (Figure 2, B).

Among bacterial bio-agents treated-*Meloidogyne incognita* infested plants, *Bacillus pumilus* treated plants showed the highest activity (Susceptible: 0.094 and Resistant: 0.216 changes in absorbance  $\text{min}^{-1} \text{g}^{-1}$ ) and it was on par with *Pseudomonas fluorescens* (Susceptible: 0.090 and Resistant: 0.207 changes in absorbance  $\text{min}^{-1} \text{g}^{-1}$ ).

### Studies on Phenylalanine Ammonia Lyase (PAL) Activity

The polyphenol oxidase activity was assayed according to the procedure given by Ross and Senderoff (1992)

spectrophotometrically at 290 nm. There was an increased enzymatic activity in *Meloidogyne incognita* infested plants (Susceptible: 0.337 and Resistant: 0.741  $\mu\text{M trans-cinnamic acid min}^{-1} \text{g}^{-1}$ ) compared to the healthy plants (0.147 and 0.323  $\mu\text{M trans-cinnamic acid min}^{-1} \text{g}^{-1}$ , respectively). However, the activity was significantly higher in bacterized-*M. incognita* infested plants.

The susceptible plants treated with *Bacillus pumilus* along with *Meloidogyne incognita* (0.510  $\mu\text{M trans-cinnamic acid min}^{-1} \text{g}^{-1}$ ) showed higher activity of phenylalanine ammonia lyase, followed by *Pseudomonas fluorescens* treated- *M. incognita* infested susceptible plants (0.503  $\mu\text{M trans-cinnamic acid min}^{-1} \text{g}^{-1}$ ). Resistant plants followed a similar course (*B. pumilus*: 1.275 and *P. fluorescens*: 1.156  $\mu\text{M trans-cinnamic acid min}^{-1} \text{g}^{-1}$ ), but they were on par with each other (Figure 2, C).

### Studies on Catalase (CAT) Activity

The activity of catalase was assayed spectrophotometrically at 240 nm by following the procedure given by Masia (1998). The increased activity of catalase was recorded in *Meloidogyne incognita* inoculated samples of susceptible and resistant tomato varieties (32.01 and 21.340  $\mu\text{g H}_2\text{O}_2 \text{g}^{-1}$ , respectively) compared to the respective healthy plants (24.964 and 16.643  $\mu\text{g H}_2\text{O}_2 \text{g}^{-1}$ ), but enzymatic activity was higher in susceptible variety (Figure 3, A). The activity of catalase significantly decreased in bacterized nematode challenged tomato varieties (Susceptible: 15.251- 18.614 and Resistant: 9.500-11.740  $\mu\text{g H}_2\text{O}_2 \text{g}^{-1}$ ).


Among bacterial bio-agents, *Bacillus pumilus* treated plants showed the least catalase activity (Susceptible: 15.251 and Resistant: 9.500  $\mu\text{g H}_2\text{O}_2 \text{g}^{-1}$ ), followed by *Pseudomonas fluorescens* (Susceptible: 16.754 and Resistant: 10.503  $\mu\text{g H}_2\text{O}_2 \text{g}^{-1}$ ) and both were on par and no significant difference was found.



**Table 3.** Effect of bacterial bio-agents on final nematode population.

Treatments	Susceptible variety				Resistant variety			
	Soil (200 cc)	Per cent reduction over control	Root (5 g)	Per cent reduction over control	Soil (200 cc)	Per cent reduction over control	Root (5 g)	Per cent reduction over control
T <sub>1</sub> : <i>Meloidogyne incognita</i> +Carbofuran 3G (Positive control)	253.33 <sup>f</sup>	35.86	50.00 <sup>e</sup>	64.36	15.00 <sup>e</sup>	16.67	5.33 <sup>e</sup>	33.37
T <sub>2</sub> : <i>M. incognita</i> + <i>Bacillus pumilus</i>	268.33 <sup>e</sup>	32.06	54.33 <sup>d</sup>	61.28	15.67 <sup>a</sup>	12.94	6.67 <sup>ab</sup>	16.62
T <sub>3</sub> : <i>M. incognita</i> + <i>B. megaterium</i>	269.67 <sup>d</sup>	31.72	57.33 <sup>c</sup>	59.14	16.33 <sup>b</sup>	9.27	7.00 <sup>ab</sup>	12.50
T <sub>4</sub> : <i>M. incognita</i> + <i>B. subtilis</i>	273.33 <sup>b</sup>	30.80	61.33 <sup>b</sup>	56.29	17.67 <sup>bc</sup>	1.83	7.67 <sup>b</sup>	4.12
T <sub>5</sub> : <i>M. incognita</i> + <i>Pseudomonas fluorescens</i>	271.67 <sup>c</sup>	31.22	61.67 <sup>b</sup>	56.05	16.00 <sup>bc</sup>	11.11	7.33 <sup>ab</sup>	8.37
T <sub>6</sub> : Control with <i>M. incognita</i> inoculation	395 <sup>a</sup>	-	140.33 <sup>a</sup>	-	18.00 <sup>a</sup>	-	8.00 <sup>a</sup>	-
T <sub>7</sub> : Absolute control without <i>M. incognita</i> inoculation	0 <sup>g</sup>	-	0 <sup>f</sup>	-	0 <sup>d</sup>	-	0 <sup>d</sup>	-
<b>SEM±</b>	<b>0.52</b>		<b>0.38</b>		<b>0.57</b>		<b>0.47</b>	
<b>CD @ 1%</b>	<b>1.75</b>		<b>1.50</b>		<b>1.83</b>		<b>1.67</b>	





**Figure 1.** Effect of *Bacillus pumilus* galling of susceptible (a) and resistant (b) tomato plants infested with *Meloidogyne incognita*.

### Studies on Total Phenols


Total phenols were estimated by spectrophotometric method using Folin Ciocalteu Reagent (FCR) at an absorbance of 700 nm. Higher accumulation of phenol was observed in bacterial-bio-agents treated plants than untreated plants. In general, the inoculation of *Meloidogyne incognita* and bacterial-bio-agents simultaneously recorded significantly higher phenol (Susceptible: 65.987-68.593 and Resistant: 79.184-82.311 mg gallic acid equivalent 100 g<sup>-1</sup> FW) content than the control with *M. incognita* inoculation (Susceptible: 48.747 and Resistant: 58.496 mg gallic acid equivalent 100 g<sup>-1</sup> FW).

Among the bacterial bio-agents treated-*Meloidogyne incognita* infested tomato plants, the plants treated with *Bacillus pumilus* showed the highest phenol content (Susceptible: 68.593 and Resistant: 82.311


mg gallic acid equivalent 100 g<sup>-1</sup> FW) and it was on par with *Pseudomonas fluorescens* (Susceptible: 67.413 and Resistant: 80.895 mg gallic acid equivalent 100 g<sup>-1</sup> FW) (Figure 3, B).

### DISCUSSION

The influence of different *Bacillus* spp. on the physiology and biochemical status of *Meloidogyne incognita*-infested tomato plants was investigated. The significant increment in the shoot and root growth of tomato plants proves that the biocontrol agents do act as plant growth-promoting agents. From the above observations, it is evident that bacterial bio-agents have a positive effect on root and shoot growth. This may be attributed to one or more of the following factors; production of phosphatases by *Bacillus* spp. facilitates the conversion of insoluble phosphorus to



**Figure 2.** Status of bio-chemicals incited by bacterial bio-agents on *Meloidogyne incognita*-infested tomato plants (A) Peroxidase (PO), (B) Polyphenol Oxidase (PPO), and (C) Phenylalanine Ammonia-Lyase (PAL) activities. **Note:** T<sub>1</sub>= *M. incognita*+carbofuran 3G, T<sub>2</sub>= *M. incognita*+*B. pumilus*, T<sub>3</sub>= *M. incognita*+*B. megaterium*, T<sub>4</sub>= *M. incognita*+*B. subtilis*, T<sub>5</sub>= *M. incognita*+*P. fluorescens*, T<sub>6</sub>= *M. incognita* only and T<sub>7</sub>= Healthy control.



**Figure 3.** Status of bio-chemicals incited by bacterial bio-agents on *Meloidogyne incognita*-infested tomato plants (A) Catalase (CAT) activity and (B) Total phenols content. **Note:** T<sub>1</sub>= *M. incognita*+carbofuran 3G, T<sub>2</sub>= *M. incognita*+*B. pumilus*, T<sub>3</sub>= *M. incognita*+*B. megaterium*, T<sub>4</sub>= *M. incognita*+*B. subtilis*, T<sub>5</sub>= *M. incognita*+*P. fluorescens*, T<sub>6</sub>= *M. incognita* only, and T<sub>7</sub>= Healthy control.

available one for the use of plants (Abdelmoteleb and Gonzalez-Mendoza, 2020), production of growth-promoting phytohormones viz., Indole Acetic Acid (IAA), Gibberellic Acid (GA), cytokine (Calvo *et al.*, 2010), improvement of water and nutrients uptake, production of antibiotic metabolites effective against soil-borne pathogens and production of B-group vitamins that promote rooting capacity and affect the population of the microbial community (Wu *et al.*, 2005; Rai, 2006).

Furthermore, the reduced galling in bacterial bio-agents treated plants might be owing to the ability of the bio-agents to modify root exudates, thus hindering the feeding site recognition by the nematodes in the soil (Siddiqui and Mahmood, 1999; Zhou *et al.*, 2019). In addition, the bacterial bio-agents produce nematocidal acids that decrease the nematode population in the soil and reduce the level of infection and, hence, the galling of roots (Iatsenko *et al.*, 2014; Lee and Kim, 2016).

However, when nematode-susceptible and resistant varieties are compared, the average number of galls formed per root system and root-knot index were less in the case of the resistant variety. These results were adjacent to the results obtained by Kumari *et al.* (2016), who made a comparative study on *M. graminicola* susceptible (Pusa 1121) and resistant (Vandana) cultivars of rice. It was found that, after 15 dpi, all the growth parameters of nematode were recorded low in the resistant variety and were significantly different from the susceptible variety. However, while comparing the response of susceptible versus resistant varieties upon infection with *M. incognita*, many contradictions arose, probably due to the nematode effects (nematode biology varies with resistant and susceptible hosts), systemic hormone signalling effects or tissue-specific differential expression of selected genes (Cabasan *et al.*, 2012).

The bio-agent activity was clearly noticed in susceptible plants infested with *M. incognita*. The genetic background of the resistant variety (Hisar Lalit) contains *Mi 1.2*

gene, which shows resistance to *M. incognita* (Reddy *et al.*, 2018) and this might be the reason for variations observed in nematode infection in the resistant genotypes, and it might attract nematodes meagerly less than susceptible ones (Peacock, 1959). Because of the ability of a resistant variety to overcome nematode infection on its own, the treatments were established to be non-significant in many of the nematode growth parameters.

Studies on biochemical levels suggest that defence-related enzymes (peroxidase, phenylalanine ammonia-lyase, polyphenol oxidase and catalase) and biomolecules (phenols) get enhanced when the plants were subjected to biotic stress and abiotic stress. The above observations on biochemical studies revealed that there was a significant accumulation of all the tested enzymes in bacterized tomato plants (susceptible and resistant) challenged with *Meloidogyne incognita*. Among different *Bacillus* spp., *B. pumilus* treated-*M. incognita* infested plants showed supreme activity of bio-chemicals, followed by *B. subtilis* in susceptible as well as in resistant varieties compared to *Pseudomonas fluorescence* (standard check).

When compared to the susceptible and resistant varieties, the activities of enzymes were superior in the resistant variety (except catalase), probably due to avoidance of nematode invasion (Rao *et al.*, 2017), which was because of its early and enhanced accumulation of enzymes in response to invading pathogens. Enhanced levels of defence enzymes and phenolic content in plants treated with bio-agents may contribute to resistance development, safeguarding the plants without causing harm.

The present investigation revealed the potentiality of *Bacillus* as a bio-agent that could reduce gall formation, root-knot index and nematode population, thus improving the shoot and root growth under pot experiments. *Bacillus* spp. could be included in integrated disease management strategies to reduce the damage caused by *M. incognita* in tomatoes. These findings



revealed that *B. pumilus* was the most potential *Bacillus* bio-agent in reducing *M. incognita* infection by embellishing the plant growth and strengthening the plants by enhancing defence-related enzymes, followed by *B. subtilis* and *B. megaterium*.

## ACKNOWLEDGEMENTS

The authors are thankful to the ICAR-Indian Institute of Horticultural Research, Bengaluru for assisting to carry out the research. They also acknowledge the support from the Head, AICRP (Nematodes), Department of Plant Pathology, University of Agricultural Sciences, GKVK, Bengaluru for the necessary facilities to carry out this work smoothly.

## REFERENCES

1. Abdelmoteleb, A. and Gonzalez-Mendoza, D. 2020. Isolation and Identification of Phosphate Solubilizing *Bacillus* spp. from *Tamarix ramosissima* Rhizosphere and Their Effect on Growth of *Phaseolus vulgaris* under Salinity Stress. *Geomicrobiol. J.*, **37(10)**: 901-908.
2. Anita, B., Rajendran, G. and Samiyappan, R. 2004. Induction of Systemic Resistance in Tomato against Root-Knot Nematode, *Meloidogyne incognita* by *Pseudomonas fluorescens*. *Nematol. Mediterr.*, **32**: 47-51.
3. Ayoub, S. M. 1977. *Plant Nematology- An Agricultural Training Aid*. Laboratory Services--Nematology, Division of Plant Industry, Department of Food and Agriculture, State of California, 157 PP.
4. Back, M. A., Haydock, P. P. J. and Jenkinson, P. 2002. Disease Complexes Involving Plant Parasitic Nematodes and Soil Borne Pathogens. *Plant Pathol.*, **51(6)**: 683-697.
5. Cabasan, M. T. N., Kumar, A. and De-Waele, D. 2012. Comparison of Migration, Penetration, Development and Reproduction of *Meloidogyne graminicola* on Susceptible and Resistant Rice Genotypes. *Nematology*, **14(4)**: 405-415.
6. Calvo, P., Ormeno-Orrillo, E., Martinez-Romero, E. and Zuniga, D. 2010. Characterization of *Bacillus* Isolates of Potato Rhizosphere from Andean Soils of Peru and Their Potential PGPR Characteristics. *Braz. J. Microbiol.*, **41**: 899-906.
7. Chander, M. S. 1990. Enzymic Associations with Resistance to Rust and Powdery Mildew in Peas. *Indian J. Hort.*, **47(3)**: 341-345.
8. Chitwood, B. G. 1949. 'Root-Knot Nematodes'. Part I. A Revision of the Genus *Meloidogyne* Goeldi, 1887. *Proc. Helminthol. Soc. Wash.*, **16(2)**: 90-114.
9. Collange, B., Navarrete, M., Peyre, G., Mateille, T. and Tchamitchian, M. 2011. Root-Knot Nematode (*Meloidogyne*) Management in Vegetable Crop Production: The Challenge of an Agronomic System Analysis. *Crop. Prot.*, **30(10)**: 1251-1262.
10. Elling, A. A. 2013. Major Emerging Problems with Minor *Meloidogyne* Species. *Phytopathol.*, **103(11)**: 1092-1102.
11. Hallmann, J., Davies, K. G. and Sikora, R. 2009. Biological Control Using Microbial Pathogens, Endophytes and Antagonists. In: "Root-Knot Nematodes". CABI, Wallingford, UK, PP. 380-411.
12. Hou, S., Liu, Z., Hen, H. and Wu, D. 2019. Damage-Associated Molecular Pattern-Triggered Immunity in Plants. *Front. Plant Sci.*, **10**: 646.
13. Iatsenko, I., Boichenko, I. and Sommer, R. J. 2014. *Bacillus thuringiensis* DB27 Produces Two Novel Protoxins, Cry21Fa1 and Cry21Ha1, Which Act Synergistically against Nematodes. *Appl. Environ. Microbiol.*, **80(10)**: 3266-3275.
14. Koenning, S. R., Overstreet, C., Noling, J. W., Donald, P. A., Becker, J. O. and Fortnum, B. A. 1999. Survey of Crop Losses in Response to Phytoparasitic Nematodes in the United States for 1994. *J. Nematol.*, **31(4S)**: 587.
15. Kumari, C., Dutta, T. K., Banakar, P. and Rao, U. 2016. Comparing the Defense-Related Gene Expression Changes upon Root-Knot Nematode Attack in Susceptible versus Resistant Cultivars of Rice. *Sci. Rep.*, **6(1)**: 1-13.
16. Lee, Y. S. and Kim, K. Y. 2016. Antagonistic Potential of *Bacillus pumilus* L1 against Root-Knot Nematode, *Meloidogyne arenaria*. *J. Phytopathol.*, **164(1)**: 29-39.

17. Masia, A., 1998. Superoxide Dismutase and Catalase Activities in Apple Fruit during Ripening and Post-Harvest and with Special Reference to Ethylene. *Physiol. Plant.*, **104**(4): 668-672.
18. Narasimhamurthy, H. B., Ravindra, H. and Mukesh Sehgal, R. N., 2017. GIS/GPS Based Survey on Incidence and Distribution of Rice Root-Knot Nematode (*Meloidogyne graminicola*) in Southern Transition Zone of Karnataka. *J. Entomol. Zool. Stud.*, **5**(2): 410-413.
19. Ogwulumba, S. I., Ugwuoke, K. I. and Ogbuji, R. O. 2011. Reaction of Tomato cv. Roma vf (*Solanum lycopersicum*) to *Meloidogyne javanica* treub Infestation in an Ultisol Treated with Aqueous Leaf Extracts of Bitter Leaf (*Vernonia Amygdalina* L.) and Mango (*Mangifera Indica* L.). *J. Plant Prot. Res.*, **51**(1): 1-4.
20. Peacock, F. C. 1959. The Development of a Technique for Studying the Host/Parasite Relationship of the Root-Knot Nematode *Meloidogyne incognita* under Controlled Conditions. *Nematologica*, **4**(1): 43-55.
21. Poveda, J., Abril-Urias, P. and Escobar, C. 2020. Biological Control of Plant-Parasitic Nematodes by Filamentous Fungi Inducers of Resistance: Trichoderma, Mycorrhizal and Endophytic Fungi. *Front. Microbiol.*, **11**:530260.
22. Prabu, P., Umamaheswari, R. and Rao, M. S. 2019. Deciphering the Biocontrol Mechanism of *Bacillus amyloliquefaciens* IIHR BA2: Detection of Nematicidal, Fungicidal and Bactericidal Lipopeptides. *Asian J. Microbiol. Biotechnol. Environ. Sci.*, **21**(3): 731-739.
23. Radwan, M. A., Farrag, S. A. A., Abu-Elamayem, M. M. and Ahmed, N. S. 2012. Biological Control of the Root-Knot Nematode, *Meloidogyne incognita* on Tomato Using Bioproducts of Microbial Origin. *Appl. Soil Ecol.*, **56**: 58-62.
24. Rai, M. K. 2006, Handbook of Microbial Biofertilizers. Food Products Press, an Imprint of The Haworth Press, Inc, Binghamton, New York, **17**: 137-170.
25. Ramirez-Prado, J. S., Abulfaraj, A. A., Rayapuram, N., Benhamed, M. and Hirt, H., 2018. Plant Immunity: From Signaling to Epigenetic Control of Defense. *Trends Plant Sci.*, **23**(9): 833-844.
26. Rao, M. S., Umamaheswari, R., Prabu, P., Kamalnath, M., Grace, G. N. and Chaya, M. K. 2017. Induction of Systemic Resistance in Bell Pepper (*Capsicum annum* L.) Mediated through *Bacillus subtilis* IIHR BS-2 against Root-Knot Nematode, *Meloidogyne incognita*. *Pest Manage. Horti. Ecosyst.*, **23**(1): 64-71.
27. Reddy, Y. S., Sellaperumal, C., Prasanna, H. C., Yadav, A., Kashyap, S. P., Singh, S., Rai, N., Singh, M. and Singh, B. 2018. Screening of Tomato Genotypes against Root-Knot Nematode and Validation of *Mi 1* Gene-Linked Markers. *Proc. Natl. Acad. Sci. India Sect. B Biol. Sci.*, **88**: 65-72.
28. Ross, W. W. and Senderoff, R. R. 1992. Phenylalanine Ammonia Lyase from Loblolly Pine; Purification of the Enzyme and Isolation of Complementary DNA Clones. *Plant Physiol.*, **98**: 380-386.
29. Selvaraj, Y. and Kumar, R. 1995. Enzymatic Regulation in Ripening Mango Fruit. *Indian J. Hortic.*, **51**(4): 316-323.
30. Siddiqui, Z. A. and Mahmood, I. 1999. Role of Bacteria in the Management of Plant Parasitic Nematodes: a review. *Bioresour. Technol.*, **69**(2): 167-179.
31. Singleton, V. L. and Rossi, J. A. 1965. Colorimetry of Total Phenolics with Phosphomolybdic-Phosphotungstic Acid Reagents. *Am. J. Enol. Vitic.*, **16**(3): 144-158.
32. Whetten, R.W. and Sederoff, R. R. 1992. Phenylalanine Ammonia-Lyase from Loblolly Pine: Purification of the Enzyme and Isolation of Complementary DNA Clones. *Plant Physiol.*, **98**(1): 380-386.
33. Wu, S. C., Cao, Z. H., Li, Z. G., Cheung, K. C. and Wong, M. H. 2005. Effects of Biofertilizer Containing N-Fixer, P and K Solubilizers and AM Fungi on Maize Growth: A Greenhouse Trial. *Geoderma*, **125**(1-2): 155-166.
34. Zhou, D., Feng, H., Schuelke, T., De-Santiago, A., Zhang, Q., Zhang, J., Luo, C. and Wei, L. 2019. Rhizosphere Microbiomes from Root-Knot Nematode Non-Infested Plants Suppress Nematode Infection. *Microb. Ecol.*, **78**: 470-481.



## ارزیابی گونه‌های باسیلوس به عنوان عوامل کنترل زیستی علیه *Meloidogyne incognita* آلوده کننده گوجه فرنگی

بومناهایی ساتیش کاویا، تیتور روپلا-نایاک کاویتا، آراتی، و ناگاوات لوهیت کومار

### چکیده

گوجه‌فرنگی میزبان جهانی نماتد گره ریشه (*Meloidogyne incognita*) است که خسارت قابل توجهی ایجاد می‌کند. استفاده از عوامل کنترل زیستی (bio-control agents) برای کنترل نماتدهای انگلی گیاهی، رویکردی سبز و جافته، پیش از استفاده از نماتدکشی‌های مصنوعی است. نقش گونه‌های باسیلوس در تحریک تغییرات فیزیولوژیکی و بیوشیمیایی در آلودگی نماتد در این مطالعه بحث شده است. ارقام حساس (PKM-1) و مقاوم (Hisar Lalit) گوجه‌فرنگی که با باسیلوس پومیلوس تیمار شدند، طول ساقه، طول ریشه و زیست‌توده گیاهان را در مقایسه با شاهد استاندارد، *Pseudomonas fluorescens* و پس از آن *B. megaterium* افزایش دادند. بر این اساس، تمام گیاهان حساس (susceptible) تحت درمان با عامل کنترل زیستی، کاهش گال‌زایی و شاخص گال ریشه ۳ (نسبتاً مقاوم) را نشان دادند و جمعیت نماتد را در خاک و ریشه‌ها کاهش دادند. برعکس، تمام گیاهان مقاوم واکنش‌های بسیار مقاومی نشان دادند. *B. pumilus* بالاترین بیان را در تمام آنزیم‌های بیوشیمیایی مانند پراکسیداز (PO)، پلی‌فنل اکسیداز (PPO)، کاتالاز (CAT)، فنیل‌آلانین آمونیالیز (PAL) و فنل‌های کل نشان داد. در نهایت، مشخص شد که *B. pumilus* با بهبود رشد گیاه و افزایش آنزیم‌های مرتبط با دفاع در گوجه‌فرنگی، بیشترین پتانسیل را در کاهش آلودگی نماتد دارد.