

Effect of Canola Physical Mutation on *Plutella xylostella* (L.) Life Table

M. Akandeh¹, F. Kocheili¹, M. Soufbaf^{2*}, A. Rasekh¹, and K. Mozafari²

ABSTRACT

Life history of insect herbivores is unclear after feeding on mutant lines of many crops. To shed light, demographic parameters of *Plutella xylostella* on three canola cultivars ("Zar", "RGS", "Talaye") and their physical mutation-derived lines ("Zar 9-9", "RGS 8-1", "RGS 10-2", "RGS 8-13" and "Talaye 8-3") were determined under greenhouse condition. Methods of life table including the female age specific life table and the age-stage, two-sex life table were applied. According to two-sex life table, there was no significant difference between demographic parameters of *P. xylostella* on "Talaye" and "Zar" in comparison with their mutant lines, but significant differences were observed between these parameters on mutant lines of "RGS". Having suitable cultural traits, "RGS 8-1" was more susceptible than the other two mutant lines and its control cultivar "RGS" in terms of population growth of the pest. According to the two-sex life table, the net reproductive rate (R_0), intrinsic rate of increase (r), and finite rate of increase (λ) were the highest on RGS 8-1 (98.63 offspring per individual, 0.208 and 1.231 day⁻¹, respectively). Also, population projection showed the rapid growth of the pest on the latter line. There was a little difference between the same population parameters estimated by two methods of life table. Investigating some consequences of plant breeding using radiation techniques on insect fitness not only leads plant breeders to do more unfailing selections but also provides some enlightenment in pest management programs effectively when plantation of such crops is prioritized.

Keywords: Age-stage two-sex life table, Diamondback moth, Female age-specific life table, Gamma ray, Insect fitness.

INTRODUCTION

Canola, *Brassica napus* L., is an important economic oilseed crop highly impacted by one of its important pests, diamondback moth, *Plutella xylostella* L. (Lep.: Plutellidae) (Furlong *et al.*, 2013) that causes more than 90% crop loss annually (Talekar and Shelton, 1993). Mutation breeding is an important branch of plant breeding that has been reported to improve self-pollinated crops such as canola. Producing plant varieties through this technique causes random, multiple genetic

modifications in plants (Novak and Brunner, 1992). These changes can be within the gene, losses or additions of genes or groups of genes and, probably, changes in the relations of genes to each other (Ahloowalia and Maluszynski, 2001). Moreover, interaction of gamma rays with atoms or molecules of plants may produce free radicals in their cells; thereby cellular structure of plants and their metabolism are impacted depending on the irradiation level (Dhanavel *et al.*, 2012). The effects of all these alterations on plant quality and higher trophic levels such as herbivores and their natural enemies, are unknown. To discover

¹ Department of Plant Protection, College of Agriculture, Shahid Chamran University of Ahvaz, Ahvaz, Islamic Republic of Iran.

*Corresponding author; e-mail: msoufbaf@nrcam.org

² Agricultural, Medical and Industrial Research School, Karaj, P. O. Box: 31485/498, Islamic Republic of Iran.



the effect of host plant quality on survival rate and reproduction of an arthropod population, calculation of demographic parameters can be applied as an important tool (Southwood and Henderson, 2000). Accordingly, evaluation of host plants antibiotic resistance to pest insects will be conceivable (Maia *et al.*, 2000).

Chi and Liu (1985) pointed out that the traditional age-specific life-table does not consider male populations, the stage differentiation and the variable developmental rate among individuals. Disregarding these items in the calculations of demographic parameters may cause errors. These errors are related to errors in calculating the l_x and m_x (Huang and Chi, 2011). In other words, it is generally assumed that all female adults emerge at the same age. If m_x and l_x are constructed based on adult age and female individuals, they may also cause errors in population parameters (Chi and Su, 2006; Huang and Chi, 2011). Therefore, they constructed an age-stage, two-sex life table to consider the attributes in question.

To highlight the potential effects of physical mutation on fitness of insect pests, we aimed to study demographic parameters of *P. xylostella* as a model insect on some mutant lines of canola ("Zar 9-9", "RGS 8-1", "RGS 10-2", "RGS 8-13" and "Talaye 8-3") contrasting to their control cultivars ("Zar", "RGS", "Talaye"), and to analyze the data using age-stage, two-sex life table and the female age-specific life table procedures.

MATERIALS AND METHODS

Modification of Seeds by Physical Mutation

Gamma irradiation of canola seeds was carried out at Nuclear Science and Technology Research Institute (NSTRI), Karaj, Iran. The mutant canola lines, "RGS 8-1" and "RGS 8-13" were produced from cultivar "RGS" seeds irradiated at 800 Gy.

"RGS 10-2" also resulted from the same cultivar but it was irradiated at 1000Gy. "Talaye 8-3" was obtained by irradiation of seeds from cultivar "Talaye" at 800Gy and "Zar 9-9" mutant line was derived from cultivar "Zar" seeds irradiated at 900 Gy. Physical mutation aimed at induction of high yield and early maturity in experimental varieties.

Plants and Insects Rearing

Seeds of canola cultivars and their mutant lines were sown in a 1:1:1 mix of peat moss: perlite: soil in 30 cm diameter plastic pots. Five-week-old plants were used in the present experiments. The colony of *P. xylostella* was established two months prior to initiation of the experiments with 30 adults collected from a cabbage field located in Karaj (35° 48' N, 51° 00' E), Iran. Colonies of *P. xylostella* were kept in ventilated cages (80×80×60 cm). Both plants and insects were reared under greenhouse condition (27±5°C, 65±10 RH, and a photoperiod of 16:8 L:D).

Demographic Parameters

All experiments were carried out under greenhouse conditions simultaneously. Thirty pairs of both sexes of *P. xylostella* were transferred from the colony to ventilated Plexiglas prepared as oviposition cages (28×23×15cm) for 24 hours. Synchronized eggs with longevity of less than 24 hours were glued to the upper surface of a leaf of the potted plant individually; then, each leaf containing an egg was confined with a clip cage (9×7×2.5 cm). Fifty two to 81 clip cages were used in each treatment. All experimental units (clip cages) were observed daily until adults emerged. Then, newly emerged females were coupled with males from the same cohort in ventilated Plexiglas cylindrical cages (10 cm in diameter and 9 cm in height). The fecundity of females, longevity

and mortality of both sexes was determined daily during the mature stage. Experiments were continued until all individuals died.

Statistical Analysis

Female Age-specific Life Table: Age-specific fecundity (m_x) and survivorship (l_x) were determined for *P. xylostella* cohorts on each host plant based on female individuals. The iterative bisection method and the Euler-Lotka Equation with the age indexed from 0 to maximum age (ω) were applied to estimate the intrinsic rate of natural increase (r).

$$\sum_{x=0}^{\omega} e^{-r(x+1)} l_x m_x = 1 \quad (1)$$

Population growth parameters (net Reproductive rate (R_0), mean generation Time (T), finite rate of increase (λ) and intrinsic rate of natural increase (r) were estimated based on female population according to Carey (1993) and Maia *et al.* (2000). Data on “RGS” and its mutant lines were analyzed with unbalanced one-way ANOVA (Proc GLM, SAS Institute 2003 (ver. 9.1) and their means were compared using the Tukey-Kramer procedure ($P < 0.05$) while the same values on two other cultivars in contrast with their mutant lines was compared using Student's *t*-test ($P < 0.05$) (Minitab Inc., 2007).

Age-stage, Two-sex Life Table: The life history data of *P. xylostella* including males, females and immature individuals that died before the adult stage were analyzed according to the age-stage, two-sex life table (Chi and Liu, 1985; Chi, 1988) using the TWOSEX-MSChart (Chi, 2015). The age-specific survivorship (l_x) including both male and female, the age-specific fecundity (m_x), and R_0 were calculated according to Chi and Liu (1985) as below:

$$l_x = \sum_{j=1}^{\beta} s_{xj} \quad (2)$$

$$m_x = \frac{\sum_{j=1}^{\beta} s_{xj} f_{xj}}{\sum_{j=1}^{\beta} s_{xj}} \quad (3)$$

$$R_0 = \sum_{x=0}^{\infty} \sum_{j=1}^{\beta} s_{xj} f_{xj} \quad (4)$$

Where, β is the number of stages. The intrinsic rate of increase was calculated by using the bisection method and Euler-Lotka Equation with age indexed from 0 (Goodman, 1982):

$$\sum_{x=0}^{\infty} e^{-r(x+1)} l_x m_x = 1 \quad (5)$$

Moreover, T and λ were calculated as:

$$T = \frac{\ln R_0}{r} \text{ and } \lambda^T = R_0, \text{ respectively.}$$

All parameters were estimated based on both sexes and the variable developmental rates among individuals (Naseri *et al.*, 2014; Safuraie-Parizi *et al.*, 2014; Mahmoudi *et al.*, 2015). The variability of life table parameters was estimated in bootstrap procedure ($B = 10,000$). The bootstrap values on “RGS” and its mutant lines were compared using the paired bootstrap test ($P < 0.05$). The same values on “Talaye” and “Zar” as opposed to their mutant lines were compared using *t*-test ($P < 0.05$).

The program TIMING-MSChart was used to project population growth (Chi 1990, 2015). For the projection, we assumed an unlimited growth with an initial population of 10 newly laid eggs of *P. xylostella*.

RESULTS

Female Age-Specific Life Table of *P. xylostella*

Survivorship (l_x) and fecundity (m_x) curves were demonstrated based on the female age-specific life table (Figure 1). The l_x schedule of the pest on all examined treatments showed a similar pattern in general. High mortality occurred during the first 10 days in periods of incubation and first-instar larvae. Age-specific survivorship of *P. xylostella* on “RGS 8-1”, “RGS 10-2”, “RGS 8-13”, “RGS”, “Talaye”, “Talaye 8-3”, “Zar” and “Zar 9-9” at the time of adult female emergence were 0.583, 0.471, 0.44,



0.6, 0.4, 0.39, 0.69 and 0.64, respectively. According to the curve of m_x , the oviposition period of females began on days 17, 21, 23, 18, 21, 22, 16, 19 on the above mentioned varieties, respectively. Also, the highest daily fecundity of *P. xylostella* on the above mentioned varieties was 22.53, 10.19, 14.46, 14.31, 10.62, 11, 18.06 and 14.38 female offspring, respectively (Figure 1).

Based on the demographic parameters obtained from female age-specific life table, there were significant differences between r values of *P. xylostella* on "RGS" and its mutant lines (Table 1). The cohort reared on "RGS 8-1" had the highest r ($F=39.73$; $df=3, 66$; $P<0.0001$), R_0 ($F=34.49$; $df=3, 65$; $P<0.001$) and λ values ($F=38.20$; $df=3, 65$; $P<0.0001$) and the lowest T value ($F=19.74$; $df=3, 66$; $P<0.0001$). Also, there was no significant difference between the parameters on cultivar "Talaye" in comparison with "Talaye 8-3" (Table 1). In addition, the R_0 value of *P. xylostella* on "Zar" was significantly higher ($T=-2.57$; $df=24$; $P=0.017$) compared to that on "Zar 9-9". Furthermore, there was no significant difference among other population parameters on "Zar 9-9" (Table 1).

Age-Stage, Two-Sex Life Table of *P. xylostella*

The age-stage specific survival rates (s_{xj}) of *P. xylostella* on different treatments are shown in Figure 2. The curves of s_{xj} demonstrated the survival rate, stage differentiation, and the variable developmental rate among individuals. The probability that a newborn egg survived to the adult female was 0.31, 0.21, 0.17, 0.26, 0.21, 0.11, 0.31 and 0.30 and the probability to the adult male was 0.22, 0.19, 0.19, 0.25, 0.16, 0.13, 0.29 and 0.28 on "RGS 8-1", "RGS 10-2", "RGS 8-13", "RGS", "Talaye", "Talaye 8-3", "Zar" and "Zar 9-9", respectively. The results revealed that the lowest survival rate of a newborn egg to female adult stage was related to *P. xylostella* reared on "Talaye 8-3" (Figure 2). The age-

Table 1. Population parameters (Means \pm SE) of *Plutella xylostella* on canola cultivars and their mutant lines under greenhouse conditions, estimated by two methods of life table.^a

Parameter	RGS (Control)	RGS 8-1 (Mutant line)	RGS 10-2 (Mutant line)	RGS 8-13 (Mutant line)	Talaye (Control)	Talaye 8-3 (Mutant line)	Zar (Control)	Zar 9-9 (mutant line)
r	0.1677 \pm 0.010 ^b 0.197 \pm 0.007 ^b	0.2077 \pm 0.01 ^a 0.245 \pm 0.006 ^a	0.1397 \pm 0.012 ^c 0.163 \pm 0.009 ^c	0.1317 \pm 0.009 ^c 0.158 \pm 0.004 ^c	0.1222 \pm 0.012 ^a 0.146 \pm 0.010 ^a	0.1230 \pm 0.013 ^a 0.143 \pm 0.009 ^a	0.1828 \pm 0.014 ^a 0.218 \pm 0.010 ^a	0.1622 \pm 0.009 ^a 0.196 \pm 0.006 ^a
R_0	71.52 \pm 14.90 ^{ab} 73.40 \pm 4.26 ^b	98.67 \pm 17.52 ^a 98.59 \pm 4.48 ^a	46.04 \pm 12.10 ^b 46.29 \pm 5.36 ^c	43.65 \pm 10.55 ^b 47.09 \pm 2.76 ^c	33.56 \pm 9.23 ^a 34.81 \pm 5.45 ^a	32.56 \pm 9.94 ^a 32.54 \pm 2.99 ^a	78.75 \pm 17.90 ^a 86.51 \pm 8.24 ^a	58.17 \pm 12.02 ^a 61.96 \pm 4.81 ^b
λ	1.182 \pm 0.0120 ^b 1.218 \pm 0.009 ^b	1.2308 \pm 0.012 ^a 1.278 \pm 0.008 ^a	1.150 \pm 0.014 ^c 1.177 \pm 0.010 ^c	1.140 \pm 0.010 ^c 1.171 \pm 0.005 ^c	1.130 \pm 0.0140 ^a 1.157 \pm 0.011 ^a	1.130 \pm 0.0162 ^a 1.153 \pm 0.011 ^a	1.200 \pm 0.0163 ^a 1.243 \pm 0.012 ^a	1.180 \pm 0.0113 ^a 1.216 \pm 0.007 ^a
T	25.47 \pm 0.718 ^b 21.78 \pm 0.65 ^{bc}	22.11 \pm 0.410 ^c 18.70 \pm 0.34 ^d	27.40 \pm 1.03 ^a 23.52 \pm 0.86 ^{bc}	28.66 \pm 0.646 ^a 24.28 \pm 0.56 ^a	28.75 \pm 1.081 ^a 24.37 \pm 0.85 ^a	28.31 \pm 1.35 ^a 24.34 \pm 1.28 ^a	23.89 \pm 1.02 ^a 20.49 \pm 0.68 ^a	25.05 \pm 0.564 ^a 21.06 \pm 0.44 ^a

^a Values in the first and second rows for each parameter are related to the age-stage two-sex life table and the age-specific female life table, respectively. Life table parameters obtained from two methods were compared separately. Values on each cultivar were compared with those on mutant lines of the same cultivar at $p=0.05$ level. Paired bootstrap test was used to compare the values on RGS and its mutant lines. Student's t-test was used to compare the values on Zar and Talaye in comparison with their mutant lines. Values obtained from two methods of life table were not compared statistically.




Figure 1. Age-specific survivorship (l_x) and fecundity (m_x) and age-specific maternity ($l_x m_x$) of *Plutella xylostella* on canola mutant lines and their controls, using female age-specific life table under greenhouse conditions.




Figure 2. Age-stage survival rate (s_{xj}) of *Plutella xylostella* on canola mutant lines and their controls under greenhouse conditions.

specific survivorship (l_x) and age-specific fecundity (m_x) and age-specific maternity ($l_x m_x$) of *P. xylostella* on mutant lines and their controls are shown in Figure 3. To calculate l_x , individuals of both sexes were considered. This curve is a simple form of the curves in Figure 2. Based on the curve of $f(x, \text{female})$, first oviposition of females on the above-mentioned varieties occurred at the age of 16.5, 19.5, 21.5, 17.5, 19.5, 20.5, 14.5 and 17.5 days, respectively. The peak of this curve depicting the highest daily fecundity of *P. xylostella* on the experimental varieties was 89.4, 59, 36.3, 38.7, 56, 37.7, 80 and 29.5 eggs, respectively. According to these results, the highest and the lowest daily fecundity was observed on “RGS 8-1” and “Talaye 8-3”, respectively (Figure 3). The age-stage reproductive value (v_{xj}) for a new egg (v_{01}) of *P. xylostella* on all mutant lines of canola and their controls is equal to the finite rate of increase (λ) on the respective varieties. The highest reproductive value of *P. xylostella* was at the age of 17.5, 20.5, 22.5, 21.5, 20.5, 21.5, 15.5 and 22.5 days on the aforementioned varieties, respectively. This indicates that, as opposed to other ages, female individuals of the above ages had the highest contribution to the future population on the experimental varieties, respectively (Figure 4).

Population parameters in this method of life table indicated a significant difference among population parameters of *P. xylostella* on cultivar “RGS” and its mutant lines (Table 1). The highest and lowest r , R_0 and λ values were obtained on “RGS 8-1” and “RGS 8-13”, respectively. Furthermore, the lowest and highest T values of *P. xylostella* were observed on two recent mentioned mutant lines, respectively. Moreover, no significant difference was observed among the population parameters of *P. xylostella* reared on “Zar” and “Talaye” in contrast with their mutant lines (Table 1).

Simulated population growth of *P. xylostella* fed on canola cultivars and their mutant lines demonstrated the best growth of population on mutant line, RGS 8-1 (Figure 5).

DISCUSSION

Induction of desired plant attributes that have been lost through evolution is possible via mutation. There are random, multiple genetic modifications in plants which are produced through this option (Novak and Brunner, 1992). There is a noticeable variation in the parameters, particularly in the intrinsic rate of natural increase (r), among different studies. This parameter is known as the most valued item to evaluate the suitability of different host plants to an insect (Soufbaf *et al.*, 2010; Goodarzi *et al.*, 2015). Our obtained r values for all three cultivars were much less than those reported by Soufbaf *et al.* (2010), but were more similar to the values obtained on different manipulated canola genotypes by Nikooei *et al.* (2015). Intrinsic rate of increase values obtained on canola by Golizadeh *et al.* (2009) were in the range of the current study. Also, our obtained R_0 values were higher than those reported by Soufbaf *et al.* (2010) and Nikooei *et al.* (2015). The differences among studies could be attributed to an array of allelochemicals activity in excised leaves (dead plant) and in those that are still attached to the host plant (living plant) (Soufbaf *et al.*, 2012). These dissimilarities may be explained by various qualities among host plants, genetic differences of the *P. xylostella* geographic populations, different rearing techniques, and differences in environmental factors (Awmack and Leather, 2002). This study revealed no significant difference between demographic parameters of *P. xylostella* on “Talaye” and “Zar” in comparison with those on their mutant lines but these parameters had significant difference on mutant lines of “RGS”. Intrinsic rate of increase, R_0 and λ of *P. xylostella* were the highest on “RGS 8-1”, “RGS” and “RGS 10-2”, “RGS 8-13”, respectively. Furthermore, simulated population projection of *P. xylostella* fed on canola cultivars and their mutant lines confirms the susceptibility of mutant line, RGS 8-1 to population growth




Figure 3. Age-specific survivorship (l_x), age-stage fecundity of female (f_{xj}) (offspring), age-specific fecundity (m_x) and age-specific maternity ($l_x m_x$) of *Plutella xylostella* on canola mutant lines and their controls using the age-stage, two-sex life table under greenhouse conditions.




Figure 4. Age-stage-specific reproductive value (v_{xj}) of *Plutella xylostella* on canola mutant lines and their controls under greenhouse conditions.




Figure 5. Simulated population growth of *Plutella xylostella* fed on on canola mutant lines and their controls under greenhouse conditions.

of the pest. Hence, having suitable cultural traits, "RGS 8-1" was more susceptible compared with the control to population growth of *P. xylostella* while the other two mutant lines i. e. "RGS 10-2" and "RGS 8-13" were more resistant to population growth of this pest. Similarly, Nikooei *et al.* (2015) by using survival rate and r values suggested that canola mutant genotype (RGS₀₀₃) was more resistant compared to its control cultivar. These results confirm findings of Ahloowalia and Maluszynski (2001), who stated that many factors such as species or genotypes of plants and irradiation doses can affect mutagenesis. In the debate upon two different methods of life table, Chi (1988) expressed that the variation in development rates, primary sex ratio, and pre-adult mortality can cause differences between the results of the age-stage two-sex life table and the female age-specific life table. In addition, this may be partly due to the difference between l_x and m_x in the two methods of life table. Also, Chi and Su (2006) discussed the possible problem of l_x and m_x based on adult age in female-based life table. Farhadi *et al.* (2011) showed that there was no significant difference between r , R_0 , λ and T calculated by either female age-specific life table or age-stage two-sex life table. Their results showed that the male individuals and the variable developmental rate had no obvious effects on the population growth. They stated that this result was achieved because of the sex ratio (close to 1:1) and the stage overlapping which was short in comparison to the entire life span. Similar to our results, Khanamani *et al.* (2013) used both methods of life table to analyze their data and showed little difference among the same life table parameters.

The achieved survival curves of two-sex and female-based life tables showed differences due to ignoring male individuals in l_x calculation in the female age-specific life table. Moreover, the daily fecundity is multiplied by the same sex ratio in calculating m_x in the method of female age-specific life table. Huang and Chi (2011)

discussed the problem of applying female age-specific life table to insect populations and illustrated that when the age-specific female life table is applied to an age-stage-structured two-sex population, both survival and fecundity curves will be improperly manipulated because the variation in pre-adult development time among individuals is disregarded. Moreover, in this method of life table, male survival curve could not be calculated and the stage overlap would not be shown because of the utilization of the mean developmental time of each stage to partition the life history into different stages and it would ultimately result in errors in the survival curves (Chi, 1988). Since the mean duration of pre-adult stages is always greater than the first adult emergence, the m_x curve in female age specific life table starts later than the true one (Chi, 1988; Huang and Chi, 2011). Furthermore, if l_x and m_x are constructed based on adult age in age-specific female life table, they may also cause errors in population parameters (Chi and Su, 2006; Huang and Chi, 2011). Regarding the differences between the two methods of life table, it seems that age-stage, two-sex life table provides more data on the life history of pests theoretically. Although the method of female age specific life table is still applied in various studies, its application in population projection and biological control will be limited, because female life table cannot take male population and stage differentiation into consideration.

CONCLUSIONS

The present study showed that *P. xylostella* can develop on mutant lines of three commonly grown canola cultivars. It revealed that population parameters of *P. xylostella* were influenced by mutant varieties but mutation impact was dependent not only on plant cultivar but also different lines of one cultivar. The differences between the two methods of life table can be related to factors such as male population,



pre-adult mortality and variable developmental rate among individuals in age-stage, two-sex life table method. Besides, our results may provide considerable information to design a program for integrated management of *P. xylostella* in mutant canola fields. Nevertheless, related field studies should be conducted in future to attain more practical information successfully.

ACKNOWLEDGEMENTS

This research is a part of the PhD. dissertation of first author that was funded by Shahid Chamran University.

REFERENCES

- Ahloowalia, B. and Maluszynski, M. 2001. Induced Mutations: A New Paradigm in Plant Breeding. *Euphytica*, **118**: 167–173.
- Awmack, C. S. and Leather, S. R. 2002. Host Plant Quality and Fecundity in Herbivorous Insects. *Annu. Rev. Entomol.*, **47**: 817–844.
- Carey, J. R. 1993. Applied Demography for Biologists with Special Emphasis on Insects. Oxford University Press, New York, 206 PP.
- Chi, H. 1988. Life-Table Analysis Incorporating Both Sexes and Variable Development Rate among Individuals. *Environ. Entomol.*, **17**: 26–34.
- Chi, H. 1990. Timing of Control Based on the Stage Structure of Pest Population: A Simulation Approach. *J. Econ. Entomol.*, **83**: 1143–1150.
- Chi, H. 2015. *TWOSEX-MSChart: A Computer Program for the Age-Stage, Two-Sex Life Table Analysis*. URL: <http://140.120.197.173/Ecology/Download/TwoSEXMSChart.zip>.
- Chi, H. 2015. *Timing-MSChart: A Computer Program for the Population Projection Based on Age-Stage, Two-sex Life Table*. (<http://140.120.197.173/Ecology/Download/TIMING-MSChart.rar>).
- Chi, H. and Liu, H. 1985. Two New Methods for the Study of Insect Population Ecology. *Bull. Inst. Zool. Acad. Sin.*, **24**: 225–240.
- Chi, H. and Su, H. Y. 2006. Age-Stage, Two-sex Life Tables of *Aphidius gifuensis* (Ashmead) (Hymenoptera: Braconidae) and Its Host *Myzus persicae* (Sulzer) (Homoptera: Aphididae) with Mathematical Proof of the Relationship between Female Fecundity and the Net Reproductive Rate. *Environ. Entomol.*, **35**: 10–21.
- Dhanavel, D., Gnanamurthy, S. and Girija, M. 2012. Effect of Gamma Rays on Induced Chromosomal Variation in Cowpea *Vigna unguiculata* (L.) Walp. *IJCS*, **2012**: 245–250.
- Farhadi, R., Allahyari, H. and Chi, H. 2011. Life Table and Predation Capacity of *Hippodamia variegata* (Coleoptera: Coccinellidae) Feeding on *Aphis fabae* (Homoptera: Aphididae). *Biol. Control.*, **59**: 83–89.
- Furlong, M. J., Wright, D. J. and Dosdall, L. M. 2013. Diamondback Moth Ecology and Management: Problems, Progress, and Prospects. *Annu. Rev. Entomol.*, **58**: 517–541.
- Golizadeh, A., Kamali, K., Fathipour, Y. and Abbasipour, H. 2009. Life Table of the Diamondback Moth, *Plutella xylostella* (L.) (Lepidoptera: Plutellidae) on Five Cultivated Brassicaceous Host Plants. *J. Agr. Sci. Tech.*, **11**: 115–124.
- Goodarzi, M., Fathipour, Y. and Talebi, A. A. 2015. Antibiotic Resistance of Canola Cultivars Affecting Demography of *Spodoptera exigua* (Lepidoptera: Noctuidae). *J. Agr. Sci. Tech.*, **17**: 23–33.
- Goodman D, 1982. Optimal Life Histories, Optimal Notation, and the Value of Reproductive Value. *Am Nat.*, **119**: 803–823.
- Huang, Y. B. and Chi, H. 2011. Age-stage, Two-sex Life Tables of *Bactrocera cucurbitae* (Coquillett) (Diptera: Tephritidae) with a Discussion on the Problem of Applying Female Age Specific Life Tables to Insect Populations. *Insect Sci.*, **19**: 263–273.
- Khanamani, M., Fathipour, Y. and Hajiqaanbar, H. 2013. Population Growth Response of *Tetranychus urticae* to Eggplant Quality: Application of Female Age-specific and Age-stage, Two-sex Life Tables. *Int. J. Acarol.*, **39**: 638–648.

18. Mahmoudi, M. Sahragard, A., Pezhman, H. and Ghadamyari, M. 2015. Demographic Analyses of Resistance of Five Varieties of Date Palm, *Phoenix dactylifera* L. to *Ommatissus lybicus* De Bergevin (Hemiptera: Tropiciduchidae). *J. Agr. Sci. Tech.*, **17**: 263-273.
19. Maia, A. D. H., Alfredo, J. B. and Campanhola, C. 2000. Statistical Influence on Associated Fertility Life Table Parameters Using Jackknife Technique: Computational Aspects. *J. Econ. Entomol.*, **93**: 511-518.
20. Minitab Inc. 2007. *MINITAB User's Guide, Version 15*. Minitab Inc., State College, PA.
21. Naseri, B., Golparvar, Z., Razmjou, J. and Golizadeh, A. 2014. Age-stage, Two-sex Life Table of *Helicoverpa armigera* (Lepidoptera: Noctuidae) on Different Bean Cultivars. *J. Agr. Sci. Tech.*, **16**: 19-32.
22. Nikooei, M., Fathipour, Y., Jalali-Javaran, M. and Soufbaf, M. 2015. How Different Genetically Manipulated Brassica Genotypes affect Life Table Parameters of *Plutella xylostella* (Lepidoptera: Plutellidae). *J. Econ. Entomol.*, **108**: 515-524.
23. Novak, F. J. and Brunner, H. 1992. Plant Breeding: Induced Mutation Technology for Crop Improvement. *IAEA Bull.*, **4**: 25-33.
24. Safuraie-Parizi, S., Fathipour, Y. and Talebi, A. A. 2014. Evaluation of Tomato Cultivars to *Helicoverpa armigera* Using Two-Sex Life Table Parameters in Laboratory. *J. Asia-Pac. Entomol.*, **17**: 837-844.
25. SAS Institute. 2003. *SAS Statistics and Graphics Guide, Release 9.1, User's Guide*. SAS Institute Inc., Cary, NC, USA.
26. Soufbaf, M., Fathipour, Y., Karimzadeh, J. and Zalucki, M. P. 2010. Bottom-Up Effect of Different Host Plants on *Plutella xylostella* (Lepidoptera: Plutellidae): a Life-table Study on Canola. *J. Econ. Entomol.*, **103**: 2019-2027.
27. Soufbaf, M., Fathipour, Y., Zalucki, M. P. and Hui, C. 2012. Importance of Primary Metabolites in Canola in Mediating Interactions between a Specialist Leaf-feeding Insect and Its Specialist Solitary Endoparasitoid. *Arthropod Plant Interact.*, **6**: 241-250.
28. Southwood, T. and Henderson, A. 2000. *Ecological Methods*. Blackwell, Oxford.
29. Talekar, N. S. and Shelton, A. M. 1993. Biology, Ecology and Management of Diamondback Moth. *Annu. Rev. Entomol.*, **38**: 275-301.

تأثیر موتاسیون فیزیکی کلزا بر جدول زندگی *Plutella xylostella* L.

م. آکنده، ف. کچیلی، م. سوف باف، آ. راسخ، و ک. مظفری

چکیده

تاریخچه زیستی حشرات گیاهخوار پس از تغذیه روی لاین‌های موتانت بسیاری از محصولات نامعلوم است. به این منظور، پارامترهای دموگرافیک *Plutella xylostella* L. روی سه رقم کلزا ("Zar"، "RGS"، "Talaye" و لاین‌های موتانت ایجاد شده از آن‌ها به طریق موتاسیون فیزیکی ("RGS 8-1"، "RGS 10-2"، "RGS 8-13"، "Talaye 8-3"، "Zar 9-9") در شرایط گلخانه ارزیابی گردید. روش‌های جدول زندگی شامل جدول زندگی ویژه‌ی سنی ماده و جدول زندگی سنی-مرحله رشدی، دوجنسی استفاده شدند. بر اساس جدول زندگی دو جنسی تفاوت معنی‌داری بین پارامترهای



دموگرافیک بید کلم روی دو رقم “Zar” و “Talaye” در مقایسه با لاین‌های موتانت آن‌ها مشاهده نشد ولی بین پارامترهای مذکور روی رقم “RGS” و لاین‌های موتانت آن تفاوت معنی‌دار وجود داشت. لاین موتانت “RGS 8-1” علی‌رغم داشتن ویژگی‌های مطلوب زراعی در مقایسه با رقم شاهد “RGS” و سایر لاین‌های موتانت آن نسبت به افزایش جمعیت آفت حساس‌تر بود به طوریکه بر اساس جدول زندگی دوجنسی بیشترین نرخ ذاتی افزایش جمعیت (r)، نرخ خالص تولیدمثل (R_0) و نرخ متناهی افزایش جمعیت (λ) روی این لاین بود (به ترتیب ۰/۲۰۸ بر روز، ۹۸/۶۳ تخم به ازای هر فرد و ۱/۲۳۱ بر روز). همچنین تصویر جمعیت، رشد سریع این آفت را روی لاین مذکور نشان داد. بین پارامترهای رشد جمعیت آفت که به هر دو روش جدول زندگی محاسبه شد تفاوت‌های بسیار جزئی وجود داشت. بررسی پیامدهای اصلاح رقم‌های گیاهی با استفاده از روش پرتوتابی بر سازگاری حشره گیاهخوار نه تنها باعث هدایت متخصصین اصلاح نباتات به سمت گزینش‌های موفق‌تر می‌شود بلکه به نحو مؤثری روشنگری‌هایی را در حوزه برنامه‌های مدیریت آفت در زمانیکه این گیاهان در اولویت کشت قرار دارند فراهم می‌کند.