## Natural Deoxynivalenol Contamination of Corn Produced in Golestan and Moqan Areas in Iran

R. Karami-Osboo<sup>1</sup>, M. Mirabolfathy<sup>1</sup>\* and F. Aliakbari<sup>2</sup>

#### ABSTRACT

Deoxynivalenol contamination was determined in corn produced in Golestan and Ardabil (Moqan) Provinces, Iran, in 2004-2005. Samples were collected from different stages of production, including before harvest, at harvest, post harvest as well as after drying. Ground sub samples were extracted with water; each extracted sample was cleaned up through an immunoaffinity column. Deoxynivalenol was estimated through reversed-phase High Performance Liquid Chromatography (HPLC). The linearity of standard curve for 50 -10,000 ng ml<sup>-1</sup> of standard solutions was proved (R<sup>2</sup>= 0.9999). Detection limit was 10 ng g<sup>-1</sup>. Recovery of the method for 1,000 and 500 ng g<sup>-1</sup> spiked samples was 73.5% and 93.5% (n= 5). Deoxynivalenol contamination was found in 76.7% of samples in the range of 54.4-518.4 ng g<sup>-1</sup>. The mean of contamination was 116.25 ng g<sup>-1</sup>. This is the first report of natural DON contamination of corn from Iran.

Keywords: Corn, Deoxynivalenol, HPLC, Iran, Natural contamination.

#### **INTRODUCTION**

Deoxynivalenol is one of the trichothecenes produced Fusarium by species. Fusarium graminearum (Teleomorph: Gibberella zeae) isolates are considered as major producers of deoxynivalenol (DON), nivalenol (NIV), and their derivatives including 3acetyldeoxynivalenol (3-ADON), 15acetyldeoxynivalenol (15-ADON), 4acetylnivalenol (4-ANIV), in wheat, barley and corn grains (Kimura et al., 2003). Many studies account for the deleterious effects of DON on animal and human health (CAST, 2003). DON ingestion causes acute and chronic toxicity and affecting animal feeding, behavior, as well as immune function (Rotter et al., 1996; Lautraite et al., 1997; Schlatter, 2004; Pestka et al., 2004). In domestic or laboratory animals, high

DON doses of caused feed refusal, decreased weight gain, vomiting, gastrointestinal and dermal irritation and immunological alterations. DON reduces growth and feed consumption (anorexia) at low concentrations in the diet and induces vomiting at higher doses (Rotter et al., 1996). DON is known to be clastogenic (Knasmuller et al., 1997). DON is reported to bind to the ribosomal peptidyl-transferase site, inhibiting protein and DNA synthesis, consequently, the exposure resulting in decreased cell proliferation (Shifrin and Anderson, 1999). This mycotoxin is a potent protein synthesis inhibitor that can significantly alter humoral immunity, cellmediated immunity, and host resistance in a variety of experimental animal models (Pestka and Bondy, 1990).

Golestan and Moqan areas are located in the North and Northwestern Iran with favorite conditions for *Fusarium* growth.

<sup>&</sup>lt;sup>1</sup> Mycotoxins Research Laboratory, Plant Pests and Diseases Research Institute, Tehran, Islamic Republic of Iran.

<sup>&</sup>lt;sup>2</sup> Islamic Azad University, Tehran, Islamic Republic of Iran.

<sup>&</sup>lt;sup>\*</sup> Corresponding author, e-mail: mmirab2000@yahoo.com

Fusarium head blight of wheat and barley are two important diseases in Golestan Province (Golzar et al., 1998). Different Fusarium species were reported as the causal agents of corn ear rot and as the mycoflora of corn seeds. F. oxysporum, F. semitectum and F. subglutinans have been isolated from maize in Iran (Gerlach and Ershad, 1970). F. proliferatum and F. verticillioides were also among the components of the mycoflora of corn seed in Iran (Boujari and Ershad, 1993; Ershad, 1995). F.monilifrome and F.prolifratum were isolated from rotted ears collected from experimental as well as from seed production fields in Sari and Karaj areas that artificially caused ear rot (Zamani and Alizadeh, 2000). A study on mycoflora of maize harvested in the main production areas of Iran in 2000 showed that Fusarium species were predominant (38.5%), F. verticillioides being the most predominant (52% of the total isolates). Following F. verticillioides, F. proliferatum was the most prevalent among the species. Other species present at low incidence levels were F. acuminatum, F. scirpi, F.equiseti,  $F_{\cdot}$ semitectum, F. nygamai and F. culmorum (Ghiasian et al., 2004). The research reported here was conducted to determine the DON contamination of corn crop in the above mentioned areas.

#### MATERIALS AND METHODS

#### **Reagents and Standards**

DON standard was purchased from Sigma (St. Louis, MO, USA). Stock solution was prepared by dissolving the solid standard in methanol (1 mg ml<sup>-1</sup>). More diluted standard solutions for calibration (50, 100, 500, 1,000, 2,000, 3,000, 5,000 and 10,000 ng ml<sup>-1</sup>) were prepared through evaporation of aliquots (under N<sub>2</sub> stream) followed by appropriate dilution with the mobile phase or methanol. The standard solutions were kept in tightly sealed vials at -20°C. They were brought to room temperature before

use. Acetonitrile, methanol and water were analytical or HPLC grade purchased from MERCK Company (Germany), DONPREP immunoaffinity columns were obtained from R-BIOPHARM RHONE LTD (Glasgow, Scotland). Filter papers (Whatman No.1) and glass microfibre (GF/A) were from Whatman (Maidstone, UK.).

# Sample Preparation, Extraction and Clean-up

Sixty corn samples were collected from different locations of Golestan and Mogan areas in the North and Northwestern Iran, during 2004-2005. Each 5-10 kg sample was finely ground, using Romer Series IITM mill, (MO, USA) and well mixed. A 100 g subsample was taken from each main (ground) sample and stored at -20°C. Two hundred ml deionized water plus 2 g of NaCl was added to each sample and then shaken for 30 minutes at 150 rpm by means of a Gallenkamp shaker. Samples were filtered through Whatman No.1 filter paper, 10 ml of the filtrate was filtered through GF/A glass microfibre filter, then 2 ml of the extract (equivalent to 0.25 g sample) was passed through the DONPREP immunoaffinity column at a flow-rate of about 1 drop min<sup>-1</sup>. The column was rinsed with 5 ml de-ionized water at the same flow rate; the analyte was eluted with 1.5 ml methanol and collected in a clean 4 ml dark glass vial. The eluant was evaporated under a nitrogen stream at 40°C and dissolved in 1 ml of the HPLC mobile phase solution (acetonitrile: water, 1: 9) (MacDonald et al. 2005).

#### **HPLC Analysis**

Chromatographic separation and detection was performed using a Waters 616 pump and a Waters 486 tunable UV absorbance detector set at 218 nm and equipped with an automatic injector (Waters 717). Signals were processed using the Millennium

Downloaded from jast.modares.ac.ir on 2025-06-06

software (version 3.2). The reverse phase column was a Waters Nova-pak® C-18, 3.9 mm×250 mm, 4 $\mu$ m particle size (Waters Milford, MA, USA). The mobile phase was acetonitrile: water mixture (1:9; v/v) and was used isocratic at 1.0 ml min<sup>-1</sup> flow rate. The mobile phase was degassed through a vacuum–degassing device (Waters). The injection volume was 200 µl (equivalent to 0.05 g of sample). The quantification of DON was estimated by integrating the area

under the curve for DON at its retention time and was compared to a standard calibration curve for this analyte by using standard, spike and blank samples (Figure 1). DON retention time was 6.7 min under the above conditions.

#### **Recovery Test**

To prepare 500 and 1,000 ng g<sup>-1</sup> spiked corn samples, appropriate amount of



**Figure 1.** HPLC chromatograms of (A) Deoxynivalenol standard 1000 ng ml<sup>-1</sup>, (B) Corn spiked sample with 1.0 ppm (C) Blank corn sample. HPLC condition, column:  $250\times3.9$  mm, C18 reverse phase ODS, 4 µm; mobile phase: acetonitrile: water (1:9; v/v); flow rate: 1.0 ml min<sup>-1</sup>; UV detector: 218 nm.

standard solution (10 µg ml<sup>-1</sup>) of DON was added to 25 g of the blank corn samples and left for one hour to have the solvent evaporated prior to extraction. Extraction and clean up were carried out as for the other samples. The detection limit was 10 ng g<sup>-1</sup>. A positive correlation was detected among all points of the standard curve in the range of 50 to 10,000 ng ml<sup>-1</sup>; with a linearity of R<sup>2</sup>= 0.9999. Mean recoveries for 500 and 1,000 ng g<sup>-1</sup> spiked sample were 93.5 and 73.5% (n= 5), respectively.

#### **RESULTS AND DISCUSSION**

The analysis of 60 corn samples, which were collected during 2004-2005 from Golestan and Moqan areas, revealed that 76.6% of samples were contaminated with DON (Table 1), the range of contamination was 54.4 to 518.4 ng  $g^{-1}$  which was under recommended levels for DON in the world. Since 80% of analyzed samples (48 samples) were collected during post harvest stages, a validated statistical regression contamination levels between or contamination incidence percents of samples and sampling stages was not found. The following presented results are the first reported determination of DON in corn from Iran.

*Fusarium graminearum* benefits from a broad host range and can cause ear disease of corn often called red ear rot or *Gibberella* ear rot. Red ear rot has been reported in corn–growing areas worldwide but is especially prevalent in temperate climates where relatively cool temperatures and weather coincide with silk emergence.

Natural epidemics are often localized and sporadic and thus difficult to predict. During early stages of symptom development, infected ears are not easily detected in fields because husks may show few symptoms. In severe infections, however, red or pink mold can progress throughout the entire ear, colonizing kernels, cobs, and husks. If wet weather persists, blue-black perithecia and a pinkish mass of mycelia and macroconidia can form on husk surfaces. Spores landing on the emerged silks can infect the ear through the silk channel. Controlled field inoculation into the silk channel indicates that corn ears are most susceptible to F. graminearum within 4 to 7 days after silk emergence (Reid et al., 1992). Wound caused by birds, insects, or extreme weather can provide an opportunity for fungal invasion; such ear damage is positively correlated with ear rot (Sutton 1982; Vigier et al., 1997). Ascospores and macroconidia produced by F. graminearum growing on corn stalks and other crop residues are the major inocula for both wheat head scab and corn ear rot. In addition, there was a positive correlation observed between F graminearum ear rot and rainfall in July and August when corn is in silking (Vigier *et al.*, 1997). Epidemics also appear to be associated with wet weather late in the growing season (Miller, 1994). Studies on corn infected with F. graminearum in the field indicate that corn can be contaminated with DON and zearalenone. Corn cob tissue often contains the highest concentrations of DON in the ear (Reid et al., 1996). DON levels in kernels of corn are highly correlated with disease severity ratings and

**Table1.** Total contaminant samples (%), range and mean of DON contamination (ng/g) for each region at harvest and before silo

Location	Contaminant samples%	DON (ng g <sup>-1</sup> ) range	DON (ng g <sup>-1</sup> ) mean
Golestan (Ali Abbad)	80	57.6-518.4	161.3
Golestan (Aq-Qala)	64.7	54.4-357.3	104.23
Moqan	85.7	61.6-117.2	73.31
Total mean	76.7	54.4-518.4	116.25

with various measures of fungal load, including colony- forming units and ergosterol levels (Atanassov *et al.*, 1994; Bennett *et al.*, 1988; Miller *et al.*, 1983; Trigo-Stockli *et al.*, 1995). However, even visibly healthy grain can contain significant amounts of DON. In an Ontario study, 52% of normal-appearing corn kernels collected from an infected field contained 0.28 to 5µg g<sup>-1</sup> DON by ELISA (Sinha and Savard, 1997).

In spite of the wide studies conducted on F. graminearum as the causal agent of Fusarium head blight diseases of wheat and barley throughout wide areas in Iran (Safaie et al., 2005; Safaie and Alizadeh, 2001; Sanjarian et al., 2005), it has not yet been isolated from corn. F. culmorum, which was known as a DON producer (Birzele, 2002), was isolated from Iranian maize mycoflora (Ghiasian et al., 2004). It seems that the reason for low isolation frequency of F. graminearum and F. culmorum from corn seed in Iran is due to the dominant population of F. verticillioides and F. proliferatum among the other Fusarium species on maize at harvest time, in which case both compete with F. graminearum growth in artificial media in in vitro experiments. Moreover fumonisin **B**1 contamination of the Golestan's corn seed samples which were used in this research and evaluated using HPLC+IAC was quite high as all the samples were contaminated ( Mirabolfathy et al., 2007) in the range of  $261-6,891 \text{ ng g}^{-1}$  with a mean of 2,658.35 ng g<sup>-1</sup>. If selective media had been used for isolating F. graminearum, among the other Fusarium species, with altering temperature during incubation period and collecting samples at early harvest time, DON producing Fusarium species could possibly be more frequently isolated. Since the residues of infected wheat and barley are the main inoculum sources of infection, especially whereby wheat and barley are planted in rotation with corn in a majority of fields in Iran, it seems monitoring of Fusarium species during corn growth stages from silking through harvest would help in identifying DON producing *Fusarium* species. A monitoring of *Fusarium* graminearum conidia on corn ears in corn growing fields would help to prevent and allow for inhibition of DON incidence in corn, while employing the integrated pest management principles.

#### REFERENCES

- Atanassov, Z., Nakamura, C., Mort, N., Kaneda, C., Kato, H., Jin, Y. Z., Yoshizawa, T. and Mural, K. 1994. Mycotoxin Production and Pathogenicity of *Fusarium* Species and Wheat Resistance to *Fusarium* Head Blight. *Can. J. Botany*, **72**: 161-167.
- Bennett, G. A., Wicklow, D. T., Caldwell, R. W. and Smalley, E. B. 1988. Distribution of Trichothecenes and Zearalenone in *Fusarium graminearum* Rotted Corn Ears Grown in a Controlled Environment. J. Agric. Food Chem., 36: 639 – 642.
- Birzele, B., Meier, A., Hindorf, H., Kr¨amer, J. and Dehne, H. W. 2002. Epidemiology of *Fusarium* Infection and Deoxynivalenol Content in Winter Wheat in the Rhineland, Germany. *Euro. J. Plant Pathol.*, **108**: 667– 673.
- Boujari, J. and Ershad, D. 1993. An Investigation on Corn-seed Mycoflora. *Iran. J. Plant Pathol.*, 29:23-35.
- 5. CAST, 2003. Mycotoxins—risks in Plant, Animal and Human Systems: Task Force
- 6. Report No. *139*. Council for Agricultural Science and Technology, Ames, Iowa, PP. 1-191.
- 7. Ershad, D. 1995. *Fungi of Iran*. Ministry of Agriculture Research, Education and Extension Organization, No. 10, 556 PP.
- 8. Gerlach, W. and Ershad, D. 1970. Beitrag zur Kenntnis *der Fusarium*- und *Cylindrocarpon*-Arten in Iran. *Nova Hedwigia*, **20**: 725-784.
- Ghiasian, S. A., Kord-Bacheh, P., Rezayat, S. M., Maghsood, A. H. and Taherkhani, H. 2004. Mycoflora of Iranian Maize Harvested in the Main Production Areas in 2000. *Mycopathologia*, **158**: 113-121.
- Golzar, H., Foroutan, A. and Ershad, D. 1998. Studies on *Fusarium* Species Causing Head Blight of Wheat and Sources of Resistance of *F. graminearum* in Golestan

and Mazandaran. *Iran. J. Plant Pathol.*, **34**: 48-52.

- Knasmuller, S., Bresgen, N., Kassie, F., Mersch-Sundermann, V., Gelderblom, W., Zohrer, E. and Eckl, P. M. 1997. Genotoxic Effects of Three *Fusarium* Mycotoxins, Fumonisin B1, Moniliformin and Vomitoxin in Bacteria and in Primary Cultures of Rat Hepatocytes. *Mutat. Res.*, **391**: 39–48.
- Lautraite, S., Parent-Massin, D., Rio, B. and Hoellinger, H. 1997. *In vitro* Toxicity Induced by Deoxynivalenol (DON) on Human and Rat Granulomonocytic Progenitors. *Cell Biol. Toxicol.*, 13: 175-183.
- Kimura, M., Tokai, T. O'Donnell, K., Ward, T. J., Fujimura, M., Hamamoto, H., Shibata, T. and Yamaguchi, I. 2003. Trichothecene Biosynthesis Gene Cluster of *Fusarium* graminearum F15 Contains a Limited Number of Essential Pathway Genes and Expressed Non-essential Genes, *FEBS* Letters, 539: 105-110.
- 14. MacDonald, J., Chan, D., Brereton, P., Damant, A. and Wood, R. 2005. Determination of Deoxynivalenol in Cereals and Cereal Products by Immunoaffinity Column Chromatography with Liquid Chromatography. *J. of Aoac. Int.*, **88(4)**: 1197-1204.
- Miller, J. D., Young, J. C. and Trenholm, H. L. 1983. *Fusarium* Toxins in Field Corn. 1. Time Course of Fungal Growth and Production of Deoxynivalenol and Other Mycotoxins. *Can. J. Botany*, **61**: 3080 – 3087.
- Miller, J. D. 1994. Epidemiology of *Fusarium* Ear Diseases of Cereals. In: "Mycotoxins in Grain; Compounds Other than Aflatoxin", Miller, J. D. and Trenholm, H. L. (Eds.). Eagan Press, St, Paul, Minnesota. PP. 19-36.
- Mirabolfathy, M., Karami-Osboo, R and Amini, H. 2007. Fumonisin B1 Contamination of Golestan Corn Product. *Iran. J. Plant Pathol.*, 42: 359-374.
- Pestka, J. J. and Bondy, G. S. 1990. Alteration of Immune Function Following Dietary Mycotoxin Exposure. *Can. J. Physiol. Pharmacol.*, **66**: 1009–1016.
- Pestka, J. J., Zhou, H. R., Moon, Y. and Chung, Y. J. 2004. Cellular and Molecular Mechanisms for Immune Modulation by Deoxynivalenol and Other Trichothecenes:

Unraveling a Paradox. *Toxicol. Lett.*, **153**: 61–73.

- Reid, L. M., Bolton, A. T., Hamilton, R. I., Wolddermariam, T. and Mather, D. E. 1992.
  Effect of Silk Age on Resistance of Maize to *Fusarium graminearum. Can. J. Plant Pathol.*, 14: 293-298.
- Reid, L. M., Mather, D. E. and Hamilton, R. I. 1996. Distribution of Deoxynivalenol in *Fusarium graminearum*–infected Maize Ears. *Phytopathol.*, 86: 110 – 114.
- Rotter, B. A., Prelusky, D. B. and Pestka, J. J. 1996. Toxicology of Deoxynivalenol (vomitoxin). *J. Toxicol. Environ. Health*, 48: 1–34.
- Safaie, N., Alizadeh, A., Saidi, A., Rahimian, H. and Adam, G. 2005. Molecular Characterization and Genetic Diversity among Iranian Populations of *Fusarium graminearum*, the Causal Agent of Wheat Head Blight. *Iran. J. Plant Pathol.*, 41(2): 171-191.
- Safaie, N. and Alizadeh, A. 2001. Phenotypic Diversity in *Fusarium* gramineum Isolates and Introduction of a New Genetic Marker for the Species. *Iran. J. Plant Pathol.*, 37(3-4): 197-209.
- 25. Sanjarian, F., Mousavi, A., Alizadeh, A., Abolmaali, Sh., BuschA, B., Lucyshin, D. and Adam, G. 2005. Identification of Wheat Ribosomal Protein L3 (*RPL3*) Genes and Their Possible Role in Conferring Resistance to Fusarium Head Blight. Iran. J. Plant Pathol., **41(2)**: 257-271.
- Schlatter, J. 2004. Toxicity Data Relevant for Hazard Characterization. *Toxicol. Lett.*, 153: 83–89.
- Shifrin, V. I. and Anderson, P. 1999. Trichothecene Mycotoxins Trigger a Ribotoxic Stress Response that Activates *c*-Jun N-Terminal Kinase and *p38* Mitogenactivated Protein Kinase and Induces Apoptosis. J. Biol. Chem., 274: 13985– 13992.
- 28. Sinha, R.C. and Savard, M.E. 1997. Concentration of deoxynivalenol in single kernels and various tissues of wheat heads. *Can. J. Plant Pathol.*, **19**: 8–12.
- 29. Sutton, J. C. 1982. Epidemiology of Wheat Head Blight and Maize Ear Rot Caused by *Fusarium graminearum. Can. J. Plant Pathol.*, **4**: 195-209.

JAST

- Trigo-Stockli, D. S., Curran, P. and Pedersen J. R. 1995. Distribution and Occurrence of Mycotoxins in 1993 Kansas Wheat. *Cereal Chem.*, 72: 470–474.
- Vigier, B., Reid, L. M., Seilfert, K. A., Stewart, D. W. and Hamilton, R. I. 1997. Distribution and Prediction of *Fusarium*

Species Associated with Maize Ear Rot in Ontario. *Can. J. Plant Pathol.*, **19:** 60-65.

 Zamani, M. and Alizadeh, A. 2000. Identification of *Fusarium* Ear Rot of Corn in Sari and Karaj. *Iran. J. Plant Pathol.*, 36(1-2): 15-31

### آلودگی طبیعی محصول ذرت مناطق گلستان و مغان به داکسی نیوالنول

## ر. كرمي اسبو، م. ميرابوالفتحي وف. على اكبري

## چکیدہ

محصول ذرت سالهای ۸۵–۱۳۸۴استان گلستان و مغان از نظر آلودگی به داکسی نیوالنول مورد بررسی قرار گرفت. نمونه ها از مراحل مختلف شامل قبل از برداشت، در مرحله برداشت، پس از برداشت و بعد از خشک نمودن دانه های ذرت جمع آوری گردید. زیر نمونه های آسیاب شده با استفاده از آب عصاره گیری و با ستون های ایمونوافینیتی تصفیه گردید. میزان داکسی نیوالنول نمونه ها با استفاده از کروماتوگرافی مایع فاز معکوس ارزیابی شد. منحنی استاندارد برای محلول های استاندارد در غلظت های ۲۰۰۰۱ – ۵۰ نانوگرم در میلی لیتر کاملا خطی بود، حد تشخیص روش ۱۰ نانوگرم درگرم و بازیافت آن برای نمونه های غنی شده حاوی ۵۰۰ و ۱۰۰۰ نانوگرم درگرم داکسی نیوالنول به ترتیب ۹/۸۳ و ۲۸ درصد بود. آلودگی به داکسی نیوالنول در ۷۶/۷ نمونه ها مشاهده گردید و گستره آلودگی برابر ۵/۴۰ تا