# Sublethal Effects of Commonly Used Insecticides in Tomato Crop on Functional Response and Biological Parameters of *Macrolophus pygmaeus* Rumber (Hemiptera: Miridae)

Y. Safari Monjezi<sup>1</sup>, F. Yarahmadi<sup>1\*</sup>, and N. Zandi Sohani<sup>1</sup>

#### ABSTRACT

The hemipteran, Macrolophus pygmaeus Rumber, is an effective biocontrol agent against many crop pests including tomato leaf miner, Tuta absoluta Meyrick. Sublethal effects of azadirachtin (Neem Azal<sup>®</sup>), indoxacarb (Avaunt<sup>®</sup>), and emamectin benzoate+lufenuron (Proclim Fit®) were studied on biological characteristics and functional response of the predatory bug in laboratory conditions at  $25\pm1^{\circ}$ C,  $6\pm5\%$  RH, and a photoperiod of 16:8 hour (L:D). For this purpose, females of the predatory bug were exposed to sublethal residues (10% of field concentration) of the insecticides. Twosex life table and Roger's model were used to investigate effects of the insecticides on biological parameters and functional response of the predator, respectively. Results indicated that sublethal residues of indoxacarb and azadirachtin had the highest and lowest side effects on life table parameters of M. pygmaeus. Whereas azadirachtin has no significant effects on the bug biological parameters, total fecundity and longevity of the bug in indoxacarb treatment were significantly lower (14.6%) and higher (6.9%) than the control. The predator showed type III functional response. The type of functional response was not affected by the insecticide residues. However, the coefficient of attack rate (b) for the indoxacarb treatment (0.1521) was significantly lower than for other treatments. However, there was no significant difference between the handling Times (T<sub>h</sub>) of the treatments. In conclusion, the experiments proved that azadirachtin is a low risk insecticide to M. pygmaeus, which can be integrated with biological control by the predator.

Keywords: Biological control, Chemical control, Foraging behavior, Life table, Mirid bug.

#### INTRODUCTION

The mired bug, *Macrolophus pygmaeus* Rumber (Heteroptera: Miridae), is an important polyphagous predator that has been reported to be effective in controlling many crop pests including whiteflies, thrips, aphids, mites and larvae of lepidopteran pests in field and greenhouse conditions (Margaritopoulos, 2003; Sylla *et al.*, 2016). This predator has been used for biological control of tomato leaf miner, *Tuta absoluta* Meyrik (Lepidoptera: Gelechiidae) (Bompard *et al.*, 2013; Zappalà *et al.*, 2013; Jaworski *et al.*, 2015; Biondi *et al.*, 2018), and actively feeds on eggs and all larval stages of *T. absoluta* (Jaworski *et al.*, 2013; Sylla *et al.*, 2016).

Insecticide application is the most commonly method used for control of T. absoluta (Arnó and Gabarra, 2011). Concern about adverse effect of the chemicals on nontarget organisms and the environment has prompted scientists to develop Integrated Pest Management (IPM) programs (Smith, 1991). Biological control is the main preventive control strategy in an IPM program (Pedigo, 2002). Therefore, studying the impact of the insecticides, as another component of IPM, on the biocontrol agents is necessary (Wright and Verkerk, 1995; Nazarpour et al., 2016). The

<sup>&</sup>lt;sup>1</sup> Department of Plant Protection, Faculty of Agriculture, Agricultural Sciences and Natural Resources University of Khuzestan, Mollasani, Ahvaz, Islamic Republic of Iran.

<sup>\*</sup>Corresponding author; e-mail: yarahmadi@asnrukh.ac.ir

sublethal effects of insecticides on physiology, biology and behavior of natural enemies must be considered for a complete analysis of their impact (Desneux *et al.*, 2007).

Indoxacarb is an oxadiazine insecticide with a great effect on lepidopteran pests (Wing et al., 2000) such as T. absoluta (Nazarpour et al., 2016). The insecticide blocks the voltage dependent sodium canals, group 22A according to IRAC MoA classification version (IRAC, 2020). Another effective 9.4 insecticide that is applied against the pest is azadirachtin (AZ) (Nazarpour et al., 2016). Products with the active ingredient AZ, a tetranortriterpenoids, group of have antifeedant, deterrent and insect growth regulatory properties. The products have been obtained from seeds and fruits of Azadirachta indica Juss and Melia azaderach L. trees, respectively (Isman, 2017; Nazarpour et al., 2016). Proclaim Fit<sup>®</sup> (Syngenta Agro AG, Basel, Switzerland) is a new insecticidal product that contains 10% emamectin benzoate, a GABA and glutamate-gated chloride channel agonist (group 6 according to IRAC 2020), and 40% lufenuron, which acts by disrupting the formation of the endocuticle, group 15 according to IRAC 2020 (Copping and Menn, 2000).

Sublethal residues of insecticides can adversely affect life parameters (developmental rate, longevity, fecundity, oviposition, sex ratio), and behavior, including functional responses to prey density) of a predatory insect (Desneux et al., 2007; Rahmani and Bandani, 2013). Estimation of demographic parameters through life table analysis is an essential approach to evaluate population growth and thus these studies are valuable for assessment of sublethal effects of a pesticide on both pest and natural enemies (Rahmani and Bandani, 2013). Moreover, functional and numerical responses are two main components of prey-predator interactions that can be significantly affected by insecticide residues (Martinou and Stavrinides, 2015). Effects of some insecticidal residues including AZ (Tedeschi et al., 2001; Arnó and Gabarra, 2011), indoxacarb, and spinosad (Arnó and Gabarra, 2011) on biological parameters (eg.

survival rate and fecundity) as well as thiacloprid, chlorantraniliprole (Martinou *et al.*, 2014; Martinou *et al.*, 2015), abamectin, imidacloprid and chlorpyrifos (Sharifian *et al.*, 2017) on functional responses and predation rate of *M. pygmaeus* were previously studied.

Using study of life table parameters, a comprehensive description of population dynamics including the multiple sub-lethal effects of insecticides on insects can be achieved (Zhang et al., 2014). Due to some limitations of traditional female age-specific life table, two-sex life table method was developed. The method can be applied to agestage-structure two-sex populations, and is able to include variation in pre-adult developmental time, therefore, survival and fecundity curves will be more accurate (Chi and Getz, 1988). Moreover, functional response models and parameters have been studied for the predator foraging and exploring their dynamics and provide a conceptual understanding of prey-predator relationships. The response can be affected by sublethal residues of insecticides (Martinou and Stavrinides, 2015).

There has been limited information the effects of AZ, indoxacarb and emamectin benzoate+lufenuron on functional response and biological characteristics of *M. pygmaeus*. Therefore, this study aimed to evaluate effects of these three insecticides on life table parameters and functional responses of this predatory bug.

# MATERIALS AND METHODS

# **Rearing of Macrolophus pygmaeus**

The initial colony of *M. pygmaeus* was provided from Koppert Biological System Co. representative in Iran, Giah Co., Tehran. The insects were reared on flour moth eggs, *Ephestia kuehniella* Zeller (Lepidoptera: Pyralidae), and date palm pollen as daily diet, and bean pod as oviposition substrate at  $25\pm1^{\circ}$ C,  $60\pm5^{\circ}$  RH, and 16:8 hours (Light:Dark) in an incubator. The insects (30-70 individuals) were kept in Plexiglas cylinders (18 cm high, 7.5 cm diameter) covered with a fine gauze, 100 meshes, lid on the top and margin for ventilation. Pieces of paper were placed in the rearing cylinder for resting the predator and reducing cannibalism.

#### **Experimental Design**

The experimental insecticides were azadirachtin A (Neem Azal® 1%EC, Trifoli Co., Germany), indoxacarb (Avaunt<sup>®</sup>15%SC, emamectin Giah Corp., Iran) and benzoate+lufenuron (Proclim Fit® 50% G, Syngenta Co., Austria). The insecticides were selected because they are currently applied to control T. absoluta in greenhouses of tomato (Derbalah et al., 2012). Standard methods to study side effects of insecticides on beneficial developed by **IOBC/WPRS** organism, Working Group, were used for the trials. The insects were exposed to 10% of field concentration in the experiments according to preliminary tests. The tests showed that the concentration caused about 18-20% mortality in the predator population. The concentrations were 400, 250, and 50 PPM for indoxacarb, azadirachtin and emamectin benzoate+lufenuron (Proclim Fit<sup>®</sup>). respectively. It should be noted that each treatment was separately carried out at 25±1°C, 65±5% RH, and a photoperiod of 16:8 hour (L:D) in an incubator that was well ventilated to avoid interference of different treatments.

## Effect of Insecticides on Life Table Parameters

For each treatment, fifty glass Petri dishes (10 cm in diameter) were directly treated with the insecticides and allowed to dry (approximately 1-hour after application). Fifty pairs (male and female) of *M. pygmaeus*, which were obtained from first progeny (G1) of reared insects, were separately placed in each dish, which was then covered with a piece of Parafilm  $M^{\oplus}$  to avoid escaping of the bugs. To facilitate ventilation, a hole was made

on top of the dishes (2.5 cm diameter) and covered with fine net. After 24 hours, 20 pairs of the bug were chosen and transferred to a new rearing cylinder, with bean pods, for oviposition. After 24 hours, the bugs were removed. Females oviposit in plant tissues, leaf and stem. Due to difficult detection of the bug eggs in the plant tissues, the first twelve instar nymphs, 1 day old, were used for experiments as cohort. The life table was constructed based on 60 individuals of 1st nymphs as cohort. Each of the first instar nymphs was placed in Plexiglas Petri-dish, 9 cm in diameter, at 25±1°C, 60±5% RH, and 16:8 hours (Light:Dark) in an incubator. The insect was daily checked and number of alive bugs and their life stages were recorded. Upon adult emergence, bean pods were removed to record the laid eggs and replaced with fresh one. The observation was continued until death of the last individual. In control, the males and females were not exposed to any insecticides. Each treatment had 12 replications.

Theory of age-stage (two sex life table), developed by Chi and Getz (1988) was used to analyze the raw life history data of *M. pygmaeus* by "TWOSEX-MS Chart for the Windows operating system" software, available at http://140.120.197.173/Ecology/prod02.htm

(Chi, 2017). According to the theory, the parameters of age-specific survival rate  $(l_x)$ , age-stage specific survival rate ( $s_{xi}$ ; where, x= Insect age and j= Life stage), life expectancy (  $e_{xi}$ ; where, x= Insect age and j= Life stag), mean generation Time (T), age-specific fecundity (mx), age-stage specific fecundity (f<sub>xi</sub>), gross reproductive rate ( $GRR = \sum m_{\chi}$ ), net Reproductive rate (R<sub>0</sub>), intrinsic rate of increase (r), as well as finite rate of increase  $(\lambda = e^{r})$  were estimated. Also, the Adult Pre-Oviposition Period (APOP: The period between the emergence of an adult female and her first oviposition), Total Pre-Oviposition Period (TPOP: The time interval from birth to the beginning of oviposition) were also calculated using the experimental data. Bootstrap technique, using 1,000,000 resampling, was used for estimating standard

errors and variance of the life table parameters (Tibshirani and Efron, 1993). Moreover, the software was used for estimating the population consumption projected by using the life tables and predation rate of the original cohort, the cohorts constructed based on the 2.5 and 97.5% percentiles of finite rate ( $\lambda$ ) and net Reproductive rate (R) in different experimental treatments. Sigma plot 12.5 was used to generate graphs.

## Effect of Insecticides on Functional Responses

Each treatment was separately carried out. The experimental arena consisted of a Plexiglass Petri dish (10 cm diameter×1 cm deep). For ventilation, a hole was created on top of the dishes (2 cm diameter) and covered with fine gauze. The dishes were dipped in concentrations of 400, 250 and 50 PPM of indoxacarb, AZ and emamectin benzoate+lufenuron 30 seconds, for respectively. Distillated water was used in the control. One hundred females were exposed to the insecticide residues during 24 hours and they were used for functional responses experiments.

Densities of 8, 16, 32, 64 and 128 eggs of *E. kuehniella* per Petri dish were used in the trials. A female bug  $(3\pm 2 \text{ days old})$  was located to each Petri-dish. Before beginning the experiments, the predator was starved for 24 hours. After 24 hours of the introduction, the predators were eliminated and the numbers of preyed eggs were recorded. All experiments were done in completely randomized design with nine replications.

Logistic regression analyses were used to determine type of *M. pygmaeus* functional responses (Juliano, 2001). A polynomial function that describes the relationship between N<sub>e</sub> and N<sub>0</sub> was used for fitting the data (Equation 1): All experiments were carried out at  $25\pm1^{\circ}$ C,  $65\pm5^{\circ}$  RH, and a photoperiod of 16:8 hours (L:D) in an incubator.

$$\frac{N_a}{N_0} = \frac{\exp(P_0 + P_1 N_0 + P_2 N_0^2 + P_3 P_0^3)}{(1 + \exp P_0 + P_1 N_0 + P_2 N_0^2 + P_3 N_0^3)}$$
(1)

Where, Ne is the Number of consumed eggs, N<sub>0</sub> is the initial Number of eggs, and  $P_0$ ,  $P_1$ ,  $P_2$ , and  $P_3$  are the intercepts of the linear, quadratic, and cubic coefficients, respectively. These parameters were calculated using the method of maximum likelihood (PROC CATMOD, SAS Institute 2001). If  $P_1 > 0$  and  $P_2 < 0$ , the proportion of consumed eggs is positively densitydependent, Therefore, the data describes type III of functional response, but if  $P_1 < 0$ and  $P_2 > 0$ , the proportion of consuming eggs decreases gradually as the initial number of egg offered increases and the data indicate type II of functional response (Juliano, 2001). Because our data implicated type III functional response (see result section), in the next step, fitting Roger's type III random predator (Equation 2) was done by using nonlinear least squares regression (PROC NLIN; SAS Institute Inc. 2001) for parameters estimating of functional responses (Rogers, 1972).

 $N_a = N_0 [1 - e^{(bN_0)(T_h N_e - T)}]$  (2) Where, N<sub>a</sub> is the Number of preyed egg, N<sub>0</sub> is the initial Number of egg, b is coefficient for change in attack rate (a) with prey density (a= bN<sub>0</sub>), T is the total available Time available for searching (in this experiment T= 24 hours), and T<sub>h</sub> is the handling Time. The functional response parameters were not considered statistically different when their confidence limits (95%) overlapped (Juliano, 2001).

#### RESULTS

### Effect of Insecticides on Life Table Parameters

Total fecundity, sex ratio, TPOP and APOP of *M. pygmaeus* in different experimental treatments are shown in Table 1. The results indicated significant differences in TPOP, naymphal periods and APOP of the bug among different treatments. TPOP in indoxacarb treatment (26.41 days) was significantly more than

<u> </u>				
Treatment	Total fecundity	Sex ratio	APOP (Days)	TPOP (Days)
Azadirachtin	$54.5 \pm 5.8^{a^*}$	$4.07 \pm 0.54^{b}$	$10.12 \pm 0.23^{b}$	26.02±0.19 <sup>a</sup>
Indoxacarb	$47.2 \pm 5.5^{b}$	$7.06 \pm 0.54^{a}$	$10.35 \pm 0.16^{a}$	$26.41 \pm 0.2^{b}$
Emamectin benzoate	$55.19 \pm 4.6^{a}$	$6.1 \pm 0.53^{a}$	$9.52 \pm 0.3^{\circ}$	$26.06 \pm 0.2^{a}$
Control	55.3±7.1 <sup>a</sup>	$6.7 \pm .063^{a}$	$10.47 \pm 0.23^{a}$	$26.12 \pm 0.18^{a}$

**Table 1.** Effects of different experimental treatments on biological parameters (mean $\pm$ SE) of *Macrolophus* pygmeus.<sup>*a*</sup>

**Table 2.** Effects of sublethal dosage of azadirachtin, indoxacarb and emamectin benzoate+lufenuron on life table parameters (mean $\pm$ SE) of *Macrolophus pygmaeus.<sup>a</sup>* 

	λ	GRR	R	$R_0$	Т
Azadirachtin	$1.086 \pm 0.006^{a^*}$	$49.424{\pm}11.088^{a}$	$0.082 \pm 0.006^{a}$	21.379±4.733 <sup>a</sup>	$37.18 \pm 1.014^{a}$
Indoxacarb	$1.085 \pm 0.005^{a}$	$55.969 \pm 7.694^{a}$	$0.091 \pm 0.004^{a}$	$28.539 \pm 4.819^{a}$	$37.256 \pm 0.697^{a}$
Emamectin	$1.076 \pm 0.007^{b}$	$43.068 \pm 8.092^{a}$	$0.074 \pm 0.007^{b}$	$16.049 \pm 4.215^{b}$	$36.749 \pm 0.686^{a}$
benzoate+Lufenuron					
Control	$1.089 \pm 0.006^{a}$	44.043±8.573a	$0.085 \pm 0.005^{a}$	$23.522 \pm 4.777^{a}$	$36.73 \pm 0.745^{a}$
P-value	0.003	0.24	0.003	0.002	0.354

<sup>*a*</sup>  $\lambda$ : Finite rate of increase, *GRR*: Finite Rate of increase; r: Intrinsic rate of increase;  $R_0$ : Net Reproductive rate, T: Mean generation time (Tukey–Kramer Procedure). (a-b) The same letters in each column indicate non-significant difference (P< 0.05) using paired-bootstrap.

that of the control (26.12 days). However, APOP in emamectin benzoate+lufenuron (9.52 days) was significantly less than other treatments. Sex ratio of *M. pygmaeus* was azadirachtin. by affected Moreover, indoxacarb had the least total fecundity (47.2 eggs) among all treatments. Total fecundity in azadirachtin (54.5 eggs) and emamectin benzoate+lufenuron (55.19 eggs) treatments did not differ significantly from the control (55.3 eggs). However, total fecundity in indoxacarb treatment (47.2 eggs) was significantly lower than the control.

The intrinsic rate of increase (r), net Reproduction rate (R<sub>0</sub>), Gross Reproductive Rate (GRR), finite rate of increase ( $\lambda$ ), and the mean generation Time (T) were estimated using the bootstrap method and are shown in Table 2. No significant dereference was observed between the GRR and T values of *M. pygmaeus* in insecticidal treatments and control. In indoxacarb treatment, the  $\lambda$ , r and R<sub>0</sub> values were 10.09, 12.94% and 31.77% lower than those in the control, respectively.

The adult, female and male longevity as well as nymphal and adult durations are shown in Table 3. The longest and shortest longevities were observed adult in indoxacarb (46.9 days) and control (43.9 days), respectively. The insecticide treatments had no effects on male and female longevities. Similarly, nymphal and adult duration times were not affected by the insecticidal treatments. The lowest nymphal duration (25.02) was observed in emamectin benzoate+lufenuron.

The curves for the age-stage specific Survival rate  $(S_{xi})$  indicate the probability of a newborn individual surviving to age x and stage j, are presented in Figure 1 for different treatments. The  $S_{xj}$  curves of the control, azadirachtin and indoxacarb are somewhat similar. but emamectin benzoate+lufenuron causes reduction in survival rate of all developmental stages. Age-specific survival rate  $(l_x)$ , age-specific fecundity of total population (m<sub>x</sub>) and agespecific net maternity  $(l_x m_x)$ , specific survival rate for different treatments are shown in Figure 2. Curves of  $l_x$ ,  $m_x$  and  $l_x m_x$ in AZ treatment are similar to those in

<sup>&</sup>lt;sup>*a*</sup> TPOP: Total PreOviposition Period, APOP: The Adult PreOviposition Period. (a-b) The same letters in each column indicate non-significant difference (P < 0.05) using paired-bootstrap method.



Table 3. Developmental time (mean±SE) of different stages of Macrolophus pygmaeus in different treatments.<sup>a</sup>

Treatment	Adult	Female	Male	Nymph	Adult
	longevity	longevity	longevity		
	(Days)	(Days)	(Days)		
Azadirachtin	$45.8 \pm 1.6^{ab^*}$	$44.89 \pm 2.36^{a}$	$47.09 \pm .18^{a}$	25.63±0.21 <sup>ab</sup>	$19.87 \pm 1.56^{a}$
Indoxacarb	$46.9 \pm 1.6^{b}$	$45.52 \pm 2.63^{a}$	$44.57 \pm 2.63^{a}$	$26.16 \pm 0.16^{a}$	$18.75 \pm 1.51^{a}$
Emamectin	$45.1 \pm 1.54^{a}$	$43.3 \pm 2.68^{a}$	45±3.01 <sup>a</sup>	$25.02 \pm 0.22^{b}$	$17.94{\pm}1.88^{a}$
benzoate+Lufenuron					
Control	$43.9 \pm 1.98^{a}$	44.74±2.24a	47.04±2.23a	$27.08 \pm 0.19^{a}$	19.68±1.51 <sup>a</sup>
P-value	0.002	0.78	0.92	0.001	0.83

 $^{a}$  (a-b) The same letters in each column indicate non-significant difference (P< 0.05) using paired-bootstrap method.

**Table 4.** Maximum-likelihood estimates from logistic regressions of the proportion of *E. kuehniella* eggs eaten by *M. pygmaeus* female on initial prey density.

Treatment	Туре	Coefficient	Estimate	SE	$\chi^2$	P-value
Azadirachtin		Constant	-0.2516	0.605	17.3	< 0.0001
	п	Linear	0.00641	0.00209	9.44	0.0021
	11	Quadratic	-0.00007	0.000028	6.93	0.0085
		Cubic	2.92×10 <sup>-7</sup>	$1.2 \times 10^{-7}$	5.63	0.0179
		Constant	-0.2508	0.604	17.27	< 0.0001
Indoxacarb	П	Linear	0.00637	0.00208	9.34	0.0022
		Quadratic	-0.00006	0.000028	6.8	0.0091
		Cubic	$2.87 \times 10^{-7}$	1.23×10 <sup>-7</sup>	5.45	0.0196
		Constant	-0.2501	0.0602	17.28	< 0.0001
Emamectin	П	Linear	0.000639	0.00208	9.41	0.0022
benzoate+Lufenuron		Quadratic	-0.00007	0.000028	6.78	0.0088
		Cubic	2.9×10 <sup>-7</sup>	1.23×10 <sup>-7</sup>	5.51	0.0189
		Constant	-0.2528	0.0605	17.48	< 0.0001
Control	П	Linear	0.00646	0.00209	9.6	0.0019
	-	Quadratic	-0.00007	0.000028	7.04	0.008
		Cubic	2.93×10 <sup>-7</sup>	1.23×10 <sup>-7</sup>	5.67	0.0173

control. The highest reduction in  $l_x$  and l<sub>x</sub>m<sub>x</sub> values were observed in emamectin benzoate treatment. The curves of the life expectancies (exi) of *M. pygmaeus* for the azadirachtin, indoxacarb control. and emamectin benzoate+lufenuron are given in Figure 3. Values of  $e_{xj}$  in all experimental treatments decreased with time. The reduction was gradual in the control and AZ treatment. There was no difference between  $e_{xj}$  curves in AZ and control. However, the age-e<sub>xi</sub> curves rapidly fall down in emamectin benzoate and indoxacarb treatments, especially during days 10-20.

There was no difference between the reproductive value  $(v_{xj})$  curves of nymphal and female in the different experimental treatments (Figure 4). However,  $(v_{xj})$  of nymphal stage in emamectin benzoate+lufenuron was relatively more than the other treatments.

### Effect of Insecticides on Functional Responses

Functional response curves of the predatory bug to the various densities of *E*.



Figure 1. Age-stage survival rate (sxj) of Macrolophus pygmaeus in different experimental treatments.



**Figure 2.** Curves of age-specific survival rate  $(l_x)$ , fecundity  $(m_x)$  and maternity  $(l_xm_x)$  of *Macrolophus pygmaeus* in different treatments.



**Figure 3.** (*e<sub>xj</sub>*) of *Macrolophus pygmaeus* in different experimental treatments.



Figure 4. The age-stage life reproductive value  $(v_{xj})$  of *Macrolophus pygmaeus* in different treatments.

() HILLS

*kuehniella* eggs are presented in Figure 5. The consumed eggs were increased by increasing the initial prey density.

The results of logistic regressions revealed a type III functional response of M. pygmaeus to eggs of E. kuehniella in the control and all insecticide treatments (Table 4). The total number of prey eggs consumed by M. pygmaeus increased linearly with increase in prey density. The coefficients of attack rate (b) and handling Time  $(T_h)$  for different treatments are shown in Table 5. The lowest value of the coefficient of attack rate (b) of M. pygmaeus females were observed in indoxacarb (0.1521). The handling Times  $(T_h)$  of the bug were 2.6471, 2.6845, 3.2226, and 2.6345 hours in AZ, indoxacarb, emamectin benzoate+lufenuron and control, respectively. There was no significant difference between the handling time values of various treatments.

#### DISCUSSION

indicated sublethal Our data that concentrations of indoxacarb and AZ had the highest and lowest negative effects on life table parameters of M. pygmaeus, respectively. The finding is in agreement with Arno and Gabarra (2011) who showed that sublethal residue of indoxacarb causes significant mortality to M. pygmaeus in comparison with AZ under field condition at days after treatment. However, seven indoxacarb residue did not adversely affect the bug fecundity. The conflict results may be due to different experimental conditions and the insecticidal concentration. For instance, temperature can significantly influence insect susceptibility to sublethal residues due to physiological effects of temperature on the insect hormone systems (Glunt et al., 2014; Michalko and Košulič,



Figure 5. Number of consumed eggs of *Ephestia keuhniella* at different prey densities in various experimental treatments.

Treatment	Parameters	Estimate	Asymptotic 95% CI	
			Lower	Upper
Azadinashtin	b	0.2854	0.2753	0.2955
Azadıracının	$T_{h}$	2.6471	2.6470	2.6472
To to see t	b	0.1521	0.1016	0.2025
Indoxacarb	$T_h$	2.6845	2.5703	2.7987
Emamectin	b	0.2121	0.1120	0.3121
benzoate+Lufenuron	$T_h$	3.2226	0.1372	3.4988
Control	b	0.2911	0.2910	0.2912
Control	$T_{h}$	2.6349	2.6348	2.6349

**Table 5.** Estimated ( $\pm$ SE) attack rate (*a*) and handling Time ( $T_h$ ) of *Macrolophus pygmaeus* female on eggs of *Ephestia kuehniella* in different experimental treatments.

2016). Effect of temperature on susceptibility of natural enemies to insecticide residues is well known (Desneux et al., 2007). Also, using various types of substrate (petri dish or plant) may influence insecticide residue effects. Petri dish is inert substance and plant has waxy surface, The differences trichomes, etc. can significantly affect the residue effects (Yarahmadi et al., 2009). Also, the results are in line with Nazarpour et al. (2016) who showed that indoxacarb had the highest adverse effects on population of coexisting generalist predators of Τ. absoluta. Coccinella septempunctata L. and Chrysopa carnea Stephens, in comparison with AZ Bacillus thuringiensis Berlinier. and Similarly, residues of neem products had fewer side effects on fecundity and survivals of Macrolophus caliginosus Wagner at five days after treatment (Tedeschi et al., 2001). Also, it is demonstrated that fecundity and survival rate of Orius laevigatus Fiber (Het., Anthocoridae) were not significantly affected by AZ and indoxacarb residues (Angeli et al., 2005). Moreover, indoxacarb did not show significant toxicity on Orius insidiosus Say (Het., Anthocoridae).

In many cases, deleterious effects of AZ on beneficial arthropods, e.g. Natural enemies, were reported. Therefore, its safety has been a matter of debate (Arno *et al.*, 2009; Gontijo *et al.*, 2015). Moreover, the bio-pesticide exhibits some side effects including sterilant activity, in addition to anti-feedant and growth regulator activity on natural enemies and pollinators. However, different side effects on non-target arthropods can be observed based on concentration of AZ residues (Lima et al., 2015). Side effect of AZ on immature stages of predatory arthropods may occur due to growth regulatory properties of biopesticide and it can also affect hormone physiology of the organisms (Pilar Marco et al., 1990; Gontijo et al., 2015).

Our data implicated the type III functional response of M. pygmaeus. Therefore, the type of functional response type was not exposure affected by to sublethal concentration of the insecticides. However, the coefficient of attack rate (b) of the predator was affected by exposure to indoxacarb residue, and no significant differences were observed among the estimated handling Times  $(T_h)$  of the treatments.

The total time of functional response includes attack rate+handling time. Therefore, increase in handling time causes decrease in available time for prey searching. It was previously reported that abiotic factors, such as pesticides, can enhance the handling time due to their behavioral effects (Li *et al.*, 2006; He *et al.*, 2012; Malaquias *et al.*, 2014; Martinou *et al.*, 2015).

Similarly, Sharifian *et al.* (2017) showed that sublethal concentration of abamectin, imidacloprid and chlorpyrifos did not significantly affect functional response type of *M. pygmaeus*. Their study revealed that

abamectin has no effect on handling time and attack rate of the predatory bug. parameters However, these were significantly influenced by imidaclopride and chlorpyrifos residues. In addition, Martinou et al. (2014) demonstrated that thiacloprid residue causes significant reduction in the predation rate of M. pygmaeus. However, no significant effect was observed for chlorantraniliprole residue on the insect predation rate. Also, it is reported that functional response parameters of Acanthaspis pedestris Stål (Het., Reduviidae) are negatively affected by cypermetrin (Claver et al., 2003).

One of the best methods to control T. absoluta in the greenhouses is integrating usage of predators and selective insecticides (Arno *et al.*, 2009). Therefore, we recommend application of AZ in integration with M. *pygmaeus* release to control T. *absoluta* in greenhouse tomatoes.

## CONCLUSIONS

Our laboratory study showed that biological parameters (fecundity, survival and longevities) of M. pygmaeu were not significantly affected by AZ residues. Moreover, functional response type and parameters were not significantly affected in AZ treatment. Therefore, the study suggests that integration of *M. pygmaeus* with AZ is a compatible and effective strategy for management of greenhouse pests. However, further studies on the compatibility and effectiveness of AZ and the predator are required under greenhouse and field conditions.

# ACKNOWLEDGEMENTS

The research was supported by grant from Agricultural Sciences and Natural Resources University of Khuzestan [Grant no. 9428403]. We acknowledge Professor Hsin Chi for his technical assistance.

#### REFERENCES

- (IRAC) Insecticide Resistance Action Committee. 2020. IRAC MoA Classification Scheme. Version 9.4 (https://iraconline.org/documents/moaclassification/)(Accessed March 2020).
- Angeli, G. I. N. O., Baldessari, M. A. R. I. O., Maines, R. O. M. A. N. O. and Duso, C. 2005. Side-Effects of Pesticides on the Predatory Bug Orius laevigatus (Heteroptera: Anthocoridae) in the Laboratory. Biocontr. Sci. Tech., 15(7): 745-754.
- Arnó, J. and Gabarra, R. 2011. Side Effects of Selected Insecticides on the *Tuta absoluta* (Lepidoptera: Gelechiidae) Predators *Macrolophus pygmaeus* and *Nesidiocoris tenuis* (Hemiptera: Miridae). J. Pest Sci., 84(4), 513-520.
- Arno, J., Gabarra, R., Estopa, M., Gorman, K., Peterschmitt, M., Bonato, O. and Albajes, R. 2009. *Implementation of IPM Programs in European Greenhouse Tomato Production Areas: Tools and Constraints*. UDL Editions and Publications, Lleida, Spain.
- Biondi, A., Guedes, R. N. C., Wan, F. H. and Desneux, N. 2018. Ecology, Worldwide Spread, and Management of the Invasive South American Tomato Pinworm, *Tuta absoluta*: Past, Present, and Future. *Ann. Rev. Entomol.*, 63: 239-258.
- Bompard, A., Jaworski, C. C., Bearez, P. and Desneux, N. 2013. Sharing a Predator: Can an Invasive Alien Pest Affect the Predation on a Local Pest?. *Popul Ecol.*, 55(3): 433-440.
- Bostanian, N. J. and Akalach, M. 2006. The Effect of Indoxacarb and Five Other Insecticides on *Phytoseiulus persimilis* (Acari: Phytoseiidae), *Amblyseius fallacis* (Acari: Phytoseiidae) and Nymphs of *Orius insidiosus* (Hemiptera: Anthocoridae). *Pest Manag. Sci.*, 62(4): 334-339.
- Chi, H. 2017. TWOSEX-MSChart: A Computer Program for the Age-Stage, Two-Sex Life Table Analysis. National Chung Hsing University, Taichung, Taiwan. (http://140.120.197.173/Ecology/Download/ Twosex-MSChart.rar) (Accessed 1 May 2017).
- 9. Chi, H. and Getz, W. M. 1988. Mass Rearing and Harvesting Based on an Age-

Stage, Two-Sex Life Table: A Potato Tuber Worm (Lepidoptera: Gelechiidae) Case Study. *Environ. Entomol.*, **17**(1): 18-25.

- Chi, H. and Su, H. Y. 2006. Age-Stage, Two-Sex Life Tables of *Aphidius gifuensis* (Ashmead) (Hymenoptera: Braconidae) and Its Host *Myzus persicae* (Sulzer) (Homoptera: Aphididae) with Mathematical Proof of the Relationship between Female Fecundity and the Net Reproductive Rate. *Environ. Entomol.*, **35**: 10–21.
- Claver, M. A., Ravichandran, B., Khan, M. M. and Ambrose, D. P. 2003. Impact of Cypermethrin on the Functional Response, Predatory and Mating Behavior of a Non-Target Potential Biological Control Agent *Acanthaspis pedestris* (Stål)(Het., Reduviidae). J. Appl. Entomol., 127(1): 18-22.
- Copping, L. G. and Menn, J. J. 2000. Biopesticides: A Review of Their Action, Applications and Efficacy. *Pest Manag. Sci.*, 56(8): 651-676.
- Derbalah, A. S., Morsey, S. Z. and El-Samahy, M. 2012. Some Recent Approaches to Control *Tuta absoluta* in Tomato under Greenhouse Conditions. *Afr. Entomol.*, 20(1): 27-34.
- Desneux, N., Decourtye, A. and Delpuech, J. M. 2007. The Sublethal Effects of Pesticides on Beneficial Arthropods. *Ann. Rev. Entomol.*, 52: 81-106.
- Glunt, K. D., Paaijmans, K. P., Read, A. F. and Thomas, M. B. 2014. Environmental Temperatures Significantly Change the Impact of Insecticides Measured Using WHOPES Protocols. *Malar. J.*, 13(1): 350.
- Gontijo, L. M., Celestino, D., Queiroz, O. S., Guedes, R. N. C. and Picanço, M. C. 2015. Impacts of Azadirachtin and Chlorantraniliprole on the Developmental Stages of Pirate Bug Predators (Hemiptera: Anthocoridae) of the Tomato Pinworm *Tuta absoluta* (Lepidoptera: Gelechiidae). *Fla. Entomol.*, **98(1)**: 59-64.
- Goodman, D. 1982. Optimal Life Histories, Optimal Notation, and the Value of Reproductive Value. Am. Nat., 119: 803-823
- Hassan, S.A., Bigler, F., Bogenschütz, H., Boller, E., Brun, J., Calis, J. N. M. and Helyer, N. 1994. Results of the Sixth Joint Pesticide Testing Programme of the IOBC/WPRS-Working Group «Pesticides

and Beneficial Organisms». *Entomophaga*, **39(1)**: 107-119.

- He, Y., Zhao, J., Zheng, Y., Desneux, N. and Wu, K. 2012. Lethal Effect of Imidacloprid on the Coccinellid Predator *Serangium japonicum* and Sublethal Effects on Predator Voracity and on Functional Response to the Whitefly *Bemisia tabaci. Ecotoxicol.*, 21: 1291–1300.
- 20. Isman, M. B. 2017. Bridging the Gap: Moving Botanical Insecticides from the Laboratory to the Farm. *Indus. Crops Product.*, **110**: 10-14.
- Jaworski, C.C., Bompard, A., Genies, L., Amiens-Desneux, E. and Desneux, N. 2013. Preference and Prey Switching in a Generalist Predator Attacking Local and Invasive Alien Pests. *PLoS One*, 8(12): e82231.
- Jaworski, C.C., Chailleux, A., Bearez, P. and Desneux, N. 2015. Apparent Competition between Major Pests Reduces Pest Population Densities on Tomato Crop, But Not Yield Loss. J. Pest Sci., 88(4): 793-803.
- Juliano, S. A. 2001. Non-Linear Curve Fitting: Predation and Functional Response Curves. In: "Design and Analysis of Ecological Experiments", (Eds.): Scheiner, S. M. and Gurevitch J. 2<sup>nd</sup> Edition, Chapman and Hall, New York, USA, PP. 178–196.
- 24. Li, D.X., Tian, J. and Shen, Z.R. 2006. Effects of Pesticides on the Functional Response of Predatory Thrips, *Scolothrips takahashii* to *Tetranychus viennensis*. J. *Appl. Entomol.*, **130**: 314–322.
- 25. Lima, D. B., Melo, J. W. S., Guedes, N. M. P., Gontijo, L. M., Guedes, R. N. C. and Gondim Jr, M. G. C., 2015. Bioinsecticide-predator Interactions: Azadirachtin Behavioral and Reproductive Impairment of the Coconut Mite Predator *Neoseiulus baraki. P. Lo. S. One*, **10**(2): e0118343.
- Malaquias, J. B., Ramalho, F. S., Omoto, C., Godoy, W. A. C. and Silveira, R. F. 2014. Imidacloprid Affects the Functional Response of Predator *Podisus nigrispinus* (Dallas) (Heteroptera: Pentatomidae) to Strains of *Spodoptera frugiperda* (J. E. Smith) on Bt Cotton. *Ecotoxicol.*, 23: 192– 200.
- 27. Margaritopoulos, J. T., Tsitsipis, J. A. and Perdikis, D. C. 2003. Biological Characteristics of the Mirids *Macrolophus costalis* and *Macrolophus pygmaeus* Preying on the Tobacco form of *Myzus persicae*

[ Downloaded from jast.modares.ac.ir on 2025-07-30 ]

(Hemiptera: Aphididae). *Bul. Entomol. Res.*, **93(1)**: 39-45.

- Martinou, A. F. and Stavrinides, M. C. 2015. Effects of Sublethal Concentrations of Insecticides on the Functional Response of Two Mirid Generalist Predators. *PLoS One*, 10(12): e0144413.
- Martinou, A. F. and Stavrinides, M. C. 2015. Effects of Sublethal Concentrations of Insecticides on the Functional Response of Two Mirid Generalist Predators. *PloS One*, **10(12)**, e0144413.
- Martinou, A.F., Seraphides, N. and Stavrinides, M. C. 2014. Lethal and Behavioral Effects of Pesticides on the Insect Predator *Macrolophus pygmaeus*. *Chemosphere*, 96: 167-173.
- Michalko, R. and Košulič, O. 2016. Temperature-Dependent Effect of Two Neurotoxic Insecticides on Predatory Potential of *Philodromus* Spiders. *J. Pest Sci.*, 89(2): 517-527.
- Nazarpour, L., Yarahmadi, F., Saber, M. and Rajabpour, A. 2016. Short and Long Term Effects of Some Bio-Insecticides on *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) and Its Coexisting Generalist Predators in Tomato Fields. *J. Crop Protect.*, 5(3): 331-342.
- Pilar Marco, M., Pascual, N., Belles, X., Camps, F. and Messeguer, A. 1990. Ecdysteroid Depletion by Azadirachtin in *Tenebrio molitor* Pupae. *Pestic. Biochem. Physiol.*, 38: 60-65.
- 34. Rahmani, S. and Bandani, A. R. 2013. Sublethal Concentrations of Thiamethoxam Adversely Affect Life Table Parameters of the Aphid Predator, *Hippodamia variegata* (Goeze) (Coleoptera: Coccinellidae). *Crop Protec.*, 54: 168-175.
- Rogers, D. 1972. Random Search and Insect Population Models. J. Anim. Ecol., 41: 369-383.
- 36. Sharifian, I., Sabahi, Q. and Bandani, A. R. 2017. Effect of Some Conventional Insecticides on Functional Response Parameters of *Macrolophus pygmaeus* (Hem.: Miridae) on *Tuta absoluta* (Lep.: Gelechiidae). *Biharean Biol.*, **11**(1): 10-14.
- Smith, R. J. 1991. Integration of Biological Control Agents with Chemical Pesticides. In: "Microbial Control of Weeds", (Eds.):

Charudattan R. and Walker, H. L. Springer, Boston, USA, PP. 189-208.

- Sylla, S., Brévault, T., Diarra, K., Bearez, P. and Desneux, N. 2016. Life-History Traits of *Macrolophus pygmaeus* with Different Prey Foods.: *PLoS One*, **11(11)**: e0166610. <u>https://doi.org/10.1371/journal.pone.016661</u> <u>0</u>.
- Tedeschi, R., Alma, A. and Tavella, L. 2001. Side-Effects of Three Neem (*Azadirachta indica* A. Juss) Products on the Predator *Macrolophus caliginosus* Wagner (Het., Miridae). J. Appl. Entomol., 125(7): 397-402.
- Tibshirani, R. J. and Efron, B. 1993. An Introduction to the Bootstrap. *Monogr. Statis. Appl. Probab.* 57: 1-436.
- 41. Wing, K. D., Sacher, M., Kagaya, Y., Tsurubuchi, Y., Mulderig, L., Connair, M. and Schnee, M. 2000. Bioactivation and Mode of Action of the Oxadiazine Indoxacarb in Insects. *Crop Protect.*, **19(8-10)**: 537-545.
- 42. Wright, D. J. and Verkerk, R. H. 1995. Integration of Chemical and Biological Control Systems for Arthropods: Evaluation in a Multitrophic Context. *Pest Manag. Sci.*, **44(3)**: 207-218.
- 43. Yarahmadi, F., Mossadegh, M.S., Soleymannejadian, E., Saber, M. and Shishehbor, P. 2009. Assessment of Acute Toxicity of Abamectin, Spinosad and Chlorpyrifos to *Thrips tabaci* Lindeman (Thysanoptera: Thripidae) on Sweet Pepper by Using Two Bioassay Techniques. *Asian J. Biol. Sci.*, 2(3): 81-87.
- Zappalà, L., Biondi, A., Alma, A., Al-Jboory, I. J., Arnò, J., Bayram, A., Chilleux, A., El-Arnaouty, A., Gerling, D., Guenaoui, Y., Shaltiel-Harpaz, L/. Siscaro, G., Stavrinides, M., Tavella, L., Aznar, R.V., Urbenja, A. and Desneux, N., 2013. Natural Enemies of the South American Moth, *Tuta absoluta*, in Europe, North Africa and Middle East, and Their Potential Use in Pest Control Strategies. J. Pest Sci., 86: 635–647.
- 45. Zhang, P., Liu, F., Mu, W., Wang, Q., Li, H. and Chen, C. 2014. Life Table Study of the Effects of Sublethal Concentrations of Thiamethoxam on *Bradysia odoriphaga* Yang and Zhang. *Pesticide Biochem. Physiol.*, **111:** 31-37.

#### - Safari Monjezi et al.



# اثرات زیر کشنده سموم رایج مورد استفاده در گوجهفرنگی روی واکنش تابعی و پارامترهای زیستی سن Macrolophus pygmaeus Rumber ( (Miridae)

ی. صفری منجزی، ف. یاراحمدی، و ن. زندی سوهانی

چکیدہ

سن Macrolophus pygmaeus Rumber عامل موثر کنترل بیولوژیک علیه آفات بسیاری از محصولات از جمله مينوز برگ گوجه فرنگی Tuta absoluta Meyrick می باشد. اثر ات زیر کشنده حشره کش های آزادیراختین (نیم آزال<sup>®</sup>)، ایندو کساکارب (آوانت<sup>®</sup>) و امامکتین بنزو آت+لوفنو رون (پروکلیم فیت<sup>®</sup>) روی خصوصیات زیستی و واکنش تابعی این سن شکارگر در شرایط آزمایشگاهی در دمای ۲۵±۱، رطوبت نسبی ۴۵±۱٪ و دوره روشنایی تاریکی ۱۶: ۸ ساعت، مورد بررسی قرار گرفت. برای این منظور، مادههای این سن شکارگر در معرض بقایای زیرکشنده (۱۰٪ غلظت مزرعهای) این حشره کش ها قرار گرفتند. روش جدول زندگی دوجنسی و مدل راجرز به ترتیب برای بررسی تاثیرات این حشره کش ها روی یارامترهای زیستی و واکنش تابعی شکارگر مذبور مورد استفاده قرار گرفت. نتایج نشان داد که بقایای زیر کشنده ایندو کساکارب و آزادیراختین به ترتیب بیشترین و کمترین اثرات جانبی را روی پارامترهای جدول زندگی M. pygmaeus داشتند. در حالیکه که آزادیراختین تاثیر معنی داری روی یارامترهای جدول زندگی این سن نداشت، زادآوری کل و طول عمر این شکارگر در تيمار ايندوكساكارب به ترتيب به صورت معنىدارى كمتر (۱۴/۶٪) و بيشتر (۶/۹٪) از تيمار شاهد بود. این شکارگر واکنش تابعی نوع دوم را نشان داد. نوع واکنش تحت تاثیر بقایای حشرهکشرها قرار نگرفت. با این حال، زمان دسترسی به طعمه در تیمار امامکتین بنزوات+لوفنورون (۱/۷۴ ساعت) به صورت معنیداری طولانی تر از سایر تیمارها بود؛ ولی اختلاف معنیداری بین نرخ حمله (a) در تیمارهای مختلف موجود نداشت. در کل، این آزمایشات ثابت نمود که آزادیراختین یک حشره کش کم خطر برای M. pygmaeus است که می توان با کنترل بیولوژیک توسط این سن تلفیق گردد.