

## Effect of Deficit Irrigation Technique on Black Cumin (*Nigella sativa* L.) Water Use Efficiency

T. Yaghi<sup>1\*</sup>, A. Arslan<sup>2</sup>, and H. Saeed<sup>3</sup>

### ABSTRACT

To increase the agricultural profit, an experiment was conducted at the Scientific Agricultural Research Center (SARC) at Salamiyah, Syria. The experiment aimed to estimate the optimum irrigation level that maximizes productivity as well as the black cumin's quality local quality standards under experimental conditions. The irrigation treatments were 25, 50, 75 and 100% of the Potential Evapotranspiration ( $ET_p$ ) based on Class-A pan evaporation. The treatments were denoted as B, C, D and E, respectively. The no-irrigation treatment (A) was considered as control. Irrigation water was applied by a drip system. The A-E treatments were designed and implemented using complete randomized block design (CRBD) with four replications. The actual Evapotranspiration ( $ET_a$ ) was calculated after calibration of soil using SPAW software. Furthermore, crop yield, Irrigation Water Use Efficiency (IWUE), yield response factor ( $k_y$ ) and some parameters (period of growth stage, date of flowering...etc.) were studied during 2018-2019 successive growth seasons. Results showed that irrigation could be scheduled using some equations and Class-A pan evaporation. Moreover, the vegetative growth parameters flourished virtually and significantly by comparing higher and lower irrigation levels. However, Irrigation Water Use Efficiency (IWUE) values increased by reducing the applied irrigation water. Treatment B recorded the highest IWUE value (2.4 kg ha<sup>-1</sup> mm<sup>-1</sup>), but key values remained less than 1.0, indicating that the plant tolerates drought. Furthermore, treatment C scored the top-seed (130.6%) concerning the profit/total costs ratio. Finally, based on the results, we recommend that irrigation should not be more than 75% and not less than 25% of the  $ET_p$ .

**Keywords:** Class-A pan evaporation, Potential evapotranspiration, Yield response factor ( $k_y$ ).

### INTRODUCTION

Since spreading the Sustainable Development (SD) concept all over the world, many agricultural researchers have called for achieving the highest productivity of the least water unit declaring the following motto: "more crop per drop" (Molden *et al.*, 2010). Salamiyah is a rural district located in the Al-Harron river sub-basin at the Orontes Basin, Syria, and has a semi-arid climate with limited annual precipitation ranging from 150 to 420 mm (Yaghi *et al.*, 2016). Due to insufficient

water resources, dry farming is practiced in most of the aforementioned areas. Therefore, efficient use of scarce water resources for sustainable agricultural water management in Salamiyah is a serious challenge. So, there is an increasing need to replace crops with high irrigation requirement (cotton, maize and alfalfa) with the low irrigation requirement; therefore, high economic return crops such as black cumin, chamomile, and thyme have become popular (Serman *et al.*, 2021; Mauget *et al.*, 2022).

Black cumin (*Nigella sativa*, L.) has been used as traditional natural medicine for

<sup>1</sup> General Commission for Scientific Agricultural Research (GCSAR), Syria.

<sup>2</sup> Department of Natural Resources and Environmental Science, University of Nevada, Reno, NV 89512, USA.

<sup>3</sup> Scientific Agricultural Research Center (SARC), Salamiyah, Hama, Syria.

\* Corresponding author; e-mail: tammam.yaghi@gmail.com



centuries. The crude oil and Thymoquinone (TQ) extracted from its seeds and oil are effective against many diseases like cancer, cardiovascular complications, diabetes, asthma, kidney disease, etc. It is effective against cancer in the blood system, lung, kidney, liver, prostate, breast, cervix, skin, with much safety (Khan *et al.*, 2011). Identifying critical growth stages for water of a particular cultivar under local conditions of climate and soil fertility allows irrigation scheduling to maximize crop yield and increase the Water Use Efficiency (WUE) in areas with scarce water resources (Mahal and Sidhu, 2006; Kumar *et al.*, 2019). There are different methods to estimate the irrigation requirements. One of them is by using potential evapotranspiration based on Class-A Pan (Allen *et al.*, 1998; Ruhi *et al.*, 2006). Prolonging the irrigation intervals reduces the growth and yield of various medicinal and aromatic plants (Hassan and Ali, 2014).

The increase in irrigation water increased the number of primary, secondary and tertiary branches, number of capsules per plant, capsule length, diameter of capsule, number of seeds per capsule, fresh and dry seed yield per plant, 1,000-seed weight, and seed yield of black cumin (Karim *et al.*, 2017).

On the contrary, Elsafi (2003) found that the black cumin yield increased by prolonging the irrigation intervals and water stress applications, while the lower WUE was associated with a higher amount of irrigation water (Bondok and El-Sharkawy, 2014). Deficit Irrigation (DI) is one of the water management practices for increasing WUE. Ghamarnia *et al.* (2010) have shown that black cumin tolerates drought. They have found that irrigation at the rate of 50% of (crop Evapotranspiration, ET) achieved high crop productivity in Iran. Likewise, for the cumin crop, Bondok and El-Sharkawy (2014) have proved that as water decreased from 1.0 to 0.8 and 0.6 of ET<sub>c</sub>, the cumin yield decreased by (14.05 and 14.25%) and by (25.6 and 26.89%) in two seasons, respectively. In the same study, WUE

increased by (5.26% and 19.30%) and by (10.17% and 16.9%) compared with the full irrigation treatment, respectively.

Due to the consumptive needs of the rapidly increasing world population, black cumin has a quite large market, which is about \$60 billion (Kumar, 2009). Furthermore, because of the great importance of black cumin (*Nigella sativa* L.) plants as natural sources for producing fixed and volatile oils, more investigations for improving the growth and productivity of this plant are still needed (Ghamarnia *et al.*, 2014). In this context, there are some investigations in the field of sustainable irrigation in Syria concerning the application of virtual water concept and alternative crops. These include replacement of crops of low economic return (such as alfalfa, wheat, and cotton) with crops of high economical return (like garlic, grapes and medical plants: black cumin, thyme...etc.) to cope with water shortage, irregular demographic issues (localizing people according to safer areas because of the Syrian war), and high costs of farming (Mourad, 2012).

On the grounds of the previous scientific reports, it will be beneficial to determine the optimum irrigation level of black cumin plant to increase its productivity per unit of applied water. Therefore, the goal of this study was to estimate the optimum irrigation level, which maximizes the productivity as well as the crop quality under experimental conditions.

## MATERIALS AND METHODS

The experiment was carried out at SARC at Salamiyah district (35° 00' N, 37 02' E and 480.8 m altitude), Hama Governorate, Syria. It forms about 37.1% of available arable land of Hama governorate; the average annual precipitation throughout the growing season (Feb-Jun, 2018-2019) was only 143 mm, while the total evaporation during that period was about 645.7 mm. Therefore, irrigation is essential for optimal crop yield and good quality. In this region,

the largest percentage of water consumption in agriculture is supplied by groundwater, which are decreasing rapidly, and groundwater quality is deteriorating. Since the experimental area is far from the sea and close to Syrian Badia, it has a semi-arid climate. The physical and chemical properties of the soil are presented in Tables 1 and 2. Available soil water holding capacity was 109 mm at 0-60 cm depth. Furthermore, hand Beerkan infiltration methods have been used and their results in the same type of soil were compared with SPAW software results by Saxton and Willey (2006), similar data were reported by (Mubarak *et al.*, 2009; Angulo-Jaramillo *et al.*, 2019).

Thus, for accurate selection of the irrigation time, Management Allowable Depletion (MAD) value of about 65% was used, based on some previous researches that indicated that this crop could tolerate water shortage (Ghamarnia *et al.*, 2010; Ghamarnia and Jalili, 2013).

Daily climate data were collected from Feb. 10 until Jun 29 during 2018-2019 growing seasons. Furthermore,  $ET_0$  was calculated by using two programs (ET0 Calculator and New\_LocClim software), and monthly values are shown in Table 3.

### Experimental Treatments, Planting, Cultural Practices and Design of Drip Irrigation System

The study was carried out using Complete Randomized Block Design (CRBD) with four replications. Experimental treatments were A (without irrigation), B (25% of  $ET_p$ ), C (50%), D (75%), and E (100%). The experimental area consisted of 20 plots. The area of each plot was 5 m<sup>2</sup> (2.5\*2 m), and the total area was 157.3 m<sup>2</sup> (14.3\*11 m). In order to prevent the interaction between irrigation treatments, 1 m space was left between the plots. Besides, two rows in each plot were left out of the assessment due to the edge's effect, and the remaining area formed the harvest plots, as shown in Figure 1.

Figure 1 shows that there are 4 drip lines for 5 cultivation lines in each plot. Each line was equipped with external drippers, except treatment A. The distance between two lines was 0.5 m. Based on technical advice received from soil lab staff, K (Potassium oxide, K<sub>2</sub>O) was not needed while the fertilizers of N (100 kg ha<sup>-1</sup>) and P (Super phosphate, P<sub>2</sub>O<sub>5</sub>) (60 kg ha<sup>-1</sup>) were needed. Half of the total nutrient amount (fertilizers) was given during soil preparation, and the rest was given with the fertigation technique

**Table 1.** Physical properties of the soil.

Soil depth (cm)	Bulk density (g cm <sup>-3</sup> )	Texture <sup>a</sup>	Field capacity (%v)	Wilting point (%v)	Water holding capacity (%)	(mm)	Clay (%)	Silt (%)	Sand (%)
0-30	1.41	CL	38.1	25.7	12.4	52.45	42	20	38
30-60	1.29	SCL	39.9	25.3	14.6	56.5	45	42.3	12.7
60-90	1.24	SCL	42.1	28.8	13.3	49.47	49	43	8

<sup>a</sup> CL: Clay, SCL: Silty Clay.

**Table 2.** Chemical properties of the soil.<sup>a</sup>

Soil depth (cm)	pH (1:5)	EC (1:5) (dS m <sup>-1</sup> )	Avail-P (mg kg <sup>-1</sup> )	Avail-K (mg kg <sup>-1</sup> )	Mineral-N (mg kg <sup>-1</sup> )	CaCO <sub>3</sub> (%)	OM (%)
0-30	7.75	0.37	38.2	504	1.34	7.2	1.19
30-60	8.03	0.32	39.2	430	1.89	4.3	1.38
60-90	7.43	0.49	19.4	380	2.1	3.3	1.34


<sup>a</sup> 1:5: Saturated soil paste; P: Phosphorus; K: Potassium, N: Nitrogen.

**Table 3.** Climate data records taken monthly for short and long-term periods.

Month	Solar (h d <sup>-1</sup> )	Temp (°C)		Rainfall (mm)	Relative humidity (%)	Wind (m s <sup>-1</sup> )	E <sub>pan</sub> (mm)	ET <sub>0</sub> <sup>a</sup> (mm)
		Max	Min					
Feb	5.1	17	2.7	76	83	3.3	36.7	20.3
Long-term <sup>b</sup>	4.13	12	2.4	32.3	87	3.7	-	19.4
Mar	8.4	18.1	5.8	18.4	81	3.4	60.7	60.7
Long-term	5.59	16.7	3.9	56.7	89	3	-	62.7
Apr	8.7	26.3	10.4	45	77	4.2	108.9	89.9
Long-term	8.12	26.1	11.9	42.4	76	4.5	-	82.7
May	11.4	33.1	15.9	19.6	64	3.1	185.2	114.4
Long-term	11.1	33.6	15	22	58	4.3	-	112.6
Jun	12.2	33.2	17.7	1.4	53	2.2	190	114.9
Long-term	12.1	34.1	17.2	1.2	48	1.7	-	115.8
Total average				160.4			581.5	400.2

<sup>a</sup> FAO Penman-Monteith method, Feb. 10-28. . . Until Jun 29, using ET<sub>0</sub> Calculator software (2018-2019).

<sup>b</sup> Between the years 1970 and 2020 using New\_LocClim software.

**Figure 1.** Layout of black cumin experiment and irrigation system at SARC, Hama, Syria.

as equal amounts during the irrigation period by dividing into the number of irrigations. The black cumin seeds were planted with a density of 4 kg ha<sup>-1</sup>. Farming operations such as reseeding, weeding, and irrigation were done as required. Moreover, pesticides were not used during the two 2018-2019 growing seasons.

### Irrigation Water, Evapotranspiration, and Water-Yield Functions

Evapotranspiration was estimated from the water balance Equation (1):

$$ET = I + R \pm \Delta S_m - D_r \quad (1)$$

Where, ET is crop Evapotranspiration (mm), I is the amount of applied Irrigation water

(mm), R is precipitation (mm),  $\Delta S_m$  is any change in Soil moisture content (mm) and Dr is Deep percolation water (mm), which was assumed to be zero. The method given by Allen *et al.* (1998) was chosen for calculating the amount of irrigation water:

$$ET_p = K_p \times E_{pan} \quad (2)$$

$$ET_c = K_c \times ET_p \quad (3)$$

Where,  $ET_p$  is Potential Evapotranspiration (mm),  $K_p$  is pan coefficient ranged 0.74-0.77 based on the principles given by Doorenbos and Pruitt (1977),  $E_{pan}$  is pan Evaporation (mm),  $ET_c$  is crop Evapotranspiration (mm) and  $K_c$  is crop coefficient. Four crop coefficients were picked up (0.52, 0.91, 1.27 and 0.77) for different black cumin growth stages (initial, development, mid-stage, and late), respectively, based on the scientific report of Ghamarnia *et al.* (2014).

The volume of irrigation water requirement for each treatment was calculated according to the squared area:

$$I = ET_c \times S_d \quad (4)$$

Where,  $S_d$  is irrigation water ratio ( $B = 25\%$ ;  $C = 50\%$ ;  $D = 75\%$ ;  $E = 100\%$ ).

$$WUE = Y / (I + P_e) \quad (5)$$

$$IWUE = (Y - Y_0) / IWU \quad (6)$$

Where,  $I$  is Irrigation water amount (mm),  $P_e$  is effective rainfall (mm) =  $0.6 P$  (Precipitation, mm) if  $P < 75\%$ ,  $P_e = 0.8 P$  if  $P > 75$  mm. month<sup>-1</sup>,  $Y$  is seed Yield obtained from the treatments (kg ha<sup>-1</sup>),  $Y_0$  is seed Yield obtained from treatment (A) (kg ha<sup>-1</sup>), and  $IWU$  is Irrigation Water Use (mm).

The yield response factor ( $k_y$ ) representing the decrease in black cumin seed yield, which may occur as a result of the decrease in the unit evapotranspiration was determined using Equations (2), (3), and (4) (Doorenbos and Kassam, 1986):

$$(1 - Y_a / Y_m) = k_y (1 - ET_a / ET_m) \quad (7)$$

Where,  $Y_a$  is actual seed Yield regarding the treatments (kg ha<sup>-1</sup>),  $Y_m$  is maximum seed Yield (kg ha<sup>-1</sup>),  $ET_a$  is actual seasonal Evapotranspiration (mm), and  $ET_m$  is maximum seasonal Evapotranspiration (mm).

The  $k_y$  values are crop specific and vary over the growing season according to growth stages of the report by Steduto *et al.* (2012):  $k_y < 1$ : Crop is more tolerant to water

deficit, and recovers partially from stress, exhibiting less than proportional reductions in yield with reduced water use.

### The Reproductive and Vegetative Parameters

The collected data for this item were the plant height, number of branches per plant, dry matter, harvest index, number of the mature and dry follicles per plant, the follicle's diameter, number of the seeds per follicle, weight of the 1000 seeds, depth of the roots, yield of the seeds, and Water Use Efficiency (WUE). Furthermore, five plants were used to determine the dry weight per plant. The plants were dried and weighed using a sensitive balance. Harvest index was calculated as follow:

$$HI = [Y / (Ps + Y)] \times 100 \quad (8)$$

Where,  $HI$  is Harvest Index (%),  $Y$  is an economical seed Yield (kg ha<sup>-1</sup>),  $Ps$  is straw yield (kg ha<sup>-1</sup>).

We had employed the univariate analysis of variance using a statistical package for social science (SPSS software) to analyze the collected data. The difference between treatment means was compared by Duncan's Multiple Range Test (DMRT) at 0.05 statistical significance level. Furthermore, an economical study was performed by determining the ratio between the profit and the total costs for each treatment.

## RESULTS AND DISCUSSION

### Impact of Water Stress on Black Cumin Growth

#### Black cumin Water Consumptive Use, $ET_a$

Daily black cumin water consumptive use values (actual water requirements) have differed according to the phenological phases (local quality standards compared to



what area farmer does) and treatments. For example, in treatment B, the  $ET_a$  value was  $0.9 \text{ mm d}^{-1}$  at initial stage and increased to  $3.8 \text{ mm d}^{-1}$  at mid-season stage, then, lowered to  $3.33 \text{ mm d}^{-1}$  again at the late and the harvest stage in the first season. The same thing in the second season but with simple differences because of the changes concerning climate and soil. Furthermore, the period of black cumin phenological phases has differed and recorded the best quality standard at the treatment B (23 days for initial phase, 42 days for development phase, 40 days for Mid-season phase and 25 days for late and harvest phase) compared to the rest of the other treatments (E, 140 days and A, 120 days) as shown in Figure 2.

The sum of black cumin water consumptive use values during two seasons

were [488.4 (1) and 392.2 mm (2) at (E); 369.5 and 376.3 mm at (D); 290.4 and 300.6 mm at (C); 285.6 and 284.9 mm at (B); and 275.8 and 273.2 mm at (A)] as given in Table 4. These differences can be attributed to the impact of irrigation on roots depth and the corresponding ground canopy above that differed significantly between the treatments. It can be clarified through the water stress, which has led to a lack of absorption of nutrients in plant due to a lack of movement of nutrients in dry land, an inadequate water absorption and lack of extension and spread of roots (Beudez and Doussan, 2013).

That has been accompanied with a significant difference for the other studied plant attributes (plant height, branches number, dry matter and harvest




Figure 2. Black cumin water consumptive use ( $ET_a$ ) at the optimal treatment (B) which achieves quality standards represented in phenological phases compared to the rest of the treatments during two seasons.

**Table 4.** Vegetative growth characters of black cumin as affected by different water regimes based on class-A pan during the two experimental seasons.<sup>a</sup>

ET <sub>p</sub> (%)	Water consumptive use (mm)	Plant height (cm)	Branches number	Dry matter per plant (g)	Roots depth (cm)	Harvest Index (HI) (%)
First season-2018						
No irrigation (A)	275.8	10.32 d	6.50 e	0.57 e	8.30 d	33.95 d
25 (B)	285.6	27.92 c	11.30 d	1.31 d	18.03 b	48.50 c
50 (C)	290.4	28.20 c	18.73 c	1.75 c	19.43 b	53.00 b
75 (D)	369.5	33.90 ba	24.48 a	2.13 ba	17.23 c	56.25 a
100 (E)	488.4	38.35 a	23.00 ba	2.29 a	29.78 a	48.40 c
Second season-2019						
No irrigation (A)	273.2	13.38 e	8.90 d	0.93 d	7.90 d	36.05 dc
25 (B)	284.9	21.73 d	17.30 c	1.61 c	19.38 c	38.00 c
50 (C)	300.6	35.00 c	24.12 b	2.414 a	18.68 c	48.75 b
75 (D)	376.3	39.15 ba	32.58 a	2.57 a	21.13 ba	52.50 a
100 (E)	392.2	43.20 a	32.75 a	2.366 ba	31.98 a	49.10 b

<sup>a</sup> Means followed by different letter (s) within a parameter differed significantly by DMRT at  $P \leq 5\%$ .

index) at E and D compared to the rest of the other treatments as given in Table 4.

The gradual reduction in the dry weight under DI could be a result of a reduction in the chlorophyll content, and consequently, photosynthesis efficiency. This result was supported by Khalid (2006). Water is a very important factor, affects the growth of the black cumin plant when the water content of the plant decreases, its cells shrink and the cell walls relax which results in lower turgor pressure (Csonka, 1989). The principles that underlie the two processes are similar because leaf expansion depends mostly on cell expansion (Taiz and Zeiger, 2002). The larger the leaf area is the more the transpiration will be. Growth reduction because of DI has been widely reported (Karim *et al.*, 2017; Mehriya *et al.*, 2020). We cannot ignore the effect of precision irrigation like DI on the flowering stage that has shown multiplying of the follicles and seeds in each one as given in Table 5. These results are also in agreement with those of Haj Seyed Hadi *et al* (2012).

The results of the study indicated that DI treatments remarkably increased the yield, whereas the weight of one thousand seeds increased from A to E ( $A < B < C < D < E$ ) as given in Table 6. Similar data were reported by Banon *et al* (2003).

On the other hand, the efficiency of sustainable irrigation does not agree with the aforementioned results. The maximum IWUE (2.4 and 2.28  $\text{kg ha}^{-1} \text{mm}^{-1}$ ) and (2.21 and 2.25  $\text{kg ha}^{-1} \text{mm}^{-1}$ ) were obtained by B and C treatments, respectively, while the lowest values in this respect (1.83 and 1.89  $\text{kg ha}^{-1} \text{mm}^{-1}$ ) were recorded for treatment (E) in both seasons, respectively. The lower WUE associated with a higher amount of irrigation water could be attributed to a greater loss of water by ET than the corresponding increase in seed yield (Al-Kayssi *et al.*, 2011). These results are also in agreement with those of Senyigit and Arslan (2018), who concluded that the DI technique based on Class-A pan remarkably affects applied water and water use efficiency.

#### Relation between the Yield and Irrigation Water

Results have shown that there are linear relations between the amount of irrigation water and the yield ( $R^2 = 0.820$ ), and between the evapotranspiration and the yield ( $R^2 = 0.804$ ) at 0.05 significance level. Although the black cumin yield could be achieved even in the non-irrigated conditions, the equations have shown that



**Table 5.** Black cumin seeds yield components as affected by different water regimes based on Class-A pan during the two experimental seasons.<sup>a</sup>

ET <sub>p</sub> (%)	Diameter of follicle (cm)	Follicles per plant	Seeds per follicle	Empty follicles per plant	1000 seeds weight (g)	Seed yield (kg ha <sup>-1</sup> )	Irrigation Water Use Efficiency (IWUE) (kg ha <sup>-1</sup> mm <sup>-1</sup> )
First season-2018							
No irrigation (A)	0.54 d	4.53 d	12.60 e	6.10 a	1.02 d	170.3 e	-
25 (B)	0.72 c	8.08 c	31.50 d	4.20 b	1.47 c	346.0 d	2.4 a
50 (C)	0.82 b	13.80 b	60.00 c	3.00 c	1.76 b	603.8 c	2.21 b
75 (D)	0.82 b	14.73 a	79.63 b	1.40 d	1.89 b	689.2 ba	1.99 c
100 (E)	1.01 a	14.30 a	81.45 a	1.20 d	2.18 a	725.5 a	1.83 dc
Second season-2019							
No irrigation (A)	0.59 e	5.13 d	16.70 d	5.50 a	1.30 d	207.8 e	-
25 (B)	0.82 d	8.03 c	41.05 c	2.40 b	2.40 c	341.8 d	2.28 a
50 (C)	0.97 c	14.50 b	66.35 b	1.20 c	2.42 c	634.0 c	2.25 a
75 (D)	1.27 b	16.13 a	87.98 a	0.82 d	2.91 a	696.9 ba	2.03 b
100 (E)	1.05 a	15.35 a	88.35 a	0.82 d	2.61 b	723.1 a	1.89 c


<sup>a</sup> Means followed by different letter (s) within a parameter differed significantly by DMRT at P ≤ 5 %.

when irrigation water and ET increased, yield also increased. These results were in agreement with those found by Yousuf *et al* (2018), Zapotoczny *et al* (2019) on black cumin and Oweis and Hachum (2012) on the grain crop. Knowing these relations between the yield and irrigation water will give us a real perspective about the significance of precision irrigation in agriculture compared to traditional one. A thing which would lead us to save more water (Molden *et al.*, 2010).

The calculated values of yield response factor ( $k_y$ ) confirmed that the DI technique

was effective on black cumin. The  $k_y$  mean achieved about 0.43 in two seasons. Its highest values were obtained by the treatment (B) replicators recording (0.69, 0.72, 0.75 and 0.71). They gradually decreased to (0.17, 0.23, 0.21 and 0.14) in the treatment (D) as shown in Figure 3. Similar data were reported by Ozer *et al.* (2020).

Considering the scientific reports issued by FAO 36 (Steduto *et al.*, 2012), we

**Figure 3.** Average black cumin ( $k_y$ ) values under (DI) technique during two seasons.



**Table 6.** The economical study of the field trials of black cumin during 2018-2019 seasons.<sup>a</sup>

ET <sub>p</sub> (%)	Yield (kg. ha <sup>-1</sup> )	Product value (\$. ha <sup>-1</sup> )	Irrigation amount (mm. ha <sup>-1</sup> )	Irrigation n costs (\$. ha <sup>-1</sup> )	Uniform costs (\$. ha <sup>-1</sup> )	Harvest costs (\$. ha <sup>-1</sup> )	Total costs (\$. ha <sup>-1</sup> )	Profit (\$. ha <sup>-1</sup> )	Profit to total cost (%)
First season-2018									
No irrigation (A)	170.3 e	459.8 e	-	-	119.0	100.0	219	240.8 d	109.9
25 (B)	346.0 d	934.2 d	145.7	123.6	210.0	100.0	433.6	500.6 c	115.5
50 (C)	603.8 c	1630.3 c	273.0	231.5	345.0	130.4	706.9	923.4 b	130.6
75 (D)	689.2 ba	1860.8 b	346.2	293.5	390.0	148.9	832.4	1028.4 a	123.5
100 (E)	725.5 a	1958.8 a	395.6	335.5	410.0	156.7	902.2	1056.6 a	117.1
Second season-2019									
No irrigation (A)	207.8 e	602.5 d	-	-	180.5	110.0	290.5	312.0 d	107.4
25 (B)	341.8 d	991.1 c	150.0	144.0	210.3	115.0	469.4	521.8 c	111.2
50 (C)	634.0 c	1839.0 b	281.7	270.4	381.6	147.1	799.1	1039.9 b	130.1
75 (D)	696.9 ba	2012.0 a	343.9	330.2	415.3	161.7	907.2	1104.8 a	121.8
100 (E)	723.1 a	2097.0 a	381.0	365.8	431.9	167.8	965.5	1131.5 a	117.2

<sup>a</sup> Means followed by different letter (s) within a parameter differed significantly by DMRT at P ≤ 5 %.



noticed that the crop yield response factor was less than one, which means that the crop can tolerate water shortage and is suitable to dry areas such as our study site. Consequently, we recommend that the deficit irrigation programs should be applied in the regions with limited water resources in order to increase the yield for black cumin plant, which is widely cultivated under rainfed (non-irrigated) conditions.

### Black Cumin Economical Feasibility under the Impact of DI

Results of statistical analyses have indicated that the black cumin was successful economically and beneficial in dry environments, but the best profit will be achieved only after the application of deficit irrigation. The ratio (profit/total costs) value was higher at C and D than the rest of the treatments. Treatment C excelled all the treatments achieving 130.6 and 130.1% in both seasons, respectively, as given in Table 6. Therefore, we recommend not to fully irrigate the crop in this type of environments.

### CONCLUSIONS

With growing water demand and increasing signs of water scarcity, there is an urgent need to achieve higher output per unit of water consumed. Fortunately, there is an extensive range to improve crop water productivity, particularly in areas where water shortage aggravates such as Syria. In the present study, effects of the deficit irrigation technique (DI) on water applied, crop quality standards and WUE were investigated. The results of the study indicated that DI treatments remarkably decreased the applied water in the order of  $B < C < D < E$  and simultaneously increased WUE in the order of  $B > C > D > E$ .

The B treatment has attained the highest WUE of  $2.4 \text{ kg ha}^{-1} \text{ mm}^{-1}$ . The lowest

WUE ( $1.85 \text{ kg ha}^{-1} \text{ mm}^{-1}$ ) was at the treatment (E).

Reference evapotranspiration ( $ET_0$ ) value was calculated based on class-A pan method ( $ET_0 = ET_p$ ), which was recorded as the maximum value during June. Water consumptive started to increase from the date of sowing in February till Midseason stage and reached maximum in May then again reduced in the last Maturity and harvest stage in June. The yield response factor ( $k_y$ ) values decreased with increasing the irrigation water application. The average of ( $k_y$ ) was 0.43 during both seasons. That certainly indicates that black cumin tolerates water shortage.

Concerning the water-yield functions, we recommend the 50% of  $ET_p$  treatment because there has not been a significant reduction in productivity compared to the savings in water consumption. Furthermore, the results of economical analyses have confirmed that there is a high economical return after the application of DI.

### ACKNOWLEDGEMENTS

The authors are thankful for the support and encouragement of Dr. Majeda Mohammed Mofleh, the DG of the General Commission for Scientific Agricultural Research in Syria (GCSAR), and the director of Natural Resources Research Administration (ANRR).

### REFERENCES

1. Al-Kayssi, A., Shibab, R. M. and Mustafa, S. H. 2011. Impact of Soil Water Stress on Nigella Oil Content of Black Cumin Seed Grown in Calcareous-Gypsiferous Soils. *Agric. Water Manage.*, **100**: 46-57.
2. Allen, R. G., Pereira, L. S., Raes, D. and Smith, M. 1998. *Crop Evapotranspiration. Guidelines for Computing Crop Water Requirements*. FAO Irrigation and Drainage Paper **56**, Food and Agriculture Organization of the United Nations. Rome, Italy, 49 PP.

3. Angulo-Jaramillo, R., Bagarello, V., Prima, S. D. and Gosset, A. 2019. Beerkan Estimation of Soil Transfer Parameters (BEST) Across Soils and Scales. *J. Hydrol.*, **576**: 239-261.
4. Banon, S., Ochoa, J., Franco, J. A., Sanchez-Blanco, M. J. and Alarco'n, J. J. 2003. Influence of Water Deficit and Low Air Humidity in the Nursery on Survival of *Rhamnus Alaternus* Seedlings Following Planting. *J. Hort. Sci. Biotech.*, **78**: 518 - 522.
5. Beudez, N. and Doussan, C. 2013. Influence of Three Root Spatial Arrangements on Soil Water Flow and Uptake. *Proc. Environ. Sci.*, **19**: 37- 46.
6. Bondok, M. Y. and El-Sharkawy, A. F. 2014. Management of the Sprinkler Irrigation System for Cumin in the Old Valley. Egypt. *J. Agric. Res.*, **92**: 1-23.
7. Csonka, L., 1989. Physiological and genetic responses of bacteria to osmotic stress. *Microbiological Reviews*, **53**, 121-147.
8. Doorenbos, J. and Pruitt, W. O. 1977. *Crop Water Requirements*. FAO Irrigation and Drainage Paper **24**, Food and Agriculture Organization of the United Nations. Rome, Italy.
9. Doorenbos, J. and Kassam, A. H. 1986. Yield Response to Water. *Irrigation and Drainage Paper*, **33**. FAO, Rome. 93 P.
10. Elsafi, H. H., 2003. Effect of Water Quantity and Irrigation- Intervals on Vegetative Growth and Yield of Black Cumin (*Nigella sativa* L.) Thesis of fulfillment of the requirements for the degree of Master of Science in horticulture. College of Agricultural Studies. Sudan University of Science and Technology. 70 P.
11. Elsafi, H. H. 1988. Effect of Water Quantity and Irrigation- Intervals on Vegetative Growth and Yield of Black Cumin (*Nigella sativa* L.). Thesis of Fulfillment of the Requirements for the Degree of Master of Science in Horticulture, College of Agricultural Studies, Sudan University of Science and Technology, 70 PP.
12. Erteck, A. 2011. Importance of Pan Evaporation for Irrigation Scheduling and Proper Use of Crop Pan Coefficient ( $K_{cp}$ ), *Crop. Afri. J. of Agri. Res.*, **32**: 6706-6718.
13. Ghamarnia, H., Khosravy, H. and Sepehri, S. 2010. Yield and Water Use Efficiency of (*Nigella sativa* L.) under Different Irrigation Treatments in A Semi-Arid Region in the West of Iran. *J. Med. Plant. Res.*, **16**: 1612-1616.
14. Ghamarnia, H. and Jalili, Z. 2013. Water Stress Effects on Different Black Cumin (*Nigella sativa* L.) Components in a Semi-Arid Region. *Int. J. Agron. Plant Prod.*, **4**: 753-762.
15. Ghamarnia, H., Miri, E. and Ghobadei, M. 2014. Determination of Water Requirement, Single and Dual Crop Coefficients of Black Cumin (*Nigella sativa* L.) in a Semi-Arid Climate. *Irrig. Sci.*, **32**:67-76.
16. Haj Seyed H., M. R., Darzi, M. T. and Ghandehari, Z. 2012. Effects of Irrigation Treatment and Azospirillum Inoculation on Yield and Yield Component of Black Cumin (*Nigella sativa* L.). *J. Med. Plant. Res.*, **30**: 4553-4561.
17. Hassan, F. A. S. and Ali, E. F. 2014. Impact of Different Water Regimes Based on Class-A Pan on Growth, Yield and Oil Content of *Coriandrum sativum* L. *Plant. J. Saudi Soc. Agri. Sci.*, **13**: 155-161.
18. Karim, M., Himel, R. M., Ferdush, J. and Zakaria, M. 2017. Effect of Irrigation Levels on Yield Performance of Black Cumin. *Int. J. Environ. Agric. Biotech. (IJEAB)*, **22(2)**: 960-966.
19. Khalid, K. A. 2006. Influence of Water Stress on Growth, Essential Oil and Chemical Composition of Herbs (*Ocimum* sp.). *Int. Agrophys.* **20**: 289-296.
20. Khan, M. A., Chen, H. C., Tania, M. and Zhang, D. Z. 2011. Anticancer Activities of *Nigella Sativa* (Black Cumin). *Afr. J. Tradit. Comp. Altern Med.* **8**: 226- 232.
21. Kumar, S. A. 2009. *Plants Based Medicines in India*. Retrieved on February, 15, 2013: [Http://pib.nic.in/feature/feyr2000/fmay2000/f240520006.html](http://pib.nic.in/feature/feyr2000/fmay2000/f240520006.html).
22. Kumar, R., Pareek, N. K., Rathore, V. S. and Nangia, V. 2019. Effect of Irrigation and Nitrogen Application on Water Productivity and Performance of Cotton (*Gossypium*. sp). *J. Soil Water Conserv.*, **18**: 282-287.
23. Mahal, S. S. and Sidhu, A. S. 2006. Effect of Different Irrigation Schedules and Nitrogen Levels on Growth Yield



- Attributes and Seed Yield of Hybrid Canola (*Brassica Napus* L.). *Environ. Ecol.*, **4**: 1108-1111.
24. Mauget, S., Ulloa, M. and Mitchell, M. D. 2022. Simulated Irrigation Water Productivity and Related Profit Effects in U.S. Southern High Plains Cotton Production. *Agric. Water Manage. J.*, **266**: 107549.
25. Mehriya, M. L., Geat, N., Sarita Singh, H., Mattar, M. A. and Elansary, H. O. 2020. Response of Drip Irrigation and Fertigation on Cumin Yield, Quality, and Water Use Efficiency Grown under Arid Climatic Conditions. *Agron.*, **10(11)**: 1-16.
26. Michael, A. M. 1978. *Irrigation Theory and Practice*. Viskas Publishing House PVT LTD, New Delhi, PP. 448-584.
27. Molden, D., Oweis, T., Steduto, P., Bindraban, P., Hanjra, M. and Kijne, J. 2010. Improving Agricultural Water Productivity: Between Optimism and Caution. *Agric. Water Manage. J.*, **97(4)**: 528-535.
28. Mourad, K. 2012. Marginal and Virtual Water for Sustainable Water Resources Management in Syria. Academic Thesis Submitted in Partial Fulfillment of the Requirements for the Degree of Doctor of Philosophy (Ph.D. Engineering), Lund University, 157 PP.
29. Mubarak, I., Mailhol, J. C., Angulo-Jaramillo, R. and Ruelle, P. 2009. Temporal Variability in Soil Hydraulic Properties under Drip Irrigation, *Geoderma* **150**: 158-165.
30. Oweis, T. and Hachum, A. 2012. *Supplemental Irrigation: A Highly Efficient Water Use Practice*. ICARDA, Aleppo, Syria, 28 PP.
31. Ozer, H., Sahin, U., Coban, F. and Ors, S. 2020. Response of Black Cumin (*Nigella Sativa* L.) to Deficit Irrigation in a Semi-Arid Region: Growth, Yield, Quality, and Water Productivity. *J. Ind. Crops Prod.*, **144**: 112048.
32. Ruhi, B., Osman, K., Koksall, A. and Dursun, B. 2006. The Effects of Drip Irrigation on Flowering and Flower Quality of Glasshouse Gladiolus Plant. *Agric. Water Manage.*, **81**: 132-144.
33. Saxton, K. E. and Willey, P. H. 2006. The SPAW Model for Agricultural Field and Pond Hydrologic Simulation. Chapter 17. In: "Mathematical Modeling of Watershed Hydrology", (EDs.): Singh, V. P. and Frevert, D. CRC Press, PP. 401-435.
34. Senyigit, U. and Arslan, M. 2018. Effects of Irrigation Programs Formed by Different Approaches on the Yield and Water Consumption of Black Cumin (*Nigella sativa* L.) under Transition Zone in the West Anatolia Conditions. *Tar. Bil. Der. (J. Agric. Sci.)*, **24**: 22-32.
35. Serman, F. V., Orgaz, F., Starobinsky, G., Capraro, F. and Fereres, E. 2021. Water Productivity and Net Profit of High-Density Olive Orchards in San Juan, Argentina. *Agric. Water Manage. j.*, **252**: 106878.
36. Steduto, P., Hsiao, T. C., Fereres, E. and Raes, D. 2012. *Crop Yield Response to Water*. FAO Irrigation and Drainage Paper **66**, Food and Agriculture Organization of the United Nations Rome, 516 PP.
37. Taiz, L., and Zeiger, E., 2002. *Plant Physiology*, Third ed. Sunderland: Sinauer Associates. \$104'95. 690 pp. doi:10.1093/aob/mcg079.
38. Yaghi, T., Aldarir, A. N., Nangia, V., Oweis, T. and Arslan, A. 2016. Impact of Climate Changes on Water Resources Availability in the Orontes River Watershed: Case of Homs Governorate in Syria. *Jord. J. Agric. Sci.*, **12(2)**: 499-519.
39. Yousuf, M., Karim, A. J. M., Solaiman, A. R., Islam, M. and Zakaria, M. 2018. Nutrient Management on the Growth and Yield of Black Cumin (*Nigella sativa* L.). Bangladesh. *J. Agric. Res.*, **43(2)**: 205-210.
40. Zapotoczny, P., Zuk-Golaszewska, K. and Ropelewska, E. 2019. Impact of Cultivation Methods on Properties of Black Cumin (*Nigella sativa* L.) Seeds. *J. Cent. Eur. Agric.*, **20(1)**: 353-364.

اثر روش کم آبیاری بر کارایی مصرف آب زیره سیاه (*Nigella sativa* L.)

ت. یاغی، ع. ارسلان، و ح. سعید

## چکیده

برای افزایش سود کشاورزی، آزمایشی در مرکز تحقیقات علمی کشاورزی (SARC) در Salamiyah در سوریه انجام شد. هدف این آزمایش برآورد مقدار آبیاری بهینه که بهره‌وری را به حداکثر می‌رساند و همچنین کیفیت زیره سیاه در شرایط آزمایشی بود. تیمارهای آبیاری ۰.۲۵٪، ۰.۵۰٪، ۰.۷۵٪ و ۱.۰۰٪ تبخیر و تعرق پتانسیل (ET<sub>p</sub>) بر اساس تبخیر تشتک کلاس A بود. تیمارها به ترتیب B، C، D و E نشان داده شد. تیمار بدون آبیاری (A) به عنوان شاهد در نظر گرفته شد. آب آبیاری با سامانه قطره ای اعمال شد. تیمارهای A-E با استفاده از طرح بلوک‌های کامل تصادفی (CRBD) با چهار تکرار طراحی و اجرا شد. تبخیر و تعرق واقعی (ET<sub>a</sub>) پس از کالیبراسیون خاک با استفاده از نرم افزار SPAW محاسبه شد. افزون بر این، عملکرد محصول، کارایی مصرف آب آبیاری (IWUE)، ضریب پاسخ عملکرد (k<sub>y</sub>) و برخی پارامترها (دوره رشد، تاریخ گل‌دهی و غیره) طی فصل‌های رشد سال‌های ۲۰۱۸-۲۰۱۹ تعیین شد. نتایج نشان داد که می‌توان برنامه آبیاری را با استفاده از برخی معادلات و تبخیر تشتک کلاس A انجام داد. نیز، در مقایسه سطوح آبیاری بالاتر و پایین‌تر، پارامترهای رشد رویشی به طور مجازی و قابل توجهی شکوفا شدند. با اینهمه، با کاهش آب آبیاری، مقادیر کارایی مصرف آب آبیاری (IWUE) افزایش یافت. تیمار B بالاترین مقدار IWUE را ثبت کرد (۲.۴ کیلوگرم در هکتار برای هر میلی متر)، اما مقادیر کلیدی کمتر از ۱.۰ باقی ماند که نشان می‌دهد زیره سیاه خشکسالی را تحمل می‌کند. افزون بر این، در رابطه با نسبت سود به هزینه کل، تیمار C امتیاز برتر (۱۳۰.۶٪) را به دست آورد. در نتیجه، بر پایه نتایج، توصیه می‌کنیم که آبیاری بیش از ۰.۷۵٪ و کمتر از ۰.۲۵٪ ET<sub>p</sub> نباشد.