

## Effects of Aloe Vera Gel Based Active Coating Functionalized with Lemon Peel Essential Oil on Shelf Life and Quality Attributes of Cheese

S. S. Nejad Sajjadi<sup>1</sup>, L. Mansouri Najand<sup>1</sup>, and F. Shahdadi<sup>2\*</sup>

### ABSTRACT

In this study, the effect of an edible aloe vera gel-coating containing lemon peel essential oil (0, 100, and 150 ppm) on the qualitative characteristics of cheese samples was examined. Treatments included 4 groups: control (without coating), Aloe Vera Gel (AVG), AVG+100 ppm lemon peel Essential Oil (EO), and AVG+150 ppm lemon peel EO. These treatments were evaluated for 60 days in terms of physicochemical, textural, sensory, and microbial counting properties. The findings revealed that, as storage duration increased, the acidity and salt increased while pH and moisture content decreased. In evaluating the sensory properties, the effect of treatments on all sensory properties, except color scores, was significant. Samples coated with AVG and 100 ppm lemon peel EO received the highest flavor scores (4.97). As the storage time increased, the hardness, chewiness and springiness of the cheese samples increased. The samples' adhesiveness was not affected by the storage duration. At the end of the storage time, the highest total microbial, mold, and yeast counts were associated with the control cheese samples (5.37 and 4.62 log CFU g<sup>-1</sup>, respectively). The lowest amount was related to the samples coated with AVG and 150 ppm of lemon peel EO (3.92 and 3.76 log CFU g<sup>-1</sup>, respectively). In general, the use of edible coating produced with AVG and lower concentrations of lemon peel EO (100 ppm and less) improved the appearance and the flavor of cheese samples during 60 days of storage.

**Keywords:** *Aloe barbadensis* miller, Cheese sensory evaluation, Cheese textural properties, Edible coating.

### INTRODUCTION

Nowadays, chemical preservatives have been proved to be harmful and consumers tend to use foods without preservatives or containing natural preservatives (Sambu *et al.*, 2022). Cheese is a nutrient-dense dairy food, providing protein, fats, and minerals (Yerlikaya and Ozer, 2014). Cheese can be used as a main ingredient in meals, as a dessert, and as a component of foods. The rapid growth of cheese consumption in the world, particularly in European countries, is due to its use in various foods (Gomes da

Cruz *et al.*, 2009).

Due to its nutrients content, cheese provides a favorable environment for the growth of several bacteria. Globally, there are vast selections of different cheese varieties, and each one has a unique microbiological profile. The high nutritional value of cheese has led to extensive studies to improve the quantitative and qualitative properties of this product and the production of more marketable products (Trmčić *et al.*, 2016). Mold growth during the ripening and manufacturing of cheese is one of the common issues faced by cheese makers. This problem is also true for sellers and

<sup>1</sup> Department of Food Hygiene, Faculty of Veterinary Medicine, Shahid Bahonar University of Kerman, Islamic Republic of Iran.

<sup>2</sup> Department of Food Science and Technology, Faculty of Agriculture, University of Jiroft, Jiroft, Islamic Republic of Iran.

\*Corresponding author; e-mail: fatemeh.shahdadi@ujiroft.ac.ir



consumers of this product during refrigeration. The use of herb materials has been considered for many years to prevent the growth of various microorganisms and molds. (Sengun *et al.*, 2008).

Some natural and edible film-forming materials can be used to preserve foods such as cheese. One of these natural ingredients is Aloe Vera Gel (AVG). A clear and firm gel that is extracted from the inner parts of the leaves of the aloe vera (*Aloe barbadensis* Miller). AVG is odorless, non-sticky and has a high absorption strength. More than 98% of aloe vera gel is made up of water, followed by polysaccharides (pectin, cellulose, hemicellulose, glucomannan, and acemannan), the acemannan being considered as the main functional component of AVG (Bozzi *et al.*, 2007). Aloe vera is also an excellent source of antioxidants, Vitamin C, Vitamin A, Vitamin E, Beta-carotene, Folic acid, Calcium, and Magnesium (Suriati, 2018). Aloe gel has high potential to be used in the food industry, one of which is an edible coating material (Suriati *et al.*, 2020). This gel is a polysaccharide coating and can prevent moisture loss of the product. Due to the presence of different chemicals including aloin, acemannan (Martinez-Romero *et al.*, 2018), anthraquinone, saponins (Ergun and Satici, 2012) and phenolic compounds such as chatechin hydrate, caffeic acid, ferulic acid, ellagic acid, and quercetin (Sumi *et al.*, 2019), AVG has antifungal and antimicrobial characteristics and inhibits the growth and proliferation of fungus.

Essential Oils (EO) are another herbal component that can be utilized in edible films and coatings. Citrus peel EO is a mixture of more than 100 compounds, divided into two volatile parts (99-85% of the total essential oil), and the non-volatile part (1-15%). The volatile parts include monoterpenes (such as limonene), sesquiterpene hydrocarbons and oxygenated derivatives [aldehydes (such as citral), ketones and acids (along with linear aldehydes), alcohols (such as linalool)] and esters. The non-volatile parts include

hydrocarbons, fatty acids, sterols, carotenoids, waxes, coumarins, and flavonoids (Bennici and Tani, 2004). Limonene is the main monoterpene compound of citrus essential oil and has antioxidant, antibacterial and antiviral properties (Espina *et al.*, 2011; Roy *et al.*, 2007). In general, citrus peel EOs have potent antioxidant and antibacterial properties (Raspo *et al.*, 2020).

In a reported study, Shenbagam *et al.* (2023) investigated the effects of aloe vera gel-based edible coating (with or without incorporation of orange peel essential oil) on the postharvest shelf life and qualitative properties of button mushroom. The results showed that maximum concentration of orange peel essential oil (1,500  $\mu\text{L L}^{-1}$ ) incorporated in the 50% aloe vera gel significantly improved the postharvest quality attributes of mushrooms and helped extend the shelf life of mushrooms up to 4 days as compared to the control.

This study's objective was to determine the effect of an edible coating made of AVG and various concentrations of lemon peel EO on physicochemical, sensory and textural properties as well as microbial profile (total microbial count and total mold and yeast counts) of cheese.

## MATERIALS AND METHODS

### Preparation of Lemon Peel Essential Oil

Twenty kg of Mexican lemon peel (*Citrus aurantifolia*) was dried at an ambient temperature (25-38°C) and in the shade. The dried lemon peel was grounded and passed through a sieve (mesh 40). The EO was extracted by steam distillation over a Clevenger system (Aria Exir, Iran) for 4h. The obtained EO was dehydrated using sodium sulfate and stored at 4°C (Chanthaphon *et al.*, 2018). Lemon peel EO was yellow in color and yield of extracted EOs was 1.1% (w/w).

### Preparation of AVG

Aloe vera leaves were collected from University of Jiroft Research Farm. The leaves were washed and their jagged edges were cut with a knife. The top layer of the leaf was removed lengthwise and the gel was carefully separated from the leaf. The gel parts were blended thoroughly and put through a clean metal sieve (mesh 20) to form a homogeneous solution, and the extract was finally pasteurized at 65°C for 15 minutes (Martinez-Romero *et al.*, 2018). In this study, Aloe vera extract at a 100% concentration was used.

### Coating Formulations and Application

The cheese samples were prepared in Kerman Pegah Milk Factory. To produce cheese, milk was pasteurized after fat standardization (3.5%) by HTST method, and then concentrated at 50°C in ultra-filtration system until reaching 34% dry matter. Starter inoculation was done at 32-35°C. Then, rennet (12 mL per 400 g) was added and mixed well. The mixture was poured into containers and, after passing through the coagulation tunnel (30°C for 20 minutes), salt (3%) was added. Later, it was sealed and placed in an incubator at 28°C until reaching a pH of 4.7. Then, it was transferred to the cold room and kept at 4°C until the experiments (Khani and Roufegari Nejad, 2018). The cheese samples were cut into cubic specimens (3×3×3 cm<sup>3</sup>) and coated by immersion method. During this step, the cheese samples were immersed in the coating mixture (AVG with various concentrations of lemon peel EO (0, 100 and 150 ppm, which was homogenized by a homogenizer at 1000 rpm) for 1 minute. The samples were incubated for about 8 hours under controlled temperature (12°C) and humidity (relative humidity of 85%) to dry all coatings (Henriques *et al.*, 2013). AVG and lemon peel EO created a colorless coating on the samples. The samples were then placed in sealed polypropylene containers, stored in the refrigerator (4°C), and evaluated at 15 days intervals throughout the 60 days of storage period.

### Determination of Acidity

The acidity of cheese samples was determined in terms of lactic acid and by titration with sodium hydroxide (0.1N) using Equation (1) (Iranian International Standard No. 2852, 2006).

$$\text{Acidity (\%)} = N \cdot 0.009 \cdot 100 / M \quad (1)$$

Where, N is the amount of sodium hydroxide 0.1N consumed (mL) and M is the weight of the sample.

### pH Measurement

A digital pH meter (Metrohm, model 827, Switzerland) was used to determine the samples' pH levels (Iranian International Standard No. 2852, 2006)

### Measurement of Moisture Content

Cheese samples were placed in an oven at 102°C until they reached a constant weight (about 5 h). The dried samples were weighed after cooling, and the amount of moisture loss was estimated using the Equation (2) (Roy *et al.*, 2007):

$$\text{Moisture loss rate} = \frac{\text{Weight}_{\text{before drying}} - \text{Weight}_{\text{after drying}}}{\text{Weight}_{\text{before drying}}} \times 100 \quad (2)$$

### Measurement of Salt Content

Mohr method was used to determine the amount of salt. Titration was performed using silver nitrate solution (0.1 N) until an orange precipitate appeared. The percentage of salt was calculated as Equation (3) (Dorosti *et al.*, 2011).

$$\text{Salt (\%)} = \frac{\text{Consumed silver nitrate (mL)} \times \text{Silver nitrate N} \times 0.585}{\text{Weight}} \quad (3)$$

### Sensory Evaluation

Sensory properties of cheese samples were evaluated using a five-point hedonic test (Very bad: 1 to Very good: 5) in the first and



60<sup>th</sup> days of storage. The evaluators were 50 people who were selected from the experts working in Pegah Kerman Factory and students familiar with the characteristics of cheese. Samples (100 g packages) were removed from the refrigerator before the test, and after reaching the ambient temperature, they were given to the evaluators in 30 g pieces.

Samples were assessed for their characteristics including flavor, odor, color, texture and overall acceptance. Mean data of the first and 60<sup>th</sup> days were reported (Beigomi *et al.*, 2013).

### Texture Analysis Test

A texture analyzer equipment (model QTS25, FARNEL CNS, UK) and a cylindrical probe with a diameter of 36 mm were utilized for the Texture Profile Analysis (TPA) test. The cheese samples were removed from the refrigerator before the test and, after slicing (20×20×20 mm) up to 50% of the initial height (10 mm depth), were compressed by the machine. Each test was performed in at least three replications. The measured traits were hardness, cohesiveness, adhesiveness, chewiness, springiness, and gumminess. It should be noted that the TPA test was a two-step test, and these traits were defined according to the standard TPA curve (Hosseini *et al.*, 2013).

### Microbial Tests

#### *Total Microbial Count*

The total microbial count was performed using a PCA (Plate Count Agar) at 37°C for 48 hours. The number of bacteria in cheese samples was calculated as follows (Rezaei *et al.*, 2010).

Microbial content  $\text{g}^{-1}$  of cheese = Number of colonies  $\times$  Inverse dilution coefficient  $\times 10$  (4)

#### *Mold and yeast count*

YGC (Yeast Extract Glucose Chloramphenicol) medium was used for mold and yeast (fungi) count at 25°C for 48-72 hours. After incubation, the obtained colonies were counted using the Equation (4) (Rezaei *et al.*, 2010).

### Statistical Analysis

The experiments were conducted in a factorial experiment based on completely randomized design and the experimental data were analyzed with SPSS: 21 software. Factors included treatments (4 levels) and storage time (5 levels). The means were compared using the Duncan's multiple range test with a 5% confidence level. All experiments were carried out in triplicate.

## RESULTS AND DISCUSSION

### Effect of Treatments on pH

The findings in Table 1 demonstrate that the pH of cheese samples significantly reduced as storage time was increased. The lowest pH reduction was observed for cheese samples coated with AVG and 150 ppm of lemon peel EO. The pH of the control treatment was found to be the lowest at the end of the maintenance time, whereas the other treatments were not significantly different ( $p > 0.05$ ).

### Effect of Treatments on Acidity

Table 2 shows that the acidity of the cheese samples was significantly influenced by the type of coating used as well as the storage time. The treatment coated with AVG and 150 ppm EO and the control had the greatest and lowest acidity, respectively, on the 60<sup>th</sup> day. The acidity of the treatments increased as storage duration increased, and

**Table 1.** The effect of treatments on the pH of samples.<sup>a</sup>

Treatments	Day 1	Day 15	Day 30	Day 45	Day 60
Control	5.50±0.1 <sup>Ab</sup>	5.43±0.05 <sup>ABb</sup>	5.28±0.12 <sup>Cb</sup>	5.09±0.1 <sup>CDb</sup>	4.86±0.14 <sup>Ec</sup>
AVG	5.86±0.09 <sup>Aa</sup>	5.79±0.1 <sup>Aba</sup>	5.57±0.05 <sup>Ca</sup>	5.36±0.1 <sup>Da</sup>	5.10±0.06 <sup>Eab</sup>
AVG+100	5.85±0.11 <sup>Aa</sup>	5.75±0.15 <sup>Aba</sup>	5.58±0.2 <sup>Ca</sup>	5.35±0.08 <sup>Da</sup>	5.16±0.1 <sup>Eab</sup>
ppm EO					
AVG+150	5.83±0.1 <sup>Aa</sup>	5.75±0.08 <sup>Aba</sup>	5.57±0.05 <sup>Ca</sup>	5.57±0.05 <sup>CDa</sup>	5.25±0.12 <sup>Da</sup>
ppm EO					
P value	0.013	0.004	0.003	0.004	0.049

<sup>a</sup> Mean values in each column that have different lower-case letters have a significant difference ( $P < 0.05$ ). Numbers in each row that have different capital letters have a significant difference ( $P < 0.05$ ).

**Table 2.** The effect of treatments on the samples acidity.<sup>a</sup>

Treatments	Day 1	Day 15	Day 30	Day 45	Day 60
Control	1.46±0.05 <sup>Ea</sup>	1.77±0.09 <sup>Da</sup>	2.11±0.2 <sup>Ca</sup>	2.66±0.09 <sup>Ba</sup>	3.26±0.1 <sup>Aa</sup>
AVG	1.26±0.1 <sup>Eb</sup>	1.56±0.1 <sup>Db</sup>	1.80±0.18 <sup>Cb</sup>	2.16±0.1 <sup>Bb</sup>	2.70±0.05 <sup>Ab</sup>
AVG + 100	1.23±0.08 <sup>Eb</sup>	1.55±0.1 <sup>Db</sup>	1.81±0.1 <sup>Cb</sup>	2.05±0.06 <sup>Bbc</sup>	2.66±0.11 <sup>Abc</sup>
ppm EO					
AVG + 150	1.26±0.08 <sup>Eb</sup>	1.50±0.1 <sup>CDb</sup>	1.67±0.15 <sup>Cb</sup>	1.93±0.1 <sup>Bc</sup>	2.52±0.15 <sup>Ac</sup>
ppm EO					
P value	0.004	0.004	0.033	0.044	0.004

<sup>a</sup> Mean values in each column that have different lower-case letters have a significant difference ( $P < 0.05$ ). Numbers in each row that have different capital letters have a significant difference ( $P < 0.05$ ).

this increase was significant in all investigated treatments on all storage days. In cheese samples coated with AVG and 150 ppm of lemon peel EO, minimal acidity changes were observed at the end of storage period.

By increasing the storage time, pH values of all samples decreased, which may be related to the activity of lactic acid bacteria species owing to the metabolization of lactose to lactate and the produced acid (Dermiki *et al.*, 2008). Ramos *et al.* (2012) found that the pH of cheeses coated with whey protein isolate, guar gum, and antimicrobial substances decreased with increasing storage time, and the coated cheeses had a higher pH than the control. Jamshidi *et al.* (2018), used a coating of AVG and Persian gum in Iranian white cheese, and reported that, during storage, the pH decreased significantly while the acidity increased. Over time, the acidity of the various treatments increased, indicating that an increase in lactic acid production by the bacteria may be the main reason for this trend, definitely consistent with the

decreasing trend observed in pH during storage.

El-Sisi *et al.* (2015) showed that the acidity of chitosan-coated cheeses increased during storage. A study also revealed that the acidity of cheddar cheese samples coated with whey protein increased during ripening (Wagh *et al.*, 2013).

On the 60<sup>th</sup> day, the lowest amount of acidity was observed in cheese samples coated with AVG and lemon peel EO. This could indicate the low activity of lactic acid bacteria (starter and non-starter) in these samples. The more activity of lactic acid bacteria leads to more decomposition of lactate and production of organic acids such as lactic acid and acetic acid, and AVG and lemon peel EO probably due to antimicrobial activity decreased growth of these bacteria in cheese samples (Wagh *et al.*, 2013)

#### Effect of Treatments on the Salt Content

**Table 3.** The effect of treatments on the salt content (%) of cheese samples.<sup>a</sup>

Treatments	Day 1	Day 15	Day 30	Day 45	Day 60
Control	3.26±0.25 <sup>Ba</sup>	3.30±0.2 <sup>ABa</sup>	3.41±0.1 <sup>ABa</sup>	3.50±0.2 <sup>Aa</sup>	3.57±0.1 <sup>Aa</sup>
AVG	2.76±0.1 <sup>Bb</sup>	2.81±0.5 <sup>ABb</sup>	2.86±0.25 <sup>Ab</sup>	2.96±0.1 <sup>Ab</sup>	3.06±0.15 <sup>Ab</sup>
AVG+100	2.73±0.3 <sup>Bb</sup>	2.80±0.5 <sup>ABb</sup>	2.86±0.21 <sup>Ab</sup>	2.93±0.09 <sup>Ab</sup>	3.07±0.15 <sup>Abc</sup>
ppm EO					
AVG+150	2.73±0.17 <sup>ABb</sup>	2.80±0.44 <sup>ABb</sup>	2.87±0.23 <sup>Ab</sup>	2.94±0.15 <sup>Ab</sup>	2.98±0.17 <sup>Ac</sup>
ppm EO					
P value	0.007	0.002	0.001	0.001	0.001

<sup>a</sup> Mean values in each column that have different lower-case letters have a significant difference ( $P < 0.05$ ). Numbers in each row that have different capital letters have a significant difference ( $P < 0.05$ ).

According to the results in Table 3, the control had the most salt content at all storage times, while the other treatments were not significantly different ( $P > 0.05$ ). The salt content of the treatments enhanced with increasing storage period, although this was not significant in samples coated with AVG and 150 ppm lemon peel EO ( $P > 0.05$ ).

It can be seen that all the coated samples have less salt than the control, which is consistent with the results of other researchers who studied the effect of coating on the properties of cheese (Ramos *et al.*, 2012; Yilmaz and Dagdemir, 2012). On the other hand, during the 60 days of storage, the salt content of samples increased slightly as a result of weight loss due to the removal of moisture from the cheese texture.

#### Effect of Treatments on the Moisture Content

Effect of the treatments on moisture content of cheese samples is shown in Table

**Table 4.** The effect of treatments on the moisture content (%) of samples.<sup>a</sup>

Treatments	Day 1	Day 15	Day 30	Day 45	Day 60
Control	63.30±4.2 <sup>Aa</sup>	61.73±2.9 <sup>Ab</sup>	60.76±2.1 <sup>ABb</sup>	58.66±3.3 <sup>Bb</sup>	58.73±4.1 <sup>Bb</sup>
AVG	64.15±3.1 <sup>Aa</sup>	64.66±4.3 <sup>Aa</sup>	62.70±2.5 <sup>ABab</sup>	61.67±3.1 <sup>Ba</sup>	61.33±4.2 <sup>Ba</sup>
AVG+100 ppm	64.60±4.0 <sup>Aa</sup>	64.73±4.2 <sup>Aa</sup>	63.00±2.1 <sup>Aa</sup>	62.08±3.5 <sup>ABa</sup>	61.40±3.8 <sup>Ba</sup>
EO					
AVG+150 ppm	64.20±4.1 <sup>Aa</sup>	64.84±3.9 <sup>Aa</sup>	63.43±2.4 <sup>ABa</sup>	62.64±3.4 <sup>Ba</sup>	61.44±4.3 <sup>Ba</sup>
EO					
P value	0.006	0.032	0.007	0.000	0.033

<sup>a</sup> Mean values in each column that have different lower-case letters have a significant difference ( $P < 0.05$ ). Numbers in each row that have different capital letters have a significant difference ( $P < 0.05$ ).

4. The results reveal that the coating treatments and storage period had a significant effect on the moisture content. As storage time increased, the moisture content of samples decreased. Cheese samples with coatings retained moisture significantly more than the control. There was no significant difference in the moisture content of all treatments on the first day of storage ( $P > 0.05$ ).

The cheese samples' moisture gradually decreased during the storage period as a result of some moisture being released from the texture of the cheese and the packaging to the outside. The difference between the coated samples is probably due to the composition of the coating as well as the kinetics of water influence and outflow into the various coatings (Pantaleão *et al.*, 2007). Jamshidi *et al.* (2018) reported that almost all cheeses coated with AVG and Persian gum showed higher moisture content than the control, which indicates the positive effect of coating on moisture retention in cheese during storage. Coating with aloe vera gel had a barrier property for moisture

loss in several fruits such as peach (Mohammadi *et al.*, 2020), plum (Martinez-Romero *et al.*, 2018), grapes, fresh cut papaya (Farina *et al.*, 2020), and tomato fruit (Tzortzakakis *et al.*, 2019).

### Effect of Treatments on the Sensory Properties

With the exception of the color index, Table 5 shows that the effects of the tested treatments on the sensory characteristics of samples are significant. The highest taste score was related to the treatment coated with AVG and 100 ppm of lemon peel EO, while the lowest taste score was related to the control and the AVG and 150 ppm of lemon peel EO. The highest and lowest odor scores were observed in AVG with 100 ppm of lemon peel EO and the control, respectively. Samples coated with AVG and different concentrations of EO did not show significant differences in terms of texture ( $P > 0.05$ ), and the lowest texture score was assigned to the control. In terms of general acceptance, AVG with 100 ppm of lemon peel EO received the highest score.

Most sensory panelists reported a bitter taste for cheeses containing 150 ppm of lemon peel essential oil. According to research of Yilmaz and Dagdemir (2012), there were no significant differences in the color of cheese samples coated in beeswax compared to the control, which is consistent with the findings of this investigation.

Abbas *et al.* (2017) reported that adding 0.005 and 0.010  $\mu\text{L}$  of basil essential oil to UF soft cheese significantly improved the taste throughout the freshness of cheese and during the 60 days of storage. According to this report, the desirability of samples containing low concentration (0.005  $\mu\text{L}$  per 100 mL) was higher than the samples containing high concentration (0.010  $\mu\text{L}$  100  $\text{mL}^{-1}$ ). Mohammadi *et al.* (2011) reported that 100 mg/kg of basil essential oil improved the odor, taste and acceptability of white cheese during the production and storage, however, the taste and acceptability of the cheese samples were adversely affected by the essential oil concentrations of 150 and 200  $\text{mg kg}^{-1}$ .

According to Otero *et al.* (2014), sheep cheese samples covered with edible films containing antimicrobial agents had improved sensory properties. The results of Pieretti *et al.* (2019) showed that cheese samples coated with alginate and low concentrations of oregano essential oil had better sensory acceptance than the control and higher concentrations of essential oil.

### Effect of the Treatments on the Textural Characteristics

The effect of the studied treatments on the textural characteristics of cheese samples is shown in Figure 1 (A-F).

#### I- Hardness

According to Figure 1 (A), both the

**Table 5.** The effect of treatments on the sensory properties of samples.<sup>a</sup>

Treatments	Taste	Odor	Color	Texture	General acceptance
Control	4.36±0.1 <sup>c</sup>	4.45±0.05 <sup>c</sup>	4.92±0.1	4.53±0.15 <sup>c</sup>	4.42±0.08 <sup>cd</sup>
AVG	4.59±0.05 <sup>b</sup>	4.63±0.1 <sup>b</sup>	4.96±0.1	4.67±0.1 <sup>b</sup>	4.78±0.12 <sup>b</sup>
AVG+100 ppm EO	4.97±0.06 <sup>a</sup>	4.89±0.12 <sup>a</sup>	4.96±0.15	4.91±0.05 <sup>Aa</sup>	4.95±0.06 <sup>a</sup>
AVG+150 ppm EO	4.36±0.05 <sup>c</sup>	4.75±0.16 <sup>b</sup>	4.97±0.09	4.89±0.12 <sup>a</sup>	4.54±0.1 <sup>c</sup>
P value	0.001	0.001	0.56	0.000	0.003

<sup>a</sup> Mean values in each column that have different lower-case letters have a significant difference ( $P < 0.05$ ).



coating and storage duration significantly affected hardness. The hardness of the samples increased with storage time. On 60<sup>th</sup> day, the AVG with 100 and 150 ppm of lemon peel EO had the lowest hardness while the control and AVG treatments had the highest.

#### 2- Adhesiveness

Figure 1 (B), shows that although numerically the adhesiveness of the samples increased during storage, the storage time had no significant effect on the adhesiveness of the samples. At the end of storage, samples coated with AVG and 150 ppm of lemon peel EO showed the highest adhesiveness, which did not show a significant difference with the AVG and 100 ppm of lemon peel EO treatment ( $P > 0.05$ ). The lowest adhesiveness was related to the control.

#### 3- Cohesiveness

Figure 1 (C), shows that the AVG treatment had the lowest cohesiveness at the end of the storage period, with no other treatments significantly different ( $P > 0.05$ ).

#### 4- Springiness

According to Figure 1 (D), no particular trend in the springiness of samples during storage time was seen in the control. In other treatments, springiness of samples increased with increasing storage time. On the 60<sup>th</sup> day, the highest springiness was related to AVG treatments with 100 and 150 ppm of lemon peel EO and the lowest amount of springiness was related to the control.

#### 5-Gumminess

According to Figure 1 (E), it can be seen that, at the end of the storage, the highest and lowest gumminess were observed in the treatment coated with AVG+150 ppm lemon peel EO and the control, respectively. The gumminess of samples significantly increased as the storage time rose.

#### 6- Chewiness

Figure 1 (E) shows that chewiness of the samples increased as storage time increased. In the samples coated with AVG and AVG containing 100 and 150 ppm of lemon peel EO on the 45<sup>th</sup> and 60<sup>th</sup> days, this enhancement was not significant ( $P > 0.05$ ).

On the 60<sup>th</sup> day, the lowest amount of chewiness was observed in the control and samples coated with AVG and the highest amount of chewiness was observed in the samples coated with AVG and 150 ppm of lemon peel EO.

According to the findings of the textural characteristics, the hardness of the samples increased with increasing storage time, which may be related to moisture loss during storage. Another factor contributing to the samples' increased hardness during storage is likely an increase in protein-protein interactions (Bianchi *et al.*, 2021). It was also observed that the coated samples had less hardness than the control. It seems that more moisture in the coatings and more hydration may reduce the hardness of the samples (Zhong *et al.*, 2014).


Pieretti *et al.* (2019) examined how rosemary and oregano EOs and alginate-based edible coatings affected the textural characteristics of fresh cheese. They found that, at the end of the storage period, the coated samples had less hardness than the control.

At the end of the storage, the highest amount of adhesiveness was observed in the samples coated with AVG and 150 ppm lemon peel EO and the lowest adhesiveness was related to the control. In the research of Wang *et al.* (2019), cheddar cheese samples coated with isolated whey protein nanofibrils and carvacrol showed more adhesiveness than the uncoated samples.

The cohesiveness of the samples increased with increasing the storage time, and at the end of the storage, the treatment with the lowest cohesiveness was in the presence of AVG. The other treatments did not significantly differ from each other. In the study reported by Wang *et al.* (2019), the cohesiveness of coated cheese samples increased with increasing storage time, while no significant difference was observed in the other samples.

With increasing storage period, the chewiness of samples increased. This is in line with the hardness and gumminess





**Figure 1.** The effects of treatments on the hardness (A), adhesiveness (B), cohesiveness (C), springiness (D), gumminess (E), and chewiness (F) of the cheese samples. (Different lower-case letters have a significant difference ( $P < 0.05$ )).

properties. On the 60<sup>th</sup> day, the highest amount of chewiness was related to the treatment coated with AVG and 150 ppm of lemon peel EO. From a sensory point of view, it is perceived that more energy is needed to chew the coated samples. It was found that the chewiness of cheese samples coated with starch and carvacrol increased with increasing storage, and the coated samples had more chewiness than the control (López-Córdoba, 2021).

### Effect of Treatments on Microbial Count of Samples during Storage

#### 1- Total Microbial Count

Table 6 shows the effect of the treatments on the total microbial count of the samples. This table shows that the total microbial

count was significantly affected by both storage times and coatings. The total microbial count increased with increasing storage time. The sample coated with AVG and 150 ppm of lemon peel EO had the lowest microbial count. In general, the coated treatments showed less microbial count than the control.

#### 2- Total Mold and Yeast Count

Table 7 shows that there is significant variation in the total number of mold and yeast in cheese samples depending on the various treatments and storage time. The total amount of mold and yeast increased with more storage time across all treatments, with the control having the highest levels. The lowest amounts of mold and yeast were found in samples that had been coated with AVG and EO.

In general, the coated treatments showed less microbial, mold, and yeast counts than

**Table 6.** The effect of treatments on the total microbial count (log CFU g<sup>-1</sup>) of samples.<sup>a</sup>

Treatments	Day 1	Day 15	Day 30	Day 45	Day 60
Control	4.87±0.15 <sup>Ca</sup>	4.88±0.1 <sup>Ca</sup>	4.94±0.21 <sup>Ba</sup>	4.98±0.15 <sup>Ba</sup>	5.37±0.2 <sup>Aa</sup>
AVG	3.64±0.1 <sup>Db</sup>	3.73±0.14 <sup>Cb</sup>	4.60±0.15 <sup>Bb</sup>	4.64±0.11 <sup>Bb</sup>	4.82±0.1 <sup>Ab</sup>
AVG+100	3.38±0.22 <sup>Dc</sup>	3.55±0.21 <sup>Cc</sup>	3.96±0.25 <sup>Bc</sup>	4.30±0.1 <sup>Ac</sup>	4.31±0.14 <sup>Ac</sup>
ppm EO					
AVG+150	2.71±0.12 <sup>Ed</sup>	2.92±0.2 <sup>Dd</sup>	3.73±0.1 <sup>Cd</sup>	3.81±0.25 <sup>Bd</sup>	3.92±0.21 <sup>Ac</sup>
ppm EO					
P value	0.001	0.002	0.001	0.000	0.003

<sup>a</sup> Mean values in each column that have different lower-case letters have a significant difference (P< 0.05). Numbers in each row that have different capital letters have a significant difference (P< 0.05).

**Table 7.** The effect of treatments on total mold and yeast count (log CFU g<sup>-1</sup>) of cheese samples.<sup>a</sup>

Treatments	Day 1	Day 15	Day 30	Day 45	Day 60
Control	3.34±0.2 <sup>Ea</sup>	3.82±0.13 <sup>Da</sup>	3.96±0.15 <sup>Ca</sup>	4.13±0.14 <sup>Ba</sup>	4.62±0.1 <sup>Aa</sup>
AVG	3.17±0.15 <sup>Eb</sup>	3.45±0.21 <sup>Db</sup>	3.81±0.11 <sup>Cb</sup>	4.01±0.1 <sup>Bb</sup>	4.15±0.2 <sup>Ab</sup>
AVG+100	0.00 <sup>Dc</sup>	3.11±0.1 <sup>Cc</sup>	3.47±0.22 <sup>Bc</sup>	3.76±0.1 <sup>Ac</sup>	3.92±0.21 <sup>Ac</sup>
ppm EO					
AVG+150	0.00 <sup>Ec</sup>	0.00 <sup>Dd</sup>	3.06±0.1 <sup>Cd</sup>	3.47±0.12 <sup>Bd</sup>	3.76±0.15 <sup>Ad</sup>
ppm EO					
P value	0.001	0.002	0.003	0.001	0.002

<sup>a</sup> Mean values in each column that have different lower-case letters have a significant difference (P< 0.05). Numbers in each row that have different capital letters have a significant difference (P< 0.05).

the control. Numerous studies have focused on the antibacterial effects of AVG and lemon peel EO (Nielsen and Rios, 2000; Irshad *et al.*, 2011; Roy *et al.*, 2007).

Aloin and aloe-emodin are the two main components of aloe vera gel. Several researchers have confirmed the antifungal and anti-bacterial properties with improved moisture and gas barrier properties of aloe vera gel based edible coating (Ortega-Toro *et al.*, 2017).

AVG as a coating can create a physical barrier against microorganisms and reduce the occurrence of microbial spoilage (Asghari and Khalili, 2014). AVG inhibits the germination and growth of fungal mycelium and the inhibitory effect of its compounds on the activity of enzymes of pathogenic fungi has been proven (Reynolds and Dweck, 1999). Saritha *et al.* (2010) reported that the antimicrobial activity of AVG against gram-positive bacteria was higher than gram-negative bacteria. Navarro *et al.* (2011) also reported that AVG controls the *Rhizopus stolonifer*, *Botrytis cinerea* and

*Penicillium digitatum*. Leitgeb *et al.* (2021) investigated the effect of two aloe vera cultivars gel on different bacteria and fungi and reported that both aloe vera cultivars gel inhibited the growth of *Bacillus cereus*, *Staphylococcus aureus*, *E. coli*, *Pseudomonas aeruginosa*, *P. fluorescens* and *Candida albicans*, representatives of Gram-positive bacteria, Gram-negative bacteria, and fungi. The antibacterial properties of aloe vera are due to its constituents, which include saponins, acemannan, and anthraquinone derivatives. Therefore, the presence of these substances and antibacterial compounds in the AVG can reduce spread of germs in the treated samples (Ramasubramanian *et al.*, 2010).

Essential oils have different mechanisms in destroying microorganisms. These compounds enter the lipids of cell membranes and mitochondria, and this causes a difference in the structure of cells and their greater permeability, resulting in the release of ions and other cell contents. The release of large amounts of cellular

contents or the release of vital molecules and ions causes cell death (Pauli, 2006). There are several reports about the antimicrobial effect of citrus EOs and extracts (Chanthaphon *et al.*, 2018; Tan *et al.*, 2011). Antimicrobial properties of lemon peel EO are related to its active ingredients. Limonene is the main monoterpene compound of lemon peel and other citrus EOs, which has antibacterial and antiviral properties (Espina *et al.*, 2011; Roy *et al.*, 2007).

Artiga-Artigas *et al.* (2017) studied the antimicrobial effect of edible coating containing different concentrations of oregano EO on low-fat cheese. Their results showed that coatings containing oregano EO significantly reduced the microbial population during storage.

## CONCLUSIONS

The use of edible coating of AVG and lemon peel EO on cheese improved the appearance and prevented textural changes during storage. Lower concentrations (50 and 100 ppm) of lemon peel EO were suitable for obtaining cheeses with better sensory properties. The coatings maintained properties such as moisture, pH, hardness, etc. The lowest microbial, mold and yeast counts were observed in the treatments coated with AVG and 150 ppm of lemon peel EO. In general, the coated treatments showed less microbial, mold and yeast counts than the control.

## REFERENCES

1. Abbas, H. M., Assem, F. M., Zaky, W. M., Kassem, J. M. and Omer, E. A. 2017. Antioxidant, Rheological and Sensorial Properties of Ultra-Filtrated Soft Cheese Supplemented with Basil Essential Oil. *Int. J. Dairy Sci.*, **12(5)**: 301-309.
2. Artigas, M.A., Acevedo-Fani, A. and Martín-Belloso, O. 2017. Improving the Shelf Life of Low-Fat Cut Cheese using Nanoemulsion-Based Edible Coatings Containing Oregano Essential Oil and Mandarin Fiber. *Food Control.*, **76**: 1–12.
3. Asghari, M. R. and Khalili, H. 2014. The Effect of *Aloe vera* Gel on Polyphenol Oxidase Activity, Qualitative Properties and Shelf Life of Cherry Fruit. *J. Hort. Sci.*, **28 (3)**: 399-406.
4. Beigomi, M., Ghods Rohani, M., Mohammadifar, M. A., Hashemi, M., Valizadeh, M. and Ghanati, K. 2013. Comparison of Textural and Sensory Characteristics of Ultrafiltrated White Cheese Produced by Paneer Bad (*Withania coagulans*) Protease and Fungal Rennet. *Iran. J. Nutr. Sci. Food Technol.*, **8(1)**: 253-262.
5. Bennici, A. and Tani, C. 2004. Anatomical and Ultrastructural Study of the Secretory Cavity Development of Citrus Sinensis and Citrus Limon: Evaluation of Schizolysigenous Ontogeny. *Flora.*, **199**: 464 -475.
6. Bianchi, A., Mallmann, S., Gazoni, I., Cavalheiro, D. and Rigo, E. 2021. Effect of Acid Casein Freezing on the Industrial Production of Processed Cheese. *Int. Dairy J.*, **118**: 105043
7. Bozzi, A., Perrin, C., Austin, S. and Arce Vera, F. 2007. Quality and Authenticity of Commercial *Aloe vera* Gel Powders. *Food Chem.*, **103(1)**: 22–30.
8. Chanthaphon, S., Chanthachum, S. and Hongpattarakere, T. 2018. Antimicrobial Activities of Essential Oils and Crude Extract from Tropical *Citrus* spp. against Food-Related Microorganism. *J. Sci. Technol.*, **30(1)**: 125-131.
9. Dermiki, M., Ntzimani, A., Badeka, A., Savvaidis, I.N. and Kontominas, M.G. 2008. Shelf-Life Extension and Quality Attributes of the Whey Cheese. *LWT-Food Sci. Technol.*, **41(2)**: 284-294.
10. Dorosti, S., Bazmi, A., Ghanbarzadeh, B. and Ayaseh, A. 2011. Effect of Brine Concentration on the Physicochemical Properties of Iranian White Cheese. *J. Food Sci. Technol.*, **8(30)**: 1-10.
11. El-Sisi, A.S., Mohamed Gapr, E.S. and Kamaly, K.M. 2015. Use of Chitosan as an Edible Coating in RAS Cheese. *Biolife.*, **3(2)**: 564-570.
12. Ergun, M. and Satici, F. 2012. Use of Aloe Vera Gel as Bio preservative for ‘Granny



- Smith' and 'Red Chief' Apples. *J. Anim. Plant Sci.*, **22(2)**: 363-368.
13. Espina, L., Somolinos, M., Loran, S., Conchello, P., Garcia, D. and Pagan, R. 2011. Chemical Composition of Commercial Citrus Fruit Essential Oils and Evaluation of Their Antimicrobial Activity Acting Alone or in Combined Processes. *Food Control*, **22**: 896- 902.
  14. Farina, V., Passafiume, R., Tinebra, I., Scuderi, D., Saletta, F. and Gugliuzza, G. 2020. Postharvest Application of Aloe Vera Gel-Based Edible Coating to Improve the Quality and Storage Stability of Fresh-Cut Papaya. *J. Food Qual.*, **200(1)**: 8303140
  15. Gomes da Cruz, A., Buriti, F. C. A., Batista de Souza, C. H., Fonseca Faria, J. A. and Isay Saad, S. M. 2009. Probiotic Cheese: Health Benefits, Technological and Stability Aspects. *Trends Food Sci. Technol.*, **20(8)**: 344-354.
  16. Henriques, M., Santos, G., Rodrigues, A., Gomes, D., Pereira, C. and Gil, M. 2013. Replacement of Conventional Cheese Coatings by Natural Whey Protein Edible Coatings with Antimicrobial Activity. *J. Hyg. Eng. Des.*, **3**: 34-47.
  17. Hosseini, M., Habibi Najafi, M. B. and Mohebbi, M. 2013. Assessment of Physico-Chemical and Sensory Properties of Imitation Cheese Containing Whey Protein Concentrate and Enzyme-Modified Lighvan Cheese. *Iran. J. Nutr. Sci. Food Ind.*, **8(2)**: 91-102.
  18. Iranian International Standard No. 2852. 2006. *Milk and Milk Products: Determination of Titratable Acidity and Value pH –Test Method*. 1st Revision, The Institute of Standards and Industrial Research of Iran, Iran.
  19. Irshad, S., Butt, M. and Younus, H. 2011. In vitro Antibacterial Activity of *Aloe Barbadosis* Miller (*Aloe vera*). *Pharma.*, **1(2)**: 59-64 .
  20. Jamshidi, F., Rahimi, S. and Fadaei Noghani, V. 2018. The Effect of Edible *Aloe vera* Gel-Persian Gum Film on Iranian White Cheese Properties. *Iran. J. Nutr. Sci. Food Technol.*, **13(1)**: 63-74.
  21. Leitgeb, M., Kupnik, K., Knez, Ž. and Primožič, M. 2021. Enzymatic and Antimicrobial Activity of Biologically Active Samples from *Aloe arborescens* and *Aloe barbadensis*. *J. Biol.*, **10(8)**: 765-771.
  22. Khani, A. and Roufegari Nejad, L. 2018. Low Fat UF-Feta Cheese Production Containing Xanthan Gum. *J. Food Ind. Res.*, **29(1)**: 155-167.
  23. López-Córdoba, A. 2021. Feasibility of Using Carvacrol/Starch Edible Coatings to Improve the Quality of Paipa Cheese. *Polym.*, **13(15)**: 1-11.
  24. Martínez-Romero, D., Paladines, D., Valverde, M., Guillén, F., Zapata, P. J. and Valero, D. 2018. Rosehip Oil Added to *Aloe vera* Gel as Postharvest Coating of 'Songría' Plums and 'President' Prunes. *Acta Hortic.*, **1194(4)**: 321–325..
  25. Mohammadi, K., Karim, G., Hanifian, Sh., Tarinejad, A. and Gaseemnezhad, R. 2011. Antimicrobial effect of *Zataria multiflora* Boiss. Essential oil on *Escherichia coli* O157:H7 during manufacture and ripening of white brined cheese. *J. Food Hyg.*, **1(2)**: 69-78.
  26. Mohammadi, L., Hassanzadeh Khankahdani, H. and Tanaka, F. 2020. Effect of *Aloe vera* Gel Combined with Basil (*Ocimum basilicum* L.) Essential Oil as a Natural Coating on Maintaining Post-Harvest Quality of Peach (*Prunus persica* L.) during Storage. *IOP Conference Series: Earth and Environmental Science*, **594(1)**: 012008
  27. Navarro, D., Díaz-Mula, H. M., Guillén, F., Zapata, P. J., Castillo, S., Serrano, M., Valero, D. and Martínez-Romero, D. 2011. Reduction of Nectarine Decay Caused by *Rhizopus stolonifer*, *Botrytis cinerea* and *Penicillium digitatum* with *Aloe vera* Gel Alone or with the Addition of Thymol. *Int. J. Food Microbiol.*, **151**: 241–246.
  28. Nielsen, V. and Rios, R. 2000. Inhibition of Fungal Growth on Bread by Volatile Components from Spices and Herbs and the Possible Application in Active Packaging with Special Emphasis on Mustard Essential Oil. *J. Food Microbiol.*, **60(2-3)**: 219-29.
  29. Otero, V., Raquel, B., Santosa, J., odríguez-Calleja, M. R., Nerín, C. and García-López, M. 2014. Evaluation of Two Antimicrobial Packaging Films against *Escherichia coli* O157:H7 Strains *in Vitro* and during Storage of a Spanish Ripened Sheep Cheese (Zamorano). *Food Control.*, **42**: 296-302.
  30. Ortega-Toro, R., Collazo-Bigliardi, S., Roselló, J., Santamarina, P. and Chiralt, A.

2017. Antifungal Starch-Based Edible Films Containing *Aloe vera*. *Food Hydrocoll.*, **72(2)**: 1–10
31. Pantaleão, I., Pintado, M. E. and Poças, M. F. 2007. Evaluation of Two Packaging Systems for Regional Cheese. *Food Chem.*, **102 (2)**: 481–487.
  32. Pauli, A. 2006.  $\alpha$ -Bisabolol from Chamomile- A Specific Ergosterol Biosynthesis Inhibitor. *J. Aromathe.*, **16**:5-21.
  33. Pieretti, G. G., Pinheiro, M. P., Scapim, M. R. D. S., Mikcha, J. M. G. and Madrona, G. S. 2019. Effect of an Edible Alginate Coating with Essential Oil to Improve the Quality of a Fresh Cheese. *Acta. Sci. Technol.*, **41**: 36402.
  34. Ramasubramanian, T. S., Sivakumar, V. T. and Thirumalai, A. V. 2010. Antimicrobial Activity of *Aloe vera* (L.) Burm. against Pathogenic Microorganisms. *J. Bio. Sci. Res.*, **4(2)**: 251–258.
  35. Ramos, O. L., Pereira, J. O., Silva, S. I., Fernandes, J. C., Franco, M. I., Lopes-da-Silva, J. A., Pintado, M. E. and Malcata, F. X. 2012. Evaluation of Antimicrobial Edible Coatings from a Whey Protein Isolate Base to Improve the Shelf Life of Cheese. *J. Dairy Sci.*, **95(11)**: 6282-92.
  36. Raspo, M. A., Vignola, M. B., Andreatta, A. E. and Juliani, H. R. 2020. Antioxidant and Antimicrobial Activities of Citrus Essential Oils from Argentina and the United States. *Food Biosci.*, **36(3)**: 27-38.
  37. Rezaei, M., Yahyaei, M., Parviz, M. and Khodaei motlagh, M. 2010. A Survey of Microbial Contamination in Traditional Cheese Distributed in Markazi Province in Iran. *J. Health Environ.*, **7(1)**: 115-121.
  38. Reynolds, T. and Dweck, A.C. 1999. *Aloe vera* Leaf Gel. A Review Update. *J. Ethnopharmacol.*, **21**: 68- 89.
  39. Roy, B.C., Hoshino, M., Ueno, H., Sasaki, M. and Goto, M. 2007. Supercritical Carbon Dioxide Extraction of the Volatiles from the Peel of Japanese Citrus Fruits. *J. Essent. Oil Res.*, **19**: 78-84.
  40. Sambu, S., Hemaram, U., Murugan, R. and Alsofi, A. A. 2022. Toxicological and Teratogenic Effect of Various Food Additives: An Updated Review. *Biomed. Res. Int.*, **24(1)**: 1-11.
  41. Saritha, V., Anilakumar, K. R. and Khanum, F. 2010. Antioxidant and Antibacterial Activity of *Aloe vera* Gel Extracts. *Int. J. Pharm. Biol. Sci.*, **1**: 376–384.
  42. Sengun, I., Yaman, D. and Gonul, S. 2008. Mycotoxins and Mould Contamination in Cheese: A Review. *World Mycotoxin J.*, **1(3)**: 291 – 298
  43. Shenbagam A., Kumar, N., Rahul, K., Upadhyay, A., Gniewosz, M. and Kieliszek, M. 2023. Characterization of Aloe Vera Gel-Based Edible Coating with Orange Peel Essential Oil and Its Preservation Effects on Button Mushroom (*Agaricus bisporus*). *Food Bioproc. Tech.*, **131(4)**: 1-22.
  44. Sumi, F. A., Sikder, B., Rahman, M. M., Lubna, S. R., Ulla, A., Hossain, M. H., Jahan, I. A., Alam, M. A., and Subhan, N. 2019. Phenolic Content Analysis of Aloe vera Gel and Evaluation of the Effect of Aloe Gel Supplementation on Oxidative Stress and Fibrosis in Isoprenaline-Administered Cardiac Damage in Rats. *Prev. Nutr. Food. Sci.*, **24(3)**: 254-264.
  45. Suriati, L., Utama, I. M. S., Harjosuwono, B. A. and Gunam, B. W. 2020. Stability Aloe Vera Gel as Edible Coating. *Earth Environ. Sci.*, **411(2)**: 1-6.
  46. Suriati, L. 2018. Studies the Resistance to Oxidation and the Changes Phases against the Characteristics of Physicochemical *Aloe vera* Gel. *J. Bio. Chem. Res.*, **35(2)**: 670-679.
  47. Tan, Q., Ai, M. and Minh, N. 2011. Volatile Constituents of Essential Oil from *Citrus sinensis* Grown in Tine Giant Province, Vietnam. *Asian J. Food Agro. Ind.*, **4(3)**: 183-186
  48. Trmčić, K., Chauhan, A., Kent, D. J., Ralyea, R. D., Martin, N. H., Boor, K. J. and Wiedmann, M. 2016. Coliform Detection in Cheese Is Associated with Specific Cheese Characteristics, But No Association Was Found with Pathogen Detection. *J. Dairy Sci.*, **99(8)**: 6105-6120.
  49. Tzortzakis, N., Xylia, P. and Chrysargyris, A. 2019. Sage Essential Oil Improves the Effectiveness of *Aloe vera* Gel on Postharvest Quality of Tomato Fruit. *Agron.*, **9(10)**: 635-643.
  50. Wagh, Y. R., Pushpadass, H. A., Magdaline Eljeeva Emerald, F. and Surendra Nath, B. 2013. Preparation and Characterization of Milk Protein Films and Their Application for Packaging of Cheddar Cheese. *Food Sci. Technol.*, **51(12)**: 3767-3775.



51. Wang, Q., Yu, H., Tian, B., Jiang, B., Xu, J., Li, D., Feng, Z. and Liu, C. 2019. Novel Edible Coating with Antioxidant and Antimicrobial Activities Based on Whey Protein Isolate Nanofibrils and Carvacrol and Its Application on Fresh-Cut Cheese. *J. Coat*, **9(9)**: 583-591.
52. Yerlikaya, O. and Ozer, E. 2014. Production of Probiotic Fresh White Cheese Using Co-Culture with *Streptococcus thermophilus*. *Food Sci. Technol.*, **34(3)**: 1-10.
53. Yilmaz, F. and Dagdemir, E. 2012. The Effects of Beeswax Coating on Quality of Kashar Cheese during Ripening. *Int. J. Food Sci. Tech.*, **47**: 2582-2589.
54. Zhong, Y., Cavender, G. and Zhao, Y. 2014. Investigation of Different Coating Application Methods on the Performance of Edible Coatings on Mozzarella Cheese. *LWT Food Sci. Technol.*, **56**: 1-8.

## تأثیر پوشش فعال مبتنی بر ژل آلونته ورا با اسانس پوست لیمو بر ماندگاری و ویژگی های کیفی پنیر

س.س. نژاد سجادی، ل. منصورى نژند، و ف. شهدادی

### چکیده

در این مطالعه تأثیر پوشش خوراکی ژل آلونته ورا حاوی اسانس پوست لیمو (۰، ۱۰۰ و ۱۵۰ پی پی ام) بر ویژگی های کیفی نمونه های پنیر مورد بررسی قرار گرفت. تیمارها شامل ۴ گروه کنترل (بدون پوشش)، ژل آلونته ورا، ژل آلونته ورا و ۱۰۰ پی پی ام اسانس پوست لیمو و ژل آلونته ورا و ۱۵۰ پی پی ام اسانس پوست لیمو بود. این تیمارها به مدت ۶۰ روز از نظر خواص فیزیکوشیمیایی، بافتی، حسی و شمارش میکروبی مورد ارزیابی قرار گرفتند. یافته ها نشان داد که با افزایش مدت نگهداری، اسیدیته و نمک افزایش و pH و رطوبت کاهش یافت. در ارزیابی ویژگی های حسی، تأثیر تیمارها بر تمامی ویژگی های حسی به جز امتیاز رنگ معنی دار بود. نمونه های پوشش داده شده با ژل آلونته ورا و ۱۰۰ پی پی ام پوست لیمو بالاترین امتیاز طعم (۴/۹۷) را دریافت کردند. با افزایش زمان نگهداری، سختی، قابلیت جویدن و فنری بودن نمونه های پنیر افزایش یافت. چسبندگی نمونه ها تحت تأثیر مدت زمان نگهداری قرار نگرفت. در پایان زمان نگهداری، بیشترین شمارش کل میکروبی، کپک و مخمر مربوط به نمونه های پنیر شاهد (به ترتیب ۵/۳۷ و ۵/۶۲ log CFU/g) و کمترین مقدار مربوط به نمونه های پوشش داده شده با ژل آلونته ورا و ۱۰۰ پی پی ام پوست لیمو (به ترتیب ۳/۹۲ و ۳/۷۶ log CFU/g) بود. به طور کلی استفاده از پوشش خوراکی تهیه شده با یل آلونته ورا و غلظت های کمتر اسانس پوست لیمو (۱۰۰ پی پی ام و کمتر) باعث بهبود ظاهر و طعم نمونه های پنیر طی ۶۰ روز نگهداری گردید.