# Abundance and Diversity of AM Fungi across a Gradient of Land Use Intensity and Their Seasonal Variations in Niligiri Biosphere of the Western Ghats, India

R. Lakshmipathy<sup>1\*</sup>, A. N. Balakrishna<sup>2</sup>, and D. J. Bagyaraj<sup>2</sup>

# ABSTRACT

The impact of land use intensity on the abundance and diversity of arbuscular mycorrhizal fungi (AMF) was investigated at six land use types viz., natural forest, grassland, acacia plantations, cardamom plantations, coffee plantations and paddy fields in the Niligiri Biosphere of the Western Ghats in South India. There was no significant difference in AMF root colonization in different land use types during pre-monsoon but there was a significant difference in AMF root infection ratings between different land use types during post-monsoon season, where it was higher in natural forests and grasslands. The AMF spore density and infective propagules were significantly higher in grasslands and acacia plantations compared to all other land use types during both seasons. Except for paddy fields, the spore density and number of infective propagules were higher in post-monsoon season compared to premonsoon season in other land use types. The numbers of AMF species identified were 56 during pre-monsoon and 67 species during post-monsoon season suggesting seasonal variations in diversity. During both seasons Glomus fasciculatum was recorded in maximum number of sampling points across the landscape followed by G. geosporum during premonsoon and G. mosseae during post-monsoon season. The species diversity was highest in natural forests and grasslands as compared to other land use types in both seasons. The species richness index for AMF was highest in natural forests and least in paddy fields during both study periods. The sand content, bulk density, total N, organic C, alkaline and acid phosphatases positively correlated with AMF activity while clay, silt, K, total P and available P were negatively correlated.

Keywords: AMF, Infective propagules, Land use types, Root colonization, Spore density.

# INTRODUCTION

A major component of the soil mycobiota in most agro-ecosystems is the arbuscular mycorrhizal fungi (AMF). AMF exist inside the root and in the soil. The intraradical mycelium consists of hyphae and other fungal structures, such as arbuscules and vesicles; the extraradical mycelium forms spores, explores soil and new areas for colonization and absorbs nutrients (Tommerup and Sivasithamparam, 1990). In the recent system of classification, all the AM fungal species are placed in four orders *i.e.* Archaeosporales, Diversisporales, Glomerales and Paraglomerales which comprise 13 families and 19 genera that belong to class Glomeromycetes of the phylum Glomeromycota (Sieverding and Oehl, 2006; Oehl *et al.*, 2008; and Palenzuela *et al.*, 2008).

Since these fungi are obligate symbionts, their population and diversity may be determined by the plant species present in the given ecosystem. Apart from plant species, human activities also affect these fungi which play an important role in

<sup>&</sup>lt;sup>1</sup> Post Harvest Technology Center, Agricultural College Campus, Bapatla-522 101, India.

<sup>\*</sup> Corresponding author; e-mail: lakshmipatty@yahoo.com

<sup>&</sup>lt;sup>2</sup> Department of Agricultural Microbiology, University of Agricultural Sciences, GKVK, Bangalore-560 065, India

carbon allocation, nutrient cycling and maintenance of diversified ecosystems (Doss and Bagyaraj, 2001). The presence of these fungi and their genetic and functional diversities are important for both plant community and ecosystem productivity. Improved plant growth due to inoculation of plants with AM fungi has been demonstrated especially under P deficient condition. The growth improvement is mainly because of enhanced P uptake. AMF can also enhance tolerance or resistance to root pathogens and abiotic stresses such as drought and metal toxicity (Clark and Zeto, 2000; Auge, 2004).

Western Ghats of India is identified as one of the biodiversity hotspots of the world (Myers et al., 2000). Information on the distribution and frequency of occurrence of specific AM fungi in the Nilgiri biosphere of the Western Ghats area is very scarce. A few studies performed so far have recorded the occurrence of Glomus species in the root zone soils of different tree species (Muthukumar and Manian. 1993: Vasanthakrishna et al., 1994). Similarly, Lakshmipathy et al. (2004) recorded Glomus etunicatum in the root zone soil of cashew. The mycorrhizal activity in terms of root colonization was examined in 59 different forest tree species and the intensity of colonization was found to be high in four species, moderate in 23 species and low in 32 species (Byra Reddy et al., 1994). Patricia et al. (2009) recovered a total number of 24 AM fungal species in pristine forest ecosystem of Amazon. Santhaguru et al. (1995) observed mycorrhizal root colonization in twenty species of tree legumes in the Eastern Ghats and found altogether 21 species of AMF belonging to six genera viz. Acaulospora, Entrophospora, Glomus, **Sclerocystis** Gigaspora, and Scutellospora. Therefore, appears it that occurrence of AM fungi in different forest tree species defines ecological niches.

In tropical soils, application of organic matter stimulated the proliferation of AMF. This was attributed to the low organic matter content in tropical soils (Harinikumar and Bagyaraj, 1989). The addition of organic amendments such as paddy straw, maize straw and pongamia leaf increased the mycorrhizal activity. Of the three amendments used, addition of pongamia leaf improved AM fungal infection to the maximum, followed by maize straw (Harinikumar and Bagyaraj, 1988). High P availability is reported to be negatively correlated with AM fungal activity (Krishna and Bagyaraj, 1982). Balakrishna et al. (2001) reported that application of inorganic fertilizers negatively affected native AM fungal spore density, infective propagule numbers and per cent root colonization in a finger millet- maize-fallow crop rotation system. Furthermore, studies conducted on the effect of mono and mixed cropping systems on AMF population in soil have revealed that mixed cropping of soybean and maize stimulated the proliferation of AM fungi compared to mono-cropping of maize or soybean (Harinikumar et al., 1990).

Seasonal fluctuations also seem to have a considerable impact on the AM fungal development in the soil. Harinikumar and Bagyaraj (1988) and Mallesh and Bagyaraj (1991) showed that AM fungi sporulate during winter in the tropics. The optimum temperature for sporulation by mycorrhizal fungi appears to be around  $25^{\circ}$ C. To our knowledge, the AMF abundance and diversity occurring over broad land use types and in different seasons under tropical conditions have not yet been investigated in India. Thus, the present study was undertaken in the Niligiri Biosphere of the Western Ghats of Karnataka in South India.

#### MATERIALS AND METHODS

#### **Study Area**

The benchmark area, Koothy village of Somwarpet taluk is located in the Kodagu district of Karnataka (Figure 1). A wide range of land use types with a diversified plant species composition is found in the region. This benchmark area is situated in the Nilgiri Biosphere Reserve of the northern region and lies between 12°40' 03" N-12° 42' 19" N and 75° 47<sup>°</sup> 10<sup>°</sup> E–75<sup>°</sup> 79<sup>°</sup> 14<sup>°</sup> E. The annual rainfall of the area ranges from 2,000 mm to 3,500 mm. Most of the rainfall is drawn from southwest monsoon during June-August period. Four seasons can be clearly distinguished; summer from March to May, monsoon from June to September, post-monsoon during October and November, and winter during December to February (Temp.  $6.2-12.5^{\circ}$ C). The temperature begins to increase from March to April, with a



Figure 1. Location of the study area-Koothy, Somvarpet taluk, Kodagu, Karnataka.

mean daily maximum of  $28.6^{\circ}C$  and mean daily minimum of  $17.8^{\circ}C$  and reaching as high as 32 to  $35^{\circ}C$  during April or May.

Coffee and cardamom plantations cover a major part of the study area. The natural forests at the periphery of the plantations are evergreen with varying levels of degradation. A few patches of *Acacia auriculiformis* plantations (monoculture) and grassy patches are found adjacent to the forests. Rain-fed agriculture is practiced in the valleys with one paddy crop every year during the rainy season. Additionally, crops like chilly and short duration grain legumes are also grown in the summer season utilizing the residual moisture and sparse rainfall of northeast monsoon.

#### Land Use Pattern

Six land use–land cover types could be distinguished in the study area using a satellite database of the year. A brief description of these land use types is presented in Tables 1 and 2.

The soil samples were collected in premonsoon season (February) and post-monsoon season (October) for the analysis of soil physical, chemical, biochemical and microbiological properties.

## **Soil Sampling**

A triangle of  $50 \times 50 \times 50$  m was laid at each sampling point. From the center of the triangle at



Figure 2. Soil sampling procedure for studying soil properties and AM fungi.

Table 1.	Dominant	vegetation	types	and	management	practices	of the	different	land u	ise t	ypes	in tl	ne
study are	a.												

Land use type	Major vegetation	Fertilization (Kg ha <sup>-1</sup> )	Plant protection	Cultivation intensity
Natural forest	These forests are little disturbed and are adjacent to coffee and cardamom plantations. There are as many as 153 species of plants. Dominated by <i>Caryota urens, Olea dioica, Canthium dicoccum,</i> <i>Artocarpus heterophyllus</i> and <i>Dimoacarpus longan</i> trees, shrubs like <i>Leea indica, Dichapetalum</i> gelonioides, <i>Flacourtia indica, Nilgirianthus</i> <i>heyneanus</i> etc. and herbs like <i>Brachiaria</i> <i>milliformis, Justisia trinervia, Blumea barabata,</i> <i>Alysicarpus vagnailis, Archidendron</i> <i>monodelphum, Derris indica, Derris scandens,</i> <i>Pongamia pinnata</i> were the dominant on the forest floor.	None	None	None
Grasslands	These are the open patches found in slightly elevated regions of the Ghats and adjoining agricultural fields. Dominated by grasses like Panicum repens, Sporobolus diander and herbs like Stachytarpheta indica, Borreria articularis, Oldenalnadia corymbosa, Derris seandens, Desmodium triquetrum, Mimosa pudica, Zornia gibbosa etc.	None	None	None
Acacia plantations	These are the monoculture plantations of <i>Acacia</i> <i>auriculiformis</i> of 15-25 years old raised in the open and disturbed areas. The ground flora is mainly dominated by species like <i>Stachytarpheta indica</i> , <i>Brachiaria milliformis</i> , <i>Centella asiatica</i> and shrubs like <i>Lantana camara</i> , <i>Maesa indica</i> , <i>Randia</i> <i>dumatorum</i> .	None	None	None
Coffee plantations	About 106 species of plants belonging to trees, shrubs and herbs were recorded in addition to coffee plantations. The most dominant tree species are <i>Grevilia robusta</i> , <i>Caryota urens</i> , <i>Artocarpus</i> <i>heterophyllus</i> , <i>Acrocarpus fraxinifolius and Derris</i> <i>scanden</i> , <i>Mucuna prurienss</i> . Most of the plantations have naturally occurring species maintained as shade trees and in a few plantations <i>Grevilia robusta</i> is planted as shade trees. The <i>Brachiaria molliformis</i> grass is the most dominant ground flora.	Mineral and farmyard manure as per the University recommendat ion	Chemical: Fungicides and Insecticide s twice a year	Very low, since it is a perennial crop
Cardamom plantations	One hundred and thirty four species of plants belonging to trees, shrubs and herbs were recorded. The plant species like Olea dioica, Litsea floribunda, Caryota urens, Cinnamomum zeylanicum, Acacia concinna, Acrocarpus fraxinifolius, Albizia lebbeck, Albizia odoratissima, Dalbergia latifolia, Derris scandens and Mucuna prurienss were the most dominant flora with Brachiaria molliformis grass as the dominant ground flora.	Mineral and farmyard manure as per the University recommendat ion	Chemical: Fungicides and Insecticide s twice a year	Very low, since it is a perennial crop
Paddy fields	The agricultural crop grown in that region is mainly rain fed rice. The ground flora varies with the season and as many as 47 species of herbs are recorded. The dominant herbaceous weeds present in the area are <i>Panicum repens</i> , <i>Grangea</i> maderaspatana, <i>Centella asiatica</i> , <i>Blumea</i> barbata	Mineral and farmyard manure	Chemical: Fungicides and Insecticide s 2 times for each crop.	High:2-3 crops/year i.e. paddy followed by pulses

Land use types pH		Organic C	Total N	Total K	Total P	Avail. P	
-	$H_2O$	KCl	(g kg')	$(mg kg^{-1})$	(mg kg ')	$(mg kg^{-1})$	(mg kg')
Natural forests	6.20	5.24	15.5	46.90	232.0	1.9	0.63
Grasslands	5.57	4.37	11.8	40.88	175.5	2.47	0.80
Acacia plantations	5.51	4.43	13.8	42.35	240.0	1.74	1.04
Cardamom plantation	6.16	5.20	10.4	95.80	261.0	3.58	1.38
Coffee plantations	6.16	5.03	11.5	47.90	283.5	3.46	1.23
Paddy fields	5.30	4.21	7.7	37.24	189.0	3.86	1.48

Table 2. Chemical soil parameters of the different land use types differing in land use intensity.

a distance of three and six meters two concentric circles were drawn as shown in the figure. At equidistance from each other three soil cores (6.25 cm dia.) of 0-20 cm depth were collected on the circumference of each of the circle as shown in the (Figure 2) avoiding the litter above the ground. The six soil samples thus collected were mixed together and a composite soil sample was drawn using quaternary technique. The soil sample was collected along with the root bits, the root bits were separated and soil was divided into two parts. The root bits collected were immediately transferred to a fixative. One part was air dried and used for chemical analysis and the other part was stored at 5°C in a refrigerator. The soil samples were collected at nine sampling points from natural forests, eight from grasslands, six from acacia plantation, thirteen from cardamom plantation, sixteen from coffee plantations and eight from paddy fields.

# Soil Physical, Chemical and Biological Properties

The clay, silt and sand were quantified by mechanical analysis as outlined in the 'TSBF Hand Book of Methods' by Anderson and Ingram (1989). The organic carbon was determined by the modified Walkley-Black method (Anderson and Ingram, 1989). The total nitrogen in the soil samples was analyzed by semi micro-Kjeldhal method (Jackson, 1973) using a Gerhardt auto analyzer. The total phosphorus content of the soil samples was determined by vanado-molybdo phosphoric yellow colour method in nitric acid system (Jackson, 1973). The soil samples used in this study were acidic and hence Bray's extracting solution containing 0.03 N NH<sub>4</sub>F in 0.025 N HCl was used (Jackson, 1973). The available potassium was determined by following the ammonium acetate method as described by Merwin and Peach (1951). The bulk density was determined following standard procedures (Anderson and Ingram, 1989). The acid and alkaline phosphatase activities were estimated as per the procedure (Eivazi and Tabatabai, 1977). The results of the soil analysis of different sites are given in the Table 2.

#### Assessment of the Mycorrhizal Parameters

The following mycorrhizal structures were determined: root colonization was determined by gridline intersection method (Giovannetti and Mosse, 1980) by staining the root bits with trypon blue. Spore density was 0.05% determined by extracting spores from soil by wet sieving and decantation method (Gerdemann and 1963) and infective Nicolson, propagule numbers 10 fold MPN method by (Sieverding, 1991).

# Identification of the Diversity of AM Fungi

Spores extracted directly from soil can be used for identification, but better identification of AMF is possible by "host baiting technique" or "trap pot culturing". Because spores from baits are in a better state, they can be identified to species level by using spore morphological charecteristics (Bagyaraj and Sturmer, 2008).

The composite samples obtained from the sampling points were transferred to the

laboratory. The soil test sample was mixed with sterile sand soil (1:1) mix (50% soil test sample+50% sterile sand soil mix) and planted with suitable trap crops such as a mixture of sorghum and cowpea under glass house conditions. Watering was done every other day. After 3-4 months, the potting mix was wet sieved and the spores were observed under a compound microscope. Morphologically similar spores were picked and the population of each spore type was enumerated from a respective field sample. The spores of each type were brought into pot culture by funnel technique after surface sterilization of the spores (Nicolson, 1967) with aqueous solution containing 200-ppm an streptomycin sulphate and 2% chloramine T. The spores were mounted on a glass slide in lactoglycerol. They were later identified with the help of "Manual for identification of VA mycorrhizal fungi" by Schenck and Perez (1990) and the Morton: INVAM Joe website by http://invam.caf.wvu.edu.

#### **Statistical Analysis**

The data collected from the field experiments were subjected to statistical analysis following the statistical package Excel and SYSTAT version 10.2. Arc sin transformations were done (Snedecor and Cochran, 1968) wherever necessary. The treatment means were separated by the Duncan's Multiple Range Test and by Probability Matrix. The Shannon-Weiner diversity index, Jakknife's species richness index, Sorenson similarity index for AMF between different land use types were calculated (Krebs, 1989).

#### RESULTS

#### **Mycorrhizal Root Colonization**

AMF root colonization was studied during two different seasons *viz*. pre and post-monsoon. Significant differences in root colonization between different land use types were observed only during post-monsoon season (Table 3). AMF root colonization was significantly higher in roots collected from natural forests and a grassland compared to other land use types and was lowest in paddy fields. The root colonization was significantly higher in post-monsoon season compared to pre-monsoon in natural forests, grasslands and acacia plantations while it was more during pre-monsoon season in coffee plantations, cardamom plantations and paddy fields.

#### **Spore Density and Infective Propagules**

In both seasons, the spore density in soils collected from acacia plantations and grasslands was significantly higher as compared to all other land use types but it was the lowest in case of paddy fields (Table 3). The average spore density of all land use types in two different seasons was higher during post-monsoon season compared to pre-monsoon season.

The infective propagule numbers in soils collected from acacia plantations and grasslands were on par with each other and significantly higher as compared to the other land use types

**Table 3.** AMF root colonization, Spore load and Infective propagules in different land use types in pre and post-monsoon seasons.

	AMF root colonization (%)		Spore density	(No 50 g <sup>-1</sup> soil)	Infective propagules (No g <sup>-1</sup> soil)		
	Pre-	Post-	Pre-	Post-	Pre-	Post-	
Land use types	monsoon	monsoon	monsoon	monsoon	monsoon	monsoon	
Natural forests	59	72 <sup>a</sup>	201 <sup>cd</sup>	374 <sup>c</sup>	305 <sup>c</sup>	559°	
Grasslands	64	$68^{ab}$	641 <sup>ab</sup>	654 <sup>a</sup>	$690^{ab}$	839 <sup>a</sup>	
Acacia plantations	57	64 <sup>b</sup>	767 <sup>a</sup>	616 <sup>ab</sup>	775 <sup>a</sup>	737 <sup>ab</sup>	
Cardamom plantations	64	49 <sup>d</sup>	176 <sup>cdef</sup>	275 <sup>d</sup>	177 <sup>cd</sup>	407 <sup>d</sup>	
Coffee plantations	66	57 <sup>cd</sup>	258 <sup>d</sup>	316 <sup>d</sup>	260 <sup>cd</sup>	443 <sup>d</sup>	
Paddy fields	48	33 <sup>e</sup>	184 <sup>cde</sup>	201 <sup>e</sup>	215 <sup>cd</sup>	184 <sup>e</sup>	
F test at 0.05%	NS	*	*	*	*	*	
Seasons mean	60	57	371	405	404	528	

\* Significant.

Mean values followed by the same superscript in each column do not differ significantly at P=0.05 level by DMRT.

during pre-monsoon season (Table 3). The paddy fields recorded the least. A similar trend was observed in the post-monsoon season too. The average numbers of infective propagules in soils collected from all land use types were higher during post-monsoon season compared to premonsoon season.

# Effect of Soil Properties on AM Fungi

The influence of soil physical, chemical and bio-chemical properties on different mycorrhizal parameters was calculated. Among the soil physical properties, sand and bulk density had a positive influence on root colonization, spore density and infective propogule numbers, while the clay content was negatively correlated to root colonization, spore density and infective propagules (Table 4). The chemical properties of the soils, such as organic C, had a significant positive influence on the formation of AMF in soil while the total P and available P had a negative influence on AMF formation. The biochemical properties of soil such as the acid and alkaline phosphatase activities were positively correlated with mycorrhizal root infection.

# Species Abundance and Diversity of AM Fungi in Different Land Use Types

Altogether, a total of 67 species of AM fungi were isolated from all six different land use types across the sampling site, out of which 56 were collected during pre-monsoon and 67 species during post-monsoon season (Table 5). There were 12 species belonging to the genus Acaulospora, one to Archeospora, 48 to Glomus, one to Intraspora, one to Pacispora and 3 to Gigaspora. G. mosseae, A. lacunosum, G. fasciculatum, A. bireticulata and G. geosporum were found virtually in all land use types either in the pre- or post monsoon seasons. Some species such as G. geosporum, G. citricolum, G. heterosporum and G. halonatum were common to natural forests, grasslands, acacia plantations and coffee plantations in both seasons. G. fulvum was associated with only cardamom plantations in both seasons and was present in natural forests only in the pre-monsoon season. The species Arccheospora trappei was recorded only in the grasslands in both seasons. G. australe, G. intraradices, A. mellea, G. hoi, A. spinosa, G. magnicaulis and A. morrowe, were recorded both in the pre- and post monsoons. The species, G. phansihalos, Gi. albida, G. pachycaulis and P. scintillans were recorded only during premonsoon while others such as G. reticulatum, G. caledonium, A. delicata, I. schenckii, G. diaphanum and G. pustulatum were recorded only during post-monsoon season and almost confined to grasslands.

#### AMF Species Abundance across the Landscape Over Two Different Seasons

The abundance of different AMF species over different land use types indicated a higher incidence during post-monsoon season than in pre-monsoon season (Table 6). During postmonsoon season, the majority of the species produced spores abundantly in field soil compared to pre-monsoon season. A few species

**Table 4.** Correlation matrix between soil properties and components of AM fungi in different land uses types.

Sol colonization	Spore density	Infective propagules
(%)	50 g⁻¹ soil	$(\text{IP g}^{-1} \text{ soil})$
-0.039	-0.067	-0.086
-0.203	-0.237	-0.142
0.122	0.151	0.091
0.068	0.302*	0.446*
0.647**	0.456*	0.400*
0.836**	0.486*	0.465*
0.796**	0.379*	0.465*
0.115	0.058	0.103
0.065	-0.073	-0.079
-0.279*	-0.308*	-0.232
-0.441*	-0.366*	-0.224
	(%) -0.039 -0.203 0.122 0.068 0.647** 0.836** 0.796** 0.115 0.065 -0.279* -0.441*	$\begin{array}{c ccccc} (\%) & 50 \ g^{-1} \ {\rm soil} \\ \hline -0.039 & -0.067 \\ -0.203 & -0.237 \\ 0.122 & 0.151 \\ 0.068 & 0.302* \\ 0.647** & 0.456* \\ 0.836** & 0.486* \\ 0.796** & 0.379* \\ 0.115 & 0.058 \\ 0.065 & -0.073 \\ -0.279* & -0.308* \\ -0.441* & -0.366* \\ \hline \end{array}$

\* Significant at P= 0.05 level.

\*\* Highly significant at P= 0.01.



Table 5. Occurrence and spore numbers (No 50 g	<sup>1</sup> soil) of AMF species in different land use types in pre- (PRM) and post-
monsoon (POM) seasons.	

Sl. No	AMF species	Nat fo	ural rest	Gras	slands	Ac: plant	acia ations	Carda planta	amom ations	Co plant	ffee tations	Pa fie	ddy elds
	-	PRM	POM	PRM	POM	PRM	POM	PRM	POM	PRM	POM	PRM	POM
1	Glomus mosseae	15	526	69	366	44	620	84	319	527	854	38	215
2	Acaulospora lacunosa	36	132	306	491	138	203	250	55	178	198	91	27
3	G. fasciculatum	219	776	401	1013	696	897	278	881	573	795	50	
4	A. bireticulata	65 82	180	230	749 407	383 67	293	251	612	166	384 612	24 114	
6	G. geospoi uni G. citricolum	78	403	230	119	90	209	251	52	2	128	114	
7	G. heterosporum	29	117	419	273	20	207		02	42	77		
8	G. halonatum	67	101	202	456					121	181		
9	A. scrobiculata	18	88					53	241	210	523		24
10	G. aggregatum	3	150				114			151	845	48	544
11	G. globiferum	8	44			34	73						
12	G. maculosum	92	155	149	157	154	402	197	152			78	182
13	G. multicaulis	26	187			96	293	128	329	21	410		
14	A. nicolsoni	41	133			276	67						
15	G. manihotis	59	141	242	260	85	81					23	75
10	G. albidum	101	101	542	368	272	288	171	100				
18	G. etunicatum G. monosporum	39 124	280	51	95		151	30	30	80	68		
19	G. monosporum G. canadense	67	71					50	46	00	00		
20	G. delhiense	25	366						10				
21	G. boreale	100	180										
22	G. multisbstensum	27	58										
23	G. dimorphicum	17	42										
24	G. versiforme	27	146										
25	S. calospora	12	105										
26	G. verrucosa	23	129										
27	G. fulvum	19						228	259				
28	G. australe		96	454	178								
29	G. intraradices		120	88	459						137	20	24
30 21	A .appendicula		22 87	115	70				22		50	28	34
22	A. idevis C. algraidaum		07		78				33		38	17	41
32	G. lentotichum		24									47	41
34	G. ambisporum		80				377						
35	A. mellea			225	217		118	319	590				
36	G. hoi			121	225	77	145	334	123	14	98	7	
37	A. spinosa			60	124	459	311	3	43				
38	G. magnicaule			116	102					213	160		
39	A. morrowae			188	231								
40	Ar. trappei			475	160	05	127	26	0.1				
41	G. phansinalos G. aaspora albida			99 58		28 272	137	30	81				
42	G gaspora albiaa G. pachycaulis			218		212		28	103				
44	pacispora scintillans			58									
45	G. pustulatum			17	109								
46	G. reticulatum				554								
47	G. caledonium				78				504				
48	A. delicate				44			244	39				
49	Intraspora shenkii				40							_	
50	G. diaphanum				56	51	96			96	50	7	66
51	G. segmentatus					90	98					21	
52 52	S. persica					158	238					21	
55	G. geruemunnu G. macrocarpum					45	380						
55	G. radiatum					138	138		41	113	386		
56	G. lacteum					230	128	75	234	28	42	69	194
57	A. dilatata							16	131	234	263		
58	G. invermaium							34	159	252	153		
59	G. constrictum								21	375	325		
60	G. clavispora									13	58		
61	G. tenebrosum									209			
62	G. deserticola									73			
63	Scutellospora										52		
64	heterogama										10	144	60
04 65	Gi .margarlia G. tortuosum										48	144 ⊿1	51
66	G. tortuosum Gi rosea											+1	30
67	G. clarum				97								37
					~ .								

**Table 6**. Abundance of spores of different species of AM fungi across sampling site during pre- and postmonsoon seasons in different land use types.

	Spore	abund	lance (N	No 50 $g^{-1}$ s	oil)	
-	Pre-	-mons	soon	Post n	nons	oon
A.bireticulata	1140	±	216	1678	±	269
A.lacunosa	749	±	110	1106	±	167
A.scrobiculata	28	±	83	852	±	206
A.nicolsoni	317	±	111	200	±	56
A.appendicula	143 ND <sup>a</sup>	±	27	89	±	24
A.laevis	ND 544	± +	ND 142	250	±	38 221
A. meenea A. marrowae	188	± +	145	0.04 231	± +	231 04
A.marrowae A dilatata	250	+	94	394	+ +	110
A.delicata	244	+	100	83	±	28
A.spinosa	522	±	109	478	±	115
Ar.trappei	475	±	64	160	±	64
G.albidum	515	±	137	817	±	151
G. ambisporum	457	±	63	ND	±	ND
G.australe	454	±	63	274	±	82
G.aggregatum	202	±	61	1653	±	343
G.borrealis	100	±	41	180	±	73
G. caledonium	ND	±	ND	502	±	202
G.canedense	113	±	27	/1	±	31
G.citricolum C. alamai daumi	1/9	±	42	911	±	142
G.clarolaeum C. alamum	47 ND	±	19 ND	156	± +	40
G. clavispora	13	- -	5	97 58	т +	40 24
G.cuvisporu G.constrictum	375	÷ +	153	346	- +	131
G.delhiens	25	+	10	366	÷ +	149
G.deserticola	73	±	30	ND	±	ND
G.diaphanum	154	±	40	268	±	38
G.dimorphicum	17	±	7	42	±	17
G.etinicatum	261	±	66	729	±	113
G.fulvum	247	±	92	259	±	106
G.geosporum	1123	±	121	2030	±	256
G.gerdimanii	45	±	18	61	±	25
G.globiferum	42	±	14	117	±	32
G.halonatum	390	±	83	738	±	179
G.heterosporum	490	±	166	46/	±	108
G.hoi C inturnadiana	223	± +	127	591 716	± +	8/
G.intraraaices G.invermaians	402	± +	52 101	508	± +	1/0
G.invermations G.lacteum	179	+	86	588	⊥ +	02 00
G.leptotichum	ND	∸ +	ND	24	÷ +	10
G.macrocarpum	99	±	203	380	±	155
G.magnicaule	329	±	90	262	±	70
G.manihotene	167	±	36	297	±	59
G.monosporum	234	±	52	290	±	75
G.mosseae	777	±	251	2900	±	233
G.multisubstensum	27	±	11	58	±	24
G.multicaulis	27	±	46	1219	±	141
G.pachycaulis	317	±	11	103	±	42
G.phansihalos	220	±	59	218	±	57
G.pustulatum	17	±	1	109	±	44
G.radiatum	251	±	65 142	202 ND	±	155 ND
G.reticulatum	554 00	± +	145	ND 08	±	ND 40
G.segmentatus G.tenebrosum	200	± +	37 85	96 ND	± +	40 ND
G.tenebrosum G.tortuosum	117	+	17	51	⊥ +	88
G.tortuosum G versiforme	27	∸ +	11	146	÷ +	60
G.verrucosa	23	±	9	129	±	53
Gi.albida	715	+	109	817	_ +	115
Gi.margarita	144	±	59	117	±	31
Gi.rosea	ND	±	ND	39	±	16
I.schenckeii	ND	±	ND	40	±	16
P.scintillans	58	±	24	ND	±	20
S.calospora	15	±	5	105	±	43
S.heterogama	ND	±	ND	52	±	21
S.persica	158	±	55	258	±	105

such as A. delicata, A. morrowae, G. albidum, G. fulvum and G. magnicaule produced more spores during pre-monsoon than in post-monsoon season. Spores of A. laevis, Intraspora. schenckii, G. caledonium, G. clarum, G. leptotichum, G. reticulatum, G. ambisporum, Gi. Rose and, S. heterogama were noticed only during post-monsoon season while spores of G. deserticola and G. tenebrosum were observed only during pre-monsoon season.

# Diversity of AM Fungi in Different Land Use Types

Shannon-Wiener diversity index during premonsoon season was higher in natural forests compared to grasslands, acacia plantations, coffee plantations, cardamom plantations and paddy fields. A similar trend also existed in the post-monsoon season and the diversity was significantly higher in natural forests compared to acacia plantations, cardamom plantations, coffee plantations and paddy fields. The diversity index was significantly higher in post monsoon as compared to pre-monsoon season. Even within land use types, the diversity index was significantly higher in post-monsoon as compared to pre-monsoon season except in paddy fields (Table 7).

Jacknife's species richness index for AMF in different land use types during pre-monsoon season was higher in natural forests and the least in cardamom plantations. During post-monsoon season the AMF species richness index followed a similar trend. Seasonal variations in species richness index were higher during post-monsoon as compared to pre-monsoon season in all land use types except in paddy fields (Table 8).

#### DISCUSSION

The present investigation in the Nilgiri Biosphere of the Western Ghats in South India has shown that more intensive land use can have a negative impact on AMF, and thus in principle, some earlier studies on similar subjects e.g. Oehl *et al.* (2003) confirmed a decrease in AMF species richness with increasing land use intensity in Central Europe. Our studies also considered the effects of monsoon periods in different land use systems.

<sup>a</sup> Not Detected.

	No. of AMF species		Shannon-Weiner diversity index		
Land use types	Pre-monsoon	Post-monsoon	Pre-monsoon	Post-monsoon	
Natural Forests	27	33	4.41 <sup>a</sup>	4.59 <sup>a</sup>	
Grasslands	24	27	$4.20^{ab}$	4.45 <sup>ab</sup>	
Acacia Plantations	22	26	3.99 <sup>b</sup>	4.26 <sup>b</sup>	
Cardamom Plantations	19	26	3.57 <sup>cd</sup>	$4.08^{bc}$	
Coffee Plantations	23	25	3.80 <sup>c</sup>	4.04 <sup>bc</sup>	
Paddy Fields	16	13	3.26 <sup>d</sup>	3.01 <sup>d</sup>	
F test at 0.05%	-	-	*	*	
Season mean			3.87	4.07	

**Table 7.** AM fungal species diversity in different land use types during pre and post-monsoon seasons.

Note: Values in parenthesis represent Error Mean Sum.

\* Significant.

Mean values followed by the same superscript in each column do not differ significantly at P=0.05 level by DMRT.

Table 8. Jakknife's species richness index for AM fungi in different land use types during pre and postmonsoon seasons.

	Species richness index				
Land use types	Pre-monsoon	Post-monsoon			
Natural Forest	43.00	51.67			
Grasslands	37.13	48.50			
Acacia plantations	38.67	41.83			
Cardamom plantations	24.38	37.08			
Coffee plantations	30.44	34.37			
Paddy fields	25.63	20.87			

Although there were no significant differences in mycorrhizal root colonization between land use types during pre-monsoon season, significant differences were observed during post-monsoon season. The relative high root colonization in natural forests and grasslands may be related to the low disturbance of these ecosystems as compared to the other land use types. Also, the P contents may play an important role in mycorrhizal colonization and AMF spore production. Several studies have indicated that high available P-content (> 9 ppm) reduced mycorrhizal colonization and spore production (Koide, 1991; Lakshmipathy et al., 2002, 2003). The available-P in soils of the Indian land use types (Table 1) is lower compared to other land uses. This could be one of the reasons for variations in mycorrhizal root colonization in different land use types in this study. Pande and Tarafdar (2004) suggested that increased rainfall and relative air humidity increased AMF colonization. With the onset of rains in the postmonsoon season significant moisture prevailed in the soil that favoured better root growth of plant species and enhanced AMF root colonization.

## AMF Spore Density and Infective Propagules

The spore density and infective propagules were significantly more in grasslands and acacia plantations compared to natural forests, cardamom and coffee plantations with the lowest in paddy fields in both seasons. The floor of acacia plantations in the study area is also covered with patches of graminaceous species and acacia itself being a leguminous plant, could have caused higher spore density and infective propagules. Mycorrhizal population can also be affected by several agricultural practices such as ploughing, application of fertilizers, pesticides and herbicides which reduced the mycorrhizal spores and infective propagules (Boddington and Dodd, 2000; Balakrishna *et al.*, 2001; Kabir, 2005). These factors may be responsible for the lower number of spores and infective propagules in paddy fields, cardamom and coffee plantations.

Seasonal variations may also influence spore density and species diversity. Several workers (Mallesh and Bagyaraj, 1991; Picone, 2000) have reported earlier that the spore density and infective propagules were higher in winter than in summer suggesting that the mycorrhizal activity is favored during winter season. In this study, also such variations were observed between land use types.

# **Influence of Soil Properties**

Soil physical, chemical and biochemical properties influenced AMF root colonization, spore density and infective propagule numbers. Sand and bulk density had a positive influence, while clay and silt contents had a negative influence on AM fungi. In sandy soils, plant roots proliferated better because of good aeration; thereby possibly enhancing spore production. Similar observations have also been reported earlier (Pande and Tarafdar, 2004). Organic C content in soils is known to favor mycorrhizal infections (Johnson and Wedin, 1997; Pande and Tarafdar, 2004). The organic-C and N content of the soils in the study area showed a positive influence on AMF. Therefore, it suggests that the status of organic matter content in soils is important for mycorrhizal activity. The available and total P contents in soil play an important role in mycorrhizal colonization and spore production. Several studies have indicated that high available Pcontent (> 9 ppm) reduced mycorrhizal colonization and spore production (Koide, 1991; Lakshmipathy et al., 2002, 2003). The total P and available P contents in soils in the study area seem to have a significant negative influence on AMF activity.

#### **AMF Species Distribution**

Variations in AMF species numbers during different seasons can potentially be attributed to changes in moisture regimes in soil and other climatic conditions. More AMF species were recorded during rainy season as compared to summer season (Lovelock *et al.*, 2003; Husband

*et al.*, 2002). There was a strong shift in mycorrhizal communities and their numbers over time. They suggested that, during wet season a large number of newly germinated plant seedlings prevail, hence there were more species present. Few plant species, which may regenerate only during wet season, may vanish during dry season. Variations in species composition in different seasons are quite obvious in this study owing to the reasons already explained.

During both seasons, higher numbers of AMF species were recorded in natural forests. grasslands and acacia plantations than in the cardamom and coffee plantations and paddy fields. Shi et al. (2007) reported higher species diversity in natural forests than in forest plantation. Oehl et al. (2003) while studying the impact of land use types have also reported that the AMF species composition was the highest in grasslands, lower in the low and moderate input arable lands and the lowest in the lands with intensive continuous maize mono-cropping. Certain other studies have also indicated that the number of species was higher in undisturbed areas than the disturbed areas. Picone (2000) while studying the AMF species composition in Nicaragua and Costa-Rican forests and pastures, found similar species composition in both forests and pastures, while Mendez et al. (2001) noticed higher AMF species composition in pasture soils than in cultivated soils. The results of the present study follow a similar pattern as the natural forests and grasslands and to some extent acacia plantations are more or less undisturbed, because these plantations are almost 10 years old. Coffee plantations and paddy fields on the other hand are more intensively cultivated which may have led to lower AMF species numbers.

## AMF Species Spore Abundance over Landscape

AMF species spore abundance was more during post-monsoon than during pre-monsoon season. The majority of the species produced more spores during post-monsoon than during pre-monsoon season. This kind of variation in AMF species composition over different seasons could be attributed to the adaptation of specific AMF species to a particular climatic condition and soil moisture regimes. Sampling of soils over seasons has revealed that some AMF species sporulated better during wet season while some species sporulated during dry season. Lovelock (2003) also observed that the relative abundance of spores of Acaulospora was lower than that of Glomus, during wet season and found that Glomus produced relatively more spores at the highest seasonal rainfall. Further, he also suggested that during wet season a number of plant species with profuse rooting favored the sporulation of AMF species. Sanders and Fitter (1992) opined that the composition of plant community might also affect mycorrhizal fungi causing differential reproduction and survival, which will definitely act as a selective force on the composition of AM fungi. Schenck and Kinloch (1980) also noticed incidence of AMF species over different periods of time over the years; the spores of some species increased, while the spores of few species decreased. Spores of few species observed during one season disappeared in the next season.

# Diversity of AM Fungi in Different Land Use Types

Mycorrhizal fungi are likely to be affected by plant community composition (Janos, 1980; Kormanik et al., 1980). The AMF populations may readily respond to the proportion of mycotrophic plants in a community because the fungi cannot live without hosts. In the present study, variations in the Shannon-Wiener diversity index of AMF species in different land use types were noticed. This kind of variation in the diversity of AMF in different land use types and during different seasons was also noticed in earlier studies. Picone (2000) and Helgason et al. (2007) have reported that the AMF diversity indices for forest and pastures were similar. Also, in this study, the diversity index of AMF in natural forests and grasslands was on par with each other. This suggests that AMF diversity in undisturbed systems as that of forests and grasslands is less affected. In contrast, the diversity index was quite different in less undisturbed grasslands and more disturbed paddy fields. Oehl et al. (2003) in their study also recorded the highest mycorrhizal diversity index in grasslands compared to moderate and low input aerable lands and intensive continuous maize mono-cropping. Carpenter et al. (2001) in their study on spore density and diversity of AMF in different land uses found that diversity of AMF changed due to change in land use types.

The AMF species richness between different land use types and seasons followed a similar trend as that of diversity. The lower species richness index for paddy fields could be due to intensive agricultural practices. Earlier studies have shown that application of fertilizers, pesticides, tillage and other soil disturbances decreased AMF diversity and species richness (Kruckelmann, 1975; Hayman, 1980, Gemma, 1988, Attitchabi et al., 2008). Further, low fertility status of the soils in natural forests, acacia plantations and grasslands probably required an increased dependence of plants on mycorrhizae for growth and survival, which facilitated the growth of a diverse species of AMF. Blaszkowski (1994) also made similar observations with species richness for different plants grown in different places of Hel peninsula of Poland, where the species richness was more in low fertile soils sampled under Cupressaceae followed by Rosaceae and less in Gramineae. The fertility status of soil was more in soils under Rosaceae and Gramineae.

#### CONCLUSIONS

This study shows that disturbance of a particular land use type leads to the destruction of flora, which in turn alter/reduce the population and diversity of mycofauna.

#### ACKNOWLEDGEMENTS

Authors express appreciation for the financial support from the GEF/UNEP through TSBF for carrying out this research work.

#### REFERENCES

- 1. Anderson, J. M. and Ingram, J. S. I. 1993. *Tropical Soil Biology and Fertility: A Handbook of Methods.* 2<sup>nd</sup> Edition, CAB International, Wallingford, 55pp.
- 2. Attitchabi, J., Coyne, D., Hountondji, F., Lawouin, L., Wiemken, A. and Oehl, F. 2008, Arbuscular Mycorrhizal Fungi in the 'Yam Belt' of West Africa. *African J. Biotechnol.*, **56**: 256-268.

JAST

- 3. Augé, R. M. 2004. Water Relations, Drought and Vesicular-arbuscular Mycorrhizal Symbiosis. *Mycorrhiza*, **11**: 3–42.
- 4. Bagyaraj, D. J. and Manjunath, A. 1980. Selection of Suitable Host for Mass Production of VA Mycorrhizal Inoculation. *Plant Soil*, **55**: 495-498.
- Bagyaraj, D. J. and Varma, A. 1995. Interactions between Arbuscular Mycorrhizal Fungi and Plants: Their Importance in Sustainable Agriculture in Arid and Semiarid Tropics. *Adv. Microbiol. Ecol.*, 14: 119-142.
- Bagyaraj, D. J. and Sturmer, S. L. 2008. Arbuscular Mycorrhizal Fungi (AMF). In: "A Handbook of Tropical Soil Biology: Sampling and Characterization of Belowground Biodiversity", (Eds.): Moreira, F. M. S., Huising, J. E. and Bignell, D. E... Earthscan Publication, London, PP. 131-147.
- Balakrishna, A. N., Lakshmipathy, R., Sudhir, K. and Bagyaraj, D. J. 2001. Effect of Long Term Application of Inorganic Fertilizers on Native VA Mycorrhizal Fungi and Soil Microbial Biomass in a Finger Millet-maize-fallow Rotation. J.Soil Biol.Ecol., 21: 1-6.
- Bhardwaj, S., Dudeja, S. S. and Khurana, A. L. 1997. Distribution of Vesiculararbuscular Mycorrhizal Fungi in the Natural Ecosystem. *Folia-Microbiol.*, 42: 589-594.
- Blaszkowski, J. 1994. Comparative Studies on the Occurrence of Arbuscular Fungi and Mycorrhizae (Glomales) in Cultivated and Uncultivated Soils of Poland. *Acta Mycol.*, 28: 93-140.
- Boddington, C. L. and Dodd, J. C. 2000. The Effect of Agricultural Practices on the Development of Indigenous Arbuscular Mycorrhizal Fungi. II. Studies in Experimental Microcosms. *Plant Soil*, 218: 145-157.
- Byra Reddy, M. S., Krishna Naik, L., Bhaskar, V. and Bagyaraj, D. J. 1994. Status of Mycorrhizal Association in Some Tropical Tree Species. J. Soil Biol. Ecol., 14: 51-54.
- Carpenter, F. L., Palacios, M. S., Gonzalez, Q. E. and Schroeder, M. 2001. Land Use and Erosion of a Costa Rican Ultisol Affect Soil Chemistry, Mycorrhizal Fungi and Early Regeneration. *For. Ecol. Manage.*, 144: 1-17.

- Doss, D. D. and Bagyaraj, D. J. 2001. Ecosystem Dynamics of Mycorrhizae. In: "Innovative Approaches in Microbiology", (Eds.): Maheswari, D. K. and Dubey, R. C.. Bishen Singh Mahendra Pal Singh Publication, Dehra Dun, India, PP. 115-129
- Eivazi, F. and Tabatabai, M. A. 1977. Phosphatases in Soils. *Soil. Biol. Biochem.*, 9: 167-172.
- Fischer, C. R., Janos, D. P., Perry, D. A. Linderman, R. G. and Sollins, P. 1994. Mycorrhiza Inoculum Potentials in Tropical Secondary Sucession. *Biotropica*, 26: 369-377.
- Gemma, J. N. and Koske, R. E. 1988. Seasonal Variation in Spore Abundance and Dormancy of *Gigaspora gigantea* and in Mycorrhizal Inoculum Potential of a Dune Soil. *Mycologia* 80: 211-216.
- 17. Gerdemann, J. W. and Nicolson, T. H. 1963. Spores of Mycorrhizal Endogone Species Extracted from the Soil by Wet Sieving and Decanting. *Trans. Br. Mycol. Soc.*, 46: 235-244.
- Giovannetti, M. and Mosse, B. 1980. An Evaluation of Techniques to Measure Vesicular Arbuscular Infection in Roots. *New Phytol.*, 84: 489-500.
- Harinikumar, K. M. and Bagyaraj, D. J. 1988. The Effect of Season on VA Mycorrhiza of Leucaena and Mango in Semi-arid Tropic. *Arid Soil Res. Rehabil.*, 7: 139-143.
- Harinikumar, K. M., Bagyaraj, D. J. and Mallesha, B. C. 1990. Effect of Intercropping and Organic Soil Amendments on Native VA Mycorrhiza in Semi-arid Tropics. *Arid Soil Res. Rehabil.*, 4: 193-197.
- Harinikumar, K. M. and Bagyaraj, D. J. 1989. Effect of Cropping Sequences, Fertilizers and Farm Yard Manure on VA Mycorrhizal Fungi. *Biol. Fert. Soils*, 7: 173-175.
- 22. Hayman, D. S. 1978. Mycorrhizal Population of Sown Pastures and Native Vegetation in Otago, New Zealand. *NZ. J. Agric. Res.*, **21**: 271-276.
- 23. Hayman, D. S. 1980. VA Mycorrhiza and Crop Production. *Nature*, **287**: 487-488.
- Helgason, T., Merryweather, J. W., Young, J. P. and Fitter, A. H. 2007. Specificity and Resilience in the Arbuscular Mycorrhizal Fungi of a Natural Woodland Community. *J. Ecol.*, **95**: 623–630.

- Hibbett, D. S., Binder, M., Bischoff, J. F., Blackwell, M. and Cannon, P. F. 2007. A Higher Level Phylogenetic Classification of the Fungi. *Mycol. Res.*, **111**: 509-514.
- Husband, R., Herre, E. A., Turner, S. L. Gallery, R. and Young, J. P. W. 2002. Molecular Diversity of Arbuscular Mycorrhizal Fungi and Pattern of Host Association over Time and Space in a Tropical Forest. *Molecular Ecol.*, 11: 2669-2678.
- Jackson, M. L. 1973. Soil Chemical Analysis. Prentice Hall Pvt. Ltd., New Delhi, India, PP.105-110.
- Janos, D. P. and Read, D. J. 1992. Heterogenity and Scale in Tropical Vesicular-arbuscular Mycorrhiza Formation. In: "*Mycorrhiza in Ecosystems*", (Eds.): Fitter, A. H. and Alexander, I. J.. CAB international US, London, PP. 276-282.
- Janos, D. P. 1980. Mycorrhizae Influence Tropical Succession. *Biotropica*, 12: 56-64.
- Johnson, N. C. and Wedin, D. A. 1997. Soil Carbon, Nutrients, and Mycorrhizae during Conversion of Dry Tropical Forest to Grassland. *Ecol. Applications*, 7: 171-182.
- Clark, R. B. and Zeto, S. K. 2000. Mineral Acquisition by Arbuscular Mycorrhizal Plants. J. Plant. Nutr., 23: 867–902.
- Koide, R. T. 1991. Nutrient Supply, Nutrient Demand and Plant Response to Mycorrhizal infection. *New Phytol.* 117: 365-386.
- 33. Kormanik, P. P., Bryan, W. C. and Schultz, R. C. 1980. Increasing Endomycorrhizal Fungus Inoculum in Forest Nursery Soil with Cover Crops. *South. J. Appl. For.*, 4: 151-153.
- 34. Krebs, C. J. 1989, *Ecological Methodology*. Harper Collins Publishers, 58 PP.
- 35. Krishna, K. R. and Bagyaraj, D. J. 1982. Effect of Vesicular-arbuscular Mycorrhiza and Soluble Phosphate on *Abelmoscus esculentus* (L.) *Moench. Plant Soil*, **64**: 209-213.
- Kruckelman, H.W. 1975. Effects of Fertilizers, Soils, Soil Tillage and Plant Species on the Frequency of *Endogone* Chlamadospores and Mycorrhizal Infection in Arable Soils. In: "Endomycorrhizas", (Eds.): Sanders, F. E., Mosse, B. and Tinker, P. B.. Academic Press. London, PP. 511-525.
- Lakshmipathy, R., Sumana, D. A., Balakrishna, A. N., Bagyaraj, D. J. and Kumar, D. P. 2004. Evaluation, Grafting

Success and Field Establishment of Cashew Rootstock as Influenced by VAM Fungi. *Ind. J. Expt. Biol.*, **42**: 1132-1135

- Lakshmipathy. R., Balakrishna, G. and Bagyaraj, D. J. 2003. VA Mycorhizal Colonization Pattern in RET Medicinal Plants (*Mammea suriga, Saraca asoca, Garcinia spp., and Embelia ribes and Calamus sp.*) in Different Parts of Karnataka. *Asian J. Microbiol. Biotech. Env. Sci.*, 5: 505-508.
- 39. Lakshmipathy. R., Balakrishna G., Chandrika, K. and Bagyaraj, D. J. 2002. VA Mycorrhizal Colonization Pattern in *Hemidesmus indicus* and *Dioscorea bulbifera* in Different Parts of Karnataka. *Giobios*, 29: 209-212.
- Lovelock, C. E., Andersen, K. and Morton, J. B. 2003. Arbuscular Mycorrhizal Communities in Tropical Forests Are Affected by Host Tree Species and Environment. *Oceologia*, 132: 268-279.
- 41. Mallesha, B. C. and Bagyaraj, D. J. 1991. Season Favouring Sporulation of VA Mycorrhizal Fungi in Cardamom Plantations. *Soil Boil. Ecol.*, **11**: 75-78.
- 42. Mendez, A. B., Scervine, J. M. and Fodeas, A. M. 2001. Arbuscular Mycorrhizal Population Associated with Natural and Cultivated Vegetation on a Site of Buenos Aires Province, Argentina. *Biol. Fertil. Soils*, 33: 373-381.
- 43. Mervin, H. D. and Peech, M. 1951. Exchangebility of Soil Potassium in the Sand, Silt and Clay Fractions as Influenced by the Nature of the Complementary Exchangeable Cation. *Proc. Soil Sci. Soc. Amer.*, **15**: 125-129.
- 44. Mohan, C. 2003. Mycorrhizae in Forest Plantations: Association, Diversity and Exploitation in Planting Stock Improvement: KFRI Research Report-252. Peechi.
- 45. Molina, R. J., Trappe, J. M. and Strickler, G. S. 1978. Mycorrhizal Fungi Associated with *Festuca* in the Western United States and Canada. *Can. J. Bot.*, **56**: 1691-1695.
- 46. Morton, J. B. and Benny, G. L. 1990. Revised Classification of Arbuscular Mycorrhizal Fungi (Zygomycetes): A New Order, Glomales, Two New Families, *Acaulosporaceae* and *Gigasporaceae* with an Emendation of *Glomaceae*. *Mycotaxon*, **37**: 471-491.
- 47. Morton, J. B. 1990. Species and Clones of Arbuscular Mycorrhizal Fungi (*Glomales*

Downloaded from jast.modares.ac.ir on 2025-05-20

*zygomycetes*): Their Role in Macro and Micro–evolutionary Process. *Mycotaxon*, **37**: 493–515.

- Muthukumar, T. and Udayan, K. 2000. Arbuscular Mycorrhizas of Plants Prowing in the Western Ghats Region, Southern India. *Mycorrhiza*, 9: 297-313.
- Muthukumar, T. K. V. and Manian, S. 1993. Vesicular Arbuscular Mycorrhizal Fungi in Western Ghats. *Ind. Bot.*, 10: 79-83.
- Myers, N., Mittermeir, R. A., Mittermeir, C. G., da Gustavo, A. B. and Kent, J. 2000. Biodiversity Hotspots for Conservation Priorities. *Nature*, 403: 853-858.
- 51. Oehl, F., Sieverding, E., Ineichen, L., Mader, P., Boller, T. and Wienmken, A. 2003. Impact of Land Use Intensity on the Species Diversity of Arbuscular Myocrrhial Fungi in Agroecosystems of Central Europe. *Appl. Env. Microbiol.*, 69: 2816-2824.
- 52. Oehl, F., Sieverding, E., Goto, T. B., Palenzeula, J., Ferrol, N., da Silva, G. A. and de Souza, F. A. 2009. The Fairytale of the Different Morphological Identification of AMF: Recent Advances in the Area. In: "Beyond the Roots". Belo Horizonte, Brazil, 2009, 17 PP.
- Palenzuela, J., Ferrol, N., Boller, T., Azcon-Aguilar, C. and Oehl, E. 2008, *Otospora bareai:* A New Fungal Species in the Glomeromycetes from a Dolomitic Shrubland in the Natural Park of Sierra de Baza (Granada, Spain). *Mycologia*, **100**: 296-305.
- Pande, M. and Tarafdar, J. C. 2004. Arbuscular Mycorrhizal Fungal Diversity in Neem Based Agroforestry Systems in Rajasthan. *App. Soil Ecol.*, 26: 233-241.
- 55. Patrícia, L.L, Stürmer S. L. and Siqueira, J. O. 2009. Occurrence and Diversity of Arbuscular Mycorrhizal Fungi in Trap Cultures from Soils under Different Land Use Systems in the Amazon, Brazil. *Brazilian J. Microbiol.*, **40**: 111-121.
- 56. Phillips, J. M. and Hayman, D. S. 1970. Improved Procedures for Clearing and Staining Parasites and Vesicular Arbuscular Mycorrhizal Fungi for Rapid Assessment of Infection. *Trans. Br. Mycol. Soc.*, 55: 158-161.
- 57. Picone, C. M. 2000. Diversity and Abundance of Arbuscular Mycorrhizal Fungus Spores in Tropical Forest and Pasture. *Biotropica*, **32**: 734-750.

- Pirozynski, K. A. and Dalphe, Y. 1989. Geological History of the Glomaceae with Particular Reference to Mycorrhizal Symbiosis. *Symbiosis*, 7: 1–36.
- 59. Porter, W. M. 1979. The Most Probable Number Method for Enumerating Infective Propagules of Vesicular Arbuscular Mycorrhizal Fungi in Soil. *Aus. J. Soil Res.*, 17: 515-519.
- Porter, W. M., Robson, A. D. and Abbott, L. K. 1987. Field Survey of the Distribution of Vesicular-arbuscular Mycorrhizal Fungi in Relation to Soil pH. *J. App. Ecol.*, 24: 659-662.
- 61. Sanders, I. R. and Fitter, A. H. 1992. Evidence for Differential Responses between Host-fungus Combinations of Vesicular-arbuscular Mycorrhizas from a Grassland. *Mycol. Res.*, **96**: 415-419.
- 62. Santhaguru, K., Ponmalar, S. B. G. and Karunakaran, R. 1995. Vesicular-arbuscular Mycorrhizae in Tree-legumes and Its Rhizospheric Soils in Alagar Hills. *Ind. For.*, **121**: 817-823.
- 63. Schenck, N. C. and Kinloch, R. A. 1980. Incidence of Mycorrhizal Fungi on Six Field Crops in Monoculture on a Newly Cleared Woodland Site. *Mycologia*, **72**: 229-443.
- Schenk, N. C. and Perez, Y. 1990. Manual for Identification of VA Mycorrhizal Fungi. 3<sup>rd</sup> Edition, Synergistic Publications, Gainesville, Florida, USA, P. 1-142.
- Shi, Z., Y., Wang, F. and WeI Y. L. 2007. Natural Forest and Forest Plantation Affect Diversity of Arbuscular Mycorrhizal Fungi in the Rhizosphere of Diptorocarpaceae. *American-Eurasian J. Agric. Environ. Sci.*, 2: 411-416.
- 66. Sieverding, E. and Leihner, D. 1984. Effect of Herbicides on Population Dynamics of VA Mycorrhiza with Cassava. *Angew Botanic*. **58**: 283-294.
- Sieverding, E. 1991. Vesicular Arbuscular Mycorrhiza Management in Tropical Agroecosystems. Technical Corporation, Federal Republic of Germany, Eschbom, PP. 307-311.
- Sieverding, E. and Oehl, F. 2006. Revision of *Entrophospora* and Description of *Kuklospora* and *Intraspora*: Two New Genera in the Arbuscular Mycorrhizal Glomeromycetes. J. App. Bot. Food Quality, 80: 69-81.



- 69. Snedecor, S. W. and Cochram, W. G. 1968. *Statistical Methods*. Oxford and IBH Publication, New Delhi, 124 PP.
- Tommerup, I. C. and Sivasithamparam, K. 1990. Zygospores and Asexual Spores of *Gigaspora decipiens*: An Arbuscular Mycorrhizal Fungus. *Mycol. Res.*, **94**: 897-900.
- Vasanthakrishna, M., Muthanna, M. B. and Bagyaraj, D. J. 1994. Succession of Vesicular Arbuscular Mycorrhizal Fungi

Associated with *Casuarina equisetifolia L.* Ann. For., **2**: 123-126.

- Walker, C., Vestberg, M., Demircik, F., Stockinger, H., Saito, M., Sawaki, H., Nishmura, I. and Schufler, A. 2007a. Molecular Phylogeny and New Taxa in the Archaeosporales. *Mycol. Res.* 111: 3-49.
- Walker, C., Vestberg, M., Demircik, F., Stockinger, H., Saito, M., Sawaki, H., Nishmura, I. and Schufler, A. 2007b. Molecular Phylogeny and New Taxa in the Archaeosporales. *Mycol. Res.* 111: 137-153.

فراوانی و تنوع قارچهای AMF در دامنهای از شدت کاربری اراضی و تغییرات فصلی آن در بیوسفر نیلیگیری در منطقه گات غربی، هندوستان

ر. لاکشمیپتی، ا. ن. بالاکریشنا، د.ج. با گیاراج

# چکیدہ

اثر شدت کاربری اراضی بر روی فراوانی و تنوع قارچهای AMF در شش نوع کاربری اراضی شامل جنگل طبیعی، علفزار، مزرعه آکاسیا، مزرعه هل، مزرعه قهوه و مزرعه برنج در بیوسفر نیلیگیری در منطقه گات غربی، در جنوب هندوستان بررسی گردید. در فصل پیش از مونسون تفاوت معنی داری بین مقدار AMF ریشه در کاربری-های مختلف مشاهده نشد در حالی که در فصل پیش از مونسون این تفاوت معنی دار بود و مقادیر در جنگل طبیعی و علفزار بیشتر بودند. تراکم هاگ AMF و اندامهای زایشی آلوده درعلفزار و مزرعه آکاسیا در هر دو فصل نسبت به دیگر کاربریها بیشتر بود. به استئنای مزرعه برنج، در بقیه کاربریها تراکم هاگ AMF و اندامهای زایشی آلوده در فصل پیش از مونسون از فصل پیش از آن بیشتر بود. تعداد گونه های AMF شناسایی شده در زایشی آلوده در فصل پس از مونسون از فصل پیش از آن بیشتر بود. تعداد گونه های AMF شناسایی شده در در هر دو فصل میش از مونسون ۶۵ و در فصل پس از مونسون ۶۷ عدد بود که نشان دهنده تغییرات فصلی در تنوع میباشد. در هر دو فصل میش از مونسون از فصل پس از مونسون ۶۷ عدد بود که نشان دهنده تغییرات فصلی در تنوع میباشد. موه دو فصل پیش از مونسون ۶۵ و در فصل پس از مونسون ۶۷ عدد بود که نشان دهنده تغییرات فصلی در تنوع میباشد. موه دو فصل میش از مونسون از فصل پس از مونسون ۶۷ عدد بود که نشان دهنده تغییرات فصلی در تنوع میباشد. مور دو فصل مناز مونسون ۶۵ و در فصل پس از مونسون ۶۰ موسون بیشترین مقدار را داشتند. در مور دو فصل، تنوع گونه ای در جنگل طبیعی و علفزار نسبت به کاربریهای دیگر بیشتر بود. در هر دو دوره زمانی مورد مطالعه، شاخص غنای گونهای برای AMF در جنگل طبیعی حداکثر و در مزرعه برنج حداقل بود. میزان مورد مطالعه، شاخص غنای گونهای برای AMF در جنگل طبیعی حداکثر و در مزرعه برنج حداقل بود. میزان موره میانسیته ظاهری، نیتروژن کل، کرین آلی و فسفاتازهای قلیایی و اسیدی با فعالیت AMF رابطه مثبت و میزان